

Experiences of Domestic Violence and Mental Disorders: A Systematic Review and Meta-Analysis

Kylee Trevillion¹*, Siân Oram¹*, Gene Feder², Louise M. Howard¹*

1 Section of Women's Mental Health, Health Service and Population Research Department, Institute of Psychiatry, King's College London, London, United Kingdom, **2** Centre for Academic Primary Care, School of Social and Community Medicine, University of Bristol, Clifton, Bristol, United Kingdom

Abstract

Background: Little is known about the extent to which being a victim of domestic violence is associated with different mental disorders in men and women. We aimed to estimate the prevalence and odds of being a victim of domestic violence by diagnostic category and sex.

Methods: Study design: Systematic review and meta-analysis. Data Sources: Eighteen biomedical and social sciences databases (including MEDLINE, EMBASE, PsycINFO); journal hand searches; scrutiny of references and citation tracking of included articles; expert recommendations, and an update of a systematic review on victimisation and mental disorder. Inclusion criteria: observational and intervention studies reporting prevalence or odds of being a victim of domestic violence in men and women (aged ≥ 16 years), using validated diagnostic measures of mental disorder. Procedure: Data were extracted and study quality independently appraised by two reviewers. Analysis: Random effects meta-analyses were used to pool estimates of prevalence and odds.

Results: Forty-one studies were included. There is a higher risk of experiencing adult lifetime partner violence among women with depressive disorders (OR 2.77 (95% CI 1.96–3.92), anxiety disorders (OR 4.08 (95% CI 2.39–6.97), and PTSD (OR 7.34 (95% CI 4.50–11.98), compared to women without mental disorders. Insufficient data were available to calculate pooled odds for other mental disorders, family violence (i.e. violence perpetrated by a non-partner), or violence experienced by men. Individual studies reported increased odds for women and men for all diagnostic categories, including psychoses, with a higher prevalence reported for women. Few longitudinal studies were found so the direction of causality could not be investigated.

Conclusions: There is a high prevalence and increased likelihood of being a victim of domestic violence in men and women across all diagnostic categories, compared to people without disorders. Longitudinal studies are needed to identify pathways to being a victim of domestic violence to optimise healthcare responses.

Citation: Trevillion K, Oram S, Feder G, Howard LM (2012) Experiences of Domestic Violence and Mental Disorders: A Systematic Review and Meta-Analysis. PLoS ONE 7(12): e51740. doi:10.1371/journal.pone.0051740

Editor: Jim van Os, Maastricht University Medical Centre, The Netherlands

Received: September 9, 2012; **Accepted:** November 9, 2012; **Published:** December 26, 2012

Copyright: © 2012 Trevillion et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The authors receive support from the NIHR Programme Grants for Applied Research scheme (RP-PG-0108-10084). Louise Howard also receives salary support from the National Institute for Health Research (NIHR) Mental Health Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. This paper presents independent research funded by the National Institute for Health Research (NIHR) under its Programme Grants for Applied Research scheme (RP-PG-0108-10084). The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. The funder had no role in study design, data collection, data analysis, decision to publish, or preparation of the manuscript.

Competing Interests: Gene Feder and Louise M. Howard are members of the WHO Guideline Development Group on Policy and Practice Guidelines for responding to Violence Against Women and the NICE/SCIE Guideline Development Group on Preventing and Reducing Domestic Violence. KT and SO declare no competing interests. This does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials.

* E-mail: louise.howard@kcl.ac.uk

† These authors contributed equally to this work.

Introduction

Domestic violence is an international public health problem, affecting the lives of hundreds of thousands of people every year. Globally, prevalence estimates of lifetime experiences of physical or sexual partner violence among women range from 15%–71%, with past year estimates ranging from 4% and 54% [1]. No such global estimates exist for men. Research on the prevalence of domestic violence within same-sex relationships is limited; however, evidence from the USA increasingly suggests that the prevalence is similar across same-sex and heterosexual relationships [2].

As a consequence of the substantial physical and psychiatric morbidity associated with domestic violence [3,4,5], victims have increased use of health services compared to those not abused [6,7]. Domestic violence is associated with substantial healthcare costs, with direct medical and mental healthcare costs approximating £1,730 million per annum in the UK and \$4.1 billion in the USA, with additional societal costs [8,9].

Prolonged exposure to threatening life events, including domestic violence, is associated with the onset, duration and recurrence of mental disorders [5,10], and men and women with mental disorders are at an increased risk of experiencing violence [11]. Recent reviews have suggested that being a victim of




Figure 1. Flow Diagram of Screened and Included Papers.
doi:10.1371/journal.pone.0051740.g001

domestic violence is common among people with mental disorders [4,5,12,13]. These reviews, however, predominantly focus on depression and PTSD (or report on “mental disorders” without diagnostic characterisation) and have not drawn upon the broader body of research on violence victimisation among people with mental disorders. Furthermore, most reviews do not critically appraise study quality [4,5,12] and do not report separately on men who experience domestic violence or on domestic violence perpetrated by family members [4,12,13]. This systematic review therefore aimed to estimate:

- The prevalence (lifetime and past year) of being a victim of domestic violence in men and women with mental disorders
- The odds of being a victim of domestic violence in men and women with mental disorders compared with non-mentally disordered controls

Methods

Search Strategy

This review followed MOOSE and PRISMA guidelines [14,15] (see Checklist S1) and the protocol is registered with the PROSPERO database of systematic reviews (<http://www.crd.york.ac.uk/prospero>); registration number CRD42011001281. A multi-stage search strategy was used, which comprised: (a) an electronic search of 18 bibliographic databases; (b) an update of a recent systematic review on violence experienced by people with mental disorders (i.e. a review which did not focus on domestic violence but may have included studies that collected data on domestic violence) [16]; (c) hand searches of three key journals (i.e. *Trauma Violence and Abuse*, *Journal of Traumatic Stress*, and

Violence Against Women); (d) screening of references lists of included studies; (e) forwards citation tracking (i.e. identifying studies that had cited the papers included in this review), and (f) expert recommendations. Medical Subject Headings (MeSH) and text words were used to search 18 biomedical and social science electronic databases, from their dates of inception up to 31st March 2011 (see Text S1 for the list of databases used). Terms for domestic violence [17] were adapted from Cochrane protocols and peer-reviewed literature reviews [12,18] and terms for mental disorders [19] were adapted from NICE guidelines [20] (see Text S2 for Medline, EMBASE and PsycINFO search strategies). When updating the victimisation review [Maniglio [16]], we used the author’s original search terms to search databases from September 2007 (the upper limit of the original review) to the 31st March 2011 [16]. No language restrictions were used.

Selection Criteria

Studies were eligible for inclusion if they: (a) included men and/or women who were 16 years or older and were diagnosed with a mental disorder using a validated diagnostic instrument (i.e. diagnostic instruments that have been validated against a gold standard measure for diagnosing mental disorder, such as the Schedules for Clinical Assessment in Neuropsychiatry [21]) (see Text S3 for the full definition of mental disorder); (b) presented the results of peer-reviewed research based on experimental studies (e.g. randomised controlled trials, non-randomised controlled trials, parallel group studies), before-and-after studies, interrupted time series studies, cohort studies, case-control studies, or cross-sectional studies; and (c) measured the prevalence or odds of lifetime/past year domestic violence (see Text S3 for the full definition of domestic violence), or reported data from which these

statistics could be calculated. In the registered protocol, we stated that studies which used validated screening instruments (i.e. that identify presence of probable mental disorder but do not diagnose mental disorders) would be included in the main text of the review but excluded from meta-analyses. However, due to the large number of screening papers identified (see Figure 1), we decided to include only studies that used validated diagnostic instruments. When we identified multiple eligible papers from the same study only the paper reporting the largest sample size with data of relevance to the objectives of the review was included.

Data Extraction and Quality Appraisal

The downloaded titles and abstracts were screened against the inclusion criteria by two reviewers (KT and SO). If it was unclear whether a reference met the inclusion criteria, it was taken forward to the next stage of screening. The full texts of potentially eligible studies were assessed by two reviewers (KT and SO). If studies collected data on the prevalence and/or odds of domestic violence but did not report it, authors were contacted for the data. Details of the 1,083 excluded papers and reasons for exclusion are available upon request.

Data from included papers were extracted into a standardised electronic database by two reviewers (KT and SO) and a random sample of 20% was independently cross-checked. Extracted data included details on: the study design; sample characteristics; measures of mental disorder and domestic violence, and the prevalence and odds of lifetime/past year domestic violence. Data were extracted separately for men and women. When reported, details on resource use, impact and severity of violence and chronicity of mental disorders were extracted.

The quality of included studies was independently appraised by two reviewers (KT and SO) using criteria adapted from validated tools [22,23,24,25]. Reviewers compared scores and resolved disagreements before allocating a final appraisal score (see Table S1). Reviewer inter-rater reliability regarding quality scores was high (i.e. for overall quality score: Pearson's r 0.98, ICC 0.95). The quality appraisal checklist includes items assessing study selection and measurement biases (see Table S2). Studies were categorised as high-quality if they scored $\geq 50\%$ on questions pertaining to selection bias. Quality scoring, particularly for observational research, is contestable [26]; yet we wanted to exclude poor studies that threatened the validity of our findings. The 50% criterion was chosen in order to maximise the number of studies eligible for inclusion in the meta-analysis, while excluding studies in which there was a high risk of selection bias.

Data Analysis

We calculated prevalence, odds ratios and 95% confidence intervals for domestic violence among men and women by type of mental disorder. If studies measured multiple disorders, odds ratios were calculated separately by type of mental disorder and for each estimate the control group were participants without any mental disorder. Prevalence and odds ratios were also calculated separately by sex and type of violence. Regarding type of violence, we report results for any violence (i.e. physical, sexual and psychological violence) and for physical violence alone. Data on the prevalence and odds of sexual and psychological violence were limited and are given in Table S1. It was not possible to adjust odds ratios for potential confounders (e.g. childhood abuse) due to the lack of data from many of the original studies; unadjusted odds ratios are therefore presented.

We calculated DerSimonian-Laird random effects odds ratio estimates (with corresponding 95% confidence intervals) for lifetime and past year domestic violence among people with

mental disorders, compared to people without a mental disorder, if reports were available from three or more high-quality studies [27]. Pooled odds ratios were calculated separately for men and women; studies for which sex-disaggregated data were not available were not eligible for inclusion in the meta-analyses. Disorder-specific summary estimates that included both high- and low-quality papers were calculated in order to assess the impact of excluding low-quality papers. We also examined the influence of individual studies on summary effect estimates by conducting influence analyses, which compute summary estimates omitting one study at a time. Unless stated, neither the inclusion of the low-quality studies nor the omission of individual studies made material differences to odds estimates. We aimed to assess the risk of small study bias with funnel plots in conjunction with Egger's tests [28]. However, due to the small number of eligible studies, statistical tests for funnel plot asymmetry were not appropriate and we were confined to visual inspection of the plots; funnel plots are presented where there were sufficient data-points to allow this (see Figure S1, Figure S2, and Figure S3). Heterogeneity among studies was estimated using the I^2 statistic (associated 95% confidence intervals were calculated with the STATA *heterogi* command using a non-central χ^2 based approach). Due to the small number of studies included in each meta-analysis, it was not possible to use meta-regression to investigate sources of heterogeneity. All analyses were conducted in STATA 11 [29].

Results

The study selection process is presented in Figure 1. The literature search yielded 29,707 unique references, of which 28,584 were excluded following title and abstract screening. Of the 1,123 references that met, or potentially met, the inclusion criteria, 59 (56 dissertations and three journal articles (in *Türkiye'de Psikiyatri*, *Psikiyatri Psikoloji* *Psikofarmakoloji Dergisi*, and *Revista de Psiquiatria Clinica*) could not be located. Following full text screening, 41 papers were included in the review: 28 were from searches of electronic databases, five from citation tracking, one from hand searching, four from re-examining and updating of an earlier systematic review of victimisation, and three from experts. Six non-English language papers were translated but were not eligible for inclusion in the review.

Key characteristics of the studies are reported in Table 1. Details of design, sample size, definition and measurement of mental disorder and domestic violence are reported in Table S1. Of the 41 included studies, 27 scored $\geq 50\%$ on quality appraisal criteria for selection bias [30,31,32,33,34,35,36,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56], and 14 scored less than 50% on quality appraisal criteria for selection bias [57,58,59,60,61,62,63,64,65,66,67,68,69,70]. Unless otherwise stated, results are reported for high-quality studies only.

Prevalence and Odds of Domestic Violence by Diagnosis

Results present data for lifetime and past year experiences of any type of partner violence (i.e. physical, sexual and/or psychological violence) across all mental disorders in women and men. Prevalence and odds estimates for all included studies are presented in Table S1; where available, data on specific types of violence (i.e. physical, sexual or psychological violence) and violence by non-intimate family members are also presented in Table S1.

Depressive Disorders

Lifetime Domestic Violence. Among women, the median prevalence of any lifetime partner violence (7 studies) was 45.8%

Table 1. Characteristics of included studies (n = 41)*.

	Total (n = 41)	Lifetime domestic violence (n = 26)	Past year domestic violence (n = 18)
Sample:			
Males only	0	0	0
Females only	25	14	14
Males and females	16†	12	4
Diagnoses:			
Schizophrenia & non-affective psychosis	3	2	1
Bipolar affective disorder	2	2	0
Depressive disorders	26	21	12
Dysthymia	5	2	3
Anxiety disorders	15	9	7
PTSD	14	9	7
OCD	2	2	0
Panic disorders	6	2	4
Phobias	3	2	1
Somatisation	1	0	1
Eating disorder	1	0	1
Personality disorder	4	3	1
Common Mental Disorder	5	4	2
Setting:			
Clinical	17	11	7
Non-clinical	24	15	11
Perpetrator:			
Partner only	38‡	24	17
Family only	0	0	0
Partner or family	3	2	1
Type of violence			
Physical violence	20	15	5
Psychological violence	9	5	4
Sexual violence	4	3	1
Physical, sexual, psychological combined	11	5	6
Recency of violence			
Lifetime domestic violence	23		
Past year domestic violence	15	-	-
Lifetime and past year domestic violence	3		
Measurement of domestic violence			
Validated measures	18∞	11	8
Non-validated measures	19◇	11	7
Trauma items from DSM/CIDI criteria	4	4	0
Region:			
North America	17	10	9
Central America	1	1	0
South America	1	1	0
Europe	6	5	2
Africa	3	3	0
Asia	8	6	2
Australasia	5	0	5

*Categories are not mutually exclusive and rows may therefore add to >40.

†Sex-disaggregated data was available for 11 of the 16 studies.

‡Five papers measured only spousal violence.

∞Four papers made modifications to validated measures and five did not use all items in the measure.

◇In 16 studies the authors developed their own measure to assess domestic violence.

doi:10.1371/journal.pone.0051740.t001

(IQR 21.3%–76.5%; range 15.6%–89.2%) [30,36,38,46,50,52,55]. The pooled odds ratio for any lifetime partner violence was 2.77 (95% CI 1.96–3.92), with high heterogeneity $I^2 = 83.9\%$ (95% CI 69.0%–92.0%) (See Figure 2) [30,36,38,46,50,52,55]. When excluding one study that used a conservative definition of partner violence, heterogeneity was considerably reduced $I^2 = 61.2\%$ (95% CI 5%–84%), and the revised pooled odds ratio for any lifetime partner violence increased to 3.21 (95% CI 2.49–4.2) [52]. The corresponding funnel plots showed some asymmetry, which may indicate publication bias (Figure S1). Two high-quality studies measured lifetime physical partner violence among men with depressive disorders and reported estimates of 5.3% and 31.3%; both studies reported that men with depressive disorders were more likely to experience domestic violence compared to men with no mental disorders [50,55].

Past Year Domestic Violence. The median prevalence of any past year partner violence (7 studies) was 35.3% (IQR 16.0%–40.1%; range 1.7%–82.5%) among women with depressive disorders [34,35,37,38,45,53,54]. The pooled odds ratio for past year partner violence was 3.31 (95% CI 2.35–4.68); $I^2 = 32.8\%$ (95% CI 0.0%–73.0%) (see Figure 3) [34,35,37,38,45,54]. Funnel plots did not indicate asymmetry (see Figure S2). Only one high-quality study reported on any past year domestic violence among men, and identified a prevalence of 80.6% [53].

Dysthymia

Lifetime Domestic Violence. Two high-quality studies reported on the prevalence of any lifetime partner violence among men and women with dysthymia [30,55]. The largest study (a nationally representative survey of 34,653 non-institutionalised American residents) reported that women with dysthymic disorder had an increased likelihood of experiencing lifetime physical partner violence (OR 5.58 95% CI 4.60–6.76); similar findings were reported among men with dysthymic disorder (OR 4.84 95% CI 2.49–8.79). The study reported lifetime prevalence estimates of 20.0% among women and 3.9% among men with dysthymic disorder [55].

Past Year Domestic Violence. A cross-sectional survey of 364 pregnant and postpartum Vietnamese women reported a prevalence of 16.7% for any past year partner violence among six women with dysthymic disorder [54]. No difference in the odds of partner violence were detected among women with dysthymic disorder and women without a mental disorder (OR 1.39 95% CI 0.03–13.00) [54].

Anxiety Disorders

Lifetime Domestic Violence. The median prevalence of any lifetime partner violence (5 studies) for women with anxiety disorders was 27.6% (IQR 24.9%–72.7%; range 22.4%–89.9%) [30,38,50,52,55]. The pooled odds ratio for any lifetime partner




Figure 2. Pooled odds estimates for lifetime intimate partner violence among women with depressive disorders.
doi:10.1371/journal.pone.0051740.g002




Figure 3. Pooled odds estimates for past year intimate partner violence among women with depressive disorders.
doi:10.1371/journal.pone.0051740.g003

violence was 4.08 (95% CI 2.39–6.97), with high heterogeneity ($I^2 = 89.0$, 95% CI 77.0%–95.0%) (see Figure 4) [30,38,50,52,55]. Heterogeneity reduced considerably upon excluding two studies that did not use a validated instrument to measure partner violence ($I^2 = 56.5\%$ (95% CI 0%–88%), and the revised pooled odds ratio for any lifetime partner violence among women with anxiety disorder was 2.92 (95% CI 1.82–4.68) [52,55]. Only two high-quality studies measured lifetime partner violence among men. These studies found that men with anxiety disorders were significantly more likely to have experienced lifetime physical partner violence than those without a mental disorder, and reported prevalence estimates of 7.4% and 27.0% [50,55]. Limited data were available on the prevalence and odds of lifetime domestic violence among men and women with phobic [30,55,69] or somatoform disorders [69] (see Table S1).

Past Year Domestic Violence. Among women with anxiety disorders, the median prevalence of any past year partner violence (4 studies) was 28.4% (IQR 25.5%–42.2%, range 20.0%–80.5%) [38,45,53,54]. The pooled odds ratio for any past year partner violence was 2.29 (95% CI 1.31–4.02); ($I^2 = 0.0\%$, 95% CI 0.0%–90.0%) (see Figure 5) [38,45,54]. One high-quality study included men, and reported a prevalence of any past year partner violence of 74.0% among men with anxiety disorders [53].

Post Traumatic Stress Disorder (PTSD)

Lifetime Domestic Violence. The median prevalence of any lifetime partner violence (4 studies) among women with PTSD was 61.0% (IQR 41.1%–80.1%; range 29.4%–89.5%) [30,38,39,55]. The pooled odds ratio for any lifetime partner violence was 7.34 (95% CI 4.50–11.98) with high heterogeneity ($I^2 = 85.1\%$, 95% CI 52.0%–92.0%) (see Figure 6) [30,38,39,55]. Funnel plots did not indicate asymmetry (see Figure S3). One high-quality study included men, and reported an increased likelihood of lifetime physical partner violence among men with PTSD compared to men without a mental disorder (OR 9.66 95% CI 6.49–14.26), with a lifetime prevalence of 7.3% among men with PTSD [55].

Past Year Domestic Violence. A survey of female welfare recipients found that 27.0% of women with PTSD had experienced physical partner violence in the past year. The study reported a greater likelihood of past year physical partner violence among women with PTSD compared to women without a mental disorder (OR 3.62; 95% CI 2.32–5.67) [38].

Obsessive Compulsive Disorder

Lifetime Domestic Violence. A cross-sectional survey of 650 women attending primary care clinics in Pakistan reported a prevalence of 93.8% for any lifetime partner violence among women with obsessive compulsive disorder (OCD). The study




Figure 4. Pooled odds estimates for lifetime intimate partner violence among women with anxiety disorders.
doi:10.1371/journal.pone.0051740.g004

found that women with OCD had an increased likelihood of experiencing any lifetime partner violence compared to women without a mental disorder (OR 6.43; 95% CI 1.95–33.23) [30].

Past Year Domestic Violence. No high-quality studies reported the prevalence or odds of past year domestic violence among men or women with OCD.

Eating Disorder

Lifetime Domestic Violence. No high-quality studies reported the prevalence or odds of lifetime domestic violence among men or women with an eating disorder.

Past Year Domestic Violence. One birth cohort study reported that at age 21, 63.6% of 11 women with eating disorders reported past year physical partner violence; women with eating disorders were more likely to report partner violence compared to women without a mental disorder (OR 7.31 95% CI 1.76–35.10) [45].

Personality Disorder

Lifetime Domestic Violence. One study, a national survey of 34,653 non-institutionalised American residents, reported an increased odds of lifetime physical partner violence among both women (OR: 6.06 95% CI 5.35–6.86) and men (OR: 7.04 95% CI 5.30–9.43) with any personality disorder, and lifetime prevalence estimates of 21.4% and 5.4% respectively [55].

Past Year Domestic Violence. One birth cohort study reported that at age 21, 100.0% of three women with an antisocial

personality disorder reported past year physical partner violence [55].

Common Mental Disorders (depressive and/or anxiety disorders identified but not disaggregated; CMD)

Lifetime Domestic Violence. The median prevalence of any lifetime partner violence among women with CMDs (3 studies) was 48.0% (IQR 35.6%–63.2%, range 23.0%–78.1%); women with CMDs were reported to be more likely to experience any lifetime partner violence compared to those without a mental disorder [42,43,44].

Past Year Domestic Violence. A national survey of 7,047 UK householders reported an increased odds of any past year partner violence among women (OR: 4.4 95% CI: 3.32–5.82) and men (OR: 3.1 95% CI 2.18–4.39) with CMDs; prevalence estimates were 15.2% and 11.7% respectively [42].

Schizophrenia and Non-Affective Psychosis

Lifetime Domestic Violence. No high-quality studies reported the prevalence or odds of any lifetime partner violence among men or women with schizophrenia and non-affective psychosis. Two lower-quality studies, both conducted with psychiatric samples, reported that the lifetime prevalence of any partner violence ranged from 43.8%–83.3% among women with schizophrenia and non-affective psychosis [59,64].




Figure 5. Pooled odds estimates for past year intimate partner violence among women with anxiety disorders.
doi:10.1371/journal.pone.0051740.g005

Past Year Domestic Violence. One birth cohort study reported a prevalence of 43.8% for past year physical partner violence among 16 women with non-affective psychosis [45]. The same study reported that women with non-affective psychosis were more likely to experience past year partner violence compared to women without a mental disorder (OR 3.25; 95% CI 0.97–10.3).

Bipolar Affective Disorder

Lifetime Domestic Violence. One study, a nationally representative survey of 34,563 non-institutionalised American residents, identified an increased odds of lifetime physical partner violence among both women (OR 8.14; 95% CI 6.99–9.47) and men (OR 9.42; 95% CI 6.57–13.50) with bipolar disorder, and lifetime prevalence estimates of 26.7% and 7.1%, respectively [55].

Past Year Domestic Violence. No high-quality studies reported the prevalence or odds of past year domestic violence among men or women with bipolar disorder.

Findings from Longitudinal Studies

Three studies presented longitudinal data on the relationship between mental disorders and domestic violence [36,51,53]. A three year cohort study of 286 women found that among 14 women who were depressed and in violent relationship during the study, only one instance of depression predated the violence; rates of depression among the 12 women who had left the violent relationship within one year was no different from that of those who never experienced violence (25% vs. 23%) [36]. Bardone et

al, reporting data from the Dunedin birth cohort, found that depression at age 15 did not predict past-year relationship violence at age 21 (OR 1.22 95% CI 0.45–3.03) [51]. Conduct disorder at age 15 was, however, associated with later partner violence (OR 3.14, 95% CI 1.47–6.64). Fergusson et al, reporting data from the Christchurch birth cohort, found that any mental disorder (i.e. depressive, anxiety, conduct, or substance use disorder) at age 14–21 years was associated with past year partner violence at age 24–25 years, and that partner violence and mental disorder at 24–25 years were significantly associated even after adjusting for prior mental disorder and other antecedent and concurrent covariates [53].

Impact and Resource Use

Limited data were available on the impact of experiences of domestic violence and victims’ resource use. Two studies reported increased odds of substance misuse problems (OR of 3.4 and 4.1) among people experiencing domestic violence [41,47]. One study reported that people experiencing domestic violence had increased odds of suicidal ideation (OR 6.3) [41] and one reported that victims of recent violence experienced greater deprivation (e.g. eviction, homelessness, and food insufficiency) compared to non-victims [38]. Few studies provided details on victims’ resource use following domestic violence. One paper reported that women who experienced physical assault (n = 28 women, of which n = 6 reported spousal assault) were significantly more likely to have used emergency services (n = 11, 43%) (p = 0.002) and mental health services (n = 7, 24%) (p < 0.001) within the past 12 months




Figure 6. Pooled odds estimates for lifetime intimate partner violence among women with post-traumatic stress disorder.
doi:10.1371/journal.pone.0051740.g006

[35]. In relation to recency of abuse, another paper found that victims reporting past year violence were twice as likely to have received treatment for mental health problems (1.6%) than victims reporting violence prior to the last twelve months (13.6%) and were twice as likely (26.8% vs.13.6%) to report currently needing treatment [38].

Discussion

Key Findings

We found consistent evidence that both men and women with all types of mental disorders report a high prevalence and increased odds of domestic violence compared to people without mental disorder, with women more likely to experience abuse than men. Due to the limited number of high-quality studies it was not possible to calculate pooled odds of partner violence among men or for men or women with disorders other than depression, anxiety or PTSD. Studies on the prevalence and odds of domestic violence by non-intimate family members were also limited. Nonetheless, across a range of diagnoses, studies indicated that men and women with a mental disorder are at an increased likelihood of experiencing domestic violence compared to those without a mental disorder. For example, data from Wave II of the large US National Epidemiologic Survey on Alcohol and Related Conditions suggests that men and women with bipolar affective disorder were more than eight times more likely to report ever having been a victim of partner violence than people with no mental disorder [55].

Although a bi-directional causal relationship between domestic violence and mental disorder seems likely [5], there were insufficient data available from which to draw conclusions about causality. Due to the paucity of longitudinal studies we were only able to make a limited assessment of the temporality of the relationship between mental disorder and domestic violence and of whether recovery from mental disorder is associated with a reduction in risk of domestic violence, or vice versa. It was also not possible, due to insufficient data, to test whether the strength of the association between specific mental disorders and domestic violence varied with severity of violence. We were not able to examine strengths of association between specific mental disorders and recency of domestic violence (i.e. past year vs. lifetime) as odds ratios were calculated from studies with different study populations and measures of violence. This limits our ability to interpret direction of causality here, as recency of mental illness (i.e. past year vs. lifetime) has been shown to affect the strength of association between mental disorder and violence; similarly measurements of lifetime diagnoses may include individuals who may not have experienced a mental disorder during the observation period for acts of violence [71].

Strengths and Limitations

We used an inclusive search strategy and followed MOOSE and PRISMA reporting guidelines [14,15]. Our review extends previous research by examining the prevalence and odds of domestic violence across all mental disorders, presenting estimates

of the prevalence and odds of domestic violence separately by sex, restricting the scope to studies that used validated diagnostic instruments, and drawing upon the related body of research on mental disorder and victimisation.

All pooled odds ratio estimates indicated that women with mental disorders are at an increased likelihood of experiencing partner violence compared to women without mental disorders. However, in light of the high heterogeneity observed between studies, caution should be exercised when interpreting these figures. Due to a lack of data it was not possible to control for confounding factors when pooling prevalence estimates. When we excluded studies that used conservative definitions of domestic violence or employed non-validated instruments to measure domestic violence, heterogeneity was reduced. However, we do not know the relative contributions of the study setting and measurement of domestic violence to the heterogeneity, and it is likely that study country and known confounding factors (e.g. age, experiences of childhood abuse and substance misuse) may also affect variations in prevalence estimates. Funnel plot asymmetry also indicated the potential for publication bias among studies of depression.

Due to the lack of consistency in the data collected by the primary studies, we were unable to adjust our pooled estimates for potential confounders (e.g. childhood abuse). Furthermore, because of a lack of primary studies, we were unable to: calculate pooled estimates of the odds of domestic violence among men with mental disorders; to assess whether the odds of violence perpetrated by family members was increased among men and women with mental disorder; to analyse whether the prevalence and odds of domestic violence among men and women with mental disorder varied according to sexual preference.

Our meta-analyses were constrained by methodological and conceptual weaknesses in the primary studies. A third of studies scored <50% on quality appraisal criteria relating to selection bias; 23 studies used non-probability sampling, 15 did not provide information on the representativeness of their samples and 14 did not report on the likely impact of non-participation. Although most studies did not score poorly in relation to measurement bias, the measurement of domestic violence varied substantially, with regards to time period (lifetime vs. past year), type of abusive behaviour (physical, sexual, psychological or a combination of behaviours), and instrument. We reported separate estimates of the prevalence and odds of lifetime and past year domestic violence, but recognise that both measures are potentially problematic: recall bias may be present in studies that measure lifetime domestic violence, while participants in studies of past year violence may have had insufficient time to acknowledge or identify their abuse experiences as such [72]. Several papers measured only experiences of physical violence, whereas others included sexual and psychological abuse within their definition of domestic violence. The Conflict Tactics Scale (CTS) was the most commonly used instrument but has been criticised for gender neutrality, measuring acts out of context (not reporting whether acts of violence were in attack or defence) which may lead to differential misclassification bias across sexes. In addition, although the revised CTS partly addresses sexual violence [73], it does not measure other forms of violence [74]. Several papers reported modifying validated instruments without detailing how, if at all, the adapted measures were validated, or reported that they developed their own measures to assess abuse. These factors are likely to reduce both the reliability and comparability of study findings; therefore greater efforts are needed for the development of methodologically robust studies examining the relationship between domestic violence and mental illness [75].

Implications of Findings

This systematic review provides strong evidence of a high prevalence and increased odds of domestic violence across all mental disorders among both men and women and draws attention to key gaps in the evidence base. The findings of this review highlight the need for healthcare professionals to recognise the increased vulnerability of men and women with mental disorders to domestic violence and to be prepared to identify and address these issues in treatment plans. Current evidence suggests that identification of domestic violence is most effective when professionals are trained to understand the nature of domestic violence and its long term impact on health, to ask about domestic violence safely if abuse is ongoing, and have clear referral and care pathways for identified victims [76,77]. New guidelines from the World Health Organisation recommend that primary care and mental health services work in partnership with the domestic violence sector to address patients' needs [78]. Further research is needed, however, to investigate which interventions are effective in reducing domestic violence experienced by men and women with mental disorders and how to improve mental health after the abuse has stopped [5].

Supporting Information

Checklist S1 PRISMA Checklist.

(DOC)

Text S1 Electronic databases searched for systematic review.

(DOC)

Text S2 Search Terms for MEDLINE, EMBASE, PsycInfo.

(DOC)

Text S3 Definitions of domestic violence and mental disorders.

(DOC)

Table S1 Characteristics and reported outcomes of included studies.

(DOC)

Table S2 Critical Appraisal Checklist for Included Studies.

(DOC)

Table S3 Quality Appraisal Form.

(DOC)

Figure S1 Funnel plot: Odds of lifetime domestic violence among women with depressive disorder.

(TIF)

Figure S2 Funnel plot: Odds of past year domestic violence among women with depressive disorder.

(TIF)

Figure S3 Funnel plot: Odds of lifetime domestic violence among women with post-traumatic stress disorder.

(TIF)

Acknowledgments

We would like to thank the experts in this area who responded so helpfully in our request for data. We also gratefully acknowledge Fraser Anderson for her assistance during summer 2011; Mauricio Moreno for technical support; Professor Michael Dewey and the Institute of Psychiatry Biostatistics team for statistical support, and Seynam Kluitse for

administrative assistance. We also wish to thank Monika Janosik, Leo Koeser, Sharron Leung, Kazuyo Machiyama, Mauricio Moreno, Michael Dewey, Oliver Schumann, and Tracy Teng for translating articles that were published in non-English languages.

References

- Garcia-Moreno C, Jansen HAFM, Ellsberg M, Heise L, Watts CH (2006) Prevalence of intimate partner violence: findings from the WHO multi-country study on women's health and domestic violence. *The Lancet* 368: 1260–1269.
- McClennen JC (2005) Domestic violence between same-gender partners: recent findings and future research. *Journal of Interpersonal Violence* 20: 149–154.
- Campbell JC (2002) Health consequences of intimate partner violence. *The Lancet* 359: 1331–1336.
- Golding MJ (1999) Intimate Partner Violence as a Risk Factor for Mental Disorders: A Meta-Analysis. *Journal of Family Violence* 14: 99–132.
- Howard LM, Trevillion K, Khalifeh H, Woodall A, Agnew-Davies R, et al. (2010) Domestic Violence and Severe Psychiatric Disorders: prevalence and interventions. *Psychological Medicine* 40: 881–893.
- MacMillan LH, Wathen CN, Jamieson E, Boyle M, McNutt AL, et al. (2006) Approaches to Screening for Intimate Partner Violence in Health Care Settings: A Randomized Trial. *JAMA* 296: 530–536.
- Rivara FP, Anderson ML, Fishman P, Bonomi AE, Reid RJ, et al. (2007) Intimate partner violence and health care costs and utilization for children living in the home. *Pediatrics* 120: 1270–1277.
- Centers for Disease Control and Prevention, National Center for Injury Prevention and Control (2003) *Costs of Intimate Partner Violence Against Women in the United States*. Atlanta: GA: Centers for Disease Control and Prevention.
- Walby S (2009) *The Cost of Domestic Violence: Up-date 2009*. Lancaster: Lancaster University.
- Brown GW, Harris TO, Hepworth C, Robinson R (1994) Clinical and Psychosocial Origins of Chronic Depressive Episodes II: A Patient Enquiry. *British Journal of Psychiatry* 165: 457–465.
- Khalifeh H, Dean K (2010) Gender and violence against people with severe mental illness. *International Review of Psychiatry* 22: 535–546.
- Friedman SH, Loe S (2007) Incidence and prevalence of intimate partner violence by and against women with severe mental illness. *Journal of Women's Health* 16: 471–480.
- Hughes K, Bellis MA, Jones L, Wood S, Bates G, et al. (2012) Prevalence and risk of violence against adults with disabilities: a systematic review and meta-analysis of observational studies. *The Lancet* 379: 1621–1629.
- Moher D, Liberati A, Tetzlaff J, Altman DG (2009) Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Medicine* 6.
- Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, et al. (2000) Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA* 283: 2008–2012.
- Maniglio R (2009) Severe mental illness and criminal victimization: a systematic review. *Acta Psychiatrica Scandinavica* 119: 180–191.
- Home Office (2005) *Domestic violence: a national report*. London: Home Office.
- Ramsay J, Carter Y, Davidson L, Dunne D, Eldridge S, et al. (2009) Advocacy interventions to reduce or eliminate violence and promote the physical and psychosocial well-being of women who experience intimate partner abuse. *Cochrane Database of Systematic Reviews*: Art. No.: CD005043. DOI: 005010.001002/14651858.CD14005043.pub14651852.
- World Health Organization (1993) *Composite International Diagnostic Interview (CIDI): Interviewer's Manual*. Geneva: World Health Organization.
- NICE (2008) *The guidelines manual*. London: National Institute for Health and Clinical Excellence.
- Wing JK (1994) *The Schedules for Clinical Assessment in Neuropsychiatry*. Geneva: World Health Organization-Division of Mental Health.
- Critical Appraisal Skills Programme (2012) *Critical Appraisal Skills Programme: making sense of the evidence*. Oxford: Critical Appraisal Skills Programme.
- Downs SH, Black N (1998) "The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions." *J Epidemiol Comm Health* 52: 377–384.
- Loney PL, Chambers LW (2000) "Critical appraisal of the health research literature: prevalence or incidence of a health problem." *Chronic Dis Canada* 19: 170–177.
- Saha S, Chant D, Welham J, McGrath J (2005) "A systematic review of the prevalence of schizophrenia." *PLoS Med* 2: 141.
- Greenland S, O'Rourke K (2001) On the bias produced by quality scores in meta-analysis, and a hierarchical view of proposed solutions. *Biostatistics* 2: 463–467.
- DerSimonian R, Laird N (1986) Meta-analysis in clinical trials. *Controlled Clinical Trials* 7: 177–188.
- Egger M, Smith GD (1998) "Meta-analysis bias in location and selection of studies." *BMJ* 316: 61.
- StataCorp LP (2009) *STATA Statistical Software: Release 11 Version*. tx: College Station.
- Ayub M, Irfan M, Nasr T, Lutufullah M, Kingdon D, et al. (2009) Psychiatric morbidity and domestic violence: A survey of married women in Lahore. *Social Psychiatry and Psychiatric Epidemiology* 44: 953–960.
- Iverson KM, Gradus JL, Resick PA, Smith KF, et al. (2011) Cognitive-behavioral therapy for PTSD and depression symptoms reduces risk for future intimate partner violence among interpersonal trauma survivors. *Journal of Consulting & Clinical Psychology* 79: 193–202.
- Saliou V, Fichelle A, McLoughlin M, Thauvin I, Lejoyeux M (2005) Psychiatric disorders among patients admitted to a French medical emergency service. *General Hospital Psychiatry* 27: 263–268.
- Zanarini MC, Frankenburg FR, Reich DB, Marino MF, Haynes MC, et al. (1999) Violence in the lives of adult borderline patients. *Journal of Nervous & Mental Disease* 187: 65–71.
- Hammen C, Brennan PA (2002) Interpersonal dysfunction in depressed women: impairments independent of depressive symptoms. *Journal of Affective Disorders* 72: 145–156.
- Thurston WE, Patten S, Lagendyk LE (2006) Prevalence of violence against women reported in a rural health region. *Canadian Journal of Rural Medicine* 11: 259–267.
- Andrews B, Brown GW (1988) Marital violence in the community: A biographical approach. *British Journal of Psychiatry* 153: 305–312.
- Hicks MHR, Li ZH (2003) Partner violence and major depression in women: A community study of Chinese Americans. *Journal of Nervous and Mental Disease* 191: 722–729.
- Tolman RM, Rosen D (2001) Domestic violence in the lives of women receiving welfare - Mental health, substance dependence, and economic well-being. *Violence against Women* 7: 141–158.
- Baker CK, Norris FH, Diaz DM, Perilla JL, Murphy AD, et al. (2005) Violence and PTSD in Mexico: Gender and Regional Differences. *Social Psychiatry and Psychiatric Epidemiology* 40: 519–528.
- Chang DF, Shen BJ, Takeuchi DT (2009) Prevalence and demographic correlates of intimate partner violence in Asian Americans. *International Journal of Law & Psychiatry* 32: 167–175.
- Jirapramukpiat T, Harpham T, Prince M (2011) Family violence and its 'adversity package': a community survey of family violence and adverse mental outcomes among young people. *Soc Psychiatry Psychiatr Epidemiol* 46: 825–831.
- Jonas S, Khalifeh H, Bebbington P, McManus S, Brugha T, et al. Intimate partner violence and psychiatric disorder in England: Results from the 2007 Adult Psychiatric Morbidity Survey. Unpublished.
- Patel V, Kirkwood BR, Pednekar S, Pereira B, Barros P, et al. (2006) Gender disadvantage and reproductive health risk factors for common mental disorders in women: a community survey in India. *Archives of General Psychiatry* 63: 404–413.
- Simsek Z, Ak D, Altindag A, Gunes M (2008) Prevalence and predictors of mental disorders among women in Sanliurfa, Southeastern Turkey. *Journal of Public Health*.
- Danielson KK, Moffitt TE, Caspi A, Silva PA (1998) Comorbidity between abuse of an adult and DSM-III-R mental disorders: Evidence from an epidemiological study. *American Journal of Psychiatry* 155: 131–133.
- Deyessa N, Berhane Y, Alem A, Mary E, Emmelin M, et al. (2009) Intimate partner violence and depression among women in rural Ethiopia: a cross-sectional study. *Clinical Practice & Epidemiology in Mental Health* 5: 1–10.
- Gass JD, Stein DJ, Williams DR, Seedat S (2011) Gender Differences in Risk for Intimate Partner Violence among South African Adults. *Journal of Interpersonal Violence* 26: 2764–2789.
- O'Leary KD, Tittle N, Bromet EJ, Gluzman SF (2008) Descriptive epidemiology of intimate partner aggression in Ukraine. *Social Psychiatry & Psychiatric Epidemiology* 43: 619–626.
- Schumacher JA, Coffey SF, Norris FH, Tracy M, Clements K, et al. (2010) Intimate partner violence and Hurricane Katrina: predictors and associated mental health outcomes. *Violence & Victims* 25: 588–603.
- Afifi TO, MacMillan H, Cox BJ, Asmundson GJ, Stein MB, et al. (2009) Mental health correlates of intimate partner violence in marital relationships in a nationally representative sample of males and females. *Journal of Interpersonal Violence* 24: 1398–1417.
- Bardone AM, Moffitt TE, Caspi A, Dickson N, Silva PA (1996) Adult mental health and social outcomes of adolescent girls with depression and conduct disorder. *Development and Psychopathology* 8: 811–829.

Author Contributions

Conceived and designed the experiments: LMH GF SO KT. Performed the experiments: KT SO. Analyzed the data: KT SO. Contributed reagents/materials/analysis tools: KT SO. Wrote the paper: KT SO GF LMH. Critical revisions of the manuscript: KT SO GF LMH.

52. DeKlyen M, Brooks-Gunn J, McLanahan S, Knab J (2006) The mental health of married, cohabiting and non-coresident parents with infants. *American Journal of Public Health* 96: 1836–1841.
53. Fergusson DM, Horwood L, Ridder EM (2005) Partner violence and mental health outcomes in a New Zealand birth cohort. *Journal of Marriage and Family* 67: 1103–1119.
54. Fisher J, Thach T, Buoi TL, Kriitmaa K, Rosenthal D, et al. (2010) Common perinatal mental disorders in northern Viet Nam: community prevalence and health care use. *Bulletin of the World Health Organization* 88: 737–745.
55. Grant BF, Goldstein RB (2011) Unpublished Data (Wave 2 of the National Epidemiologic Survey on Alcohol and Related Conditions). Supplements Pietrzak, R.H. et al “Prevalence and Axis I comorbidity of full and partial posttraumatic stress disorder in the United States: Results from Wave 2 of the National Epidemiologic Survey on Alcohol and Related Conditions”. *Journal of Anxiety Disorders* 25: 456–465.
56. Kaminer D, Grimsrud A, Myer L, Stein DJ, Williams DR (2008) Risk for post-traumatic stress disorder associated with different forms of interpersonal violence in South Africa. *Social Science & Medicine* 67: 1589–1595.
57. Dennis MF, Flood AM, Reynolds V, Araujo G, Clancy CP, et al. (2009) Evaluation of Lifetime Trauma Exposure and Physical Health in Women With Posttraumatic Stress Disorder or Major Depressive Disorder. *Violence Against Women* 15: 618–627.
58. Duran B, Oetzel J, Parker T, Malcoe LH, Lucero J, et al. (2009) Intimate partner violence and alcohol, drug, and mental disorders among American Indian women from southwest tribes in primary care. *American Indian and Alaska Native Mental Health Research* 16: 11–27.
59. Friedman S, Loue S, Goldman Heaphy E, Mendez N (2011) Intimate Partner Violence Victimization and Perpetration by Puerto Rican Women with Severe Mental Illnesses. *Community Mental Health Journal* 47: 156–163.
60. Najavits LM, Sonn J, Walsh M, Weiss RD (2004) Domestic violence in women with PTSD and substance abuse. *Addictive Behaviors* 29: 707–715.
61. Cascardi M, O’Leary K, Lawrence EE, Schlee KA (1995) Characteristics of women physically abused by their spouses and who seek treatment regarding marital conflict. *Journal of Consulting and Clinical Psychology* 63: 616–623.
62. Vahip I, Doganavsargil O (2006) [Domestic violence and female patients]. *Turk Psikiyatri Dergisi* 17: 107–114.
63. Tuten M, Jones HE, Tran G, Svikis DS (2004) Partner violence impacts the psychosocial and psychiatric status of pregnant, drug-dependent women. *Addictive Behaviors* 29: 1029–1034.
64. Wong SPY, Phillips MR (2009) Nonfatal Suicidal Behavior among Chinese Women Who Have Been Physically Abused by Their Male Intimate Partners. *Suicide and Life Threatening Behavior* 39: 648–658.
65. Cerulli C, Talbot NL, Tang W, Chaudron LH (2011) Co-Occurring Intimate Partner Violence and Mental Health Diagnoses in Perinatal Women. *Journal of Women’s Health* 0: 1–7.
66. Weizmann-Henelius G, Viemero V, Eronen M (2004) Psychological Risk Markers in Violent Female Behavior. *The International Journal of Forensic Mental Health* 3: 185–196.
67. Contreras-Pezzotti L, Arteaga-Medina J, Latorre J, Folino J, Campo-Arias A (2011) Association between Intimate Partner Violence and Posttraumatic Stress Disorder: A Case-Control Study. *Rev Colomb Psiquiat* 39: 85–92.
68. Lejoyeux M, Zillhardt P, Chieze F, Fichelle A, Mc Loughlin M, et al. (2002) Screening for domestic violence among patients admitted to French emergency service. *European Psychiatry* 17: 479–483.
69. Roberts GL, Lawrence JM, Williams GM, Raphael B (1998) The impact of domestic violence on women’s mental health. *Australian & New Zealand Journal of Public Health* 22: 796–801.
70. Yasan A, Saka G, Ozkan M, Ertem M (2009) Trauma type, gender, and risk of PTSD in a region within an area of conflict. *Journal of Traumatic Stress* 22: 663–666.
71. Van Dorn R, Volavka J, Johnson N (2012) Mental disorder and violence: is there a relationship beyond substance use? *Social Psychiatry & Psychiatric Epidemiology* 47: 487–503.
72. Feder G, Ramsay J, Dunne D, Rose M, Arsene C, et al. (2009) How far does screening women for domestic (partner) violence in different health-care settings meet criteria for a screening programme? Systematic reviews of nine UK, National Screening Committee criteria. *Health Technology Assessment* 13(16) (pp iii-113). 2009 Date of Publication: March 2009: iii-113.
73. Straus MA, Hamby LS, Boney-McCoy S, Sugarman BD (1996) The Revised Conflict Tactics Scales (CTS2) development and preliminary psychometric data. *Journal of Family Issues* 17: 283–316.
74. Loseke DR, Kurz D (2005) Men’s violence toward women is the serious social problem. In: Loseke DR, Gelles RJ, Cavanaugh MM, editors. *Current controversies on family violence*, 2nd edition. Newbury Park: Sage. pp. 79–96.
75. Van Dorn R, Desmarais S, Singh J (2011) Violence and victimization in adults with severe mental illness: Part 1 - An application of integrated data analysis. In: I . Needham, T . Palmstierna, R . Almvik, Oud N, editors. *Proceedings of the 7th European Congress on Violence in Clinical Psychiatry*. Amsterdam: Kavanah.
76. Feder G, Agnew-Davies R, Baird K, Dunne D, Eldridge S, et al. (2011) Identification and Referral to Improve Safety (IRIS) of women experiencing domestic violence with a primary care training and support programme: a cluster randomised controlled trial. *Lancet* 11: 1–8.
77. Trevillion K, Byford S, Cary M, Rose D, Oram S, et al. (Submitted) A pilot study of domestic violence advocacy integrated within community mental health services.
78. World Health Organization (forthcoming) Guideline Development Group on Policy and Clinical practice guidelines for responding to violence against women (2011–2012). Geneva: World Health Organization.