

2021

## Experimental methods to study the pathogenesis of human enteric RNA viruses

Somya Aggarwal

*Washington University School of Medicine in St. Louis*

Ebrahim Hassan

*Washington University School of Medicine in St. Louis*

Megan T Baldrige

*Washington University School of Medicine in St. Louis*

Follow this and additional works at: [https://digitalcommons.wustl.edu/open\\_access\\_pubs](https://digitalcommons.wustl.edu/open_access_pubs)

---

### Recommended Citation

Aggarwal, Somya; Hassan, Ebrahim; and Baldrige, Megan T, "Experimental methods to study the pathogenesis of human enteric RNA viruses." *Viruses*. 13,6. . (2021).  
[https://digitalcommons.wustl.edu/open\\_access\\_pubs/10378](https://digitalcommons.wustl.edu/open_access_pubs/10378)

This Open Access Publication is brought to you for free and open access by Digital Commons@Becker. It has been accepted for inclusion in Open Access Publications by an authorized administrator of Digital Commons@Becker. For more information, please contact [vanam@wustl.edu](mailto:vanam@wustl.edu).

Review

# Experimental Methods to Study the Pathogenesis of Human Enteric RNA Viruses

Somya Aggarwal <sup>1</sup>, Ebrahim Hassan <sup>1</sup>  and Megan T. Baldrige <sup>1,2,\*</sup> 

<sup>1</sup> Division of Infectious Diseases, Department of Medicine, Edison Family Center for Genome Sciences & Systems Biology, Washington University School of Medicine, St. Louis, MO 63110, USA; s.aggarwal@wustl.edu (S.A.); e.hassan@wustl.edu (E.H.)

<sup>2</sup> Department of Molecular Microbiology, Washington University School of Medicine, St. Louis, MO 63110, USA

\* Correspondence: mbaldrige@wustl.edu

**Abstract:** Every year, millions of children are infected with viruses that target the gastrointestinal tract, causing acute gastroenteritis and diarrheal illness. Indeed, approximately 700 million episodes of diarrhea occur in children under five annually, with RNA viruses norovirus, rotavirus, and astrovirus serving as major causative pathogens. Numerous methodological advancements in recent years, including the establishment of novel cultivation systems using enteroids as well as the development of murine and other animal models of infection, have helped provide insight into many features of viral pathogenesis. However, many aspects of enteric viral infections remain elusive, demanding further study. Here, we describe the different in vitro and in vivo tools available to explore different pathophysiological attributes of human enteric RNA viruses, highlighting their advantages and limitations depending upon the question being explored. In addition, we discuss key areas and opportunities that would benefit from further methodological progress.



**Citation:** Aggarwal, S.; Hassan, E.; Baldrige, M.T. Experimental Methods to Study the Pathogenesis of Human Enteric RNA Viruses. *Viruses* **2021**, *13*, 975. <https://doi.org/10.3390/v13060975>

Academic Editors: Akio Adachi and Masako Nomaguchi

Received: 15 April 2021

Accepted: 20 May 2021

Published: 25 May 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in

**Keywords:** human norovirus; human rotavirus; human astrovirus; gastroenteritis; immortalized cell lines; human enteroids

## 1. Introduction

Acute gastroenteritis, characterized by symptoms including nausea, vomiting, malaise, abdominal pain, fever, and diarrhea, is one of the most common health problems worldwide. More than 700 million cases occur annually in children under five years of age, resulting in few deaths in developed countries, but more than 2 million deaths in developing countries [1]. A diverse group of viral, bacterial, and parasitic pathogens are responsible for acute gastroenteritis, but among these, enteric viruses cause almost half of the cases affecting patients of all ages worldwide, and in the United States, viruses are the leading cause [2]. Viral gastroenteritis is usually self-limiting with symptom resolution occurring

and picobirnaviruses [1]. With progressive advances in sequencing-based approaches to identify causative pathogens, we are increasingly adept at defining those viruses causing symptoms, characterizing their prevalence during asymptomatic infection, and identifying novel viral pathogens [9,10]. However, characterization of the effects of established or novel strains on the host requires the use of both in vitro and in vivo experimental model systems, many of which have, until recently, been lacking for these pathogens.

The past decade has witnessed the advent of intestinal organoid-based studies of virus–host interactions as well as an expansion in available animal models, both of which have dramatically increased our capacity to decipher physiologically relevant viral and cellular processes during infection. Here, we discuss the in vitro and in vivo tools that are currently used to study the pathogenesis of human RNA enteric viruses, focusing on norovirus, rotavirus, and astrovirus, to provide a useful compendium for those beginning to explore the field of gastroenteritis (Tables 1 and 2).

**Table 1.** Summary of commonly employed in vitro and in vivo models to study human rotavirus, norovirus, and astrovirus, with most broadly-used approaches shown in red.

Virus	Method	Origin	Experimental Model	Viral Strain(s)	References
<i>Human Rotavirus</i>	<i>in vitro</i>		<i>Immortalized cells</i>		
		Human colon adenocarcinoma	Caco-2	Trypsin-activated HRV-Wa or SA114F	[11,12]
		Human colorectal adenocarcinoma	HT-29	Trypsin-activated SA114F	[12]
		Human colorectal adenocarcinoma	Genetically-modified HT-29 ( <i>STAG2</i> <sup>-/-</sup> , <i>STAT1</i> <sup>-/-</sup> , and <i>STING</i> <sup>-/-</sup> ) HT-29	Trypsin-activated HRV-Wa, G2P[4], G4P[6], G12P[4], G9P[8], G8P[11], G8P[10]	[13]
		Human colorectal adenocarcinoma	HT29-MTX	Trypsin-activated SA114F	[14]
		African Green Monkey Kidney	<b>MA-104 and Vero</b>	Trypsin-activated HRV	[15]
		African Green Monkey Kidney	Genetically-modified ( <i>LRGUK</i> <sup>-/-</sup> , <i>WR62</i> <sup>-/-</sup> , <i>EMX2</i> <sup>-/-</sup> ) Vero cells	Trypsin-activated RV3, CDC-9, Rotarix and 116E	[16,17]
			<i>Human Intestinal Enteroids (HIEs)</i>		
		Human intestinal biopsies	3D-HIEs	Trypsin activated G1P[8] and G9P[8]	[18,19]
		Human intestinal biopsies	Differentiated 2D-monolayer and Transwell HIEs	Trypsin activated HRV	[20]
		Human intestinal			

Table 1. Cont.

Virus	Method	Origin	Experimental Model	Viral Strain(s)	References
<i>HIEs</i>					
		Human intestinal biopsies	Differentiated 2D-monolayer & Transwell HIEs	Bile-treated GII.1, GII.3 & GII.4 stool filtrate	[20,31]
		Human intestinal biopsies	Genetically-modified ( <i>IFNAR1</i> <sup>-/-</sup> , <i>IFNLR1</i> <sup>-/-</sup> , <i>STAT1</i> <sup>-/-</sup> , <i>MAVS</i> <sup>-/-</sup> , and <i>STAT1</i> <sup>-/-</sup> <i>STAT2</i> <sup>-/-</sup> ) HIEs	GI.3 & GII.4 stool filtrate	[32]
		Human intestinal biopsies	Genetically-modified ( <i>FUT2</i> <sup>-/-</sup> ) HIEs	GI.3, GII.4 & GII.17 stool filtrate	[33]
	<i>in vivo</i>	<i>Animal Model</i>			
		Fish	Zebrafish	GI.7, GII.2, GII.3, GII.4, GII.6 from stool suspension	[34]
		Rodent	Humanized BALB/c <i>Rag</i> <sup>-/-</sup> <i>Il2rg</i> <sup>-/-</sup> mice (engrafted with human CD34+)	GI.3a, GII.4, GII.6 from stool suspension	[35]
		Pig	Gnotobiotic piglets	GI.4 stool filtrate	[36,37]
		Pig	<i>Rag2</i> <sup>-/-</sup> <i>Il2rg</i> <sup>-/-</sup> gnotobiotic piglets	GI.4 stool filtrate	[38]
		Non-human primate	Pig-tailed macaques	GI.3 from stool suspension	[39]
		Non-human primate	Chimpanzee	GI.1 stool filtrate	[40]
<i>Human Astrovirus</i>	<i>in vitro</i>	<i>Immortalized Cell line</i>			
		Human colon adenocarcinoma	<b>Caco-2</b>	HAstV-1, 2, 3, 4, 5, 6, 7	[41–43]
		Human colon adenocarcinoma	Genetically-modified Caco-2 ( <i>CLTC</i> silencing)	Trypsin-activated HAstV-8	[44]
		Human colorectal adenocarcinoma	HT-29	Trypsin-activated HAstV-1, 2, 3, 4, 5, 6, 7	[41]
<i>HIEs</i>					
		Human intestinal biopsies	3D HIEs	HAstV-1	[45,46]
		Human intestinal biopsies	Differentiated 2D-monolayer in transwell	HAstV-VA1; HAstV-MLB1 & Trypsin-activated HAstV-1	[43,47]

## 2. Human RNA Enteric Viruses

Human RNA enteric viruses include an array of viral genera that are predominantly transmitted through the fecal-oral route and can invade and replicate in the gastrointestinal tract. These viruses can be categorized as: (1) viruses predominantly infecting the small intestinal epithelium to cause acute gastroenteritis, such as rotaviruses, caliciviruses, and astroviruses; (2) viruses that briefly infect the gastrointestinal tract before causing clinical disease at extraintestinal sites, such as polioviruses, coxsackieviruses, and hepatitis A (HAV) and hepatitis E (HEV); and (3) viruses that infect the intestinal tract at late stages of disease in immunocompromised hosts such as human immunodeficiency virus. In this review, we will predominantly focus on this first category of viruses that infect the intestinal epithelium, and here will briefly describe what is known about their pathogenesis and cellular tropism.

### 2.1. Rotavirus

Human rotavirus (HRV) is a non-enveloped, segmented, double-stranded RNA virus from the family Reoviridae, first identified from duodenal biopsies and fecal samples of children suffering from diarrhea [48,49]. A leading cause of acute gastroenteritis in children under five, HRV infection can range from asymptomatic to causing severe non-bloody diarrhea with vomiting and fever lasting from 3 to 8 days [50–53]. In some cases, rapid dehydration and electrolyte imbalance can lead to death [54,55], and indeed prior to the development of effective vaccines, HRV claimed more than 500,000 lives each year and accounted for an estimated \$1 billion in health care costs in the US annually [4,56]. RV can be classified into seven major serogroups (A–G). Groups A, B, and C infect both humans and animals, while the rest have only been found in animals to date, and Group A has been established as the most common RV responsible for causing human illness [57]. These viruses possess a distinct cell tropism, predominantly infecting the mature enterocytes and enteroendocrine cells of the small intestine [19,58]. It has been recently observed that transmission of HRV can occur via vesicle-cloaked virus clusters, indicating that both free and clustered virions may contribute to disease pathogenesis [59].

HRV antigen can be detected in stool samples using ELISA or immunochromatography, but qPCR-based assays provide greater sensitivity and allow genotyping of virus isolates, and thus are routinely used in vaccine and epidemiological studies [60,61]. Currently, two live, attenuated RV vaccines are widely used worldwide: the RV5 or RotaTeq vaccine is a pentavalent vaccine composed of five bovine-human RV strains [62], while the RV1 or Rotarix vaccine is monovalent containing one HRV strain [63]. Since the introduction of these vaccines in 2006 in national vaccination programs of America and Europe [64], a massive reduction in the incidence of HRV infection has occurred [63,65,66]. Based on the performance of these vaccines, in 2009, the WHO extended this recommendation worldwide. Although the effectiveness of RV vaccines remains higher in developed countries than in developing countries, potentially due to a variety of factors including microbiota variation, implementation of these vaccines has universally translated into a substantial

protein VP1, the NoV genus is divided into ten genogroups (GI-GX) which are further sub-divided into different genotypes containing individual virus strains [74]. Among these, GI, GII and GIV strains infect humans and genotype 4 of GII (GII.4) is responsible for the majority of human outbreaks [75]. NoVs are generally species-specific, with each genogroup of NoV infecting distinct groups of hosts [76]. While the cellular tropism for HuNoV has been a subject of some controversy, the development of in vitro culture systems, discussed in detail below, have suggested mature enterocytes and B cells as targets for HuNoV [29,31,77]. A dual tropism for immune cells and intestinal epithelial cells, specifically enteroendocrine cells, is supported by histologic analyses of intestinal tissues from HuNoV-infected immunocompromised patients [78,79]. HuNoV clusters cloaked within vesicles have, as for HRV, been implicated in viral transmission [59].


Diagnosis of HuNoV can be performed using commercially available enzyme immunoassays or immunochromatographic assays on samples including diarrheal stool or vomitus. Multiple real-time qRT-PCR (qPCR) assays have also been developed for standard diagnostic use [80]. While there are no HuNoV vaccines currently available, multiple vaccine candidates designed to target the viral capsid have been evaluated in clinical trials [81].

### 2.3. *Astrovirus*

Human astroviruses (HAstVs) were first reported in 1975, after electron microscopic analysis of stool following an outbreak of pediatric diarrhea revealed virions with a characteristic star-like appearance [82,83]. HAstVs are non-enveloped, positive-sense RNA viruses from the family *Astroviridae* which have been classified into three divergent groups: the classic HAstVs, the nonclassic HAstV-MLB and HAstV-VA/HMO groups [7]. Among these, classic HAstVs include eight serotypes and are responsible for 2–9% of all acute viral gastroenteritis in children worldwide [7]. The more recently discovered nonclassic AstVs are less well-studied, and their pathological effects and prevalence worldwide remain unclear. In general, HAstV induces diarrheal disease that is milder than HRV or HuNoV, and is associated with abdominal pain, vomiting, and fever that lasts 2–3 days. Although infections are generally self-limiting, immunocompromised individuals may succumb to disseminated infection [84,85]. Asymptomatic infections have been reported in children as well as adults [86,87]. HAstV antigen has been found in the mature enterocytes of the small intestine [88–90], and an in vitro cultivation system in human intestinal enteroids supports this tropism, but also identifies goblet cells and intestinal progenitor cells as potentially permissive to infection [43]. No vaccines are currently available against HAstVs, and dehydration secondary to gastroenteritis is treated with oral or intravenous fluids. Currently, real-time RT-quantitative PCR assays are sensitive, fast, and reproducible for HAstV diagnosis [91], and next-generation sequencing continues to facilitate the identification of new emerging strains [92,93].

## 3. In Vitro Tools to Study Human Enteric RNA Viruses

practical. An important consideration for in vitro viral experimentation, applicable to enteric RNA viruses [97–100], is the potential for passaged viruses to mutate substantially as they adapt to in vitro conditions. While fundamental aspects of viral pathogenesis may remain intact even with viral genetic changes, interpretation of results should always be considered with this caveat in mind. Here, we will discuss the in vitro systems available for studying human RNA enteric viruses (Figure 1).



**Figure 1.** In vitro methods commonly used for the study of human noroviruses (HuNoVs), rotaviruses (HRVs) and astroviruses (HAstVs). Immortalized cell lines and human intestinal enteroids have been used for the study of human enteric viruses. Modifications to cells either to facilitate infections and/or interrogate the role of host genes during infection can be performed using methods including CRISPR-Cas9-based editing, gene overexpression or silencing. Created with [BioRender.com](https://www.biorender.com) (accessed on 21 January 2021).

### 3.1. Immortalized Cell Lines

Immortalized cell lines provide a pure population of cells that are easy to maintain and have the potential to divide indefinitely, and thus many host–enteric virus interaction studies have been performed using cancer-derived or immortalized cell lines. Different human intestinal cell lines, manifesting specific functions and characteristics of the gut epithelium, are widely used, and African green monkey kidney cell lines, which are more susceptible to a variety of viruses, have also served as useful lines for enteric viral research [101].

cells, but requires trypsin during the entire course of infection for efficient replication [11]. Caco-2 cells also support HAstV replication, for which the usage of serum-free media during infection is recommended [41,108]. While Caco-2 cells express histo-blood group antigens, which are key attachment factors for HuNoV [109–113], they do not consistently support HuNoV replication in monolayer conditions [96,114].

HT-29 cells are a pluripotent, heterogeneous cell line that grows as a multilayer of unpolarized, undifferentiated cells with less than 5% differentiated mucus-secreting cells and columnar absorptive cells [115]. These cells have the potential to further differentiate: in the absence of glucose, HT-29 cells can undergo a typical enterocytic differentiation [116], whereas in the absence of serum, ~50% of cells differentiate into goblet-like cells expressing mucins [117]. HT-29 cells support in vitro growth of HRV [118], and a variant, the mucus-secreting HT-29-MTX line generated by differentiating HT-29 into mature goblet cells using methotrexate [115], has been successfully employed to investigate the role of glycans in promoting HRV virulence [14]. HT-29 cells can also support efficient replication of some HAstV serotypes when grown without glucose [41]. Unlike HRV and HAstV, HuNoV does not replicate in HT-29 cells [96,114,119].

HuNoV GI and GII strains have been previously cultivated in 3-dimensional models wherein intestinal epithelial cell line INT-407 was grown on collagen-coated porous micro-carrier beads in rotating-wall vessel bioreactors [119], an approach also used for HuNoV replication with Caco-2 cells [120]. However, challenges in replicating this model have prevented its widespread implementation [121–123].

### 3.1.2. B Cell Lines

After many unsuccessful attempts to grow HuNoV in immortalized cell lines, the identification of an immune cell tropism for a strain of murine norovirus (MNoV), a prominent small animal model for HuNoV studies, suggested potential utility in exploring immune cells for HuNoV cultivation [124]. GII.4 strains of HuNoV have since been cultivated in human-derived transformed B-cell line BJABs [29,77], an infection which is enhanced in the presence of histo-blood group antigen carbohydrates that facilitate viral attachment to the B cells [77]. However, low virus yield and inconsistent results among different laboratories have been reported [29].

### 3.1.3. Non-Human Primate Cell Lines

African green monkey (*Cercopithecus aethiops*) kidney cell lines are the most common non-human primate cell lines used for enteric viral research. These cell lines are susceptible to many viruses due to the absence of type I interferon (IFN) and cyclin-dependent kinase inhibitor genes [125,126]. Indeed, African green monkey kidney cells were the first used for the growth of HRV [127].

Vero cells are the most widely accepted immortalized cell line for the development of human viral vaccines, and have been used for poliovirus, rabies virus, influenza virus, and HRV vaccine propagation [128], though yield of high-titer virus has been a frequent



HAdV replication as well [41]. These cell lines are also used for the study of enteric RNA viruses causing extraintestinal disease including HAV and HEV [135–137].

#### Advantages and Disadvantages of Using Immortalized Cell Lines

Because they are cost-effective and have the potential to grow indefinitely, immortalized cell lines offer multiple advantages over alternate approaches (Table 2). They are easy to maintain and devoid of any ethical concerns associated with the usage of animals or human tissues. Although cell lines serve as a powerful tool for viral studies, due to the homogeneity of the cell population and the lack of host factors such as immune cells and the enteric nervous system, they do not necessarily represent the *in vivo* tissue environment. These lines have been genetically manipulated, either via transformation in cancer cell lines or via artificial expression of cancer genes to drive indefinite proliferation. This manipulation can alter key phenotypes including immune responses and cellular functions. Further, serial passaging can lead to genetic drift, driving heterogeneity in cell populations that can confer genotypic and phenotypic variation, both over time within a laboratory and between research groups [138].

#### 3.2. Primary Cells

Primary cells are isolated directly from tissue and are grown *in vitro*, retaining morphological and functional properties of their tissue of origin. For viral studies, primary cells may facilitate adaptation of virus to *in vitro* conditions before transitioning to immortalized cell lines. For example, HRV is more efficiently grown in primary African green monkey kidney cells than immortalized lines [131,139], and multiple rounds of passaging in primary cells were critical to adapt virus for growth in MA-104 cells [15,139]. A potential link has been reported between RV infection and pancreatitis and subsequent autoimmune type-1 diabetes [140,141]; replication of HRV has also been demonstrated in primary monkey islet cells [142]. Human intestinal primary cells have also been recently utilized for replication of other enteric viruses such as HEV [143]. Moreover, HEV has also been cultivated successfully in extrahepatic primary cells such as hematopoietic cells, endometrial stromal cells and renal epithelium [144–146]. However, HuNoV has not been shown to replicate in primary cells [96].

#### Advantages and Disadvantages of Primary Cells

Primary cells offer advantages including maintenance of the physiological features and genetic makeup of the tissue of origin; they are also cost-effective in comparison to animal models. However, they also exhibit longer doubling times and limited growth potential compared to immortalized cells. They are difficult to maintain and may change with each passage, and cells taken from different sources can exhibit high levels of variation in responses to external stimuli [138] (Table 2).

#### 3.3. Intestinal Enteroids


nucleoside analog ribavirin against HRV, which varies by viral strain, supporting the use of these cultures towards personalized medicine approaches [149,150]. A variety of HuNoV strains have also been successfully cultivated in HIEs [20,31,151]. Studies exploring HuNoV pathogenesis in HIEs have revealed enterocytes to be a primary target for HuNoV replication, and indicated that cells derived from duodenum, jejunum, and ileum can be permissive to infection [31]. They have also identified viral strain-specific requirements for bile acids and histo-blood group antigens, regulated by FUT2 expression, for susceptibility of HIEs to HuNoV [16,31,152]. Importantly, genetic modifications of HIEs have been successfully performed using CRISPR-Cas9 approaches to explore FUT2 requirements and IFN regulation of HuNoV [32,33]; this capacity to genetically modify HIEs is likely to be increasingly applied to enteric viral cultivation studies (Figure 1). HIEs also permit efficient replication of HAstV, with HIEs derived from all intestinal segments supporting the growth of representative strains from all three clades [43,45,46]. These studies have shown that HAstV can infect multiple cell types such as goblet cells, mature enterocytes, and intestinal progenitor cells [43,45,46], and have revealed the importance of IFN-mediated antiviral responses against HAstV [45].

Because 3-dimensional enteroids have a closed structure, depending upon the pathogen being studied, it may be critical to explore whether luminal/apical entry requires derivation of a 2-dimensional monolayer, wherein the organoids are dissociated by enzymatic treatment and cells are seeded onto plates coated with matrigel [31,152,153], collagen mimicking the extracellular matrix [154], or polyethylene glycol [155]. Two-dimensional monolayers have the advantage of scaling up and can be used for high throughput screening studies [156]. Another option to provide simultaneous apical and basolateral access is to grow the 2-dimensional monolayer on a Transwell insert containing a porous membrane and coated with extracellular matrix-like proteins [156,157]. These Transwell inserts can also be used to co-culture immune and epithelial cells in the different chamber compartments, with the porous membrane permitting transport of secreted factors [158,159]. An alternate approach to provide apical access in a 3-dimensional enteroid system is to leverage a new technique of reverse enteroid development wherein manipulation of extracellular matrix proteins permits access to the apical surface [156]. These ‘api-cal-out’ enteroids can differentiate into various intestinal epithelial cell lineages, act as diffusion barriers and perform intestinal functions including nutrient absorption and mucus secretion [160], and have been recently applied to study the pathogenesis of pandemic viral strain SARS-CoV-2 [161,162], highlighting the potential for their application to other viral systems. Another consideration for these cultures is the absence of physical forces, such as fluid shear effect or peristaltic movement, characteristic of intestinal physiology. These forces regulate the behavior of intestinal cells [163] and their interaction with luminal contents [164]. Microfluidic devices for introducing shear fluid forces to generate a microenvironment similar to the gut, a model called “intestine-on-a-chip”, that employ monolayers derived from 3-dimensional organoids have thus been developed [165–167]. However, they require specialized technical expertise, microfabrication, and fluidic systems that are not universally

individuals can lead to distinct phenotypic responses [170]. HIE models generally lack components of the host microenvironment, as discussed above [171]. Finally, maintenance of HIEs is very expensive and time-consuming, requiring a substantial degree of expertise, compared to immortalized cell lines (Table 2).

#### 4. In Vivo Tools to Study Human Enteric RNA Viruses

While the study of viruses in vitro provides many key insights into virus–cell interactions, it is critical to complement these analyses with the study of infection in the context of an organism. Factors such as adaptive immune responses to and the influence of the microbiota on viral infection in the intestine can often be most readily achieved with the use of animal models [172,173] (Figure 2).



**Figure 2.** Animal models available for the study of HRV and HuNoV. Numerous animal models described to date support replication of HRV and HuNoV, including a variety of non-human primates as well as gnotobiotic pigs, zebrafish, and humanized mice. Created with BioRender.com (accessed on 21 January 2021).

##### 4.1. Non-Human Primates

Due to the genetic proximity of non-human primates (NHP), including vervet mon-

Enteric viruses isolated directly from NHPs have also been used extensively for study of viral pathogenesis. For example, SA11 is a simian rotavirus initially isolated from a vervet monkey [178] which has subsequently been used as a model for the study of RV both in vivo and in vitro [179,180]. Study of another simian rotavirus strain, rhesus RV, contributed to the formulation of RV vaccines [181,182]. Recovirus is an enteric calicivirus isolated from rhesus macaques [183] which causes diarrhea in infected animals and can be used to study pathogenesis of and immunity to caliciviruses such as HuNoV in vivo [184]. In the last several years, AstVs have been increasingly identified in NHP samples, though these have not yet been extensively characterized [185,186].

#### 4.2. Gnotobiotic Pigs

Gnotobiotic pigs, referring to pigs for whom the microbial status is well-defined and including germ-free status, have been a long-standing resource for the study of both the microbiota as well as enteric viruses due to their strong similarities to humans in pathophysiological responses [187]. Human microbiota samples can be efficiently transplanted into gnotobiotic piglets, resulting in microbial profiles similar to the donor samples [188].

Infection of gnotobiotic piglets with either HRV or HuNoV recapitulates human symptoms of acute viral gastroenteritis including diarrhea and fecal viral shedding, thus making these models extremely useful for immunological studies [25,189]. Gnotobiotic piglets have been extensively used for RV vaccine evaluation [190], and studies of HuNoV infection in gnotobiotic piglets demonstrate an association of HuNoV infection with intestinal epithelial cell apoptosis and barrier disruption, indicating a mechanism for HuNoV-induced diarrhea in humans [191]. Porcine enteric viruses, causing pathological symptoms of gastroenteritis such as diarrhea, vomiting, dehydration and even mortality in neonatal piglets, have been found which share similarity with HuNoV GII and HAstV strains [192–194].

#### 4.3. Mouse Models

One of the most widely used animal models in biomedical sciences is the laboratory mouse (*Mus musculus*). Mice have many features that make them ideal for research studies, including a relatively fast reproduction rate with large litter sizes, the existence of large and well-controlled housing facilities in the majority of research institutions, and an enormous breadth of genetically modified models permitting interrogation of the sufficiency or necessity of factors for any given phenotype [195]. In the field of enteric virus research, mice can potentially be infected directly with human viruses. BALB/c and outbred Swiss SW55 mouse pups can be orally inoculated with HRV strains, with infection causing development of symptoms and histopathological changes associated with gastroenteritis [196,197]. Similarly, BALB/c *Rag2<sup>-/-</sup>Il2rg<sup>-/-</sup>* mice engrafted with human CD34+ hematopoietic stem cells have been used for intraperitoneal infection with HuNoV GII.4, wherein viral structural and nonstructural proteins were found to be expressed in the spleen and liver [35]. Interestingly, in this infection model the genetic background of the mice may play a more important role than the engraftment of human cells [35]. These models have not yet been

revealing important regulatory aspects of the commensal microbiota, IFN signaling, and roles for nonstructural proteins in antagonism of host responses which are conserved with HuNoV [124,207–209]. A recent set of studies identified CD300LF as the host protein which serves as the MNoV receptor, and though human CD300LF does not mediate the same role for HuNoV, this finding raises intriguing questions about the potential existence of a proteinaceous receptor for HuNoV [124,210–212]. More recently, murine astrovirus (muAstV) was identified as a small animal model for HAsVs [213,214]. MuAstV has been shown to mediate viral interference against other enteric viruses in immunodeficient hosts [215], and its tropism for mucus-secreting goblet cells may also permit muAstV to regulate enteric bacterial pathogens [46,216]. Although mouse models of viral infection may not reflect all clinical outcomes of human enteric viruses, their importance in interrogating and defining key aspects of viral regulation in vivo cannot be denied.

#### 4.4. Zebrafish

Recently, zebrafish (*Danio rerio*) have gained prominence as in vitro and in vivo models for the study of viral pathogenesis. Zebrafish offer numerous advantages including being inexpensive, breeding rapidly, and being genetically tractable [217,218]. In addition, zebrafish are transparent during development, facilitating visualization of internal structures and viral infection. This model shares genetic and physiologic similarities to humans including innate and adaptive immunity, making them widely used vertebrate models of human diseases. Recently, zebrafish larvae have been used as a robust replication model for HuNoV. GI and GII viruses replicate to high titers, with virus detectable in both intestinal and hematopoietic tissues, consistent with the possible dual tropism for HuNoV [34]. In addition, an increase in expression of several immune genes was also observed [34]. The zebrafish model has also been utilized for high throughput anti-HuNoV drug screening [219,220]. Of interest, a transgenic zebrafish expressing green fluorescent protein under the influence of an IFN stimulated gene promoter has been developed, providing an in vivo tracking system for viral infections [221].

#### 4.5. Turkeys

Turkeys have proven to be useful small animal models for the study of AstV pathogenesis and immunity [222,223]. Turkey poults infected with turkey AstV (TAsV-2) show clinical symptoms including age-dependent diarrhea, and have been used to identify the viral capsid as a critical mediator of diarrhea and to test therapeutic antiviral strategies against AstVs [224,225].

## 5. Conclusions and Future Directions

Viral gastroenteritis remains a serious global health concern, as we currently lack vaccines for HuNoV and HAsV, and the efficacy of the RV vaccine may be limited in some countries due to unclear environmental factors. While the lack of suitable in vitro and in vivo model systems that could reliably recapitulate the primary aspects of viral

systems to reliably infect inexpensive and manipulatable small animal models such as mice and zebrafish with human enteric viruses would be tremendously useful and may rely on generation of transgenic animals expressing human viral receptors, similar to the introduction of the poliovirus receptor into mice [228]. Additional future advances in these model systems will almost certainly continue to yield an improved understanding of the pathogenesis of these enteric viruses, which will be key to improved vaccine and therapeutic approaches to combat the global burden of acute viral gastroenteritis.

**Author Contributions:** S.A., E.H. and M.T.B. wrote, read, and edited the manuscript. All authors have read and agreed to the published version of the manuscript.

**Funding:** This work was supported by NIH grants R01 AI127552, R01 AI139314, and R01 AI141478, a Children’s Discovery Institute of Washington University and St. Louis Children’s Hospital Interdisciplinary Research Initiative grant (MI-II-2019–790), the Pew Biomedical Scholars Program, and The Mathers Foundation.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Bishop, R.F.; Kirkwood, C. Enteric viruses. *Ref. Modul. Biomed. Sci.* **2014**. [CrossRef]
2. Chiejina, M.; Samant, H. Viral Diarrhea. *StatPearls [Internet]*. 2019. Available online: <https://www.ncbi.nlm.nih.gov/books/NBK470525/> (accessed on 21 January 2021).
3. Stuempfig, N.D.; Seroy, J. Viral gastroenteritis. *StatPearls [Internet]*. 2020. Available online: <https://www.ncbi.nlm.nih.gov/books/NBK518995/> (accessed on 21 January 2021).
4. Tate, J.E.; Burton, A.H.; Boschi-Pinto, C.; Steele, A.D.; Duque, J.; Parashar, U.D. 2008 estimate of worldwide rotavirus-associated mortality in children younger than 5 years before the introduction of universal rotavirus vaccination programmes: A systematic review and meta-analysis. *Lancet Infect. Dis.* **2012**, *12*, 136–141. [CrossRef]
5. McAtee, C.L.; Webman, R.; Gilman, R.H.; Mejia, C.; Bern, C.; Apaza, S.; Espetia, S.; Pajuelo, M.; Saito, M.; Challappa, R. Burden of norovirus and rotavirus in children after rotavirus vaccine introduction, Cochabamba, Bolivia. *Am. J. Trop. Med. Hyg.* **2016**, *94*, 212–217. [CrossRef] [PubMed]
6. Patel, M.M.; Widdowson, M.-A.; Glass, R.I.; Akazawa, K.; Vinjé, J.; Parashar, U.D. Systematic literature review of role of noroviruses in sporadic gastroenteritis. *Emerg. Infect. Dis.* **2008**, *14*, 1224. [CrossRef]
7. Bosch, A.; Pintó, R.M.; Guix, S. Human astroviruses. *Clin. Microbiol. Rev.* **2014**, *27*, 1048–1074. [CrossRef]
8. Ham, H.; Oh, S.; Jang, J.; Jo, S.; Choi, S.; Pak, S. Prevalence of human astrovirus in patients with acute gastroenteritis. *Ann. Lab. Med.* **2014**, *34*, 145. [CrossRef]
9. Fischer, T.K.; Rasmussen, L.D.; Fonager, J. Taking gastro-surveillance into the 21st century. *J. Clin. Virol.* **2019**, *117*, 43–48. [CrossRef]
10. Fernandez-Cassi, X.; Martínez-Puchol, S.; Silva-Sales, M.; Cornejo, T.; Bartolome, R.; Bofill-Mas, S.; Girones, R. Unveiling Viruses Associated with Gastroenteritis Using a Metagenomics Approach. *Viruses* **2020**, *12*, 1432. [CrossRef]
11. Jourdan, N.; Laffitte, J.C.; Forestier, F.; Servin, A.; Quero, A. Infection of cultured human intestinal cells by monkey RRV and human Wa rotavirus as a function of intestinal epithelial cell differentiation. *Res. Virol.* **1995**, *146*, 325–331. [CrossRef]
12. Casola, A.; Estes, M.K.; Crawford, S.E.; Ogra, P.L.; Ernst, P.B.; Garofalo, R.P.; Crowe, S.E. Rotavirus infection of cultured intestinal epithelial cells induces secretion of CXC and CC chemokines. *Gastroenterology* **1998**, *114*, 947–955. [CrossRef]
13. Ding, S.; Diep, J.; Feng, N.; Ren, L.; Li, B.; Ooi, Y.S.; Wang, X.; Brulois, K.F.; Yasukawa, L.L.; Li, X. STAG2 deficiency induces interferon responses via cGAS-STING pathway and restricts virus infection. *Nat. Commun.* **2018**, *9*, 1–8. [CrossRef] [PubMed]

20. Zou, W.Y.; Blutt, S.E.; Crawford, S.E.; Ettayebi, K.; Zeng, X.-L.; Saxena, K.; Ramani, S.; Karandikar, U.C.; Zachos, N.C.; Estes, M.K. Human intestinal enteroids: New models to study gastrointestinal virus infections. In *Organoids*; Springer: Berlin/Heidelberg, Germany, 2017; pp. 229–247.
21. Chang-Graham, A.L.; Danhof, H.A.; Engevik, M.A.; Tomaro-Duchesneau, C.; Karandikar, U.C.; Estes, M.K.; Versalovic, J.; Britton, R.A.; Hyser, J.M. Human intestinal enteroids with inducible neurogenin-3 expression as a novel model of gut hormone secretion. *Cell. Mol. Gastroenterol. Hepatol.* **2019**, *8*, 209–229. [[CrossRef](#)]
22. Wen, X.; Cao, D.; Jones, R.W.; Hoshino, Y.; Yuan, L. Tandem truncated rotavirus VP8\* subunit protein with T cell epitope as non-replicating parenteral vaccine is highly immunogenic. *Hum. Vaccines Immunother.* **2015**, *11*, 2483–2489. [[CrossRef](#)]
23. Wen, X.; Wen, K.; Cao, D.; Li, G.; Jones, R.W.; Li, J.; Szu, S.; Hoshino, Y.; Yuan, L. Inclusion of a universal tetanus toxoid CD4+ T cell epitope P2 significantly enhanced the immunogenicity of recombinant rotavirus ΔVP8\* subunit parenteral vaccines. *Vaccine* **2014**, *32*, 4420–4427. [[CrossRef](#)]
24. Torres, A.; Ji-Huang, L. Diarrheal response of gnotobiotic pigs after fetal infection and neonatal challenge with homologous and heterologous human rotavirus strains. *J. Virol.* **1986**, *60*, 1107–1112. [[CrossRef](#)] [[PubMed](#)]
25. Yuan, L.; Ward, L.A.; Rosen, B.I.; To, T.L.; Saif, L.J. Systematic and intestinal antibody-secreting cell responses and correlates of protective immunity to human rotavirus in a gnotobiotic pig model of disease. *J. Virol.* **1996**, *70*, 3075–3083. [[CrossRef](#)] [[PubMed](#)]
26. Bentes, G.A.; Guimarães, J.R.; Volotão, E.D.M.; Fialho, A.M.; Hooper, C.; Ganime, A.C.; Gardinali, N.R.; Lanzarini, N.M.; Da Silva, A.D.S.; Pitcovski, J. Cynomolgus monkeys (*Macaca fascicularis*) as an experimental infection model for human group A rotavirus. *Viruses* **2018**, *10*, 355. [[CrossRef](#)] [[PubMed](#)]
27. YAP, K.L.; ASRI, A. Experimental group A rotaviral infection in cynomolgus monkeys raised on formula diet. *Microbiol. Immunol.* **1990**, *34*, 153–162.
28. Majer, M.; Behrens, F.; Weinmann, E.; Mauler, R.; Maass, G.; Baumeister, H.; Luthardt, T. Diarrhea in newborn cynomolgus monkeys infected with human rotavirus. *Infection* **1978**, *6*, 71–72. [[CrossRef](#)]
29. Jones, M.K.; Grau, K.R.; Costantini, V.; Kolawole, A.O.; De Graaf, M.; Freiden, P.; Graves, C.L.; Koopmans, M.; Wallet, S.M.; Tibbetts, S.A. Human norovirus culture in B cells. *Nat. Protoc.* **2015**, *10*, 1939. [[CrossRef](#)]
30. Todd, K.V.; Tripp, R.A. Vero cells as a mammalian cell substrate for human norovirus. *Viruses* **2020**, *12*, 439. [[CrossRef](#)]
31. Ettayebi, K.; Crawford, S.E.; Murakami, K.; Broughman, J.R.; Karandikar, U.; Tenge, V.R.; Neill, F.H.; Blutt, S.E.; Zeng, X.-L.; Qu, L. Replication of human noroviruses in stem cell-derived human enteroids. *Science* **2016**, *353*, 1387–1393. [[CrossRef](#)]
32. Lin, S.-C.; Qu, L.; Ettayebi, K.; Crawford, S.E.; Blutt, S.E.; Robertson, M.J.; Zeng, X.-L.; Tenge, V.R.; Ayyar, B.V.; Karandikar, U.C. Human norovirus exhibits strain-specific sensitivity to host interferon pathways in human intestinal enteroids. *Proc. Natl. Acad. Sci. USA* **2020**, *117*, 23782–23793. [[CrossRef](#)]
33. Haga, K.; Ettayebi, K.; Tenge, V.R.; Karandikar, U.C.; Lewis, M.A.; Lin, S.-C.; Neill, F.H.; Ayyar, B.V.; Zeng, X.-L.; Larson, G. Genetic manipulation of human intestinal enteroids demonstrates the necessity of a functional fucosyltransferase 2 gene for secretor-dependent human norovirus infection. *MBio* **2020**, *11*. [[CrossRef](#)]
34. Van Dycke, J.; Ny, A.; Conceição-Neto, N.; Maes, J.; Hosmillo, M.; Cuvry, A.; Goodfellow, I.; Nogueira, T.C.; Verbeken, E.; Matthijnsens, J. A robust human norovirus replication model in zebrafish larvae. *PLoS Pathog.* **2019**, *15*, e1008009. [[CrossRef](#)] [[PubMed](#)]
35. Taube, S.; Kolawole, A.O.; Höhne, M.; Wilkinson, J.E.; Handley, S.A.; Perry, J.W.; Thackray, L.B.; Akkina, R.; Wobus, C.E. A mouse model for human norovirus. *MBio* **2013**, *4*. [[CrossRef](#)] [[PubMed](#)]
36. Cheetham, S.; Souza, M.; Meulia, T.; Grimes, S.; Han, M.G.; Saif, L.J. Pathogenesis of a genogroup II human norovirus in gnotobiotic pigs. *J. Virol.* **2006**, *80*, 10372–10381. [[CrossRef](#)] [[PubMed](#)]
37. Bui, T.; Kocher, J.; Li, Y.; Wen, K.; Li, G.; Liu, F.; Yang, X.; LeRoith, T.; Tan, M.; Xia, M. Median infectious dose of human norovirus GII.4 in gnotobiotic pigs is decreased by simvastatin treatment and increased by age. *J. Gen. Virol.* **2013**, *94 Pt 9*, 2005. [[CrossRef](#)]
38. Lei, S.; Ryu, J.; Wen, K.; Twitchell, E.; Bui, T.; Ramesh, A.; Weiss, M.; Li, G.; Samuel, H.; Clark-Deener, S. Increased and prolonged human norovirus infection in RAG2/IL2RG deficient gnotobiotic pigs with severe combined immunodeficiency. *Sci. Rep.* **2016**, *6*, 1–2. [[CrossRef](#)] [[PubMed](#)]

44. Méndez, E.; Muñoz-Yañez, C.; Sánchez-San Martín, C.; Aguirre-Crespo, G.; del Rocio Baños-Lara, M.; Gutierrez, M.; Espinosa, R.; Acevedo, Y.; Arias, C.F.; López, S. Characterization of human astrovirus cell entry. *J. Virol.* **2014**, *88*, 2452–2460. [[CrossRef](#)] [[PubMed](#)]
45. Triana, S.; Stanifer, M.L.; Shahraz, M.; Mukenhirn, M.; Kee, C.; Rueda, D.O.; Paulsen, M.; Benes, V.; Boulant, S.; Alexandrov, T. Single-cell transcriptomics reveals immune response of intestinal cell types to viral infection. *BioRxiv* **2020**. [[CrossRef](#)]
46. Ingle, H.; Hassan, E.; Gawron, J.; Mihi, B.; Li, Y.; Kennedy, E.A.; Kalugotla, G.; Makimaa, H.; Lee, S.; Desai, P. Murine astrovirus tropism for goblet cells and enterocytes facilitates an IFN- $\lambda$  response in vivo and in enteroid cultures. *Mucosal Immunol.* **2021**, *14*, 751–761. [[CrossRef](#)] [[PubMed](#)]
47. Owusu, I.A.; Mirabelli, C.; Kolawole, A.O. Intestinal Enteroid Culture for Human Astroviruses. *Bio Protoc.* **2020**, *10*, e3687. [[CrossRef](#)]
48. Flewett, T.; Bryden, A.; Davies, H. Virus particles in gastroenteritis. *Lancet* **1973**, *2*, 1497. [[CrossRef](#)]
49. Bishop, R. Discovery of rotavirus: Implications for child health. *J. Gastroenterol. Hepatol.* **2009**, *24*, S81–S85. [[CrossRef](#)]
50. Lundgren, O.; Svensson, L. Pathogenesis of rotavirus diarrhea. *Microbes Infect.* **2001**, *3*, 1145–1156. [[CrossRef](#)]
51. Morris, A.P.; Estes, M.K. Microbes and microbial toxins: Paradigms for microbial-mucosal interactions VIII. Pathological consequences of rotavirus infection and its enterotoxin. *Am. J. Physiol.* **2001**, *281*, G303–G310.
52. Chen, C.-C.; Huang, J.-L.; Chang, C.-J.; Kong, M.-S. Fecal calprotectin as a correlative marker in clinical severity of infectious diarrhea and usefulness in evaluating bacterial or viral pathogens in children. *J. Pediatric Gastroenterol. Nutr.* **2012**, *55*, 541–547. [[CrossRef](#)]
53. Wiegering, V.; Kaiser, J.; Tappe, D.; Weißbrich, B.; Morbach, H.; Girschick, H.J. Gastroenteritis in childhood: A retrospective study of 650 hospitalized pediatric patients. *Int. J. Infect. Dis.* **2011**, *15*, e401–e407. [[CrossRef](#)]
54. Gurwith, M.; Wenman, W.; Hinde, D.; Feltham, S.; Greenberg, H. A prospective study of rotavirus infection in infants and young children. *J. Infect. Dis.* **1981**, *144*, 218–224. [[CrossRef](#)] [[PubMed](#)]
55. Glass, R.I.; Kilgore, P.E.; Holman, R.C.; Jin, S.; Smith, J.C.; Woods, P.A.; Clarke, M.J.; Ho, M.S.; Gentsch, J.R. The epidemiology of rotavirus diarrhea in the United States: Surveillance and estimates of disease burden. *J. Infect. Dis.* **1996**, *174* (Suppl. 1), S5–S11. [[CrossRef](#)]
56. Desselberger, U.; Huppertz, H.-I. Immune responses to rotavirus infection and vaccination and associated correlates of protection. *J. Infect. Dis.* **2011**, *203*, 188–195. [[CrossRef](#)]
57. Luchs, A.; Timenetsky, M.d.C.S.T. Group A rotavirus gastroenteritis: Post-vaccine era, genotypes and zoonotic transmission. *Einstein* **2016**, *14*, 278–287. [[CrossRef](#)] [[PubMed](#)]
58. Lopez, S.; Arias, C. Early steps in rotavirus cell entry. *Reoviruses Entry Assem. Morphog.* **2006**, 39–66. [[CrossRef](#)]
59. Santiana, M.; Ghosh, S.; Ho, B.A.; Rajasekaran, V.; Du, W.-L.; Mutsafi, Y.; De Jesús-Díaz, D.A.; Sosnovtsev, S.V.; Levenson, E.A.; Parra, G.I. Vesicle-cloaked virus clusters are optimal units for inter-organismal viral transmission. *Cell Host Microbe* **2018**, *24*, 208–220.e8. [[CrossRef](#)] [[PubMed](#)]
60. Li, W.; Xiang, W.; Li, C.; Xu, J.; Zhou, D.; Shang, S. Molecular epidemiology of rotavirus A and adenovirus among children with acute diarrhea in Hangzhou, China. *Gut Pathog.* **2020**, *12*, 1–5. [[CrossRef](#)] [[PubMed](#)]
61. Gutiérrez-Aguirre, I.; Steyer, A.; Boben, J.; Gruden, K.; Poljšak-Prijatelj, M.; Ravnikar, M. Sensitive detection of multiple rotavirus genotypes with a single reverse transcription-real-time quantitative PCR assay. *J. Clin. Microbiol.* **2008**, *46*, 2547–2554. [[CrossRef](#)]
62. Vesikari, T.; Matson, D.O.; Dennehy, P.; Van Damme, P.; Santosham, M.; Rodriguez, Z.; Dallas, M.J.; Heyse, J.F.; Gouveia, M.G.; Black, S.B. Safety and efficacy of a pentavalent human-bovine (WC3) reassortant rotavirus vaccine. *N. Engl. J. Med.* **2006**, *354*, 23–33. [[CrossRef](#)]
63. Ruiz-Palacios, G.M.; Pérez-Schael, I.; Velázquez, F.R.; Abate, H.; Breuer, T.; Clemens, S.C.; Cheuvart, B.; Espinoza, F.; Gillard, P.; Innis, B.L. Safety and efficacy of an attenuated vaccine against severe rotavirus gastroenteritis. *N. Engl. J. Med.* **2006**, *354*, 11–22. [[CrossRef](#)]
64. World Health Organization. Rotavirus vaccines = Vaccins antirotavirus. *Wkly. Epidemiol. Rec. Relev. Épidémiol. Hebd.* **2007**, *82*, 285–295.



69. Harris, V.C.; Armah, G.; Fuentes, S.; Korpela, K.E.; Parashar, U.; Victor, J.C.; Tate, J.; de Weerth, C.; Giaquinto, C.; Wiersinga, W.J. Significant correlation between the infant gut microbiome and rotavirus vaccine response in rural Ghana. *J. Infect. Dis.* **2017**, *215*, 34–41. [[CrossRef](#)]
70. Dolin, R.; Blacklow, N.R.; DuPont, H.; Buscho, R.F.; Wyatt, R.G.; Kasel, J.A.; Hornick, R.; Chanock, R.M. Biological properties of Norwalk agent of acute infectious nonbacterial gastroenteritis. *Proc. Soc. Exp. Biol. Med.* **1972**, *140*, 578–583. [[CrossRef](#)]
71. Kapikian, A.Z.; Wyatt, R.G.; Dolin, R.; Thornhill, T.S.; Kalica, A.R.; Chanock, R.M. Visualization by immune electron microscopy of a 27-nm particle associated with acute infectious nonbacterial gastroenteritis. *J. Virol.* **1972**, *10*, 1075–1081. [[CrossRef](#)]
72. Atmar, R.L.; Opekun, A.R.; Gilger, M.A.; Estes, M.K.; Crawford, S.E.; Neill, F.H.; Graham, D.Y. Norwalk virus shedding after experimental human infection. *Emerg. Infect. Dis.* **2008**, *14*, 1553. [[CrossRef](#)]
73. Kirby, A.; Shi, J.; Montes, J.; Lichtenstein, M.; Moe, C. Disease course and viral shedding in experimental Norwalk virus and Snow Mountain virus infection. *J. Med. Virol.* **2014**, *86*, 2055–2064. [[CrossRef](#)]
74. Chhabra, P.; de Graaf, M.; Parra, G.I.; Chan, M.C.-W.; Green, K.; Martella, V.; Wang, Q.; White, P.A.; Katayama, K.; Vennema, H. Updated classification of norovirus genogroups and genotypes. *J. Gen. Virol.* **2019**, *100*, 1393. [[CrossRef](#)] [[PubMed](#)]
75. Zheng, D.-P.; Widdowson, M.-A.; Glass, R.I.; Vinjé, J. Molecular epidemiology of genogroup II-genotype 4 noroviruses in the United States between 1994 and 2006. *J. Clin. Microbiol.* **2010**, *48*, 168–177. [[CrossRef](#)] [[PubMed](#)]
76. Vinjé, J. Advances in laboratory methods for detection and typing of norovirus. *J. Clin. Microbiol.* **2015**, *53*, 373–381. [[CrossRef](#)] [[PubMed](#)]
77. Jones, M.K.; Watanabe, M.; Zhu, S.; Graves, C.L.; Keyes, L.R.; Grau, K.R.; Gonzalez-Hernandez, M.B.; Iovine, N.M.; Wobus, C.E.; Vinjé, J. Enteric bacteria promote human and mouse norovirus infection of B cells. *Science* **2014**, *346*, 755–759. [[CrossRef](#)] [[PubMed](#)]
78. Karandikar, U.C.; Crawford, S.E.; Ajami, N.J.; Murakami, K.; Kou, B.; Ettayebi, K.; Papanicolaou, G.A.; Jongwutiwes, U.; Perales, M.-A.; Shia, J. Detection of human norovirus in intestinal biopsies from immunocompromised transplant patients. *J. Gen. Virol.* **2016**, *97*, 2291. [[CrossRef](#)] [[PubMed](#)]
79. Green, K.Y.; Kaufman, S.S.; Nagata, B.M.; Chaimongkol, N.; Kim, D.Y.; Levenson, E.A.; Tin, C.M.; Yardley, A.B.; Johnson, J.A.; Barletta, A.B.F. Human norovirus targets enteroendocrine epithelial cells in the small intestine. *Nat. Commun.* **2020**, *11*, 1–10. [[CrossRef](#)]
80. Zhang, H.; Morrison, S.; Tang, Y.-W. Multiplex polymerase chain reaction tests for detection of pathogens associated with gastroenteritis. *Clin. Lab. Med.* **2015**, *35*, 461–486. [[CrossRef](#)]
81. Kocher, J.; Yuan, L. Norovirus vaccines and potential antinorovirus drugs: Recent advances and future perspectives. *Future Virol.* **2015**, *10*, 899–913. [[CrossRef](#)]
82. Appleton, H. Viruses and gastroenteritis in infants. *Lancet* **1975**, *1*, 1297. [[CrossRef](#)]
83. Madeley, C.; Cosgrove, B. 28 nm particles in faeces in infantile gastroenteritis. *Lancet* **1975**, *306*, 451–452. [[CrossRef](#)]
84. González, G.G.; Pujol, F.H.; Liprandi, F.; Deibis, L.; Ludert, J.E. Prevalence of enteric viruses in human immunodeficiency virus seropositive patients in Venezuela. *J. Med. Virol.* **1998**, *55*, 288–292. [[CrossRef](#)]
85. Wunderli, W.; Meerbach, A.; Guengoer, T.; Berger, C.; Greiner, O.; Caduff, R.; Trkola, A.; Bossart, W.; Gerlach, D.; Schibler, M. Astrovirus infection in hospitalized infants with severe combined immunodeficiency after allogeneic hematopoietic stem cell transplantation. *PLoS ONE* **2011**, *6*, e27483. [[CrossRef](#)]
86. Méndez-Toss, M.; Griffin, D.D.; Calva, J.; Contreras, J.F.; Puerto, F.I.; Mota, F.; Guiscafré, H.; Cedillo, R.; Muñoz, O.; Herrera, I. Prevalence and genetic diversity of human astroviruses in Mexican children with symptomatic and asymptomatic infections. *J. Clin. Microbiol.* **2004**, *42*, 151–157. [[CrossRef](#)] [[PubMed](#)]
87. Kurtz, J.; Lee, T.; Craig, J.; Reed, S.E. Astrovirus infection in volunteers. *J. Med. Virol.* **1979**, *3*, 221–230. [[CrossRef](#)] [[PubMed](#)]
88. Sebire, N.; Malone, M.; Shah, N.; Anderson, G.; Gaspar, H.; Cubitt, W. Pathology of astrovirus associated diarrhoea in a paediatric bone marrow transplant recipient. *J. Clin. Pathol.* **2004**, *57*, 1001–1003. [[CrossRef](#)] [[PubMed](#)]
89. Gray, E.; Angus, K.; Snodgrass, D. Ultrastructure of the small intestine in astrovirus-infected lambs. *J. Gen. Virol.* **1980**, *49*, 71–82. [[CrossRef](#)]

96. Duizer, E.; Schwab, K.J.; Neill, F.H.; Atmar, R.L.; Koopmans, M.P.; Estes, M.K. Laboratory efforts to cultivate noroviruses. *J. Gen. Virol.* **2004**, *85*, 79–87. [[CrossRef](#)] [[PubMed](#)]
97. Wobus, C.E.; Karst, S.M.; Thackray, L.B.; Chang, K.-O.; Sosnovtsev, S.V.; Belliot, G.; Krug, A.; Mackenzie, J.M.; Green, K.Y.; Virgin IV, H.W. Replication of Norovirus in cell culture reveals a tropism for dendritic cells and macrophages. *PLoS Biol.* **2004**, *2*, e432. [[CrossRef](#)]
98. Kitamoto, T.; Takai-Todaka, R.; Kato, A.; Kanamori, K.; Takagi, H.; Yoshida, K.; Katayama, K.; Nakanishi, A. Viral population changes during murine norovirus propagation in RAW 264.7 cells. *Front. Microbiol.* **2017**, *8*, 1091. [[CrossRef](#)] [[PubMed](#)]
99. Kearney, K.; Chen, D.; Taraporewala, Z.F.; Vende, P.; Hoshino, Y.; Tortorici, M.A.; Barro, M.; Patton, J.T. Cell-line-induced mutation of the rotavirus genome alters expression of an IRF3-interacting protein. *EMBO J.* **2004**, *23*, 4072–4081. [[CrossRef](#)]
100. Tsugawa, T.; Tatsumi, M.; Tsutsumi, H. Virulence-associated genome mutations of murine rotavirus identified by alternating serial passages in mice and cell cultures. *J. Virol.* **2014**, *88*, 5543–5558. [[CrossRef](#)] [[PubMed](#)]
101. Govorkova, E.; Murti, G.; Meignier, B.; De Taisne, C.; Webster, R. African green monkey kidney (Vero) cells provide an alternative host cell system for influenza A and B viruses. *J. Virol.* **1996**, *70*, 5519–5524. [[CrossRef](#)]
102. Pinto, M. Enterocyte-like differentiation and polarization of the human colon carcinoma cell line Caco-2 in culture. *Biol. Cell* **1983**, *47*, 323–330.
103. Rousset, M. The human colon carcinoma cell lines HT-29 and Caco-2: Two in vitro models for the study of intestinal differentiation. *Biochimie* **1986**, *68*, 1035–1040. [[CrossRef](#)]
104. Pintó, R.M.; Diez, J.M.; Bosch, A. Use of the colonic carcinoma cell line CaCo-2 for in vivo amplification and detection of enteric viruses. *J. Med. Virol.* **1994**, *44*, 310–315. [[CrossRef](#)]
105. Peterson, M.D.; Mooseker, M.S. Characterization of the enterocyte-like brush border cytoskeleton of the C2BBE clones of the human intestinal cell line, Caco-2. *J. Cell Sci.* **1992**, *102*, 581–600. [[CrossRef](#)] [[PubMed](#)]
106. Svensson, L.; Finlay, B.; Bass, D.; Von Bonsdorff, C.; Greenberg, H. Symmetric infection of rotavirus on polarized human intestinal epithelial (Caco-2) cells. *J. Virol.* **1991**, *65*, 4190–4197. [[CrossRef](#)]
107. Jourdan, N.; Maurice, M.; Delautier, D.; Quero, A.M.; Servin, A.L.; Trugnan, G. Rotavirus is released from the apical surface of cultured human intestinal cells through nonconventional vesicular transport that bypasses the Golgi apparatus. *J. Virol.* **1997**, *71*, 8268–8278. [[CrossRef](#)]
108. Janowski, A.B.; Wang, D. Infection and propagation of astrovirus VA1 in cell culture. *Curr. Protoc. Microbiol.* **2019**, *52*, e73. [[CrossRef](#)]
109. Harrington, P.R.; Vinjé, J.; Moe, C.L.; Baric, R.S. Norovirus capture with histo-blood group antigens reveals novel virus-ligand interactions. *J. Virol.* **2004**, *78*, 3035–3045. [[CrossRef](#)]
110. Harrington, P.R.; Lindesmith, L.; Yount, B.; Moe, C.L.; Baric, R.S. Binding of Norwalk virus-like particles to ABH histo-blood group antigens is blocked by antisera from infected human volunteers or experimentally vaccinated mice. *J. Virol.* **2002**, *76*, 12335–12343. [[CrossRef](#)]
111. Hutson, A.M.; Atmar, R.L.; Marcus, D.M.; Estes, M.K. Norwalk virus-like particle hemagglutination by binding to H histo-blood group antigens. *J. Virol.* **2003**, *77*, 405–415. [[CrossRef](#)]
112. Marionneau, S.; Ruvoën, N.; Le Moullac-Vaidye, B.; Clement, M.; Cailleau-Thomas, A.; Ruiz-Palacois, G.; Huang, P.; Jiang, X.; Le Pendu, J. Norwalk virus binds to histo-blood group antigens present on gastroduodenal epithelial cells of secretor individuals. *Gastroenterology* **2002**, *122*, 1967–1977. [[CrossRef](#)]
113. White, L.J.; Ball, J.M.; Hardy, M.E.; Tanaka, T.N.; Kitamoto, N.; Estes, M.K. Attachment and entry of recombinant Norwalk virus capsids to cultured human and animal cell lines. *J. Virol.* **1996**, *70*, 6589–6597. [[CrossRef](#)] [[PubMed](#)]
114. Oka, T.; Stoltzfus, G.T.; Zhu, C.; Jung, K.; Wang, Q.; Saif, L.J. Attempts to grow human noroviruses, a sapovirus, and a bovine norovirus in vitro. *PLoS ONE* **2018**, *13*, e0178157. [[CrossRef](#)]
115. Huet, G.; Kim, I.; De Bolos, C.; Lo-Guidice, J.; Moreau, O.; Hemon, B.; Richet, C.; Delannoy, P.; Real, F.; Degand, P. Characterization of mucins and proteoglycans synthesized by a mucin-secreting HT-29 cell subpopulation. *J. Cell Sci.* **1995**, *108*, 1275–1285. [[CrossRef](#)] [[PubMed](#)]

122. Herbst-Kralovetz, M.M.; Radtke, A.L.; Lay, M.K.; Hjelm, B.E.; Bolick, A.N.; Sarker, S.S.; Atmar, R.L.; Kingsley, D.H.; Arntzen, C.J.; Estes, M.K. Lack of norovirus replication and histo-blood group antigen expression in 3-dimensional intestinal epithelial cells. *Emerg. Infect. Dis.* **2013**, *19*, 431. [[CrossRef](#)]
123. Takanashi, S.; Saif, L.J.; Hughes, J.H.; Meulia, T.; Jung, K.; Scheuer, K.A.; Wang, Q. Failure of propagation of human norovirus in intestinal epithelial cells with microvilli grown in three-dimensional cultures. *Arch. Virol.* **2014**, *159*, 257–266. [[CrossRef](#)]
124. Wilen, C.B.; Lee, S.; Hsieh, L.L.; Orchard, R.C.; Desai, C.; Hykes, B.L.; McAllaster, M.R.; Balce, D.R.; Feehley, T.; Brestoff, J.R. Tropism for tuft cells determines immune promotion of norovirus pathogenesis. *Science* **2018**, *360*, 204–208. [[CrossRef](#)]
125. Desmyter, J.; Melnick, J.L.; Rawls, W.E. Defectiveness of interferon production and of rubella virus interference in a line of African green monkey kidney cells (Vero). *J. Virol.* **1968**, *2*, 955–961. [[CrossRef](#)]
126. Osada, N.; Kohara, A.; Yamaji, T.; Hirayama, N.; Kasai, F.; Sekizuka, T.; Kuroda, M.; Hanada, K. The genome landscape of the african green monkey kidney-derived vero cell line. *DNA Res.* **2014**, *21*, 673–683. [[CrossRef](#)]
127. Wyatt, R.G.; James, W.D.; Bohl, E.H.; Theil, K.W.; Saif, L.J.; Kalica, A.R.; Greenberg, H.B.; Kapikian, A.Z.; Chanock, R.M. Human rotavirus type 2: Cultivation in vitro. *Science* **1980**, *207*, 189–191. [[CrossRef](#)]
128. Barrett, P.N.; Mundt, W.; Kistner, O.; Howard, M.K. Vero cell platform in vaccine production: Moving towards cell culture-based viral vaccines. *Expert Rev. Vaccines* **2009**, *8*, 607–618. [[CrossRef](#)]
129. Wu, W.; Orr-Burks, N.; Karpilow, J.; Tripp, R.A. Development of improved vaccine cell lines against rotavirus. *Sci. Data* **2017**, *4*, 1–12. [[CrossRef](#)]
130. Guix, S.; Pérez-Bosque, A.; Miró, L.; Moretó, M.; Bosch, A.; Pintó, R.M. Type I interferon response is delayed in human astrovirus infections. *PLoS ONE* **2015**, *10*, e0123087. [[CrossRef](#)]
131. Hasegawa, A.; Matsuno, S.; Inouye, S.; Kono, R.; Tsurukubo, Y.; Mukoyama, A.; Saito, Y. Isolation of human rotaviruses in primary cultures of monkey kidney cells. *J. Clin. Microbiol.* **1982**, *16*, 387–390. [[CrossRef](#)]
132. Whitaker, A.; Hayward, C. The characterization of three monkey kidney cell lines. *Dev. Biol. Stand.* **1985**, *60*, 125–131.
133. Sato, K.; Inaba, Y.; Shinozaki, T.; Fujii, R.; Matumoto, M. Isolation of human rotavirus in cell cultures. *Arch. Virol.* **1981**, *69*, 155–160. [[CrossRef](#)]
134. Urasawa, T.; Urasawa, S.; Taniguchi, K. Sequential passages of human rotavirus in MA-104 cells. *Microbiol. Immunol.* **1981**, *25*, 1025–1035. [[CrossRef](#)] [[PubMed](#)]
135. Kanda, T.; Sasaki, R.; Masuzaki, R.; Matsumoto, N.; Ogawa, M.; Moriyama, M. Cell culture systems and drug targets for hepatitis A virus infection. *Viruses* **2020**, *12*, 533. [[CrossRef](#)] [[PubMed](#)]
136. Fu, R.M.; Decker, C.C.; Dao Thi, V.L. Cell culture models for hepatitis E virus. *Viruses* **2019**, *11*, 608. [[CrossRef](#)]
137. El-Mokhtar, M.A.; Sayed, I.M. Model systems for studying extrahepatic pathogenesis of hepatitis E virus. Current knowledge and future directions. *Rev. Med. Virol.* **2021**. [[CrossRef](#)] [[PubMed](#)]
138. Kaur, G.; Dufour, J.M. *Cell Lines: Valuable Tools or Useless Artifacts*; Taylor & Francis: Abingdon, UK, 2012.
139. Ward, R.L.; Knowlton, D.R.; Pierce, M.J. Efficiency of human rotavirus propagation in cell culture. *J. Clin. Microbiol.* **1984**, *19*, 748–753. [[CrossRef](#)]
140. De La Rubia, L.; Herrera, M.; Cebrero, M.; De Jong, J. Acute pancreatitis associated with rotavirus infection. *Pancreas* **1996**, *12*, 98–99. [[CrossRef](#)]
141. Honeyman, M.C.; Laine, D.; Zhan, Y.; Londrigan, S.; Kirkwood, C.; Harrison, L.C. Rotavirus infection induces transient pancreatic involution and hyperglycemia in weanling mice. *PLoS ONE* **2014**, *9*, e106560. [[CrossRef](#)]
142. Coulson, B.S.; Witterick, P.D.; Tan, Y.; Hewish, M.J.; Mountford, J.N.; Harrison, L.C.; Honeyman, M.C. Growth of rotaviruses in primary pancreatic cells. *J. Virol.* **2002**, *76*, 9537–9544. [[CrossRef](#)]
143. Marion, O.; Lhomme, S.; Nayrac, M.; Dubois, M.; Pucelle, M.; Requena, M.; Miguères, M.; Abravanel, F.; Peron, J.M.; Carrere, N. Hepatitis E virus replication in human intestinal cells. *Gut* **2020**, *69*, 901–910. [[CrossRef](#)]
144. El-Mokhtar, M.A.; Othman, E.R.; Khashbah, M.Y.; Ismael, A.; Ghaliony, M.A.; Seddik, M.I.; Sayed, I.M. Evidence of the extrahepatic replication of Hepatitis E virus in human endometrial stromal cells. *Pathogens* **2020**, *9*, 295. [[CrossRef](#)]
145. El-Mokhtar, M.A.; Seddik, M.I.; Osman, A.; Adel, S.; Abdel Aziz, E.M.; Mandour, S.A.; Mohammed, N.; Zarzour, M.A.; Abdel-

151. Ettayebi, K.; Tenge, V.R.; Cortes-Penfield, N.W.; Crawford, S.E.; Neill, F.H.; Zeng, X.-L.; Yu, X.; Ayyar, B.V.; Burrin, D.; Ramani, S. New insights and enhanced human norovirus cultivation in human intestinal enteroids. *Mosphere* **2021**, *6*, 1. [[CrossRef](#)]
152. Costantini, V.; Morantz, E.K.; Browne, H.; Ettayebi, K.; Zeng, X.-L.; Atmar, R.L.; Estes, M.K.; Vinjé, J. Human norovirus replication in human intestinal enteroids as model to evaluate virus inactivation. *Emerg. Infect. Dis.* **2018**, *24*, 1453. [[CrossRef](#)]
153. Li, L.; Xue, M.; Fu, F.; Yin, L.; Feng, L.; Liu, P. IFN-lambda 3 mediates antiviral protection against porcine epidemic diarrhea virus by inducing a distinct antiviral transcript profile in porcine intestinal epithelia. *Front. Immunol.* **2019**, *10*, 2394. [[CrossRef](#)]
154. Murakami, K.; Tenge, V.R.; Karandikar, U.C.; Lin, S.-C.; Ramani, S.; Ettayebi, K.; Crawford, S.E.; Zeng, X.-L.; Neill, F.H.; Ayyar, B.V. Bile acids and ceramide overcome the entry restriction for GII. 3 human norovirus replication in human intestinal enteroids. *Proc. Natl. Acad. Sci. USA* **2020**, *117*, 1700–1710. [[CrossRef](#)]
155. Wilson, R.L.; Swaminathan, G.; Ettayebi, K.; Bomidi, C.; Zeng, X.-L.; Blutt, S.E.; Estes, M.K.; Grande-Allen, K.J. Protein-functionalized poly (ethylene glycol) hydrogels as scaffolds for monolayer organoid culture. *Tissue Eng. Part C Methods* **2021**, *27*, 12–23. [[CrossRef](#)] [[PubMed](#)]
156. Atmar, R.L.; Ettayebi, K.; Ayyar, B.V.; Neill, F.H.; Braun, R.P.; Ramani, S.; Estes, M.K. Comparison of Microneutralization and Histo-Blood Group Antigen-Blocking Assays for Functional Norovirus Antibody Detection. *J. Infect. Dis.* **2020**, *221*, 739–743. [[CrossRef](#)] [[PubMed](#)]
157. Roodsant, T.; Navis, M.; Aknouch, I.; Renes, I.B.; Van Elburg, R.M.; Pajkrt, D.; Wolthers, K.C.; Schultsz, C.; Van Der Ark, K.C.; Sridhar, A. A human 2D primary organoid-derived epithelial monolayer model to study host-pathogen interaction in the small intestine. *Front. Cell. Infect. Microbiol.* **2020**, *10*, 272. [[CrossRef](#)]
158. Noel, G.; Baetz, N.W.; Staab, J.F.; Donowitz, M.; Kovbasnjuk, O.; Pasetti, M.F.; Zachos, N.C. A primary human macrophage-enteroid co-culture model to investigate mucosal gut physiology and host-pathogen interactions. *Sci. Rep.* **2017**, *7*, 1–13.
159. De Rudder, C.; Arroyo, M.C.; Lebeer, S.; Van de Wiele, T. Dual and Triple Epithelial Coculture Model Systems with Donor-Derived Microbiota and THP-1 Macrophages to Mimic Host-Microbe Interactions in the Human Sinonasal Cavities. *Mosphere* **2020**, *5*. [[CrossRef](#)] [[PubMed](#)]
160. Co, J.Y.; Margalef-Catala, M.; Li, X.; Mah, A.T.; Kuo, C.J.; Monack, D.M.; Amieva, M.R. Controlling epithelial polarity: A human enteroid model for host-pathogen interactions. *Cell Rep.* **2019**, *26*, 2509–2520.e4. [[CrossRef](#)]
161. Zang, R.; Castro, M.F.G.; McCune, B.T.; Zeng, Q.; Rothlauf, P.W.; Sonnek, N.M.; Liu, Z.; Brulois, K.F.; Wang, X.; Greenberg, H.B. TMPRSS2 and TMPRSS4 promote SARS-CoV-2 infection of human small intestinal enterocytes. *Sci. Immunol.* **2020**, *5*, eabc3582. [[CrossRef](#)]
162. Giobbe, G.G.; Bonfante, F.; Zambaiti, E.; Gagliano, O.; Jones, B.C.; Luni, C.; Laterza, C.; Perin, S.; Stuart, H.T.; Pagliari, M. SARS-CoV-2 infection and replication in human fetal and pediatric gastric organoids. *bioRxiv* **2020**. [[CrossRef](#)]
163. Gayer, C.P.; Basson, M.D. The effects of mechanical forces on intestinal physiology and pathology. *Cell. Signal.* **2009**, *21*, 1237–1244. [[CrossRef](#)] [[PubMed](#)]
164. Lentle, R.; Janssen, P. Physical characteristics of digesta and their influence on flow and mixing in the mammalian intestine: A review. *J. Comp. Physiol. B* **2008**, *178*, 673–690. [[CrossRef](#)]
165. Kim, H.J.; Huh, D.; Hamilton, G.; Ingber, D.E. Human gut-on-a-chip inhabited by microbial flora that experiences intestinal peristalsis-like motions and flow. *Lab A Chip* **2012**, *12*, 2165–2174. [[CrossRef](#)] [[PubMed](#)]
166. Shah, P.; Fritz, J.V.; Glaab, E.; Desai, M.S.; Greenhalgh, K.; Frachet, A.; Niegowska, M.; Estes, M.; Jäger, C.; Seguin-Devaux, C. A microfluidics-based in vitro model of the gastrointestinal human-microbe interface. *Nat. Commun.* **2016**, *7*, 1–15. [[CrossRef](#)]
167. Kasendra, M.; Tovaglieri, A.; Sontheimer-Phelps, A.; Jalili-Firoozinezhad, S.; Bein, A.; Chalkiadaki, A.; Scholl, W.; Zhang, C.; Rickner, H.; Richmond, C.A. Development of a primary human Small Intestine-on-a-Chip using biopsy-derived organoids. *Sci. Rep.* **2018**, *8*, 1–4. [[CrossRef](#)] [[PubMed](#)]
168. Ashammakhi, N.; Nasiri, R.; De Barros, N.R.; Tebon, P.; Thakor, J.; Goudie, M.; Shamloo, A.; Martin, M.G.; Khademhosseni, A. Gut-on-a-chip: Current progress and future opportunities. *Biomaterials* **2020**, 120196. [[CrossRef](#)]
169. Wilson, R.L.; Hewes, S.A.; Rajan, A.; Lin, S.-C.; Bomidi, C.; Iida, T.; Estes, M.K.; Maresso, A.W.; Grande-Allen, K.J. A Millifluidic Perfusion Cassette for Studying the Pathogenesis of Enteric Infections Using Ex-Vivo Organoids. *Ann. Biomed. Eng.* **2021**, *49*.

177. Wyatt, R.G.; Greenberg, H.B.; Dalgard, D.W.; Allen, W.P.; Sly, D.L.; Thornhill, T.S.; Chanock, R.M.; Kapikian, A.Z. Experimental infection of chimpanzees with the Norwalk agent of epidemic viral gastroenteritis. *J. Med. Virol.* **1978**, *2*, 89–96. [[CrossRef](#)]
178. Malherbe, H.; Harwin, R. The cytopathic effects of vervet monkey viruses. *S. Afr. Med. J.* **1963**, *37*, 407–411. [[PubMed](#)]
179. Yin, N.; Yang, F.-M.; Qiao, H.-T.; Zhou, Y.; Duan, S.-Q.; Lin, X.-C.; Wu, J.-Y.; Xie, Y.-P.; He, Z.-L.; Sun, M.-S. Neonatal rhesus monkeys as an animal model for rotavirus infection. *World J. Gastroenterol.* **2018**, *24*, 5109. [[CrossRef](#)] [[PubMed](#)]
180. Ludert, J.E.; Feng, N.; Yu, J.H.; Broome, R.L.; Hoshino, Y.; Greenberg, H.B. Genetic mapping indicates that VP4 is the rotavirus cell attachment protein in vitro and in vivo. *J. Virol.* **1996**, *70*, 487–493. [[CrossRef](#)]
181. McNeal, M.M.; Sestak, K.; Choi, A.H.-C.; Basu, M.; Cole, M.J.; Aye, P.P.; Bohm, R.P.; Ward, R.L. Development of a rotavirus-shedding model in rhesus macaques, using a homologous wild-type rotavirus of a new P genotype. *J. Virol.* **2005**, *79*, 944–954. [[CrossRef](#)]
182. Hoshino, Y.; Honma, S.; Jones, R.W.; Santos, N.; Nakagomi, O.; Nakagomi, T.; Kapikian, A.Z.; Thouless, M.E. A rotavirus strain isolated from pig-tailed macaque (*Macaca nemestrina*) with diarrhea bears a P6 [1]: G8 specificity. *Virology* **2006**, *345*, 1–12. [[CrossRef](#)]
183. Farkas, T.; Sestak, K.; Wei, C.; Jiang, X. Characterization of a rhesus monkey calicivirus representing a new genus of Caliciviridae. *J. Virol.* **2008**, *82*, 5408–5416. [[CrossRef](#)]
184. Sestak, K.; Feely, S.; Fey, B.; Dufour, J.; Hargitt, E.; Alvarez, X.; Pahar, B.; Gregoricus, N.; Vinjé, J.; Farkas, T. Experimental inoculation of juvenile rhesus macaques with primate enteric caliciviruses. *PLoS ONE* **2012**, *7*, e37973. [[CrossRef](#)]
185. Xin, Y.-Y.; Li, L.-L.; Ao, Y.-Y.; Xie, Z.-P.; Li, J.-S.; Duan, Z.-J.; Yu, J.-M.; Zhang, B. A novel astrovirus identified in wild rhesus monkey feces in China. *Arch. Virol.* **2019**, *164*, 2385–2388. [[CrossRef](#)] [[PubMed](#)]
186. Karlsson, E.A.; Small, C.T.; Freiden, P.; Feeroz, M.; Matsen IV, F.A.; San, S.; Hasan, M.K.; Wang, D.; Jones-Engel, L.; Schultz-Cherry, S. Non-human primates harbor diverse mammalian and avian astroviruses including those associated with human infections. *PLoS Pathog.* **2015**, *11*, e1005225. [[CrossRef](#)] [[PubMed](#)]
187. Meurens, F.; Summerfield, A.; Nauwynck, H.; Saif, L.; Gerdts, V. The pig: A model for human infectious diseases. *Trends Microbiol.* **2012**, *20*, 50–57. [[CrossRef](#)] [[PubMed](#)]
188. Vlasova, A.N.; Rajashekara, G.; Saif, L.J. Interactions between human microbiome, diet, enteric viruses and immune system: Novel insights from gnotobiotic pig research. *Drug Discov. Today Dis. Models* **2018**, *28*, 95–103. [[CrossRef](#)] [[PubMed](#)]
189. Saif, L.; Ward, L.; Yuan, L.; Rosen, B.; To, T. The gnotobiotic piglet as a model for studies of disease pathogenesis and immunity to human rotaviruses. *Viral Gastroenteritis* **1996**, *12*, 153–161.
190. Azevedo, M.P.; Vlasova, A.N.; Saif, L.J. Human rotavirus virus-like particle vaccines evaluated in a neonatal gnotobiotic pig model of human rotavirus disease. *Expert Rev. Vaccines* **2013**, *12*, 169–181. [[CrossRef](#)]
191. Troeger, H.; Loddenkemper, C.; Schneider, T.; Schreier, E.; Epple, H.-J.; Zeitz, M.; Fromm, M.; Schulzke, J.D. Structural and functional changes of the duodenum in human norovirus infection. *Gut* **2009**, *58*, 1070–1077. [[CrossRef](#)]
192. Wang, Q.-H.; Han, M.G.; Cheetham, S.; Souza, M.; Funk, J.A.; Saif, L.J. Porcine noroviruses related to human noroviruses. *Emerg. Infect. Dis.* **2005**, *11*, 1874. [[CrossRef](#)]
193. Sugieda, M.; Nakajima, S. Viruses detected in the caecum contents of healthy pigs representing a new genetic cluster in genogroup II of the genus ‘Norwalk-like viruses’. *Virus Res.* **2002**, *87*, 165–172. [[CrossRef](#)]
194. Indik, S.; Valíček, L.; Šmíd, B.; Dvořáková, H.; Rodák, L. Isolation and partial characterization of a novel porcine astrovirus. *Vet. Microbiol.* **2006**, *117*, 276–283. [[CrossRef](#)]
195. Perlman, R.L. Mouse models of human disease: An evolutionary perspective. *Evol. Med. Public Health* **2016**, *2016*, 170–176. [[PubMed](#)]
196. Gouvea, V.; Alencar, A.; Barth, O.; De Castro, L.; Fialho, A.; Araujo, H.; Majerowicz, S.; Pereira, H. Diarrhoea in mice infected with a human rotavirus. *J. Gen. Virol.* **1986**, *67*, 577–581. [[CrossRef](#)]
197. Ebina, T.; Tsukada, K.; Umezu, K.; Nose, M.; Tsuda, K.; Hatta, H.; Kim, M.; Yamamoto, T. Gastroenteritis in suckling mice caused by human rotavirus can be prevented with egg yolk immunoglobulin (IgY) and treated with a protein-bound polysaccharide preparation (PSK). *Microbiol. Immunol.* **1990**, *34*, 617–629. [[CrossRef](#)] [[PubMed](#)]

204. Zhang, B.; Chassaing, B.; Shi, Z.; Uchiyama, R.; Zhang, Z.; Denning, T.L.; Crawford, S.E.; Pruijssers, A.J.; Iskarpatyoti, J.A.; Estes, M.K. Prevention and cure of rotavirus infection via TLR5/NLRC4-mediated production of IL-22 and IL-18. *Science* **2014**, *346*, 861–865. [[CrossRef](#)]
205. Shi, Z.; Zou, J.; Zhang, Z.; Zhao, X.; Noriega, J.; Zhang, B.; Zhao, C.; Ingle, H.; Bittinger, K.; Mattei, L.M. Segmented filamentous bacteria prevent and cure rotavirus infection. *Cell* **2019**, *179*, 644–658.e13. [[CrossRef](#)] [[PubMed](#)]
206. Karst, S.M.; Wobus, C.E.; Lay, M.; Davidson, J.; Virgin, H.W. STAT1-dependent innate immunity to a Norwalk-like virus. *Science* **2003**, *299*, 1575–1578. [[CrossRef](#)] [[PubMed](#)]
207. Baldrige, M.T.; Nice, T.J.; McCune, B.T.; Yokoyama, C.C.; Kambal, A.; Wheadon, M.; Diamond, M.S.; Ivanova, Y.; Artyomov, M.; Virgin, H.W. Commensal microbes and interferon- $\lambda$  determine persistence of enteric murine norovirus infection. *Science* **2015**, *347*, 266–269. [[CrossRef](#)] [[PubMed](#)]
208. Orchard, R.C.; Wilen, C.B.; Doench, J.G.; Baldrige, M.T.; McCune, B.T.; Lee, Y.-C.J.; Lee, S.; Pruett-Miller, S.M.; Nelson, C.A.; Fremont, D.H. Discovery of a proteinaceous cellular receptor for a norovirus. *Science* **2016**, *353*, 933–936. [[CrossRef](#)]
209. Taube, S.; Perry, J.W.; McGreevy, E.; Yetming, K.; Perkins, C.; Henderson, K.; Wobus, C.E. Murine noroviruses bind glycolipid and glycoprotein attachment receptors in a strain-dependent manner. *J. Virol.* **2012**, *86*, 5584–5593. [[CrossRef](#)]
210. Lee, S.; Wilen, C.B.; Orvedahl, A.; McCune, B.T.; Kim, K.-W.; Orchard, R.C.; Peterson, S.T.; Nice, T.J.; Baldrige, M.T.; Virgin, H.W. Norovirus cell tropism is determined by combinatorial action of a viral non-structural protein and host cytokine. *Cell Host Microbe* **2017**, *22*, 449–459.e4. [[CrossRef](#)]
211. Grau, K.R.; Roth, A.N.; Zhu, S.; Hernandez, A.; Colliou, N.; DiVita, B.B.; Philip, D.T.; Riffe, C.; Giasson, B.; Wallet, S.M. The major targets of acute norovirus infection are immune cells in the gut-associated lymphoid tissue. *Nat. Microbiol.* **2017**, *2*, 1586–1591. [[CrossRef](#)]
212. Graziano, V.R.; Walker, F.C.; Kennedy, E.A.; Wei, J.; Ettayebi, K.; Strine, M.S.; Filler, R.B.; Hassan, E.; Hsieh, L.L.; Kim, A.S. CD300lf is the primary physiologic receptor of murine norovirus but not human norovirus. *PLoS Pathog.* **2020**, *16*, e1008242. [[CrossRef](#)] [[PubMed](#)]
213. Yokoyama, C.C.; Loh, J.; Zhao, G.; Stappenbeck, T.S.; Wang, D.; Huang, H.V.; Virgin, H.W.; Thackray, L.B. Adaptive immunity restricts replication of novel murine astroviruses. *J. Virol.* **2012**, *86*, 12262–12270. [[CrossRef](#)]
214. Cortez, V.; Sharp, B.; Yao, J.; Livingston, B.; Vogel, P.; Schultz-Cherry, S. Characterizing a murine model for astrovirus using viral isolates from persistently infected immunocompromised mice. *J. Virol.* **2019**, *93*. [[CrossRef](#)]
215. Ingle, H.; Lee, S.; Ai, T.; Orvedahl, A.; Rodgers, R.; Zhao, G.; Sullender, M.; Peterson, S.T.; Locke, M.; Liu, T.-C. Viral complementation of immunodeficiency confers protection against enteric pathogens via interferon- $\lambda$ . *Nat. Microbiol.* **2019**, *4*, 1120–1128. [[CrossRef](#)] [[PubMed](#)]
216. Cortez, V.; Boyd, D.F.; Crawford, J.C.; Sharp, B.; Livingston, B.; Rowe, H.M.; Davis, A.; Alsallaq, R.; Robinson, C.G.; Vogel, P. Astrovirus infects actively secreting goblet cells and alters the gut mucus barrier. *Nat. Commun.* **2020**, *11*, 1–9. [[CrossRef](#)] [[PubMed](#)]
217. Zou, P.F.; Nie, P. Zebrafish as a model for the study of host-virus interactions. In *Innate Antiviral Immunity*; Springer: Berlin/Heidelberg, Germany, 2017; pp. 57–78.
218. Sullivan, C.; Soos, B.-L.; Millard, P.J.; Kim, C.H.; King, B.L. Modeling Virus-Induced Inflammation in Zebrafish: A Balance Between Infection Control and Excessive Inflammation. *Front. Immunol.* **2021**, *12*, 1635. [[CrossRef](#)]
219. Tan, M.T.H.; Li, Y.; Eshaghi Gorji, M.; Gong, Z.; Li, D. Fucoidan But Not 2'-Fucosyllactose Inhibits Human Norovirus Replication in Zebrafish Larvae. *Viruses* **2021**, *13*, 461. [[CrossRef](#)]
220. Van Dycke, J.; Cuvry, A.; Knickmann, J.; Ny, A.; Rakers, S.; Taube, S.; de Witte, P.; Neyts, J.; Rocha-Pereira, J. Infection of zebrafish larvae with human norovirus and evaluation of the in vivo efficacy of small-molecule inhibitors. *Nat. Protoc.* **2021**, *16*, 1830–1849. [[CrossRef](#)]
221. Balla, K.M.; Rice, M.C.; Gagnon, J.A.; Elde, N.C. Linking virus discovery to immune responses visualized during zebrafish infections. *Curr. Biol.* **2020**, *30*, 2092–2103.e5. [[CrossRef](#)]
222. Schultz-Cherry, S.; Kapczynski, D.R.; Simmons, V.M.; Koci, M.D.; Brown, C.; Barnes, H.I. Identifying agent (s) associated with