

# Explicit solution to an optimal switching problem in the two-regime case\*

Vathana LY VATH

Laboratoire de Probabilités et

Modèles Aléatoires

CNRS, UMR 7599

Université Paris 7

e-mail: lyvath@math.jussieu.fr

Huyên PHAM

Laboratoire de Probabilités et

Modèles Aléatoires

CNRS, UMR 7599

Université Paris 7

e-mail: pham@math.jussieu.fr  
and CREST

August 2005

This version : June 2006

## Abstract

This paper considers the problem of determining the optimal sequence of stopping times for a diffusion process subject to regime switching decisions. This is motivated in the economics literature, by the investment problem under uncertainty for a multi-activity firm involving opening and closing decisions. We use a viscosity solutions approach combined with the smooth-fit property, and explicitly solve the problem in the two regime case when the state process is of geometric Brownian nature. The results of our analysis take several qualitatively different forms, depending on model parameter values.

**Key words :** Optimal switching, system of variational inequalities, viscosity solutions, smooth-fit principle.

**MSC Classification (2000) :** 60G40, 49L25, 62L15.

---

\*We would like to thank Xin Guo for discussions. We are also grateful to the anonymous referees for constructive suggestions and comments that helped to improve the paper.

# 1 Introduction

The theory of optimal stopping and its generalization, thoroughly studied in the seventies, has received a renewed interest with a variety of applications in economics and finance. These applications range from asset pricing (American options, swing options) to firm investment and real options. We refer to [4] for a classical and well documented reference on the subject.

In this paper, we consider the optimal switching problem for a one dimensional stochastic process  $X$ . The diffusion process  $X$  may take a finite number of regimes that are switched at stopping time decisions. For example in the firm's investment problem under uncertainty, a company (oil tanker, electricity station ....) manages several production activities operating in different modes or regimes representing a number of different economic outlooks (e.g. state of economic growth, open or closed production activity, ...). The process  $X$  is the price of input or output goods of the firm and its dynamics may differ according to the regimes. The firm's project yields a running payoff that depends on the commodity price  $X$  and on the regime choice. The transition from one regime to another one is realized sequentially at time decisions and incurs certain fixed costs. The problem is to find the switching strategy that maximizes the expected value of profits resulting from the project.

Optimal switching problems were studied by several authors, see [1] or [10]. These control problems lead via the dynamic programming principle to a system of variational inequalities. Applications to option pricing, real options and investment under uncertainty were considered by [2], [5] and [7]. In this last paper, the drift and volatility of the state process depend on an uncontrolled finite-state Markov chain, and the author provides an explicit solution to the optimal stopping problem with applications to Russian options. In [2], an explicit solution is found for a resource extraction problem with two regimes (open or closed field), a linear profit function and a price process following a geometric Brownian motion. In [5], a similar model is solved with a general profit function in one regime and equal to zero in the other regime. In both models [2], [5], there is no switching in the diffusion process : changes of regimes only affect the payoff functions. Their method of resolution is to construct a solution to the dynamic programming system by guessing a priori the form of the strategy, and then validate a posteriori the optimality of their candidate by a verification argument. Our model combines regime switchings both on the diffusion process and on the general profit functions. We use a viscosity solutions approach for determining the solution to the system of variational inequalities. In particular, we derive directly the smooth-fit property of the value functions and the structure of the switching regions. Explicit solutions are provided in the following cases :  $\star$  the drift and volatility terms of the diffusion take two different regime values, and the profit functions are identical of power type,  $\star$  there is no switching on the diffusion process, and the two different profit functions satisfy a general condition, including typically power functions. We also consider the cases for which both switching costs are positive, and for which one of the two is negative. This last case is interesting in applications where a firm chooses between an open or closed activity, and may regain a fraction of its opening costs when

it decides to close. The results of our analysis take several qualitatively different forms, depending on model parameter values, essentially the payoff functions and the switching costs.

The paper is organized as follows. We formulate in Section 2 the optimal switching problem. In Section 3, we state the system of variational inequalities satisfied by the value functions in the viscosity sense. The smooth-fit property for this problem, proved in [9], plays an important role in our subsequent analysis. We also state some useful properties on the switching regions. In Section 4, we explicitly solve the problem in the two-regimes case when the state process is of geometric Brownian nature.

## 2 Formulation of the optimal switching problem

We consider a stochastic system that can operate in  $d$  modes or regimes. The regimes can be switched at a sequence of stopping times decided by the operator (individual, firm, ...). The indicator of the regimes is modeled by a cadlag process  $I_t$  valued in  $\mathbb{I}_d = \{1, \dots, d\}$ . The stochastic system  $X$  (commodity price, salary, ...) is valued in  $\mathbb{R}_+^* = (0, \infty)$  and satisfies the s.d.e.

$$dX_t = b_{I_t} X_t dt + \sigma_{I_t} X_t dW_t, \quad (2.1)$$

where  $W$  is a standard Brownian motion on a filtered probability space  $(\Omega, \mathcal{F}, \mathbb{F} = (\mathcal{F}_t)_{t \geq 0}, P)$  satisfying the usual conditions.  $b_i \in \mathbb{R}$ , and  $\sigma_i > 0$  are the drift and volatility of the system  $X$  once in regime  $I_t = i$  at time  $t$ .

A strategy decision for the operator is an impulse control  $\alpha$  consisting of a double sequence  $\tau_1, \dots, \tau_n, \dots, \kappa_1, \dots, \kappa_n, \dots$ ,  $n \in \mathbb{N}^* = \mathbb{N} \setminus \{0\}$ , where  $\tau_n$  are stopping times,  $\tau_n < \tau_{n+1}$  and  $\tau_n \rightarrow \infty$  a.s., representing the switching regimes time decisions, and  $\kappa_n$  are  $\mathcal{F}_{\tau_n}$ -measurable valued in  $\mathbb{I}_d$ , and representing the new value of the regime at time  $t = \tau_n$ . We denote by  $\mathcal{A}$  the set of all such impulse controls. Now, for any initial condition  $(x, i) \in (0, \infty) \times \mathbb{I}_d$ , and any control  $\alpha = (\tau_n, \kappa_n)_{n \geq 1} \in \mathcal{A}$ , there exists a unique strong solution valued in  $(0, \infty) \times \mathbb{I}_d$  to the controlled stochastic system :

$$X_0 = x, \quad I_{0-} = i, \quad (2.2)$$

$$dX_t = b_{\kappa_n} X_t dt + \sigma_{\kappa_n} X_t dW_t, \quad I_t = \kappa_n, \quad \tau_n \leq t < \tau_{n+1}, \quad n \geq 0. \quad (2.3)$$

Here, we set  $\tau_0 = 0$  and  $\kappa_0 = i$ . We denote by  $(X^{x,i}, I^i)$  this solution (as usual, we omit the dependence in  $\alpha$  for notational simplicity). We notice that  $X^{x,i}$  is a continuous process and  $I^i$  is a cadlag process, possibly with a jump at time 0 if  $\tau_1 = 0$  and so  $I_0 = \kappa_1$ .

We are given a running profit function  $f : \mathbb{R}_+ \times \mathbb{I}_d \rightarrow \mathbb{R}$  and we set  $f_i(\cdot) = f(\cdot, i)$  for  $i \in \mathbb{I}_d$ . We assume that for each  $i \in \mathbb{I}_d$ , the function  $f_i$  is nonnegative and is Hölder continuous on  $\mathbb{R}_+$  : there exists  $\gamma_i \in (0, 1]$  s.t.

$$|f_i(x) - f_i(\hat{x})| \leq C|x - \hat{x}|^{\gamma_i}, \quad \forall x, \hat{x} \in \mathbb{R}_+, \quad (2.4)$$

for some positive constant  $C$ . Without loss of generality (see Remark 2.1), we may assume that  $f_i(0) = 0$ . We also assume that for all  $i \in \mathbb{I}_d$ , the conjugate of  $f_i$  is finite on  $(0, \infty)$  :

$$\tilde{f}_i(y) := \sup_{x \geq 0} [f_i(x) - xy] < \infty, \quad \forall y > 0. \quad (2.5)$$

The cost for switching from regime  $i$  to  $j$  is a constant equal to  $g_{ij}$ , with the convention  $g_{ii} = 0$ , and we assume the triangular condition :

$$g_{ik} < g_{ij} + g_{jk}, \quad j \neq i, k. \quad (2.6)$$

This last condition means that it is less expensive to switch directly in one step from regime  $i$  to  $k$  than in two steps via an intermediate regime  $j$ . Notice that a switching cost  $g_{ij}$  may be negative, and condition (2.6) for  $i = k$  prevents arbitrage by switching back and forth, i.e.

$$g_{ij} + g_{ji} > 0, \quad i \neq j \in \mathbb{I}_d. \quad (2.7)$$

The expected total profit of running the system when initial state is  $(x, i)$  and using the impulse control  $\alpha = (\tau_n, \kappa_n)_{n \geq 1} \in \mathcal{A}$  is

$$J_i(x, \alpha) = E \left[ \int_0^\infty e^{-rt} f(X_t^{x,i}, I_t^i) dt - \sum_{n=1}^\infty e^{-r\tau_n} g_{\kappa_{n-1}, \kappa_n} \right].$$

Here  $r > 0$  is a positive discount factor, and we use the convention that  $e^{-r\tau_n(\omega)} = 0$  when  $\tau_n(\omega) = \infty$ . We also make the standing assumption :

$$r > b := \max_{i \in \mathbb{I}_d} b_i. \quad (2.8)$$

The objective is to maximize this expected total profit over all strategies  $\alpha$ . Accordingly, we define the value functions

$$v_i(x) = \sup_{\alpha \in \mathcal{A}} J_i(x, \alpha), \quad x \in \mathbb{R}_+^*, \quad i \in \mathbb{I}_d. \quad (2.9)$$

We shall see in the next section that under (2.5) and (2.8), the expectation defining  $J_i(x)$  is well-defined and the value function  $v_i$  is finite.

**Remark 2.1** The initial values  $f_i(0)$  of the running profit functions received by the firm manager (the controller) before any decision are considered as included into the switching costs when changing of regime. This means that w.l.o.g. we may assume that  $f_i(0) = 0$ . Indeed, for any profit function  $f_i$ , and by setting  $\tilde{f}_i = f_i - f_i(0)$ , we have for all  $x > 0$ ,  $\alpha \in \mathcal{A}$ ,

$$\begin{aligned} J_i(x, \alpha) &= E \left[ \sum_{n=1}^\infty \int_{\tau_{n-1}}^{\tau_n} e^{-rt} f(X_t^{x,i}, \kappa_{n-1}) dt - \sum_{n=1}^\infty e^{-r\tau_n} g_{\kappa_{n-1}, \kappa_n} \right] \\ &= E \left[ \sum_{n=1}^\infty \int_{\tau_{n-1}}^{\tau_n} e^{-rt} \left( \tilde{f}(X_t^{x,i}, \kappa_{n-1}) + f_{\kappa_{n-1}}(0) \right) dt - \sum_{n=1}^\infty e^{-r\tau_n} g_{\kappa_{n-1}, \kappa_n} \right] \\ &= E \left[ \sum_{n=1}^\infty \int_{\tau_{n-1}}^{\tau_n} e^{-rt} \tilde{f}(X_t^{x,i}, \kappa_{n-1}) dt + \frac{f_{\kappa_0}(0)}{r} \right. \\ &\quad \left. - \sum_{n=1}^\infty e^{-r\tau_n} \left( g_{\kappa_{n-1}, \kappa_n} + \frac{f_{\kappa_n}(0) - f_{\kappa_{n-1}}(0)}{r} \right) \right] \\ &= \frac{f_i(0)}{r} + E \left[ \int_0^\infty e^{-rt} \tilde{f}(X_t^{x,i}, I_t^i) dt - \sum_{n=1}^\infty e^{-r\tau_n} \tilde{g}_{\kappa_{n-1}, \kappa_n} \right], \end{aligned}$$

with modified switching costs that take into account the possibly different initial values of the profit functions :

$$\tilde{g}_{ij} = g_{ij} + \frac{f_j(0) - f_i(0)}{r}.$$

### 3 System of variational inequalities, switching regions and viscosity solutions

We first state the linear growth property and the boundary condition on the value functions.

**Lemma 3.1** *We have for all  $i \in \mathbb{I}_d$  :*

$$\max_{j \in \mathbb{I}_d} [-g_{ij}] \leq v_i(x) \leq \frac{xy}{r-b} + \max_{j \in \mathbb{I}_d} \frac{\tilde{f}_j(y)}{r} + \max_{j \in \mathbb{I}_d} [-g_{ij}], \quad \forall x > 0, \forall y > 0. \quad (3.1)$$

In particular, we have  $v_i(0^+) = \max_{j \in \mathbb{I}_d} [-g_{ij}]$ .

**Proof.** By considering the particular strategy  $\tilde{\alpha} = (\tilde{\tau}_n, \tilde{\kappa}_n)$  of immediate switching from the initial state  $(x, i)$  to state  $(x, j)$ ,  $j \in \mathbb{I}_d$  (eventually equal to  $i$ ), at cost  $g_{ij}$  and then doing nothing, i.e.  $\tilde{\tau}_1 = 0$ ,  $\tilde{\kappa}_1 = j$ ,  $\tilde{\tau}_n = \infty$ ,  $\tilde{\kappa}_n = j$  for all  $n \geq 2$ , we have

$$J_i(x, \tilde{\alpha}) = E \left[ \int_0^\infty e^{-rt} f_j(\tilde{X}_t^{x,j}) dt - g_{ij} \right],$$

where  $\tilde{X}^{x,j}$  denotes the geometric brownian in regime  $j$  starting from  $x$  at time 0. Since  $f_j$  is nonnegative, and by the arbitrariness of  $j$ , we get the lower bound in (3.1).

Given an initial state  $(X_0, I_{0-}) = (x, i)$  and an arbitrary impulse control  $\alpha = (\tau_n, \kappa_n)$ , we get from the dynamics (2.2)-(2.3), the following explicit expression of  $X^{x,i}$  :

$$\begin{aligned} X_t^{x,i} &= x Y_t(i) \\ &:= x \left( \prod_{l=0}^{n-1} e^{b_{\kappa_l}(\tau_{l+1} - \tau_l)} Z_{\tau_l, \tau_{l+1}}^{\kappa_l} \right) e^{b_{\kappa_n}(t - \tau_n)} Z_{\tau_n, t}^{\kappa_n}, \quad \tau_n \leq t < \tau_{n+1}, \quad n \in \mathbb{N}, \end{aligned} \quad (3.2)$$

where

$$Z_{s,t}^j = \exp \left( \sigma_j (W_t - W_s) - \frac{\sigma_j^2}{2} (t - s) \right), \quad 0 \leq s \leq t, \quad j \in \mathbb{I}_d. \quad (3.3)$$

Here, we used the convention that  $\tau_0 = 0$ ,  $\kappa_0 = i$ , and the product term from  $l$  to  $n-1$  in (3.2) is equal to 1 when  $n = 1$ . We then deduce the inequality  $X_t^{x,i} \leq x e^{bt} M_t$ , for all  $t$ , where

$$M_t = \left( \prod_{l=0}^{n-1} Z_{\tau_l, \tau_{l+1}}^{\kappa_l} \right) Z_{\tau_n, t}^{\kappa_n}, \quad \tau_n \leq t < \tau_{n+1}, \quad n \in \mathbb{N}. \quad (3.4)$$

Now, we notice that  $(M_t)$  is a martingale obtained by continuously patching the martingales  $(Z_{\tau_{n-1}, t}^{\kappa_{n-1}})$  and  $(Z_{\tau_n, t}^{\kappa_n})$  at the stopping times  $\tau_n$ ,  $n \geq 1$ . In particular, we have  $E[M_t] = M_0 = 1$  for all  $t$ .

We set  $\tilde{f}(y) = \max_{j \in \mathbb{I}_d} \tilde{f}_j(y)$ ,  $y > 0$ , and we notice by definition of  $\tilde{f}_i$  in (2.5) that  $f(X_t^{x,i}, I_t^i) \leq yX_t^{x,i} + \tilde{f}(y)$  for all  $t, y$ . Moreover, we show by induction on  $N$  that for all  $N \geq 1$ ,  $\tau_1 \leq \dots \leq \tau_N$ ,  $\kappa_0 = i$ ,  $\kappa_n \in \mathbb{I}_d$ ,  $n = 1, \dots, N$  :

$$-\sum_{n=1}^N e^{-r\tau_n} g_{\kappa_{n-1}, \kappa_n} \leq \max_{j \in \mathbb{I}_d} [-g_{ij}], \quad a.s.$$

Indeed, the above assertion is obviously true for  $N = 1$ . Suppose now it holds true at step  $N$ . Then, at step  $N + 1$ , we distinguish two cases : If  $g_{\kappa_N, \kappa_{N+1}} \geq 0$ , then we have  $-\sum_{n=1}^{N+1} e^{-r\tau_n} g_{\kappa_{n-1}, \kappa_n} \leq -\sum_{n=1}^N e^{-r\tau_n} g_{\kappa_{n-1}, \kappa_n}$  and we conclude by the induction hypothesis at step  $N$ . If  $g_{\kappa_N, \kappa_{N+1}} < 0$ , then by (2.6), and since  $\tau_N \leq \tau_{N+1}$ , we have  $-e^{-r\tau_N} g_{\kappa_{N-1}, \kappa_N} - e^{-r\tau_{N+1}} g_{\kappa_N, \kappa_{N+1}} \leq e^{-r\tau_N} g_{\kappa_{N-1}, \kappa_{N+1}}$ , and so  $-\sum_{n=1}^{N+1} e^{-r\tau_n} g_{\kappa_{n-1}, \kappa_n} \leq -\sum_{n=1}^N e^{-r\tau_n} g_{\tilde{\kappa}_{n-1}, \tilde{\kappa}_n}$ , with  $\tilde{\kappa}_n = \kappa_n$  for  $n = 1, \dots, N - 1$ ,  $\tilde{\kappa}_N = \kappa_{N+1}$ . We then conclude by the induction hypothesis at step  $N$ .

It follows that

$$\begin{aligned} J_i(x, \alpha) &\leq E \left[ \int_0^\infty e^{-rt} \left( yx e^{bt} M_t + \tilde{f}(y) \right) dt + \max_{j \in \mathbb{I}_d} [-g_{ij}] \right] \\ &= \int_0^\infty e^{-(r-b)t} yx E[M_t] dt + \int_0^\infty e^{-rt} \tilde{f}(y) dt + \max_{j \in \mathbb{I}_d} [-g_{ij}] \\ &= \frac{xy}{r-b} + \frac{\tilde{f}(y)}{r} + \max_{j \in \mathbb{I}_d} [-g_{ij}]. \end{aligned}$$

From the arbitrariness of  $\alpha$ , this shows the upper bound for  $v_i$ .

By sending  $x$  to zero and then  $y$  to infinity into the r.h.s. of (3.1), and recalling that  $\tilde{f}_i(\infty) = f_i(0) = 0$  for  $i \in \mathbb{I}_d$ , we conclude that  $v_i$  goes to  $\max_{j \in \mathbb{I}_d} [-g_{ij}]$  when  $x$  tends to zero.  $\square$

We next show the Hölder continuity of the value functions.

**Lemma 3.2** *For all  $i \in \mathbb{I}_d$ ,  $v_i$  is Hölder continuous on  $(0, \infty)$  :*

$$|v_i(x) - v_i(\hat{x})| \leq C|x - \hat{x}|^\gamma, \quad \forall x, \hat{x} \in (0, \infty), \quad \text{with } |x - \hat{x}| \leq 1,$$

for some positive constant  $C$ , and where  $\gamma = \min_{i \in \mathbb{I}_d} \gamma_i$  of condition (2.4).

**Proof.** By definition (2.9) of  $v_i$  and under condition (2.4), we have for all  $x, \hat{x} \in (0, \infty)$ , with  $|x - \hat{x}| \leq 1$  :

$$\begin{aligned} |v_i(x) - v_i(\hat{x})| &\leq \sup_{\alpha \in \mathcal{A}} |J_i(x, \alpha) - J_i(\hat{x}, \alpha)| \\ &\leq \sup_{\alpha \in \mathcal{A}} E \left[ \int_0^\infty e^{-rt} \left| f(X_t^{x,i}, I_t^i) - f(X_t^{\hat{x},i}, I_t^i) \right| dt \right] \\ &\leq C \sup_{\alpha \in \mathcal{A}} E \left[ \int_0^\infty e^{-rt} \left| X_t^{x,i} - X_t^{\hat{x},i} \right|^{\gamma_{I_t^i}} dt \right] \\ &= C \sup_{\alpha \in \mathcal{A}} \int_0^\infty E \left[ e^{-rt} |x - \hat{x}|^{\gamma_{I_t^i}} |Y_t(i)|^{\gamma_{I_t^i}} dt \right] \\ &\leq C|x - \hat{x}|^\gamma \sup_{\alpha \in \mathcal{A}} \int_0^\infty e^{-(r-b)t} E|M_t|^{\gamma_{I_t^i}} dt \end{aligned} \tag{3.5}$$

by (3.2) and (3.4). For any  $\alpha = (\tau_n, \kappa_n)_n \in \mathcal{A}$ , by the independence of  $(Z_{\tau_n, \tau_{n+1}}^{\kappa_n})_n$  in (3.3), and since

$$E \left[ \left| Z_{\tau_n, \tau_{n+1}}^{\kappa_n} \right|^{\gamma_{\kappa_n}} \middle| \mathcal{F}_{\tau_n} \right] = E \left[ \exp \left( \gamma_{\kappa_n} (\gamma_{\kappa_n} - 1) \frac{\sigma_{\kappa_n}^2}{2} (\tau_{n+1} - \tau_n) \right) \middle| \mathcal{F}_{\tau_n} \right] \leq 1, \quad a.s.,$$

we clearly see that  $E|M_t|^{\gamma_{\kappa_n}} \leq 1$  for all  $t \geq 0$ . We thus conclude with (3.5).  $\square$

The dynamic programming principle combined with the notion of viscosity solutions are known to be a general and powerful tool for characterizing the value function of a stochastic control problem via a PDE representation, see [6]. We recall the definition of viscosity solutions for a P.D.E in the form

$$H(x, v, D_x v, D_{xx}^2 v) = 0, \quad x \in \mathcal{O}, \quad (3.6)$$

where  $\mathcal{O}$  is an open subset in  $\mathbb{R}^n$  and  $H$  is a continuous function and nonincreasing in its last argument (with respect to the order of symmetric matrices).

**Definition 3.1** *Let  $v$  be a continuous function on  $\mathcal{O}$ . We say that  $v$  is a viscosity solution to (3.6) on  $\mathcal{O}$  if it is*

(i) *a viscosity supersolution to (3.6) on  $\mathcal{O}$  : for any  $\bar{x} \in \mathcal{O}$  and any  $C^2$  function  $\varphi$  in a neighborhood of  $\bar{x}$  s.t.  $\bar{x}$  is a local minimum of  $v - \varphi$ , we have :*

$$H(\bar{x}, v(\bar{x}), D_x \varphi(\bar{x}), D_{xx}^2 \varphi(\bar{x})) \geq 0.$$

and

(ii) *a viscosity subsolution to (3.6) on  $\mathcal{O}$  : for any  $\bar{x} \in \mathcal{O}$  and any  $C^2$  function  $\varphi$  in a neighborhood of  $\bar{x}$  s.t.  $\bar{x}$  is a local maximum of  $v - \varphi$ , we have :*

$$H(\bar{x}, v(\bar{x}), D_x \varphi(\bar{x}), D_{xx}^2 \varphi(\bar{x})) \leq 0.$$

**Remark 3.1 1.** By misuse of notation, we shall say that  $v$  is viscosity supersolutin (resp. subsolution) to (3.6) by writing :

$$H(x, v, D_x v, D_{xx}^2 v) \geq (\text{resp. } \leq) 0, \quad x \in \mathcal{O}, \quad (3.7)$$

**2.** We recall that if  $v$  is a smooth  $C^2$  function on  $\mathcal{O}$ , supersolution (resp. subsolution) in the classical sense to (3.7), then  $v$  is a viscosity supersolution (resp. subsolution) to (3.7).

**3.** There is an equivalent formulation of viscosity solutions, which is useful for proving uniqueness results, see [3] :

(i) A continuous function  $v$  on  $\mathcal{O}$  is a viscosity supersolution to (3.6) if

$$H(x, v(x), p, M) \geq 0, \quad \forall x \in \mathcal{O}, \forall (p, M) \in J^{2,-} v(x).$$

(ii) A continuous function  $v$  on  $\mathcal{O}$  is a viscosity subsolution to (3.6) if

$$H(x, v(x), p, M) \leq 0, \quad \forall x \in \mathcal{O}, \forall (p, M) \in J^{2,+} v(x).$$

Here  $J^{2,+}v(x)$  is the second order superjet defined by :

$$J^{2,+}v(x) = \left\{ (p, M) \in \mathbb{R}^n \times S^n : \limsup_{\substack{x' \rightarrow x \\ x \in \mathcal{O}}} \frac{v(x') - v(x) - p \cdot (x' - x) - \frac{1}{2}(x' - x) \cdot M(x' - x)}{|x' - x|^2} \leq 0 \right\},$$

$S^n$  is the set of symmetric  $n \times n$  matrices, and  $J^{2,-}v(x) = -J^{2,+}(-v)(x)$ .

In the sequel, we shall denote by  $\mathcal{L}_i$  the second order operator associated to the diffusion  $X$  when we are in regime  $i$  : for any  $C^2$  function  $\varphi$  on  $(0, \infty)$ ,

$$\mathcal{L}_i \varphi = \frac{1}{2} \sigma_i^2 x^2 \varphi'' + b_i x \varphi'.$$

We then have the following PDE characterization of the value functions  $v_i$  by means of viscosity solutions.

**Theorem 3.1** *The value functions  $v_i$ ,  $i \in \mathbb{I}_d$ , are the unique viscosity solutions with linear growth condition on  $(0, \infty)$  and boundary condition  $v_i(0^+) = \max_{j \in \mathbb{I}_d} [-g_{ij}]$  to the system of variational inequalities :*

$$\min \left\{ rv_i - \mathcal{L}_i v_i - f_i, v_i - \max_{j \neq i} (v_j - g_{ij}) \right\} = 0, \quad x \in (0, \infty), \quad i \in \mathbb{I}_d. \quad (3.8)$$

*This means*

(1) Viscosity property : for each  $i \in \mathbb{I}_d$ ,  $v_i$  is a viscosity solution to

$$\min \left\{ rv_i - \mathcal{L}_i v_i - f_i, v_i - \max_{j \neq i} (v_j - g_{ij}) \right\} = 0, \quad x \in (0, \infty). \quad (3.9)$$

(2) Uniqueness property : if  $w_i$ ,  $i \in \mathbb{I}_d$ , are viscosity solutions with linear growth conditions on  $(0, \infty)$  and boundary conditions  $w_i(0^+) = \max_{j \in \mathbb{I}_d} [-g_{ij}]$  to the system of variational inequalities (3.8) , then  $v_i = w_i$  on  $(0, \infty)$ .

**Proof.** (1) The viscosity property follows from the dynamic programming principle and is proved in [9].

(2) Uniqueness results for switching problems has been proved in [10] in the finite horizon case under different conditions. For sake of completeness, we provide in Appendix a proof of comparison principle in our infinite horizon context, which implies the uniqueness result.  $\square$

**Remark 3.2** For fixed  $i \in \mathbb{I}_d$ , we also have uniqueness of viscosity solution to equation (3.9) in the class of continuous functions with linear growth condition on  $(0, \infty)$  and given boundary condition on 0. In the next section, we shall use either uniqueness of viscosity solutions to the system (3.8) or for fixed  $i$  to equation (3.9), for the identification of an explicit solution in the two-regimes case  $d = 2$ .



We shall also combine the uniqueness result for the viscosity solutions with the smooth-fit property on the value functions that we state below.

For any regime  $i \in \mathbb{I}_d$ , we introduce the switching region :

$$\mathcal{S}_i = \left\{ x \in (0, \infty) : v_i(x) = \max_{j \neq i} (v_j - g_{ij})(x) \right\}.$$

$\mathcal{S}_i$  is a closed subset of  $(0, \infty)$  and corresponds to the region where it is optimal for the operator to change of regime. The complement set  $\mathcal{C}_i$  of  $\mathcal{S}_i$  in  $(0, \infty)$  is the so-called continuation region :

$$\mathcal{C}_i = \left\{ x \in (0, \infty) : v_i(x) > \max_{j \neq i} (v_j - g_{ij})(x) \right\},$$

where the operator remains in regime  $i$ . In this open domain, the value function  $v_i$  is smooth  $C^2$  on  $\mathcal{C}_i$  and satisfies in a classical sense :

$$rv_i(x) - \mathcal{L}_i v_i(x) - f_i(x) = 0, \quad x \in \mathcal{C}_i.$$

As a consequence of the condition (2.6), we have the following elementary partition property of the switching regions, see Lemma 4.2 in [9] :

$$\mathcal{S}_i = \cup_{j \neq i} \mathcal{S}_{ij}, \quad i \in \mathbb{I}_d,$$

where

$$\mathcal{S}_{ij} = \{x \in \mathcal{C}_j : v_i(x) = (v_j - g_{ij})(x)\}.$$

$\mathcal{S}_{ij}$  represents the region where it is optimal to switch from regime  $i$  to regime  $j$  and stay here for a moment, i.e. without changing instantaneously from regime  $j$  to another regime. The following Lemma gives some partial information about the structure of the switching regions.

**Lemma 3.3** *For all  $i \neq j$  in  $\mathbb{I}_d$ , we have*

$$\mathcal{S}_{ij} \subset Q_{ij} := \{x \in \mathcal{C}_j : (\mathcal{L}_j - \mathcal{L}_i)v_j(x) + (f_j - f_i)(x) - rg_{ij} \geq 0\}.$$

**Proof.** Let  $x \in \mathcal{S}_{ij}$ . By setting  $\varphi_j = v_j - g_{ij}$ , this means that  $x$  is a minimum of  $v_i - \varphi_j$  with  $v_i(x) = \varphi_j(x)$ . Moreover, since  $x$  lies in the open set  $\mathcal{C}_j$  where  $v_j$  is smooth, we have that  $\varphi_j$  is  $C^2$  in a neighborhood of  $x$ . By the supersolution viscosity property of  $v_i$  to the PDE (3.8), this yields :

$$r\varphi_j(x) - \mathcal{L}_i \varphi_j(x) - f_i(x) \geq 0. \tag{3.10}$$

Now recall that for  $x \in \mathcal{C}_j$ , we have

$$rv_j(x) - \mathcal{L}_j v_j(x) - f_j(x) = 0,$$

so that by substituting into (3.10), we obtain :

$$(\mathcal{L}_j - \mathcal{L}_i)v_j(x) + (f_j - f_i)(x) - rg_{ij} \geq 0,$$

which is the required result. □

We quote the smooth fit property on the value functions, proved in [9].

**Theorem 3.2** For all  $i \in \mathbb{I}_d$ , the value function  $v_i$  is continuously differentiable on  $(0, \infty)$ .

**Remark 3.3** In a given regime  $i$ , the variational inequality satisfied by the value function  $v_i$  is a free-boundary problem as in optimal stopping problem, which divides the state space into the switching region (stopping region in pure optimal stopping problem) and the continuation region. The main difficulty with regard to optimal stopping problems for proving the smooth-fit property through the boundaries of the switching regions, comes from the fact that the switching region for the value function  $v_i$  depends also on the other value functions  $v_j$ . The method in [9] use viscosity solutions arguments and the condition of one-dimensional state space is critical for proving the smooth-fit property. The crucial conditions in this paper require that the diffusion coefficient in any regime of the system  $X$  is strictly positive on the interior the the state space, which is the case here since  $\sigma_i > 0$  for all  $i \in \mathbb{I}_d$ , and a triangular condition (2.6) on the switching costs. Under these conditions, on a point  $x$  of the switching region  $\mathcal{S}_i$  for regime  $i$ , there exists some  $j \neq i$  s.t.  $x \in \mathcal{S}_{ij}$ , i.e.  $v_i(x) = v_j(x) - g_{ij}$ , and the  $C^1$  property of the value functions is written as :  $v'_i(x) = v'_j(x)$  since  $g_{ij}$  is constant.

The next result provides suitable conditions for determining a viscosity solution to the variational inequality type arising in our switching problem.

**Lemma 3.4** Fix  $i \in \mathbb{I}_d$ . Let  $\mathcal{C}$  be an open set in  $(0, \infty)$ ,  $\mathcal{S} = (0, \infty) \setminus \mathcal{C}$  supposed to be an union of a finite number of closed intervals in  $(0, \infty)$ , and  $w, h$  two continuous functions on  $(0, \infty)$ , with  $w = h$  on  $\mathcal{S}$  such that

$$w \text{ is } C^1 \text{ on } \partial\mathcal{S} \tag{3.11}$$

$$w \geq h \text{ on } \mathcal{C}, \tag{3.12}$$

$w$  is  $C^2$  on  $\mathcal{C}$ , solution to

$$rw - \mathcal{L}_i w - f_i = 0 \text{ on } \mathcal{C}, \tag{3.13}$$

and  $w$  is a viscosity supersolution to

$$rw - \mathcal{L}_i w - f_i \geq 0 \text{ on } \text{int}(\mathcal{S}). \tag{3.14}$$

Here  $\text{int}(\mathcal{S})$  is the interior of  $\mathcal{S}$  and  $\partial\mathcal{S} = \mathcal{S} \setminus \text{int}(\mathcal{S})$  its boundary. Then,  $w$  is a viscosity solution to

$$\min \{rw - \mathcal{L}_i w - f_i, w - h\} = 0 \text{ on } (0, \infty). \tag{3.15}$$

**Proof.** Take some  $\bar{x} \in (0, \infty)$  and distinguish the following cases :

★  $\bar{x} \in \mathcal{C}$ . Since  $w = v$  is  $C^2$  on  $\mathcal{C}$  and satisfies  $rw(\bar{x}) - \mathcal{L}_i w(\bar{x}) - f_i(\bar{x}) = 0$  by (3.13), and recalling  $w(\bar{x}) \geq h(\bar{x})$  by (3.12), we obtain the classical solution property, and so a fortiori the viscosity solution property (3.15) of  $w$  at  $\bar{x}$ .

★  $\bar{x} \in \mathcal{S}$ . Then  $w(\bar{x}) = h(\bar{x})$  and the viscosity subsolution property is trivial at  $\bar{x}$ . It remains to show the viscosity supersolution property at  $\bar{x}$ . If  $\bar{x} \in \text{int}(\mathcal{S})$ , this follows

directly from (3.14). Suppose now  $\bar{x} \in \partial\mathcal{S}$ , and to fix the idea, we consider that  $\bar{x}$  is on the left-boundary of  $\mathcal{S}$  so that from the assumption on the form of  $\mathcal{S}$ , there exists  $\varepsilon > 0$  s.t.  $(\bar{x} - \varepsilon, \bar{x}) \subset \mathcal{C}$  on which  $w$  is smooth  $C^2$  (the same argument holds true when  $\bar{x}$  is on the right-boundary of  $\mathcal{S}$ ). Take some smooth  $C^2$  function  $\varphi$  s.t.  $\bar{x}$  is a local minimum of  $w - \varphi$ . Since  $w$  is  $C^1$  by (3.11), we have  $\varphi'(\bar{x}) = w'(\bar{x})$ . We may also assume w.l.o.g (by taking  $\varepsilon$  small enough) that  $(w - \varphi)(\bar{x}) \leq (w - \varphi)(x)$  for  $x \in (\bar{x} - \varepsilon, \bar{x})$ . Moreover, by Taylor's formula, we have :

$$w(\bar{x} - \eta) = w(\bar{x}) - \eta \int_0^1 w'(\bar{x} - t\eta) dt, \quad \varphi(\bar{x} - \eta) = \varphi(\bar{x}) - \eta \int_0^1 \varphi'(\bar{x} - t\eta) dt,$$

so that

$$\int_0^1 \varphi'(\bar{x} - t\eta) - w'(\bar{x} - t\eta) dt \geq 0, \quad \forall 0 < \eta < \varepsilon.$$

Since  $\varphi'(\bar{x}) = w'(\bar{x})$ , this last inequality is written as

$$\int_0^1 \frac{\varphi'(\bar{x} - t\eta) - \varphi'(\bar{x})}{\eta} - \frac{w'(\bar{x} - t\eta) - w'(\bar{x})}{\eta} dt \geq 0, \quad \forall 0 < \eta < \varepsilon, \quad (3.16)$$

Now, from (3.13), we have  $rw(x) - \mathcal{L}_i w(x) - f_i(x) = 0$  for  $x \in (\bar{x} - \varepsilon, \bar{x})$ . By sending  $x$  towards  $\bar{x}$  into this last equality, this shows that  $w''(\bar{x}^-) = \lim_{x \nearrow \bar{x}} w''(x)$  exists, and

$$rw(\bar{x}) - b_i \bar{x} w'(\bar{x}) - \frac{1}{2} \sigma_i^2 \bar{x}^2 w''(\bar{x}^-) - f_i(\bar{x}) = 0. \quad (3.17)$$

Moreover, by sending  $\eta$  to zero into (3.16), we obtain :

$$\int_0^1 t[-\varphi''(\bar{x}) + w''(\bar{x}^-)] dt \geq 0,$$

and so  $\varphi''(\bar{x}) \leq w''(\bar{x}^-)$ . By substituting into (3.17), and recalling that  $w'(\bar{x}) = \varphi'(\bar{x})$ , we then obtain :

$$rw(\bar{x}) - \mathcal{L}_i \varphi(\bar{x}) - f_i(\bar{x}) \geq 0,$$

which is the required supersolution inequality, and ends the proof.  $\square$

**Remark 3.4** Since  $w = h$  on  $\mathcal{S}$ , relation (3.14) means equivalently that  $h$  is a viscosity supersolution to

$$rh - \mathcal{L}_i h - f_i \geq 0 \quad \text{on } \text{int}(\mathcal{S}). \quad (3.18)$$

Practically, Lemma 3.4 shall be used as follows in the next section : we consider two  $C^1$  functions  $v$  and  $h$  on  $(0, \infty)$  s.t.

$$\begin{aligned} v(x) &= h(x), \quad v'(x) = h'(x), \quad x \in \partial\mathcal{S} \\ v &\geq h \quad \text{on } \mathcal{C}, \end{aligned}$$

$v$  is  $C^2$  on  $\mathcal{C}$ , solution to

$$rv - \mathcal{L}_i v - f_i = 0 \quad \text{on } \mathcal{C},$$

and  $h$  is a viscosity supersolution to (3.18). Then, the function  $w$  defined on  $(0, \infty)$  by :

$$w(x) = \begin{cases} v(x), & x \in \mathcal{C} \\ h(x), & x \in \mathcal{S} \end{cases}$$

satisfies the conditions of Lemma 3.4 and is so a viscosity solution to (3.15). This Lemma combined with uniqueness viscosity solution result may be viewed as an alternative to the classical verification approach in the identification of the value function. Moreover, with our viscosity solutions approach, we shall see in subsection 4.2 that Lemma 3.3 and smooth-fit property of the value functions in Theorem 3.2 provide a direct derivation for the structure of the switching regions and then of the solution to our problem.

## 4 Explicit solution in the two regime case

In this section, we consider the case where  $d = 2$ . In this two-regimes case, we know from Theorem 3.1 that the value functions  $v_i$ ,  $i = 1, 2$ , are the unique continuous viscosity solutions with linear growth condition on  $(0, \infty)$ , and boundary conditions  $v_i(0^+) = (-g_{ij})_+ := \max(-g_{ij}, 0)$ ,  $j \neq i$ , to the system :

$$\min \{rv_1 - \mathcal{L}_1 v_1 - f_1, v_1 - (v_2 - g_{12})\} = 0 \quad (4.1)$$

$$\min \{rv_2 - \mathcal{L}_2 v_2 - f_2, v_2 - (v_1 - g_{21})\} = 0. \quad (4.2)$$

Moreover, the switching regions are :

$$\mathcal{S}_i = \mathcal{S}_{ij} = \{x > 0 : v_i(x) = v_j(x) - g_{ij}\}, \quad i, j = 1, 2, i \neq j.$$

We set

$$\underline{x}_i^* = \inf \mathcal{S}_i \in [0, \infty] \quad \bar{x}_i^* = \sup \mathcal{S}_i \in [0, \infty],$$

with the usual convention that  $\inf \emptyset = \infty$ .

Let us also introduce some other notations. We consider the second order o.d.e for  $i = 1, 2$  :

$$rv - \mathcal{L}_i v - f_i = 0, \quad (4.3)$$

whose general solution (without second member  $f_i$ ) is given by :

$$v(x) = Ax^{m_i^+} + Bx^{m_i^-},$$

for some constants  $A, B$ , and where

$$m_i^- = -\frac{b_i}{\sigma_i^2} + \frac{1}{2} - \sqrt{\left(-\frac{b_i}{\sigma_i^2} + \frac{1}{2}\right)^2 + \frac{2r}{\sigma_i^2}} < 0$$

$$m_i^+ = -\frac{b_i}{\sigma_i^2} + \frac{1}{2} + \sqrt{\left(-\frac{b_i}{\sigma_i^2} + \frac{1}{2}\right)^2 + \frac{2r}{\sigma_i^2}} > 1.$$

We also denote

$$\hat{V}_i(x) = E \left[ \int_0^\infty e^{-rt} f_i(\hat{X}_t^{x,i}) dt \right],$$

with  $\hat{X}^{x,i}$  the solution to the s.d.e.  $d\hat{X}_t = b_i \hat{X}_t dt + \sigma_i \hat{X}_t dW_t$ ,  $\hat{X}_0 = x$ . Actually,  $\hat{V}_i$  is a particular solution to ode (4.3), with boundary condition  $\hat{V}_i(0^+) = f_i(0) = 0$ . It corresponds to the reward function associated to the no switching strategy from initial state  $(x, i)$ , and so  $\hat{V}_i \leq v_i$ .

**Remark 4.1** If  $g_{ij} > 0$ , then from (2.7), we have  $v_i(0^+) = 0 > (-g_{ji})_+ - g_{ij} = v_j(0^+) - g_{ij}$ . Therefore, by continuity of the value functions on  $(0, \infty)$ , we get  $\underline{x}_i^* > 0$ .

We now give the explicit solution to our problem in the following two situations :

- ★ the diffusion operators are different and the running profit functions are identical.
- ★ the diffusion operators are identical and the running profit functions are different

We also consider the cases for which both switching costs are positive, and for which one of the two is negative, the other being then positive according to (2.7). This last case is interesting in applications where a firm chooses between an open or closed activity, and may regain a fraction of its opening costs when it decides to close.

#### 4.1 Identical profit functions with different diffusion operators

In this subsection, we suppose that the running functions are identical in the form :

$$f_1(x) = f_2(x) = x^\gamma, \quad 0 < \gamma < 1, \quad (4.4)$$

and the diffusion operators are different. A straightforward calculation shows that under (4.4), we have

$$\hat{V}_i(x) = K_i x^\gamma, \quad \text{with } K_i = \frac{1}{r - b_i \gamma + \frac{1}{2} \sigma_i^2 \gamma (1 - \gamma)} > 0, \quad i = 1, 2.$$

We show that the structure of the switching regions depends actually only on the sign of  $K_2 - K_1$ , and of the sign of the switching costs  $g_{12}$  and  $g_{21}$ . More precisely, we have the following explicit result.

**Theorem 4.1** *Let  $i, j = 1, 2, i \neq j$ .*

1) *If  $K_i = K_j$ , then*

$$v_i(x) = \hat{V}_i(x) + (-g_{ij})_+, \quad x \in (0, \infty),$$

$$\mathcal{S}_i = \begin{cases} \emptyset & \text{if } g_{ij} > 0 \\ (0, \infty) & \text{if } g_{ij} \leq 0. \end{cases}$$

*It is always optimal to switch from regime  $i$  to  $j$  if the corresponding switching cost is nonpositive, and never optimal to switch otherwise.*

2) *If  $K_j > K_i$ , then we have the following situations depending on the switching costs :*

a)  $g_{ij} \leq 0$  : we have  $\mathcal{S}_i = (0, \infty)$ ,  $\mathcal{S}_j = \emptyset$ , and

$$v_i = \hat{V}_j - g_{ij}, \quad v_j = \hat{V}_j.$$

b)  $g_{ij} > 0$  :

- if  $g_{ji} \geq 0$ , then  $\mathcal{S}_i = [\underline{x}_i^*, \infty)$  with  $\underline{x}_i^* \in (0, \infty)$ ,  $\mathcal{S}_j = \emptyset$ , and

$$v_i(x) = \begin{cases} Ax^{m_i^+} + \hat{V}_i(x), & x < \underline{x}_i^* \\ v_j(x) - g_{ij}, & x \geq \underline{x}_i^* \end{cases} \quad (4.5)$$

$$v_j(x) = \hat{V}_j(x), \quad x \in (0, \infty) \quad (4.6)$$

where the constants  $A$  and  $\underline{x}_i^*$  are determined by the continuity and smooth-fit conditions of  $v_i$  at  $\underline{x}_i^*$ , and explicitly given by :

$$\underline{x}_i^* = \left( \frac{m_i^+}{m_i^+ - \gamma} \frac{g_{ij}}{K_j - K_i} \right)^{\frac{1}{\gamma}} \quad (4.7)$$

$$A = (K_j - K_i) \frac{\gamma}{m_i^+} (\underline{x}_i^*)^{\gamma - m_i^+}. \quad (4.8)$$

When we are in regime  $i$ , it is optimal to switch to regime  $j$  whenever the state process  $X$  exceeds the threshold  $\underline{x}_i^*$ , while when we are in regime  $j$ , it is optimal never to switch.

- if  $g_{ji} < 0$ , then  $\mathcal{S}_i = [\underline{x}_i^*, \infty)$  with  $\underline{x}_i^* \in (0, \infty)$ ,  $\mathcal{S}_j = (0, \bar{x}_j^*]$ , and

$$v_i(x) = \begin{cases} Ax^{m_i^+} + \hat{V}_i(x), & x < \underline{x}_i^* \\ v_j(x) - g_{ij}, & x \geq \underline{x}_i^* \end{cases} \quad (4.9)$$

$$v_j(x) = \begin{cases} v_i(x) - g_{ji}, & x \leq \bar{x}_j^* \\ Bx^{m_j^-} + \hat{V}_j(x), & x > \bar{x}_j^* \end{cases} \quad (4.10)$$

where the constants  $A$ ,  $B$  and  $\bar{x}_j^* < \underline{x}_i^*$  are determined by the continuity and smooth-fit conditions of  $v_i$  and  $v_j$  at  $\underline{x}_i^*$  and  $\bar{x}_j^*$ , and explicitly given by :

$$\begin{aligned} \bar{x}_j^* &= \left[ \frac{-m_j^-(g_{ji} + g_{ij}y^{m_i^+})}{(K_i - K_j)(\gamma - m_j^-)(1 - y^{m_i^+ - \gamma})} \right]^{\frac{1}{\gamma}} \\ \underline{x}_i &= \frac{\bar{x}_j^*}{y} \\ B &= \frac{(K_i - K_j)(m_i^+ - \gamma)\underline{x}_i^{\gamma - m_j^-} + m_i^+g_{ij}\underline{x}_i^{-m_j^-}}{m_i^+ - m_j^-} \\ A &= B\underline{x}_i^{m_j^- - m_i^+} - (K_i - K_j)\underline{x}_i^{\gamma - m_i^+} - g_{ij}\underline{x}_i^{-m_i^+} \end{aligned}$$

with  $y$  solution in  $\left(0, \left(-\frac{g_{ji}}{g_{ij}}\right)^{\frac{1}{m_i^+}}\right)$  to the equation :

$$\begin{aligned} &m_i^+(\gamma - m_j^-) \left(1 - y^{m_i^+ - \gamma}\right) \left(g_{ij}y^{m_j^-} + g_{ji}\right) \\ &+ m_j^-(m_i^+ - \gamma) \left(1 - y^{m_j^- - \gamma}\right) \left(g_{ij}y^{m_i^+} + g_{ji}\right) = 0 \end{aligned}$$

When we are in regime  $i$ , it is optimal to switch to regime  $j$  whenever the state process  $X$  exceeds the threshold  $\underline{x}_i^*$ , while when we are in regime  $j$ , it is optimal to switch to regime  $i$  for values of the state process  $X$  under the threshold  $\bar{x}_j^*$ .

### Economic interpretation.

In the particular case where  $\sigma_1 = \sigma_2$ , then  $K_2 - K_1 > 0$  means that regime 2 provides a higher expected return  $b_2$  than the one  $b_1$  of regime 1 for the same volatility coefficient  $\sigma_i$ . Moreover, if the switching cost  $g_{21}$  from regime 2 to regime 1 is nonnegative, it is intuitively clear that one has always interest to stay in regime 2, which is formalized by the property that  $\mathcal{S}_2 = \emptyset$ . However, if one receives some gain compensation to switch from regime 2 to regime 1, i.e. the corresponding cost  $g_{21}$  is negative, then one has interest to change of regime for small values of the current state. This is formalized by the property that  $\mathcal{S}_2 = (0, \bar{x}_2^*]$ . On the other hand, in regime 1, one has interest to switch to regime 2, for all current values of the state if the corresponding switching cost  $g_{12}$  is nonpositive, or from a certain threshold  $\underline{x}_1^*$  if the switching cost  $g_{12}$  is positive. A similar interpretation holds when  $b_1 = b_2$ , and  $K_2 - K_1 > 0$ , i.e.  $\sigma_2 < \sigma_1$ . Theorem 4.1 extends these results for general coefficients  $b_i$  and  $\sigma_i$ , and show that the critical parameter value determining the form of the optimal strategy is given by the sign of  $K_2 - K_1$  and the switching costs. The different optimal strategy structures are depicted in Figure I.

### Proof of Theorem 4.1.

1) If  $K_i = K_j$ , then  $\hat{V}_i = \hat{V}_j$ . We consider the smooth functions  $w_i = \hat{V}_i + (-g_{ij})_+$  for  $i, j = 1, 2$  and  $j \neq i$ . Since  $\hat{V}_i$  are solution to (4.3), we see that  $w_i$  satisfy :

$$rw_i - \mathcal{L}w_i - f_i = r(-g_{ij})_+ \quad (4.11)$$

$$w_i - (w_j - g_{ij}) = g_{ij} + (-g_{ij})_+ - (-g_{ji})_+. \quad (4.12)$$

Notice that the l.h.s of (4.11) and (4.12) are both nonnegative by (2.7). Moreover, if  $g_{ij} > 0$ , then the l.h.s. of (4.11) is zero, and if  $g_{ij} \leq 0$ , then  $g_{ji} > 0$  and the l.h.s. of (4.12) is zero. Therefore,  $w_i, i = 1, 2$  is solution to the system :

$$\min \{rw_i - \mathcal{L}_i w_i - f_i, w_i - (w_j - g_{ij})\} = 0.$$

Since  $\hat{V}_i(0^+) = 0$ , we have  $w_i(0^+) = (-g_{ij})_+$ . Moreover,  $w_i$  satisfy like  $\hat{V}_i$  a linear growth condition. Therefore, from uniqueness of solution to the PDE system (4.1)-(4.2), we deduce that  $v_i = w_i$ . As observed above, if  $g_{ij} \leq 0$ , then the l.h.s. of (4.12) is zero, and so  $\mathcal{S}_i = (0, \infty)$ . Finally, if  $g_{ij} > 0$ , then the l.h.s. of (4.12) is positive, and so  $\mathcal{S}_i = \emptyset$ .

2) We now suppose w.l.o.g. that  $K_2 > K_1$ .

a) Consider first the case where  $g_{12} \leq 0$ , and so  $g_{21} > 0$ . We set  $w_1 = \hat{V}_2 - g_{12}$  and  $w_2 = \hat{V}_2$ . Then, by construction, we have  $w_1 = w_2 - g_{12}$  on  $(0, \infty)$ , and by definition of  $\hat{V}_1$  and  $\hat{V}_2$  :

$$rw_1(x) - \mathcal{L}_1 w_1(x) - f_1(x) = \frac{K_2 - K_1}{K_1} x^\gamma - rg_{12} > 0, \quad \forall x > 0.$$

On the other hand, we also have  $rw_2 - \mathcal{L}_2 w_2 - f_2 = 0$  on  $(0, \infty)$ , and  $w_2 \geq w_1 - g_{21}$  since  $g_{12} + g_{21} \geq 0$ . Hence,  $w_1$  and  $w_2$  are smooth (hence viscosity) solutions to the system

### Figure I




Figure I.1.a:  $f_1 = f_2$ ,  $K_1 = K_2$ ,  $g_{12} > 0$ ,  $g_{21} > 0$




Figure I.1.b:  $f_1 = f_2$ ,  $K_1 = K_2$ ,  $g_{12} > 0$ ,  $g_{21} \leq 0$




Figure I.2.a:  $f_1 = f_2$ ,  $K_2 > K_1$ ,  $g_{12} \leq 0$




Figure I.2.bi:  $f_1 = f_2$ ,  $K_2 > K_1$ ,  $g_{12} > 0$ ,  $g_{21} \geq 0$




Figure I.2.bii:  $f_1 = f_2$ ,  $K_2 > K_1$ ,  $g_{12} > 0$ ,  $g_{21} < 0$



(4.1)-(4.2), with linear growth conditions and boundary conditions  $w_1(0^+) = V_1(0^+) - g_{12} = (-g_{12})_+$ ,  $w_2(0^+) = \hat{V}_2(0^+) = 0 = (-g_{21})_+$ . By uniqueness result of Theorem 3.1, we deduce that  $v_1 = w_1$ ,  $v_2 = w_2$ , and thus  $\mathcal{S}_1 = (0, \infty)$ ,  $\mathcal{S}_2 = \emptyset$ .

b) Consider now the case where  $g_{12} > 0$ . We already know from Remark 4.1 that  $\underline{x}_1^* > 0$ , and we claim that  $\underline{x}_1^* < \infty$ . Otherwise,  $v_1$  should be equal to  $\hat{V}_1$ . Since  $v_1 \geq v_2 - g_{12} \geq \hat{V}_2 - g_{12}$ , this would imply  $(\hat{V}_2 - \hat{V}_1)(x) = (K_2 - K_1)x^\gamma \leq g_{12}$  for all  $x > 0$ , an obvious contradiction. By definition of  $\underline{x}_1^*$ , we have  $(0, \underline{x}_1^*) \subset \mathcal{C}_1$ . We shall prove actually the equality :  $(0, \underline{x}_1^*) = \mathcal{C}_1$ , i.e.  $\mathcal{S}_1 = [\underline{x}_1^*, \infty)$ . On the other hand, the form of  $\mathcal{S}_2$  will depend on the sign of  $g_{21}$ .

• *Case* :  $g_{21} \geq 0$ .

We shall prove that  $\mathcal{C}_2 = (0, \infty)$ , i.e.  $\mathcal{S}_2 = \emptyset$ . To this end, let us consider the function

$$w_1(x) = \begin{cases} Ax^{m_1^+} + \hat{V}_1(x), & 0 < x < x_1 \\ \hat{V}_2(x) - g_{12}, & x \geq x_1, \end{cases}$$

where the positive constants  $A$  and  $x_1$  satisfy

$$Ax_1^{m_1^+} + \hat{V}_1(x_1) = \hat{V}_2(x_1) - g_{12} \quad (4.13)$$

$$Am_1^+ x_1^{m_1^+ - 1} + \hat{V}_1'(x_1) = \hat{V}_2'(x_1), \quad (4.14)$$

and are explicitly determined by :

$$(K_2 - K_1)x_1^\gamma = \frac{m_1^+}{m_1^+ - \gamma} g_{12} \quad (4.15)$$

$$A = (K_2 - K_1) \frac{\gamma}{m_1^+} x_1^{\gamma - m_1^+}. \quad (4.16)$$

Notice that by construction,  $w_1$  is  $C^2$  on  $(0, x_1) \cup (x_1, \infty)$ , and  $C^1$  on  $x_1$ .

★ By using Lemma 3.4, we now show that  $w_1$  is a viscosity solution to

$$\min \left\{ rw_1 - \mathcal{L}_1 w_1 - f_1, w_1 - (\hat{V}_2 - g_{12}) \right\} = 0, \quad \text{on } (0, \infty). \quad (4.17)$$

We first check that

$$w_1(x) \geq \hat{V}_2(x) - g_{12}, \quad \forall 0 < x < x_1, \quad (4.18)$$

i.e.

$$G(x) := Ax^{m_1^+} + \hat{V}_1(x) - \hat{V}_2(x) + g_{12} \geq 0, \quad \forall 0 < x < x_1.$$

Since  $A > 0$ ,  $0 < \gamma < 1 < m_1^+$ ,  $K_2 - K_1 > 0$ , a direct derivation shows that the second derivative of  $G$  is positive, i.e.  $G$  is strictly convex. By (4.14), we have  $G'(x_1) = 0$  and so  $G'$  is negative, i.e.  $G$  is strictly decreasing on  $(0, x_1)$ . Now, by (4.13), we have  $G(x_1) = 0$  and thus  $G$  is positive on  $(0, x_1)$ , which proves (4.18).

By definition of  $w_1$  on  $(0, x_1)$ , we have in the classical sense

$$rw_1 - \mathcal{L}_1 w_1 - f_1 = 0, \quad \text{on } (0, x_1). \quad (4.19)$$

We now check that

$$rw_1 - \mathcal{L}_1 w_1 - f_1 \geq 0, \quad \text{on } (x_1, \infty), \quad (4.20)$$

holds true in the classical sense, and so a fortiori in the viscosity sense. By definition of  $w_1$  on  $(x_1, \infty)$ , and  $K_1$ , we have for all  $x > x_1$ ,

$$rw_1(x) - \mathcal{L}_1 w_1(x) - f_1(x) = \frac{K_2 - K_1}{K_1} x^\gamma - rg_{12}, \quad \forall x > x_1,$$

so that (4.20) is satisfied iff  $\frac{K_2 - K_1}{K_1} x_1^\gamma - rg_{12} \geq 0$  or equivalently by (4.15) :

$$\frac{m_1^+}{m_1^+ - \gamma} \geq rK_1 = \frac{r}{r - b_1\gamma + \frac{1}{2}\sigma_1^2\gamma(1 - \gamma)} \quad (4.21)$$

Now, since  $\gamma < 1 < m_1^+$ , and by definition of  $m_1^+$ , we have

$$\frac{1}{2}\sigma_1^2 m_1^+(\gamma - 1) < \frac{1}{2}\sigma_1^2 m_1^+(m_1^+ - 1) = r - b_1 m_1^+,$$

which proves (4.21) and thus (4.20).

Relations (4.13)-(4.14), (4.18)-(4.19)-(4.20) mean that conditions of Lemma 3.4 are satisfied with  $\mathcal{C} = (0, x_1)$ ,  $h = \hat{V}_2 - g_{12}$ , and we thus get the required assertion (4.17).

★ On the other hand, we check that

$$\hat{V}_2(x) \geq w_1(x) - g_{21}, \quad \forall x > 0, \quad (4.22)$$

which amounts to show

$$H(x) := Ax^{m_1^+} + \hat{V}_1(x) - \hat{V}_2(x) - g_{21} \leq 0, \quad \forall 0 < x < x_1.$$

Since  $A > 0$ ,  $0 < \gamma < 1 < m_1^+$ ,  $K_2 - K_1 > 0$ , a direct derivation shows that the second derivative of  $H$  is positive, i.e.  $H$  is strictly convex. By (4.14), we have  $H'(x_1) = 0$  and so  $H'$  is negative, i.e.  $H$  is strictly decreasing on  $(0, x_1)$ . Now, we have  $H(0) = -g_{21} \leq 0$  and thus  $H$  is negative on  $(0, x_1)$ , which proves (4.22). Recalling that  $\hat{V}_2$  is solution to  $r\hat{V}_2 - \mathcal{L}_2\hat{V}_2 - f_2 = 0$  on  $(0, \infty)$ , we deduce obviously from (4.22) that  $\hat{V}_2$  is a classical, hence a viscosity solution to :

$$\min \left\{ r\hat{V}_2 - \mathcal{L}_2\hat{V}_2 - f_2, \hat{V}_2 - (w_1 - g_{21}) \right\} = 0, \quad \text{on } (0, \infty). \quad (4.23)$$

★ Since  $w_1(0^+) = 0 = (-g_{12})_+$ ,  $\hat{V}_2(0^+) = 0 = (-g_{21})_+$ , and  $w_1, \hat{V}_2$  satisfy a linear growth condition, we deduce from (4.17), (4.23), and uniqueness to the PDE system (4.1)-(4.2), that

$$v_1 = w_1, \quad v_2 = \hat{V}_2, \quad \text{on } (0, \infty).$$

This proves  $\underline{x}_1^* = x_1$ ,  $\mathcal{S}_1 = [x_1, \infty)$  and  $\mathcal{S}_2 = \emptyset$ .

- *Case* :  $g_{21} < 0$ .

We shall prove that  $\mathcal{S}_2 = (0, \bar{x}_2]$ . To this end, let us consider the functions

$$\begin{aligned} w_1(x) &= \begin{cases} Ax^{m_1^+} + \hat{V}_1(x), & x < \underline{x}_1 \\ w_2(x) - g_{12}, & x \geq \underline{x}_1 \end{cases} \\ w_2(x) &= \begin{cases} w_1(x) - g_{21}, & x \leq \bar{x}_2 \\ Bx^{m_2^-} + \hat{V}_2(x), & x > \bar{x}_2, \end{cases} \end{aligned}$$

where the positive constants  $A, B, \underline{x}_1 > \bar{x}_2$ , solution to

$$A\underline{x}_1^{m_1^+} + \hat{V}_1(\underline{x}_1) = w_2(\underline{x}_1) - g_{12} = B\underline{x}_1^{m_2^-} + \hat{V}_2(\underline{x}_1) - g_{12} \quad (4.24)$$

$$Am_1^+\underline{x}_1^{m_1^+-1} + \hat{V}_1'(\underline{x}_1) = w_2'(\underline{x}_1) = Bm_2^-\underline{x}_1^{m_2^- - 1} + \hat{V}_2'(\underline{x}_1) \quad (4.25)$$

$$A\bar{x}_2^{m_1^+} + \hat{V}_1(\bar{x}_2) - g_{21} = w_1(\bar{x}_2) - g_{21} = B\bar{x}_2^{m_2^-} + \hat{V}_2(\bar{x}_2) \quad (4.26)$$

$$Am_1^+\bar{x}_2^{m_1^+-1} + \hat{V}_1'(\bar{x}_2) = w_1'(\bar{x}_2) = Bm_2^-\bar{x}_2^{m_2^- - 1} + \hat{V}_2'(\bar{x}_2), \quad (4.27)$$

exist and are explicitly determined after some calculations by

$$\bar{x}_2 = \left[ \frac{-m_2^-(g_{21} + g_{12}y^{m_1^+})}{(K_1 - K_2)(\gamma - m_2^-)(1 - y^{m_1^+ - \gamma})} \right]^{\frac{1}{\gamma}} \quad (4.28)$$

$$\underline{x}_1 = \frac{\bar{x}_2}{y} \quad (4.29)$$

$$B = \frac{(K_1 - K_2)(m_1^+ - \gamma)\underline{x}_1^{\gamma - m_2^-} + m_1^+g_{12}\underline{x}_1^{-m_2^-}}{m_1^+ - m_2^-} \quad (4.30)$$

$$A = B\underline{x}_1^{m_2^- - m_1^+} - (K_1 - K_2)\underline{x}_1^{\gamma - m_1^+} - g_{12}\underline{x}_1^{-m_1^+}, \quad (4.31)$$

with  $y$  solution in  $\left(0, \left(-\frac{g_{21}}{g_{12}}\right)^{\frac{1}{m_1^+}}\right)$  to the equation :

$$\begin{aligned} & m_1^+(\gamma - m_2^-) \left(1 - y^{m_1^+ - \gamma}\right) \left(g_{12}y^{m_2^-} + g_{21}\right) \\ & + m_2^-(m_1^+ - \gamma) \left(1 - y^{m_2^- - \gamma}\right) \left(g_{12}y^{m_1^+} + g_{21}\right) = 0. \end{aligned} \quad (4.32)$$

Using (2.7), we have  $y < \left(-\frac{g_{21}}{g_{12}}\right)^{\frac{1}{m_1^+}} < 1$ . As such,  $0 < \bar{x}_2 < \underline{x}_1$ . Furthermore, by using (4.29) and the equation (4.32) satisfied by  $y$ , we may easily check that  $A$  and  $B$  are positive constants.

Notice that by construction,  $w_1$  (resp.  $w_2$ ) is  $C^2$  on  $(0, \underline{x}_1) \cup (\underline{x}_1, \infty)$  (resp.  $(0, \bar{x}_2) \cup (\bar{x}_2, \infty)$ ) and  $C^1$  at  $\underline{x}_1$  (resp.  $\bar{x}_2$ ).

★ By using Lemma 3.4, we now show that  $w_i$ ,  $i = 1, 2$ , is a viscosity solution to the system :

$$\min\{rw_i - \mathcal{L}_i w_i - f_i, w_i - (w_j - g_{ij})\} = 0, \quad \text{on } (0, \infty), \quad i, j = 1, 2, \quad j \neq i. \quad (4.33)$$

Since the proof is similar for both  $w_i$ ,  $i = 1, 2$ , we only prove the result for  $w_1$ . We first check that

$$w_1 \geq w_2 - g_{12}, \quad \forall 0 < x < \underline{x}_1. \quad (4.34)$$

From the definition of  $w_1$  and  $w_2$  and using the fact that  $g_{12} + g_{21} > 0$ , it is straightforward to see that

$$w_1 \geq w_2 - g_{12}, \quad \forall 0 < x \leq \bar{x}_2. \quad (4.35)$$

Now, we need to prove that

$$G(x) := Ax^{m_1^+} + \hat{V}_1(x) - Bx^{m_2^-} - \hat{V}_2(x) + g_{12} \geq 0, \quad \forall \bar{x}_2 < x < \underline{x}_1. \quad (4.36)$$

We have  $G(\bar{x}_2) = g_{12} + g_{21} > 0$  and  $G(\underline{x}_1) = 0$ . Suppose that there exists some  $x_0 \in (\bar{x}_2, \underline{x}_1)$  such that  $G(x_0) = 0$ . We then deduce that there exists  $x_3 \in (\bar{x}_2, \underline{x}_1)$  such that  $G'(x_3) = 0$ . As such, the equation  $G'(x) = 0$  admits at least three solutions in  $[\bar{x}_2, \underline{x}_1] : \{\bar{x}_2, x_3, \underline{x}_1\}$ . However, a straightforward study of the function  $G$  shows that  $G'$  can take the value zero at most at two points in  $(0, \infty)$ . This leads to a contradiction, proving therefore (4.36).

By definition of  $w_1$ , we have in the classical sense

$$rw_1 - \mathcal{L}_1 w_1 - f = 0, \quad \text{on } (0, \underline{x}_1). \quad (4.37)$$

We now check that

$$rw_1 - \mathcal{L}_1 w_1 - f \geq 0, \quad \text{on } (\underline{x}_1, \infty) \quad (4.38)$$

holds true in the classical sense, and so a fortiori in the viscosity sense. By definition of  $w_1$  on  $(x_1, \infty)$ , and  $K_1$ , we have for all  $x > \underline{x}_1$ ,

$$H(x) := rw_1(x) - \mathcal{L}_1 w_1(x) - f(x) = \frac{K_2 - K_1}{K_1} x^\gamma + m_2^- LBx^{m_2^-} - rg_{12}, \quad \forall x > \underline{x}_1, \quad (4.39)$$

where  $L = \frac{1}{2}(\sigma_2^2 - \sigma_1^2)(m_2^- - 1) + b_2 - b_1$ .

We distinguish two cases :

- First, if  $L \geq 0$ , the function  $H$  would be non-decreasing on  $(0, \infty)$  with  $\lim_{x \rightarrow 0^+} H(x) = -\infty$  and  $\lim_{x \rightarrow \infty} H(x) = +\infty$ . As such, it suffices to show that  $H(\underline{x}_1) \geq 0$ . From (4.24)-(4.25), we have

$$H(\underline{x}_1) = (K_2 - K_1) \left[ \frac{m_1^+ - m_2^-}{K_1} - (m_1^+ - \gamma)m_2^- L \right] - rg_{12} + m_1^+ m_2^- g_{12} L.$$

Using relations (4.21), (4.24), (4.25), (4.29) and the definition of  $m_1^+$  and  $m_2^-$ , we then obtain

$$H(\underline{x}_1) = \frac{m_1^+(m_1^+ - m_2^-)}{K_1(m_1^+ - \gamma)} - r \geq \frac{m_1^+}{K_1(m_1^+ - \gamma)} - r \geq 0.$$

- Second, if  $L < 0$ , it suffices to show that

$$\frac{K_2 - K_1}{K_1} x^\gamma - rg_{12} \geq 0, \quad \forall x > \underline{x}_1,$$

which is rather straightforward from (4.21) and (4.29) .

Relations (4.34), (4.37) (4.38) and the regularity of  $w_i$ ,  $i = 1, 2$ , as constructed, mean that

conditions of Lemma 3.4 are satisfied and we thus get the required assertion (4.33).

★ Since  $w_1(0^+) = 0 = (-g_{12})_+$ ,  $w_2(0^+) = -g_{21} = (-g_{21})_+$ , and  $w_1, \hat{V}_2$  satisfy a linear growth condition, we deduce from (4.33) and uniqueness to the PDE system (4.1)-(4.2), that

$$v_1 = w_1, \quad v_2 = w_2, \quad \text{on } (0, \infty).$$

This proves  $\underline{x}_1^* = \underline{x}_1$ ,  $\mathcal{S}_1 = [x_1, \infty)$  and  $\bar{x}_2^* = \bar{x}_2$ ,  $\mathcal{S}_2 = (0, \bar{x}_2]$ .

## 4.2 Identical diffusion operators with different profit functions

In this subsection, we suppose that  $\mathcal{L}_1 = \mathcal{L}_2 = \mathcal{L}$ , i.e.  $b_1 = b_2 = b$ ,  $\sigma_1 = \sigma_2 = \sigma > 0$ . We then set  $m^+ = m_1^+ = m_2^+$ ,  $m^- = m_1^- = m_2^-$ , and  $\hat{X}^x = \hat{X}^{x,1} = \hat{X}^{x,2}$ . Notice that in this case, the set  $Q_{ij}$ ,  $i, j = 1, 2$ ,  $i \neq j$ , introduced in Lemma 3.3, satisfies :

$$\begin{aligned} Q_{ij} &= \{x \in \mathcal{C}_j : (f_j - f_i)(x) - rg_{ij} \geq 0\} \\ &\subset \hat{Q}_{ij} := \{x > 0 : (f_j - f_i)(x) - rg_{ij} \geq 0\}. \end{aligned} \quad (4.40)$$

Once we are given the profit functions  $f_i, f_j$ , the set  $\hat{Q}_{ij}$  can be explicitly computed. Moreover, we prove in the next key Lemma that the structure of  $\hat{Q}_{ij}$ , when it is connected, determines the same structure for the switching region  $\mathcal{S}_i$ .

**Lemma 4.1** *Let  $i, j = 1, 2$ ,  $i \neq j$ .*

1) *Assume that*

$$\sup_{x>0} (\hat{V}_j - \hat{V}_i)(x) > g_{ij}. \quad (4.41)$$

• *If there exists  $0 < \underline{x}_{ij} < \infty$  such that*

$$\hat{Q}_{ij} = [\underline{x}_{ij}, \infty), \quad (4.42)$$

*then  $0 < \underline{x}_i^* < \infty$  and*

$$\mathcal{S}_i = [\underline{x}_i^*, \infty).$$

• *If  $g_{ij} \leq 0$  and there exists  $0 < \bar{x}_{ij} < \infty$  such that*

$$\hat{Q}_{ij} = (0, \bar{x}_{ij}], \quad (4.43)$$

*then  $0 < \bar{x}_i^* < \infty$  and*

$$\mathcal{S}_i = (0, \bar{x}_i^*].$$

2) *If there exist  $0 < \underline{x}_{ij} < \bar{x}_{ij} < \infty$  such that*

$$\hat{Q}_{ij} = [\underline{x}_{ij}, \bar{x}_{ij}]. \quad (4.44)$$

*Then  $0 < \underline{x}_i^* < \bar{x}_i^* < \infty$  and*

$$\mathcal{S}_i = [\underline{x}_i^*, \bar{x}_i^*].$$

3) *If  $g_{ij} \leq 0$  and  $\hat{Q}_{ij} = (0, \infty)$ , then  $\mathcal{S}_i = (0, \infty)$  and  $\mathcal{S}_j = \emptyset$ .*

**Proof.** 1) • Consider the case of condition (4.42). Since  $\mathcal{S}_i \subset \hat{Q}_{ij}$  by Lemma 3.3, this implies  $\underline{x}_i^* := \inf \mathcal{S}_i \geq \underline{x}_{ij} > 0$ . We now claim that  $\underline{x}_i^* < \infty$ . On the contrary, the switching region  $\mathcal{S}_i$  would be empty, and so  $v_i$  would satisfy on  $(0, \infty)$  :

$$rv_i - \mathcal{L}v_i - f_i = 0, \quad \text{on } (0, \infty).$$

Then,  $v_i$  would be on the form :

$$v_i(x) = Ax^{m^+} + Bx^{m^-} + \hat{V}_i(x), \quad x > 0.$$

Since  $0 \leq v_i(0^+) < \infty$  and  $v_i$  is a nonnegative function satisfying a linear growth condition, and using the fact that  $m^- < 0$  and  $m^+ > 1$ , we deduce that  $v_i$  should be equal to  $\hat{V}_i$ . Now, since we have  $v_i \geq v_j - g_{ij} \geq \hat{V}_j - g_{ij}$ , this would imply :

$$\hat{V}_j(x) - \hat{V}_i(x) \leq g_{ij}, \quad \forall x > 0.$$

This contradicts condition (4.41) and so  $0 < \underline{x}_i^* < \infty$ .

By definition of  $\underline{x}_i^*$ , we already know that  $(0, \underline{x}_i^*) \subset \mathcal{C}_i$ . We prove actually the equality, i.e.  $\mathcal{S}_i = [\underline{x}_i^*, \infty)$  or  $v_i(x) = v_j(x) - g_{ij}$  for all  $x \geq \underline{x}_i^*$ . Consider the function

$$w_i(x) = \begin{cases} v_i(x), & 0 < x < \underline{x}_i^* \\ v_j(x) - g_{ij}, & x \geq \underline{x}_i^* \end{cases}$$

We now check that  $w_i$  is a viscosity solution of

$$\min \{rw_i - \mathcal{L}w_i - f_i, w_i - (v_j - g_{ij})\} = 0 \quad \text{on } (0, \infty). \quad (4.45)$$

From Theorem 3.2, the function  $w_i$  is  $C^1$  on  $(0, \infty)$  and in particular at  $\underline{x}_i^*$  where  $w_i'(\underline{x}_i^*) = v_i'(\underline{x}_i^*) = v_j'(\underline{x}_i^*)$ . We also know that  $w_i = v_i$  is  $C^2$  on  $(0, \underline{x}_i^*) \subset \mathcal{C}_i$ , and satisfies  $rw_i - \mathcal{L}w_i - f_i = 0$ ,  $w_i \geq (v_j - g_{ij})$  on  $(0, \underline{x}_i^*)$ . Hence, from Lemma 3.4, we only need to check the viscosity supersolution property of  $w_i$  to :

$$rw_i - \mathcal{L}w_i - f_i \geq 0, \quad \text{on } (\underline{x}_i^*, \infty). \quad (4.46)$$

For this, take some point  $\bar{x} > \underline{x}_i^*$  and some smooth test function  $\varphi$  s.t.  $\bar{x}$  is a local minimum of  $w_i - \varphi$ . Then,  $\bar{x}$  is a local minimum of  $v_j - (\varphi + g_{ij})$ , and by the viscosity solution property of  $v_j$  to its Bellman PDE, we have

$$rv_j(\bar{x}) - \mathcal{L}\varphi(\bar{x}) - f_j(\bar{x}) \geq 0.$$

Now, since  $\underline{x}_i^* \geq \underline{x}_{ij}$ , we have  $\bar{x} > \underline{x}_{ij}$  and so by (4.42),  $\bar{x} \in \hat{Q}_{ij}$ . Hence,

$$(f_j - f_i)(\bar{x}) - rg_{ij} \geq 0.$$

By adding the two previous inequalities, we also obtain the required supersolution inequality :

$$rw_i(\bar{x}) - \mathcal{L}\varphi(\bar{x}) - f_i(\bar{x}) \geq 0,$$

and so (4.45) is proved.

Since  $w_i(0^+) = v_i(0^+)$  and  $w_i$  satisfies a linear growth condition, and from uniqueness of viscosity solution to PDE (4.45), we deduce that  $w_i$  is equal to  $v_i$ . In particular, we have  $v_i(x) = v_j(x) - g_{ij}$  for  $x \geq \underline{x}_i^*$ , which shows that  $\mathcal{S}_i = [\underline{x}_i^*, \infty)$ .

• The case of condition (4.43) is dealt by same arguments as above : we first observe that  $0 < \bar{x}_i^* := \sup \mathcal{S}_i < \infty$  under (4.41), and then show with Lemma 3.4 that the function

$$w_i(x) = \begin{cases} v_j(x) - g_{ij}, & 0 < x < \bar{x}_i^* \\ v_i(x), & x \geq \bar{x}_i^* \end{cases}$$

is a viscosity solution to

$$\min \{rw_i - \mathcal{L}w_i - f_i, w_i - (v_j - g_{ij})\} = 0 \quad \text{on } (0, \infty).$$

Then, under the condition that  $g_{ij} \leq 0$ , we see that  $g_{ji} > 0$  by (2.7), and so  $v_i(0^+) = -g_{ij} = (-g_{ji})_+ - g_{ij} = v_j(0^+) - g_{ij} = w_i(0^+)$ . From uniqueness of viscosity solution to PDE (4.45), we conclude that  $v_i = w_i$ , and so  $\mathcal{S}_i = (0, \bar{x}_i^*]$ .

2) By Lemma 3.3 and (4.40), the condition (4.44) implies  $0 < \underline{x}_{ij} \leq \underline{x}_i^* \leq \bar{x}_i^* \leq \bar{x}_{ij} < \infty$ . We claim that  $\underline{x}_i^* < \bar{x}_i^*$ . Otherwise,  $\mathcal{S}_i = \{\bar{x}_i^*\}$  and  $v_i$  would satisfy  $rv_i - \mathcal{L}v_i - f_i = 0$  on  $(0, \bar{x}_i^*) \cup (\bar{x}_i^*, \infty)$ . By continuity and smooth-fit condition of  $v_i$  at  $\bar{x}_i^*$ , this implies that  $v_i$  satisfies actually

$$rv_i - \mathcal{L}v_i - f_i = 0, \quad x \in (0, \infty),$$

and so is in the form :

$$v_i(x) = Ax^{m^+} + Bx^{m^-} + \hat{V}_i(x), \quad x \in (0, \infty)$$

Since  $0 \leq v_i(0^+) < \infty$  and  $v_i$  is nonnegative function satisfying a linear growth condition, this implies  $A = B = 0$ . Therefore,  $v_i$  is equal to  $\hat{V}_i$ , which also means that  $\mathcal{S}_i = \emptyset$ , a contradiction.

We now prove that  $\mathcal{S}_i = [\underline{x}_i^*, \bar{x}_i^*]$ . Let us consider the function

$$w_i(x) = \begin{cases} v_i(x), & x \in (0, \underline{x}_i^*) \cup (\bar{x}_i^*, \infty) \\ v_j(x) - g_{ij}, & x \in [\underline{x}_i^*, \bar{x}_i^*], \end{cases}$$

which is  $C^1$  on  $(0, \infty)$  and in particular on  $\underline{x}_i^*$  and  $\bar{x}_i^*$  from Theorem 3.2. Hence, by similar arguments as in case 1), using Lemma 3.4, we then show that  $w_i$  is a viscosity solution of

$$\min \{rw_i - \mathcal{L}w_i - f_i, w_i - (v_j - g_{ij})\} = 0. \quad (4.47)$$

Since  $w_i(0^+) = v_i(0^+)$  and  $w_i$  satisfies a linear growth condition, and from uniqueness of viscosity solution to PDE (4.47), we deduce that  $w_i$  is equal to  $v_i$ . In particular, we have  $v_i(x) = v_j(x) - g_{ij}$  for  $x \in [\underline{x}_i^*, \bar{x}_i^*]$ , which shows that  $\mathcal{S}_i = [\underline{x}_i^*, \bar{x}_i^*]$ .

3) Suppose that  $g_{ij} \leq 0$  and  $\hat{Q}_{ij} = (0, \infty)$ . We shall prove that  $\mathcal{S}_i = (0, \infty)$  and  $\mathcal{S}_j = \emptyset$ . To this end, we consider the smooth functions  $w_i = \hat{V}_j - g_{ij}$  and  $w_j = \hat{V}_j$ . Then, recalling the ode satisfied by  $\hat{V}_j$ , and inequality (2.7), we get :

$$rw_j - \mathcal{L}w_j - f_j = 0, \quad w_j - (w_i - g_{ji}) = g_{ij} + g_{ji} \geq 0.$$

Therefore  $w_j$  is a smooth (and so a viscosity) solution to :

$$\min [rw_j - \mathcal{L}w_j - f_j, w_j - (w_i - g_{ji})] = 0 \quad \text{on } (0, \infty).$$

On the other hand, by definition of  $\hat{Q}_{ij}$ , which is supposed equal to  $(0, \infty)$ , we have :

$$\begin{aligned} rw_i(x) - \mathcal{L}w_i(x) - f_i(x) &= r\hat{V}_j(x) - \mathcal{L}\hat{V}_j(x) - f_j(x) + f_j(x) - f_i(x) - rg_{ij} \\ &= f_j(x) - f_i(x) - rg_{ij} \geq 0, \quad \forall x > 0. \end{aligned}$$

Moreover, by construction we have  $w_i = w_j - g_{ij}$ . Therefore  $w_i$  is a smooth (and so a viscosity) solution to :

$$\min [rw_i - \mathcal{L}w_i - f_i, w_i - (w_j - g_{ij})] = 0 \quad \text{on } (0, \infty).$$

Notice also that  $g_{ji} > 0$  by (2.7) and since  $g_{ij} \leq 0$ . Hence,  $w_i(0^+) = -g_{ij} = (-g_{ij})_+ = v_i(0^+)$ ,  $w_j(0^+) = 0 = (-g_{ji})_+ = v_j(0^+)$ . From uniqueness result of Theorem 3.1, we deduce that  $v_i = w_i$ ,  $v_j = w_j$ , which proves that  $\mathcal{S}_i = (0, \infty)$ ,  $\mathcal{S}_j = \emptyset$ .  $\square$

We shall now provide explicit solutions to the switching problem under general assumptions on the running profit functions, which include several interesting cases for applications :

$$\begin{aligned} \text{(HF)} \quad & \text{There exists } \hat{x} \in \mathbb{R}_+ \text{ s.t the function } F := f_2 - f_1 \\ & \text{is decreasing on } (0, \hat{x}), \text{ increasing on } [\hat{x}, \infty), \\ & \text{and } F(\infty) := \lim_{x \rightarrow \infty} F(x) > 0, \quad g_{12} > 0. \end{aligned}$$

Under **(HF)**, there exists some  $\bar{x} \in \mathbb{R}_+$  ( $\bar{x} > \hat{x}$  if  $\hat{x} > 0$  and  $\bar{x} = 0$  if  $\hat{x} = 0$ ) from which  $F$  is positive :  $F(x) > 0$  for  $x > \bar{x}$ . Economically speaking, condition **(HF)** means that the profit in regime 2 is “better” than profit in regime 1 from a certain level  $\bar{x}$ , eventually equal to zero, and the improvement becomes then better and better. Moreover, since profit in regime 2 is better than the one in regime 1, it is natural to assume that the corresponding switching cost  $g_{12}$  from regime 1 to 2 should be positive. However, we shall consider both cases where  $g_{21}$  is positive and nonpositive. Notice that  $F(\hat{x}) < 0$  if  $\hat{x} > 0$ ,  $F(\hat{x}) = 0$  if  $\hat{x} = 0$ , and we do not assume necessarily  $F(\infty) = \infty$ .

**Example 4.1** A typical example of different running profit functions satisfying **(HF)** is given by

$$f_i(x) = k_i x^{\gamma_i}, \quad i = 1, 2, \quad \text{with } 0 < \gamma_1 < \gamma_2 < 1, \quad k_1 \in \mathbb{R}_+, \quad k_2 > 0. \quad (4.48)$$



In this case,  $\hat{x} = \left(\frac{k_1\gamma_1}{k_2\gamma_2}\right)^{\frac{1}{\gamma_2-\gamma_1}}$ , and  $\lim_{x \rightarrow \infty} F(x) = \infty$ .

Another example of profit functions of interest in applications is the case where the profit function in regime 1 is  $f_1 = 0$ , and the other  $f_2$  is increasing. In this case, assumption **(HF)** is satisfied with  $\hat{x} = 0$ .

The next proposition states the form of the switching regions in regimes 1 and 2, depending on the parameter values.

**Proposition 4.1** *Assume that **(HF)** holds.*

- 1) (i) If  $rg_{12} \geq F(\infty)$ , then  $\underline{x}_1^* = \infty$ , i.e.  $\mathcal{S}_1 = \emptyset$ .
- (ii) If  $rg_{12} < F(\infty)$ , then  $\underline{x}_1^* \in (0, \infty)$  and  $\mathcal{S}_1 = [\underline{x}_1^*, \infty)$ .
- 2) (i) If  $rg_{21} \geq -F(\hat{x})$ , then  $\mathcal{S}_2 = \emptyset$ .
- (ii) If  $0 < rg_{21} < -F(\hat{x})$ , then  $0 < \underline{x}_2^* < \bar{x}_2^* < \underline{x}_1^*$ , and  $\mathcal{S}_2 = [\underline{x}_2^*, \bar{x}_2^*]$ .
- (iii) If  $g_{21} \leq 0$  and  $-F(\infty) < rg_{21} < -F(\hat{x})$ , then  $0 = \underline{x}_2^* < \bar{x}_2^* < \underline{x}_1^*$ , and  $\mathcal{S}_2 = (0, \bar{x}_2^*]$ .
- (iv) If  $rg_{21} \leq -F(\infty)$ , then  $\mathcal{S}_2 = (0, \infty)$ .

**Proof.** 1) From Lemma 3.3, we have

$$\hat{Q}_{12} = \{x > 0 : F(x) \geq rg_{12}\}. \quad (4.49)$$

Since  $g_{12} > 0$ , and  $f_i(0) = 0$ , we have  $F(0) = 0 < rg_{12}$ . Under **(HF)**, we then distinguish the two following cases :

- (i) If  $rg_{12} \geq F(\infty)$ , then  $\hat{Q}_{12} = \emptyset$ , and so by Lemma 3.3 and (4.40),  $\mathcal{S}_1 = \emptyset$ .
- (ii) If  $rg_{12} < F(\infty)$ , then there exists  $\hat{x}_{12} \in (0, \infty)$  such that

$$\hat{Q}_{12} = [\hat{x}_{12}, \infty). \quad (4.50)$$

Moreover, since

$$(\hat{V}_2 - \hat{V}_1)(x) = E \left[ \int_0^\infty e^{-rt} F(\hat{X}_t^x) dt \right], \quad \forall x > 0,$$

and  $F$  is lower-bounded, we obtain by Fatou's lemma :

$$\liminf_{x \rightarrow \infty} (\hat{V}_2 - \hat{V}_1)(x) \geq E \left[ \int_0^\infty e^{-rt} F(\infty) dt \right] = \frac{F(\infty)}{r} > g_{12}.$$

Hence, conditions (4.41)-(4.42) with  $i = 1, j = 2$ , are satisfied, and we obtain the first assertion by Lemma 4.1 1).

2) From Lemma 3.3, we have

$$\hat{Q}_{21} = \{x > 0 : -F(x) \geq rg_{21}\}. \quad (4.51)$$

Under **(HF)**, we distinguish the following cases :

- (i1) If  $rg_{21} > -F(\hat{x})$ , then  $\hat{Q}_{21} = \emptyset$ , and so  $\mathcal{S}_2 = \emptyset$ .

(i2) If  $rg_{21} = -F(\hat{x})$ , then either  $\hat{x} = 0$  and so  $\mathcal{S}_2 = \hat{Q}_{21} = \emptyset$ , or  $\hat{x} > 0$ , and so  $\hat{Q}_{21} = \{\hat{x}\}$ ,  $\mathcal{S}_2 \subset \{\hat{x}\}$ . In this last case,  $v_2$  satisfies  $rv_2 - \mathcal{L}v_2 - f_2 = 0$  on  $(0, \hat{x}) \cup (\hat{x}, \infty)$ . By continuity and smooth-fit condition of  $v_2$  at  $\hat{x}$ , this implies that  $v_2$  satisfies actually

$$rv_2 - \mathcal{L}v_2 - f_2 = 0, \quad x \in (0, \infty),$$

and so is in the form :

$$v_2(x) = Ax^{m^+} + Bx^{m^-} + \hat{V}_2(x), \quad x \in (0, \infty)$$

Recalling that  $0 \leq v_2(0^+) < \infty$  and  $v_2$  is a nonnegative function satisfying a linear growth condition, this implies  $A = B = 0$ . Therefore,  $v_2$  is equal to  $\hat{V}_2$ , which also means that  $\mathcal{S}_2 = \emptyset$ .

► If  $rg_{21} < -F(\hat{x})$ , we need to distinguish three subcases depending on  $g_{21}$  :

- If  $g_{21} > 0$ , then there exist  $0 < \underline{x}_{21} < \hat{x} < \bar{x}_{21} < \infty$  such that

$$\hat{Q}_{21} = [\underline{x}_{21}, \bar{x}_{21}]. \quad (4.52)$$

We then conclude with Lemma 4.1 2) for  $i = 2, j = 1$ .

- If  $g_{21} \leq 0$  with  $rg_{21} > -F(\infty)$ , then there exists  $\bar{x}_{21} < \infty$  s.t.

$$\hat{Q}_{21} = (0, \bar{x}_{21}].$$

Moreover, we clearly have  $\sup_{x>0}(\hat{V}_1 - \hat{V}_2)(x) > (\hat{V}_1 - \hat{V}_2)(0) = 0 \geq g_{21}$ . Hence, conditions (4.41) and (4.43) with  $i = 2, j = 1$  are satisfied, and we deduce from Lemma 4.1 1) that  $\mathcal{S}_2 = (0, \bar{x}_2^*]$  with  $0 < \bar{x}_2^* < \infty$ .

- If  $rg_{21} \leq -F(\infty)$ , then  $\hat{Q}_{21} = (0, \infty)$ , and we deduce from Lemma 4.1 3) for  $i = 2, j = 1$ , that  $\mathcal{S}_2 = (0, \infty)$ .

Finally, in the two above subcases when  $\mathcal{S}_2 = [\underline{x}_2^*, \bar{x}_2^*]$  or  $(0, \bar{x}_2^*]$ , we notice that  $\bar{x}_2^* < \underline{x}_1^*$  since  $\mathcal{S}_2 \subset \mathcal{C}_1 = (0, \infty) \setminus \mathcal{S}_1$ , which is equal, from 1), either to  $(0, \infty)$  when  $\underline{x}_1^* = \infty$  or to  $(0, \underline{x}_1^*)$ .  $\square$

**Remark 4.2** In our viscosity solutions approach, the structure of the switching regions is derived from the smooth fit property of the value functions, uniqueness result for viscosity solutions and Lemma 3.3. This contrasts with the classical verification approach where the structure of switching regions should be guessed ad-hoc and checked a posteriori by a verification argument.

### Economic interpretation.

The previous proposition shows that, under **(HF)**, the switching region in regime 1 has two forms depending on the size of its corresponding positive switching cost : If  $g_{12}$  is larger than the “maximum net” profit  $F(\infty)$  that one can expect by changing of regime (case 1) (i), which may occur only if  $F(\infty) < \infty$ ), then one has no interest to switch of regime, and one always stay in regime 1, i.e.  $\mathcal{C}_1 = (0, \infty)$ . However, if this switching cost is smaller

than  $F(\infty)$  (case 1) (ii), which always holds true when  $F(\infty) = \infty$ ), then there is some positive threshold from which it is optimal to change of regime.

The structure of the switching region in regime 2 exhibits several different forms depending on the sign and size of its corresponding switching cost  $g_{21}$  with respect to the values  $-F(\infty) < 0$  and  $-F(\hat{x}) \geq 0$ . If  $g_{21}$  is nonnegative larger than  $-F(\hat{x})$  (case 2) (i), then one has no interest to switch of regime, and one always stay in regime 2, i.e.  $\mathcal{C}_2 = (0, \infty)$ . If  $g_{21}$  is positive, but not too large (case 2) (ii)), then there exists some bounded closed interval, which is not a neighborhood of zero, where it is optimal to change of regime. Finally, when the switching cost  $g_{21}$  is negative, it is optimal to switch to regime 1 at least for small values of the state. Actually, if the negative cost  $g_{21}$  is larger than  $-F(\infty)$  (case 2) (iii), which always holds true for negative cost when  $F(\infty) = \infty$ ), then the switching region is a bounded neighborhood of 0. Moreover, if the cost is negative large enough (case 2) (iv), which may occur only if  $F(\infty) < \infty$ ), then it is optimal to change of regime for every values of the state.

By combining the different cases for regimes 1 and 2, and observing that case 2) (iv) is not compatible with case 1) (ii) by (2.7), we then have a priori seven different forms for both switching regions. These forms reduce actually to three when  $F(\infty) = \infty$ . The various structures of the switching regions are depicted in Figure II.

Finally, we complete results of Proposition 4.1 by providing the explicit solutions for the value functions and the corresponding boundaries of the switching regions in the seven different cases depending on the model parameter values.

**Theorem 4.2** *Assume that (HF) holds.*

1) *If  $rg_{12} < F(\infty)$  and  $rg_{21} \geq -F(\hat{x})$ , then*

$$\begin{aligned} v_1(x) &= \begin{cases} Ax^{m^+} + \hat{V}_1(x), & x < \underline{x}_1^* \\ v_2(x) - g_{12}, & x \geq \underline{x}_1^* \end{cases} \\ v_2(x) &= \hat{V}_2(x), \end{aligned}$$

where the constants  $A$  and  $\underline{x}_1^*$  are determined by the continuity and smooth-fit conditions of  $v_1$  at  $\underline{x}_1^*$  :

$$\begin{aligned} A(\underline{x}_1^*)^{m^+} + \hat{V}_1(\underline{x}_1^*) &= \hat{V}_2(\underline{x}_1^*) - g_{12} \\ Am^+(\underline{x}_1^*)^{m^+-1} + \hat{V}_1'(\underline{x}_1^*) &= \hat{V}_2'(\underline{x}_1^*). \end{aligned}$$

*In regime 1, it is optimal to switch to regime 2 whenever the state process  $X$  exceeds the threshold  $\underline{x}_1^*$ , while when we are in regime 2, it is optimal never to switch.*

2) *If  $rg_{12} < F(\infty)$  and  $0 < rg_{21} < -F(\hat{x})$ , then*

$$v_1(x) = \begin{cases} A_1x^{m^+} + \hat{V}_1(x), & x < \underline{x}_1^* \\ v_2(x) - g_{12}, & x \geq \underline{x}_1^* \end{cases} \quad (4.53)$$

$$v_2(x) = \begin{cases} A_2x^{m^+} + \hat{V}_2(x), & x < \underline{x}_2^* \\ v_1(x) - g_{21}, & \underline{x}_2^* \leq x \leq \bar{x}_2^* \\ B_2x^{m^-} + \hat{V}_2(x), & x > \bar{x}_2^*, \end{cases} \quad (4.54)$$

### Figure II




Figure II.1:  $rg_{12} < F(\infty)$ ,  $rg_{21} \geq -F(\hat{x})$




Figure II.2:  $rg_{12} < F(\infty)$ ,  $0 < rg_{21} < -F(\hat{x})$




Figure II.3:  $rg_{12} < F(\infty)$ ,  $g_{21} \leq 0$ ,  $-F(\infty) < rg_{21} < -F(\hat{x})$




Figure II.4:  $rg_{12} \geq F(\infty)$ ,  $rg_{21} > -F(\hat{x})$




Figure II.5:  $rg_{12} \geq F(\infty)$ ,  $0 < rg_{21} < -F(\hat{x})$




Figure II.6:  $rg_{12} \geq F(\infty)$ ,  $g_{21} \leq 0$ ,  $F(\infty) < rg_{21} < -F(\hat{x})$




Figure II.7:  $rg_{12} \geq F(\infty)$ ,  $g_{21} \leq -F(\infty)$

where the constants  $A_1$  and  $\underline{x}_1^*$  are determined by the continuity and smooth-fit conditions of  $v_1$  at  $\underline{x}_1^*$ , and the constants  $A_2$ ,  $B_2$ ,  $\underline{x}_2^*$ ,  $\bar{x}_2^*$  are determined by the continuity and smooth-fit conditions of  $v_2$  at  $\underline{x}_2^*$  and  $\bar{x}_2^*$  :

$$A_1(\underline{x}_1^*)^{m^+} + \hat{V}_1(\underline{x}_1^*) = B_2(\underline{x}_1^*)^{m^-} + \hat{V}_2(\underline{x}_1^*) - g_{12} \quad (4.55)$$

$$A_1 m^+ (\underline{x}_1^*)^{m^+-1} + \hat{V}_1'(\underline{x}_1^*) = B_2 m^- (\underline{x}_1^*)^{m^- -1} + \hat{V}_2'(\underline{x}_1^*) \quad (4.56)$$

$$A_2(\underline{x}_2^*)^{m^+} + \hat{V}_2(\underline{x}_2^*) = A_1(\underline{x}_2^*)^{m^+} + \hat{V}_1(\underline{x}_2^*) - g_{21} \quad (4.57)$$

$$A_2 m^+ (\underline{x}_2^*)^{m^+-1} + \hat{V}_2'(\underline{x}_2^*) = A_1 m^+ (\underline{x}_2^*)^{m^+-1} + \hat{V}_1'(\underline{x}_2^*) \quad (4.58)$$

$$A_1(\bar{x}_2^*)^{m^+} + \hat{V}_1(\bar{x}_2^*) - g_{21} = B_2(\bar{x}_2^*)^{m^-} + \hat{V}_2(\bar{x}_2^*) \quad (4.59)$$

$$A_1 m^+ (\bar{x}_2^*)^{m^+-1} + \hat{V}_1'(\bar{x}_2^*) = B_2 m^- (\bar{x}_2^*)^{m^- -1} + \hat{V}_2'(\bar{x}_2^*). \quad (4.60)$$

In regime 1, it is optimal to switch to regime 2 whenever the state process  $X$  exceeds the threshold  $\underline{x}_1^*$ , while when we are in regime 2, it is optimal to switch to regime 1 whenever the state process lies between  $\underline{x}_2^*$  and  $\bar{x}_2^*$ .

**3)** If  $rg_{12} < F(\infty)$  and  $g_{21} \leq 0$  with  $-F(\infty) < rg_{21} < -F(\hat{x})$ , then

$$v_1(x) = \begin{cases} Ax^{m^+} + \hat{V}_1(x), & x < \underline{x}_1^* \\ v_2(x) - g_{12}, & x \geq \underline{x}_1^* \end{cases}$$

$$v_2(x) = \begin{cases} v_1(x) - g_{21}, & 0 < x \leq \bar{x}_2^* \\ Bx^{m^-} + \hat{V}_2(x), & x > \bar{x}_2^*, \end{cases}$$

where the constants  $A$  and  $\underline{x}_1^*$  are determined by the continuity and smooth-fit conditions of  $v_1$  at  $\underline{x}_1^*$ , and the constants  $B$  and  $\bar{x}_2^*$  are determined by the continuity and smooth-fit conditions of  $v_2$  at  $\bar{x}_2^*$  :

$$A(\underline{x}_1^*)^{m^+} + \hat{V}_1(\underline{x}_1^*) = B(\underline{x}_1^*)^{m^-} + \hat{V}_2(\underline{x}_1^*) - g_{12}$$

$$Am^+ (\underline{x}_1^*)^{m^+-1} + \hat{V}_1'(\underline{x}_1^*) = Bm^- (\underline{x}_1^*)^{m^- -1} + \hat{V}_2'(\underline{x}_1^*)$$

$$A(\bar{x}_2^*)^{m^+} + \hat{V}_1(\bar{x}_2^*) - g_{21} = B(\bar{x}_2^*)^{m^-} + \hat{V}_2(\bar{x}_2^*)$$

$$Am^+ (\bar{x}_2^*)^{m^+-1} + \hat{V}_1'(\bar{x}_2^*) = Bm^- (\bar{x}_2^*)^{m^- -1} + \hat{V}_2'(\bar{x}_2^*).$$

**4)** If  $rg_{12} \geq F(\infty)$  and  $rg_{21} \geq -F(\hat{x})$ , then  $v_1 = \hat{V}_1$ ,  $v_2 = \hat{V}_2$ . It is optimal never to switch in both regimes 1 and 2.

**5)** If  $rg_{12} \geq F(\infty)$  and  $0 < rg_{21} < -F(\hat{x})$ , then

$$v_1(x) = \hat{V}_1(x)$$

$$v_2(x) = \begin{cases} Ax^{m^+} + \hat{V}_2(x), & x < \underline{x}_2^* \\ v_1(x) - g_{21}, & \underline{x}_2^* \leq x \leq \bar{x}_2^* \\ Bx^{m^-} + \hat{V}_2(x), & x > \bar{x}_2^*, \end{cases}$$

where the constants  $A$ ,  $B$ ,  $\underline{x}_2^*$ ,  $\bar{x}_2^*$  are determined by the continuity and smooth-fit conditions of  $v_2$  at  $\underline{x}_2^*$  and  $\bar{x}_2^*$  :

$$A(\underline{x}_2^*)^{m^+} + \hat{V}_2(\underline{x}_2^*) = \hat{V}_1(\underline{x}_2^*) - g_{21}$$

$$\begin{aligned}
Am^+(\underline{x}_2^*)^{m^+-1} + \hat{V}_2'(\underline{x}_2^*) &= \hat{V}_1'(\underline{x}_2^*) \\
\hat{V}_1(\bar{x}_2^*) - g_{21} &= B(\bar{x}_2^*)^{m^-} + \hat{V}_2(\bar{x}_2^*) \\
\hat{V}_1'(\bar{x}_2^*) &= Bm^-(\bar{x}_2^*)^{m^- - 1} + \hat{V}_2'(\bar{x}_2^*).
\end{aligned}$$

In regime 1, it is optimal never to switch, while when we are in regime 2, it is optimal to switch to regime 1 whenever the state process lies between  $\underline{x}_2^*$  and  $\bar{x}_2^*$ .

**6)** If  $rg_{12} \geq F(\infty)$  and  $g_{21} \leq 0$  with  $-F(\infty) < rg_{21} < -F(\hat{x})$ , then

$$\begin{aligned}
v_1(x) &= \hat{V}_1(x) \\
v_2(x) &= \begin{cases} v_1(x) - g_{21}, & 0 < x \leq \bar{x}_2^* \\ Bx^{m^-} + \hat{V}_2(x), & x > \bar{x}_2^*, \end{cases}
\end{aligned}$$

where the constants  $B$  and  $\bar{x}_2^*$  are determined by the continuity and smooth-fit conditions of  $v_2$  at  $\bar{x}_2^*$  :

$$\begin{aligned}
\hat{V}_1(\bar{x}_2^*) - g_{21} &= B(\bar{x}_2^*)^{m^-} + \hat{V}_2(\bar{x}_2^*) \\
\hat{V}_1'(\bar{x}_2^*) &= Bm^-(\bar{x}_2^*)^{m^- - 1} + \hat{V}_2'(\bar{x}_2^*).
\end{aligned}$$

In regime 1, it is optimal never to switch, while when we are in regime 2, it is optimal to switch to regime 1 whenever the state process lies below  $\bar{x}_2^*$ .

**7)** If  $rg_{12} \geq F(\infty)$  and  $rg_{21} \leq -F(\infty)$ , then  $v_1 = \hat{V}_1$  and  $v_2 = v_1 - g_{12}$ . In regime 1, it is optimal never to switch, while when we are in regime 2, it is always optimal to switch to regime 1.

**Proof.** We prove the result only for the case **2)** since the other cases are dealt similarly and are even simpler. This case **2)** corresponds to the combination of cases 1) (ii) and 2) (ii) in Proposition 4.1. We then have  $\mathcal{S}_1 = [\underline{x}_1^*, \infty)$ , which means that  $v_1 = v_2 - g_{12}$  on  $[\underline{x}_1^*, \infty)$  and  $v_1$  is solution to  $rv_1 - \mathcal{L}v_1 - f_1 = 0$  on  $(0, \underline{x}_1^*)$ . Since  $0 \leq v_1(0^+) < \infty$ ,  $v_1$  should have the form expressed in (4.53). Moreover,  $\mathcal{S}_2 = [\underline{x}_2^*, \bar{x}_2^*]$ , which means that  $v_2 = v_1 - g_{21}$  on  $[\underline{x}_2^*, \bar{x}_2^*]$ , and  $v_2$  satisfies on  $\mathcal{C}_2 = (0, \underline{x}_2^*) \cup (\bar{x}_2^*, \infty) : rv_2 - \mathcal{L}v_2 - f_2 = 0$ . Recalling again that  $0 \leq v_2(0^+) < \infty$  and  $v_2$  satisfies a linear growth condition, we deduce that  $v_2$  has the form expressed in (4.54). Finally, the constants  $A_1, \underline{x}_1^*$ , which characterize completely  $v_1$ , and the constants  $A_2, B_2, \underline{x}_2^*, \bar{x}_2^*$ , which characterize completely  $v_2$ , are determined by the six relations (4.55)-(4.56)-(4.57)-(4.58)-(4.59)-(4.60) resulting from the continuity and smooth-fit conditions of  $v_1$  at  $\underline{x}_1^*$  and  $v_2$  at  $\underline{x}_2^*$  and  $\bar{x}_2^*$ , and recalling that  $\bar{x}_2^* < \underline{x}_1^*$ .  $\square$

**Remark 4.3** In the classical approach, for instance in the case **2)**, we construct a priori a candidate solution in the form (4.53)-(4.54), and we have to check the existence of a sextuple solution to (4.55)-(4.56)-(4.57)-(4.58)-(4.59)-(4.60), which may be somewhat tedious! Here, by the viscosity solutions approach, and since we already state the smooth-fit  $C^1$  property of the value functions, we know a priori the existence of a sextuple solution to (4.55)-(4.56)-(4.57)-(4.58)-(4.59)-(4.60).

## Appendix: proof of comparison principle

In this section, we prove a comparison principle for the system of variational inequalities (3.8). The comparison result in [10] for switching problems in finite horizon does not apply in our context. Inspired by [8], we first produce some suitable perturbation of viscosity supersolution to deal with the switching obstacle, and then follow the general viscosity solution technique, see e.g. [3].

**Theorem 4.3** *Suppose  $u_i$ ,  $i \in \mathbb{I}_d$ , are continuous viscosity subsolutions to the system of variational inequalities (3.8) on  $(0, \infty)$ , and  $w_i$ ,  $i \in \mathbb{I}_d$ , are continuous viscosity supersolutions to the system of variational inequalities (3.8) on  $(0, \infty)$ , satisfying the boundary conditions  $u_i(0^+) \leq w_i(0^+)$ ,  $i \in \mathbb{I}_d$ , and the linear growth condition :*

$$|u_i(x)| + |w_i(x)| \leq C_1 + C_2x, \quad \forall x \in (0, \infty), \quad i \in \mathbb{I}_d, \quad (\text{A.1})$$

for some positive constants  $C_1$  and  $C_2$ . Then,

$$u_i \leq w_i, \quad \text{on } (0, \infty), \quad \forall i \in \mathbb{I}_d.$$

**Proof.** *Step 1.* Let  $u_i$  and  $w_i$ ,  $i \in \mathbb{I}_d$ , as in Theorem 4.3. We first construct strict supersolutions to the system (3.8) with suitable perturbations of  $w_i$ ,  $i \in \mathbb{I}_d$ . We set

$$h(x) = C'_1 + C'_2x^p, \quad x > 0,$$

where  $C'_1, C'_2 > 0$  and  $p > 1$  are positive constants to be determined later. We then define for all  $\lambda \in (0, 1)$ , the continuous functions on  $(0, \infty)$  by :

$$w_i^\lambda = (1 - \lambda)w_i + \lambda(h + \alpha_i), \quad i \in \mathbb{I}_d,$$

where  $\alpha_i = \min_{j \neq i} g_{ji}$ . We then see that for all  $\lambda \in (0, 1)$ ,  $i \in \mathbb{I}_d$  :

$$\begin{aligned} w_i^\lambda - \max_{j \neq i} (w_j^\lambda - g_{ij}) &= \lambda\alpha_i + (1 - \lambda)w_i - \max_{j \neq i} [(1 - \lambda)(w_j - g_{ij}) + \lambda\alpha_j - \lambda g_{ij}] \\ &\geq (1 - \lambda)[w_i - \max_{j \neq i} (w_j - g_{ij})] + \lambda \left( \alpha_i + \min_{j \neq i} (g_{ij} - \alpha_j) \right) \\ &\geq \lambda \min_{i \in \mathbb{I}_d} \left( \alpha_i + \min_{j \neq i} (g_{ij} - \alpha_j) \right) \\ &\geq \lambda \underline{\nu} \end{aligned} \quad (\text{A.2})$$

where  $\underline{\nu} := \min_{i \in \mathbb{I}_d} \left[ \alpha_i + \min_{j \neq i} (g_{ij} - \alpha_j) \right]$  is a constant independent of  $i$ . We now check that  $\underline{\nu} > 0$ , i.e.  $\nu_i := \alpha_i + \min_{j \neq i} (g_{ij} - \alpha_j) > 0$ ,  $\forall i \in \mathbb{I}_d$ . Indeed, fix  $i \in \mathbb{I}_d$ , and let  $k \in \mathbb{I}_d$  such that  $\min_{j \neq i} (g_{ij} - \alpha_j) = g_{ik} - \alpha_k$  and set  $\underline{i}$  such that  $\alpha_i = \min_{j \neq i} g_{ji} = g_{\underline{i}i}$ . We then have

$$\nu_i = g_{\underline{i}i} + g_{ik} - \min_{j \neq k} g_{jk} > g_{\underline{i}k} - \min_{j \neq k} g_{jk} \geq 0,$$

by (2.6) and thus  $\underline{\nu} > 0$ .

By definition of the Fenchel Legendre in (2.5), and by setting  $\tilde{f}(1) = \max_{i \in \mathbb{I}_d} \tilde{f}_i(1)$ , we have for all  $i \in \mathbb{I}_d$ ,

$$f_i(x) \leq \tilde{f}(1) + x \leq \tilde{f}(1) + 1 + x^p, \quad \forall x > 0.$$

Moreover, recalling that  $r > b := \max_i b_i$ , we can choose  $p > 1$  s.t.

$$\rho := r - pb - \frac{1}{2}\sigma^2 p(p-1) > 0,$$

where we set  $\sigma := \max_i \sigma_i > 0$ . By choosing

$$C'_1 \geq \frac{2 + \tilde{f}(1)}{r} - \min_i \alpha_i, \quad C'_2 \geq \frac{1}{\rho},$$

we then have for all  $i \in \mathbb{I}_d$ ,

$$\begin{aligned} rh(x) - \mathcal{L}_i h(x) - f_i(x) &= rC'_1 + C'_2 x^p [r - pb_i - \frac{1}{2}\sigma_i^2 p(p-1)] - f_i(x) \\ &\geq rC'_1 + \rho C'_2 x^p - f_i(x) \\ &\geq 1, \quad \forall x > 0. \end{aligned} \tag{A.3}$$

From (A.2) and (A.3), we then deduce that for all  $i \in \mathbb{I}_d$ ,  $\lambda \in (0, 1)$ ,  $w_i^\lambda$  is a supersolution to

$$\min \left\{ rw_i^\lambda - \mathcal{L}_i w_i^\lambda - f_i, w_i^\lambda - \max_{j \neq i} (w_j^\lambda - g_{ij}) \right\} \geq \lambda \delta, \quad \text{on } (0, \infty), \tag{A.4}$$

where  $\delta = \underline{\nu} \wedge 1 > 0$ .

Step 2. In order to prove the comparison principle, it suffices to show that for all  $\lambda \in (0, 1)$  :

$$\max_{j \in \mathbb{I}_d} \sup_{(0, +\infty)} (u_j - w_j^\lambda) \leq 0$$

since the required result is obtained by letting  $\lambda$  to 0. We argue by contradiction and suppose that there exists some  $\lambda \in (0, 1)$  and  $i \in \mathbb{I}_d$  s.t.

$$\theta := \max_{j \in \mathbb{I}_d} \sup_{(0, +\infty)} (u_j - w_j^\lambda) = \sup_{(0, +\infty)} (u_i - w_i^\lambda) > 0. \tag{A.5}$$

From the linear growth condition (A.1), and since  $p > 1$ , we observe that  $u_i(x) - w_i^\lambda(x)$  goes to  $-\infty$  when  $x$  goes to infinity. By choosing also  $C'_1 \geq \max_i w_i(0^+)$ , we then have  $u_i(0^+) - w_i^\lambda(0^+) = u_i(0^+) - w_i(0^+) + \lambda(w_i(0^+) - C'_1) \leq 0$ . Hence, by continuity of the functions  $u_i$  and  $w_i^\lambda$ , there exists  $x_0 \in (0, \infty)$  s.t.

$$\theta = u_i(x_0) - w_i^\lambda(x_0).$$

For any  $\varepsilon > 0$ , we consider the functions

$$\begin{aligned} \Phi_\varepsilon(x, y) &= u_i(x) - w_i^\lambda(y) - \phi_\varepsilon(x, y), \\ \phi_\varepsilon(x, y) &= \frac{1}{4}|x - x_0|^4 + \frac{1}{2\varepsilon}|x - y|^2, \end{aligned}$$



for all  $x, y \in (0, \infty)$ . By standard arguments in comparison principle, the function  $\Phi_\varepsilon$  attains a maximum in  $(x_\varepsilon, y_\varepsilon) \in (0, \infty)^2$ , which converges (up to a subsequence) to  $(x_0, x_0)$  when  $\varepsilon$  goes to zero. Moreover,

$$\lim_{\varepsilon \rightarrow 0} \frac{|x_\varepsilon - y_\varepsilon|^2}{\varepsilon} = 0. \quad (\text{A.6})$$

Applying Theorem 3.2 in [3], we get the existence of  $M_\varepsilon, N_\varepsilon \in \mathbb{R}$  such that:

$$\begin{aligned} (p_\varepsilon, M_\varepsilon) &\in J^{2,+}u_i(x_\varepsilon), \\ (q_\varepsilon, N_\varepsilon) &\in J^{2,-}w_i^\lambda(y_\varepsilon) \end{aligned}$$

where

$$\begin{aligned} p_\varepsilon &= D_x \phi_\varepsilon(x_\varepsilon, y_\varepsilon) = \frac{1}{\varepsilon}(x_\varepsilon - y_\varepsilon) + (x_\varepsilon - x_0)^3 \\ q_\varepsilon &= -D_y \phi_\varepsilon(x_\varepsilon, y_\varepsilon) = \frac{1}{\varepsilon}(x_\varepsilon - y_\varepsilon) \end{aligned}$$

and

$$\begin{pmatrix} M_\varepsilon & 0 \\ 0 & -N_\varepsilon \end{pmatrix} \leq D^2 \phi_\varepsilon(x_\varepsilon, y_\varepsilon) + \varepsilon (D^2 \phi_\varepsilon(x_\varepsilon, y_\varepsilon))^2 \quad (\text{A.7})$$

with

$$D^2 \phi_\varepsilon(x_\varepsilon, y_\varepsilon) = \begin{pmatrix} 3(x_\varepsilon - x_0)^2 + \frac{1}{\varepsilon} & -\frac{1}{\varepsilon} \\ -\frac{1}{\varepsilon} & \frac{1}{\varepsilon} \end{pmatrix},$$

By writing the viscosity subsolution property (3.9) of  $u_i$  and the viscosity strict supersolution property (A.4) of  $w_i^\lambda$ , we have the following inequalities:

$$\begin{aligned} \min \left\{ r u_i(x_\varepsilon) - \left( \frac{1}{\varepsilon}(x_\varepsilon - y_\varepsilon) + (x_\varepsilon - x_0)^3 \right) b_i x_\varepsilon - \frac{1}{2} \sigma_i^2 x_\varepsilon^2 M_\varepsilon - f_i(x_\varepsilon), \right. \\ \left. u_i(x_\varepsilon) - \max_{j \neq i} (u_j - g_{ij})(x_\varepsilon) \right\} \leq 0 \quad (\text{A.8}) \end{aligned}$$

$$\begin{aligned} \min \left\{ r w_i^\lambda(y_\varepsilon) - \frac{1}{\varepsilon}(x_\varepsilon - y_\varepsilon) b_i y_\varepsilon - \frac{1}{2} \sigma_i^2 y_\varepsilon^2 N_\varepsilon - f_i(y_\varepsilon), \right. \\ \left. w_i^\lambda(y_\varepsilon) - \max_{j \neq i} (w_j^\lambda - g_{ij})(y_\varepsilon) \right\} \geq \lambda \delta \quad (\text{A.9}) \end{aligned}$$

We then distinguish the following two cases :

(1)  $u_i(x_\varepsilon) - \max_{j \neq i} (u_j - g_{ij})(x_\varepsilon) \leq 0$  in (A.8).

By sending  $\varepsilon \rightarrow 0$ , this implies

$$u_i(x_0) - \max_{j \neq i} (u_j - g_{ij})(x_0) \leq 0. \quad (\text{A.10})$$

On the other hand, we have by (A.9) :

$$w_i^\lambda(y_\varepsilon) - \max_{j \neq i} (w_j^\lambda - g_{ij})(y_\varepsilon) \geq \lambda \delta,$$

so that by sending  $\varepsilon$  to zero :

$$w_i^\lambda(x_0) - \max_{j \neq i}(w_j^\lambda - g_{ij})(x_0) \geq \lambda\delta. \quad (\text{A.11})$$

Combining (A.10) and (A.11), we obtain :

$$\begin{aligned} \theta = u_i(x_0) - w_i^\lambda(x_0) &\leq -\lambda\delta + \max_{j \neq i}(u_j - g_{ij})(x_0) - \max_{j \neq i}(w_j^\lambda - g_{ij})(x_0) \\ &\leq -\lambda\delta + \max_{j \neq i}(u_j - w_j^\lambda)(x_0) \\ &\leq -\lambda\delta + \theta, \end{aligned}$$

which is a contradiction.

(2)  $ru_i(x_\varepsilon) - \left(\frac{1}{\varepsilon}(x_\varepsilon - y_\varepsilon) + (x_\varepsilon - x_0)^3\right) b_i x_\varepsilon - \frac{1}{2}\sigma_i^2 x_\varepsilon^2 M_\varepsilon - f_i(x_\varepsilon) \leq 0$  in (A.8).

Since by (A.9), we also have :

$$rw_i^\lambda(y_\varepsilon) - \frac{1}{\varepsilon}(x_\varepsilon - y_\varepsilon)b_i y_\varepsilon - \frac{1}{2}\sigma_i^2 y_\varepsilon^2 N_\varepsilon - f_i(y_\varepsilon) \geq \lambda\delta,$$

this yields by combining the above two inequalities :

$$\begin{aligned} ru_i(x_\varepsilon) - rw_i^\lambda(y_\varepsilon) - \frac{1}{\varepsilon}b_i(x_\varepsilon - y_\varepsilon)^2 - (x_\varepsilon - x_0)^3 b_i x_\varepsilon \\ + \frac{1}{2}\sigma_i^2 y_\varepsilon^2 N_\varepsilon - \frac{1}{2}\sigma_i^2 x_\varepsilon^2 M_\varepsilon + f_i(y_\varepsilon) - f_i(x_\varepsilon) \leq -\lambda\delta. \end{aligned} \quad (\text{A.12})$$

Now, from (A.7), we have :

$$\frac{1}{2}\sigma_i^2 x_\varepsilon^2 M_\varepsilon - \frac{1}{2}\sigma_i^2 y_\varepsilon^2 N_\varepsilon \leq \frac{3}{2\varepsilon}\sigma_i^2 (x_\varepsilon - y_\varepsilon)^2 + \frac{3}{2}\sigma_i^2 x_\varepsilon^2 (x_\varepsilon - x_0)^2 (3\varepsilon(x_\varepsilon - x_0)^2 + 2),$$

so that by plugging into (A.12) :

$$\begin{aligned} r \left( u_i(x_\varepsilon) - w_i^\lambda(y_\varepsilon) \right) &\leq \frac{1}{\varepsilon}b_i(x_\varepsilon - y_\varepsilon)^2 + (x_\varepsilon - x_0)^3 b_i x_\varepsilon + \frac{3}{2\varepsilon}\sigma_i^2 (x_\varepsilon - y_\varepsilon)^2 \\ &\quad + \frac{3}{2}\sigma_i^2 x_\varepsilon^2 (x_\varepsilon - x_0)^2 (3\varepsilon(x_\varepsilon - x_0)^2 + 2) + f_i(y_\varepsilon) - f_i(x_\varepsilon) - \lambda\delta \end{aligned}$$

By sending  $\varepsilon$  to zero, and using (A.6), continuity of  $f_i$ , we obtain the required contradiction:  $r\theta \leq -\lambda\delta < 0$ . This ends the proof of Theorem 4.3.  $\square$

## References

- [1] Bensoussan A. and J.L. Lions (1982) : Contrôle impulsionnel et inéquations variationnelles, Dunod.
- [2] Brekke K. and B. Oksendal (1994) : “Optimal switching in an economic activity under uncertainty”, *SIAM J. Cont. Optim.*, 32, 1021-1036.
- [3] Crandall M., Ishii H. and P.L. Lions (1992) : “User’s guide to viscosity solutions of second order partial differential equations”, *Bull. Amer. Math. Soc.*, 27, 1-67.

- [4] Dixit A. and R. Pindyck (1994) : Investment under uncertainty, Princeton University Press.
- [5] Duckworth K. and M. Zervos (2001) : “A model for investment decisions with switching costs”, *Annals of Applied Probability*, 11, 239-250.
- [6] Fleming W. and M. Soner (1993) : Controlled Markov processes and viscosity solutions, Springer Verlag.
- [7] Guo X. (2001) : “An explicit solution to an optimal stopping problem with regime switching”, *Journal of Applied Probability*, 38, 464-481.
- [8] Ishii H. and P.L. Lions (1990) : “Viscosity solutions of fully nonlinear second order elliptic partial differential equations”, *Journal of Differential equations*, 83, 26-78.
- [9] Pham H. (2005) : “On the smooth-fit property for one-dimensional optimal switching problem”, to appear in *Séminaire de Probabilités*, Vol. XL.
- [10] Tang S. and J. Yong (1993) : “Finite horizon stochastic optimal switching and impulse controls with a viscosity solution approach”, *Stoch. and Stoch. Reports*, 45, 145-176.