

## Review Article

# Exponential Polynomials, Stirling Numbers, and Evaluation of Some Gamma Integrals

**Khristo N. Boyadzhiev**

*Department of Mathematics, Ohio Northern University, Ada, OH 45810, USA*

Correspondence should be addressed to Khristo N. Boyadzhiev, k-boyadzhiev@onu.edu

Received 15 May 2009; Accepted 4 August 2009

Recommended by Lance Littlejohn

This article is a short elementary review of the exponential polynomials (also called single-variable Bell polynomials) from the point of view of analysis. Some new properties are included, and several analysis-related applications are mentioned. At the end of the paper one application is described in details—certain Fourier integrals involving  $\Gamma(a+it)$  and  $\Gamma(a+it)\Gamma(b-it)$  are evaluated in terms of Stirling numbers.

Copyright © 2009 Khristo N. Boyadzhiev. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

We review the exponential polynomials  $\phi_n(x)$  and present a list of properties for easy reference. Exponential polynomials in analysis appear, for instance, in the rule for computing derivatives like  $(d/dt)^n e^{ae^t}$  and the related Mellin derivatives:

$$\left(x \frac{d}{dx}\right)^n f(x), \quad \left(\frac{d}{dx} x\right)^n f(x). \quad (1.1)$$

Namely, we have

$$\left(\frac{d}{dt}\right)^n e^{ae^t} = \phi_n(ae^t)e^{ae^t}, \quad (1.2)$$

or, after the substitution  $x = e^t$ ,

$$\left(x \frac{d}{dx}\right)^n e^{ax} = \phi_n(ax)e^{ax}. \quad (1.3)$$

We also include in this review two properties relating exponential polynomials to Bernoulli numbers,  $B_k$ . One is the semiorthogonality

$$\int_{-\infty}^0 \phi_n(x)\phi_m(x)e^{2x} \frac{dx}{x} = (-1)^n \frac{2^{n+m} - 1}{n+m} B_{n+m}, \quad (1.4)$$

where the right-hand side is zero if  $n+m$  is odd. The other property is (2.25).

At the end we give one application. Using exponential polynomials we evaluate the integrals

$$\int_{\mathbb{R}} e^{-it\lambda} t^n \Gamma(a+it) dt, \quad (1.5)$$

$$\int_{\mathbb{R}} e^{-it\lambda} t^n \Gamma(a+it) \Gamma(b-it) dt$$

for  $n = 0, 1, \dots$ , in terms of Stirling numbers.

## 2. Exponential Polynomials

The evaluation of the series

$$S_n = \sum_{k=0}^{\infty} \frac{k^n}{k!}, \quad n = 0, 1, 2, \dots \quad (2.1)$$

has a long and interesting history. Clearly,  $S_0 = e$ , with the agreement that  $0^0 = 1$ . Several reference books (e.g., [1]) provide the following numbers:

$$\begin{aligned} S_1 = e, & \quad S_2 = 2e, & \quad S_3 = 5e, & \quad S_4 = 15e, & \quad S_5 = 52e, \\ S_6 = 203e, & \quad S_7 = 877e, & \quad S_8 = 4140e. \end{aligned} \quad (2.2)$$

As noted by Gould in [2, page 93], the problem of evaluating  $S_n$  appeared in the Russian journal *Matematicheskii Sbornik*, 3 (1868), page 62, with solution *ibid*, 4 (1868-9), page 39. Evaluations are presented also in two papers by Dobinski and Ligowski. In 1877 Dobinski [3] evaluated the first eight series  $S_1, \dots, S_8$  by regrouping

$$\begin{aligned} S_1 &= \sum_1^{\infty} \frac{k}{k!} = 1 + \frac{2}{2!} + \frac{3}{3!} + \dots = 1 + \frac{1}{1!} + \frac{1}{2!} + \dots = e, \\ S_2 &= \sum_1^{\infty} \frac{k^2}{k!} = 1 + \frac{2^2}{2!} + \frac{3^2}{3!} + \dots = 1 + \frac{2}{1!} + \frac{3}{2!} + \frac{4}{3!} + \dots \\ &= \left\{ 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots \right\} + \left\{ \frac{1}{1!} + \frac{2}{2!} + \frac{3}{3!} + \dots \right\} = e + S_1 = 2e, \end{aligned} \quad (2.3)$$

and continuing like that to  $S_8$ . For large  $n$  this method is not convenient. However, later that year Ligowski [4] suggested a better method, providing a generating function for the numbers  $S_n$ :

$$e^{e^z} = \sum_{k=0}^{\infty} \frac{e^{kz}}{k!} = \sum_{k=0}^{\infty} \sum_{n=0}^{\infty} \frac{k^n z^n}{k! n!} = \sum_{n=0}^{\infty} S_n \frac{z^n}{n!}. \tag{2.4}$$

Further, an effective iteration formula was found

$$S_n = \sum_{j=0}^{n-1} \binom{n-1}{j} S_j \tag{2.5}$$

by which every  $S_n$  can be evaluated starting from  $S_1$ .

These results were preceded, however, by the work [5] of Grunert (1797–1872), professor at Greifswalde. Among other things, Grunert obtained formula (2.9) from which the evaluation of (2.1) follows immediately.

The structure of the series  $S_n$  hints at the exponential function. Differentiating the expansion

$$e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!} \tag{2.6}$$

and multiplying both sides by  $x$  we get

$$xe^x = \sum_{k=0}^{\infty} \frac{kx^k}{k!}, \tag{2.7}$$

which, for  $x = 1$ , gives  $S_1 = e$ . Repeating the procedure, we find  $S_2 = 2e$  from

$$x(xe^x)' = (x + x^2)e^x = \sum_{k=0}^{\infty} \frac{k^2 x^k}{k!}, \tag{2.8}$$

and continuing like that, for every  $n = 0, 1, 2, \dots$ , we find the relation

$$\left(x \frac{d}{dx}\right)^n e^x = \phi_n(x)e^x = \sum_{k=0}^{\infty} \frac{k^n x^k}{k!}, \tag{2.9}$$

where  $\phi_n$  are polynomials of degree  $n$ . Thus,

$$S_n = \phi_n(1)e, \quad \forall n \geq 0. \tag{2.10}$$

The polynomials  $\phi_n$  deserve a closer look. From the defining relation (2.9) we obtain

$$x(\phi_n e^x)' = x(\phi_n' + \phi_n)e^x = \phi_{n+1}e^x, \tag{2.11}$$

that is,

$$\phi_{n+1} = x(\phi'_n + \phi_n), \quad (2.12)$$

which helps to find  $\phi_n$  explicitly starting from  $\phi_0$ :

$$\begin{aligned} \phi_0(x) &= 1, \\ \phi_1(x) &= x, \\ \phi_2(x) &= x^2 + x, \\ \phi_3(x) &= x^3 + 3x^2 + x, \\ \phi_4(x) &= x^4 + 6x^3 + 7x^2 + x, \\ \phi_5(x) &= x^5 + 10x^4 + 25x^3 + 15x^2 + x, \end{aligned} \quad (2.13)$$

and so on. Another interesting relation, easily proved by induction, is

$$\phi_{n+1}(x) = x \sum_{k=0}^n \binom{n}{k} \phi_k(x). \quad (2.14)$$

From (2.12) and (2.14) one finds immediately

$$\phi'_n(x) = \sum_{k=0}^{n-1} \binom{n}{k} \phi_k(x). \quad (2.15)$$

Obviously,  $x = 0$  is a zero for all  $\phi_n$ ,  $n > 0$ . It can be seen that all the zeros of  $\phi_n$  are real, simple, and nonpositive. The nice and short induction argument belongs to Harper [6].

The assertion is true for  $n = 1$ . Suppose that for some  $n$  the polynomial  $\phi_n$  has  $n$  distinct real nonpositive zeros (including  $x = 0$ ). Then the same is true for the function

$$f_n(x) = \phi_n(x)e^x. \quad (2.16)$$

Moreover,  $f_n$  is zero at  $-\infty$  and by Rolle's theorem its derivative

$$\frac{d}{dx} f_n = \frac{d}{dx} (\phi_n(x)e^x) \quad (2.17)$$

has  $n$  distinct real negative zeros. It follows that the function

$$\phi_{n+1}(x)e^x = x \frac{d}{dx} (\phi_n(x)e^x) \quad (2.18)$$

has  $n + 1$  distinct real nonpositive zeros (adding here  $x = 0$ ).

The polynomials  $\phi_n$  can be defined also by the exponential generating function (extending Ligowski's formula)

$$e^{x(e^z-1)} = \sum_{n=0}^{\infty} \phi_n(x) \frac{z^n}{n!}. \tag{2.19}$$

It is not obvious, however, that the polynomials defined by (2.9) and (2.19) are the same, so we need the following simple statement.

**Proposition 2.1.** *The polynomials  $\phi_n(x)$  defined by (2.9) are exactly the partial derivatives  $(\partial/\partial z)^n e^{x(e^z-1)}$  evaluated at  $z = 0$ .*

Equation (2.19) follows from (2.9) after expanding the exponential  $e^{xe^z}$  in double series and changing the order of summation. A different proof will be given later.

Setting  $z = 2k\pi i$ ,  $k = \pm 1, \pm 2, \dots$ , in the generating function (2.19) one finds

$$e^{2k\pi i} = 1, \quad e^{x(e^{2k\pi i}-1)} = e^0 = 1, \tag{2.20}$$

which shows that the exponential polynomials are linearly dependent:

$$1 = \sum_{n=0}^{\infty} \phi_n(x) \frac{(2k\pi i)^n}{n!} \quad \text{or} \quad 0 = \sum_{n=1}^{\infty} \phi_n(x) \frac{(2k\pi i)^n}{n!}, \quad k = \pm 1, \pm 2, \dots \tag{2.21}$$

In particular,  $\phi_n$  are not orthogonal for any scalar product on polynomials. (However, they have the semiorthogonality property mentioned in Section 1 and proved in Section 4.)

Comparing coefficient for  $z$  in the equation

$$e^{(x+y)e^z} = e^{xe^z} e^{ye^z} \tag{2.22}$$

yields the binomial identity

$$\phi_n(x+y) = \sum_{k=0}^n \binom{n}{k} \phi_k(x) \phi_{n-k}(y). \tag{2.23}$$

With  $y = -x$  this implies the interesting "orthogonality" relation for  $n \geq 1$ :

$$\sum_{k=0}^n \binom{n}{k} \phi_k(x) \phi_{n-k}(-x) = 0. \tag{2.24}$$

Next, let  $B_n$ ,  $n = 0, 1, \dots$ , be the Bernoulli numbers. Then for  $p = 0, 1, \dots$ , we have

$$\int_0^x \phi_p(t) dt = \frac{1}{p+1} \sum_{k=1}^{p+1} \binom{p+1}{k} B_{p+1-k} \phi_k(x). \quad (2.25)$$

For proof see Example 4 in [7, page 51], or [9].

### *Some Historical Notes*

As already mentioned, formula (2.9) appears in the work of Grunert [5, page 260], where he gives also the representation (3.4) and computes explicitly the first six exponential polynomials. The polynomials  $\phi_n$  were studied more systematically (and independently) by S. Ramanujan in his unpublished notebooks. Ramanujan's work is presented and discussed by Berndt in [7, Part 1, Chapter 3]. Ramanujan, for example, obtained (2.19) from (2.9) and also proved (2.14), (2.15), and (2.25). Later, these polynomials were studied by Bell [10] and Touchard [11, 12]. Both Bell and Touchard called them "exponential" polynomials, because of their relation to the exponential function, for example, (1.2), (1.3), (2.9), and (2.19). This name was used also by Rota [13]. As a matter of fact, Bell introduced in [10] a more general class of polynomials of many variables,  $Y_{n,k}$ , including  $\phi_n$  as a particular case. For this reason  $\phi_n$  are known also as the single-variable Bell polynomials [14–17]. These polynomials are also a special case of the actuarial polynomials introduced by Toscano [18] which, on their part, belong to the more general class of Sheffer polynomials [19]. The exponential polynomials appear in a number of papers and in different applications—see [9, 13, 20–24] and the references therein. In [25] they appear on page 524 as the horizontal generating functions of the Stirling numbers of the second kind (see (3.4)).

The numbers

$$b_n = \phi_n(1) = \frac{1}{e} S_n \quad (2.26)$$

are sometimes called exponential numbers, but a more established name is Bell numbers. They have interesting combinatorial and analytical applications [15, 16, 18, 26–32]. An extensive list of 202 references for Bell numbers is given in [33].

We note that (2.9) can be used to extend  $\phi_n$  to  $\phi_z$  for any complex number  $z$  by the formula

$$\phi_z(x) = e^{-x} \sum_{k=0}^{\infty} \frac{k^z x^k}{k!} \quad (2.27)$$

(Butzer et al. [34, 35]). The function appearing here is an interesting entire function in both variables,  $x$  and  $z$ . Another possibility is to study the polyexponential function

$$e_s(x, \lambda) = \sum_{n=0}^{\infty} \frac{x^n}{n!(n+\lambda)^s}, \quad (2.28)$$

where  $\operatorname{Re} \lambda > 0$ . When  $s$  is a negative integer, the polyexponential can be written as a finite linear combination of exponential polynomials (see [9]).

### 3. Stirling Numbers and Mellin Derivatives

The iteration formula (2.12) shows that all polynomials  $\phi_n$  have positive integer coefficients. These coefficients are the Stirling numbers of the second kind  $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$  (or  $S(n, k)$ )—see [25, 28, 36–39]. Given a set of  $n$  elements,  $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$  represents the number of ways by which this set can be partitioned into  $k$  nonempty subsets ( $0 \leq k \leq n$ ). Obviously,  $\left\{ \begin{smallmatrix} n \\ 1 \end{smallmatrix} \right\} = 1$ ,  $\left\{ \begin{smallmatrix} n \\ n \end{smallmatrix} \right\} = 1$  and a short computation gives  $\left\{ \begin{smallmatrix} n \\ 2 \end{smallmatrix} \right\} = 2^{n-1} - 1$ . For symmetry one sets  $\left\{ \begin{smallmatrix} 0 \\ 0 \end{smallmatrix} \right\} = 1$ ,  $\left\{ \begin{smallmatrix} n \\ 0 \end{smallmatrix} \right\} = 0$ . The definition of  $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$  implies the property

$$\left\{ \begin{smallmatrix} n+1 \\ k \end{smallmatrix} \right\} = k \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} + \left\{ \begin{smallmatrix} n \\ k-1 \end{smallmatrix} \right\} \tag{3.1}$$

(see [38, page 259]) which helps to compute all  $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$  by iteration. For instance,

$$\left\{ \begin{smallmatrix} n \\ 3 \end{smallmatrix} \right\} = \frac{3^{n-1} - 2^n + 1}{2}. \tag{3.2}$$

A general formula for the Stirling numbers of the second kind is

$$\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} = \frac{1}{k!} \sum_{j=1}^k (-1)^{k-j} \binom{k}{j} j^n. \tag{3.3}$$

**Proposition 3.1.** *For every  $n = 0, 1, 2, \dots$*

$$\phi_n(x) = \left\{ \begin{smallmatrix} n \\ 0 \end{smallmatrix} \right\} + \left\{ \begin{smallmatrix} n \\ 1 \end{smallmatrix} \right\} x + \left\{ \begin{smallmatrix} n \\ 2 \end{smallmatrix} \right\} x^2 + \dots + \left\{ \begin{smallmatrix} n \\ n \end{smallmatrix} \right\} x^n = \sum_{k=0}^n \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} x^k. \tag{3.4}$$

The proof is by induction and is left to the reader. Setting here  $x = 1$  we come to the well-known representation for the numbers  $S_n$

$$S_n = e \left( \left\{ \begin{smallmatrix} n \\ 0 \end{smallmatrix} \right\} + \left\{ \begin{smallmatrix} n \\ 1 \end{smallmatrix} \right\} + \left\{ \begin{smallmatrix} n \\ 2 \end{smallmatrix} \right\} + \dots + \left\{ \begin{smallmatrix} n \\ n \end{smallmatrix} \right\} \right). \tag{3.5}$$

It is interesting that formula (3.4) is very old—it was obtained by Grunert [5, page 260] together with the representation (3.3) for the coefficients which are called now Stirling numbers of the second kind. In fact, coefficients of the form

$$\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} k! \tag{3.6}$$

appear in the computations of Euler—see [37].

Next we turn to some special differentiation formulas. Let  $D = d/dx$ .

### *Mellin Derivatives*

It is easy to see that the first equality in (2.9) extends to (1.3), where  $a$  is an arbitrary complex number, that is,

$$(xD)^n e^{ax} = \phi_n(ax) e^{ax} \quad (3.7)$$

by the substitution  $x \rightarrow ax$ . Even further, this extends to

$$(xD)^n e^{ax^p} = p^n \phi_n(ax^p) e^{ax^p} \quad (3.8)$$

for any  $a, p$  and  $n = 0, 1, \dots$  (simple induction and (2.12)). Again by induction, it is easy to prove that

$$(xD)^n f(x) = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} x^k D^k f(x) \quad (3.9)$$

for any  $n$ -times differentiable function  $f$ . This formula was obtained by Grunert [5, pages 257-258] (see also [2, page 89], where a proof by induction is given).

As we know the action of  $xD$  on exponentials, formula (3.9) can be "discovered" by using Fourier transform. Let

$$F[f](t) = \int_{\mathbb{R}} e^{-ixt} f(x) dx \quad (3.10)$$

be the Fourier transform of some function  $f$ . Then

$$\begin{aligned} f(x) &= \frac{1}{2\pi} \int_{\mathbb{R}} e^{ixt} F[f](t) dx, \\ (xD)^n f(x) &= \frac{1}{2\pi} \int_{\mathbb{R}} e^{ixt} \phi_n(ixt) F[f](t) dx \\ &= \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} x^k F^{-1}[(it)^k F[f]](x) = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} x^k D^k f(x). \end{aligned} \quad (3.11)$$

Next we turn to formula (1.2) and explain its relation to (1.3). If we set  $x = e^t$ , then for any differentiable function  $f$

$$\frac{d}{dt} f = \left( \frac{d}{dx} f \right) \frac{dx}{dt} = \left( \frac{d}{dx} f \right) e^t = (xD)f, \quad (3.12)$$



and we see that (1.2) and (1.3) are equivalent:

$$\left(\frac{d}{dt}\right)^n e^{ae^t} = (xD)^n e^{ax} = \phi_n(ax)e^{ax} = \phi_n(ae^t)e^{ae^t}. \quad (3.13)$$

*Proof of Proposition 2.1.* We apply (1.2) to the function  $f_x(z) = e^{x(e^z-1)} = e^{xe^z}e^{-x}$ :

$$\left(\frac{d}{dz}\right)^n f_x(z) = \phi_n(xe^z)f_x(z). \quad (3.14)$$

From here, with  $z = 0$

$$\left(\frac{d}{dz}\right)^n f_x(z)\Big|_{z=0} = \phi_n(x) \quad (3.15)$$

as needed. □

Now we list some simple operational formulas. Starting from the obvious relation

$$(xD)^n x^k = k^n x^k, \quad n = 0, 1, \dots, k \in \mathbb{R} \quad (3.16)$$

for any function of the form

$$f(t) = \sum_{n=0}^{\infty} a_n t^n, \quad (3.17)$$

we define the differential operator

$$f(xD) = \sum_{n=0}^{\infty} a_n (xD)^n \quad (3.18)$$

with action on functions  $g(x)$ :

$$f(xD)g(x) = \sum_{n=0}^{\infty} a_n (xD)^n g(x). \quad (3.19)$$

When  $g(x) = x^k$ , (3.16) and (3.19) show that

$$f(xD)x^k = \sum_{n=0}^{\infty} a_n k^n x^k = f(k)x^k. \quad (3.20)$$

If now

$$g(x) = \sum_{k=0}^{\infty} c_k x^k \quad (3.21)$$

is a function analytical in a neighborhood of zero, the action of  $f(xD)$  on this function is given by

$$f(xD)g(x) = \sum_{k=0}^{\infty} c_k f(k)x^k, \quad (3.22)$$

provided that the series on the right side converges. When  $f$  is a polynomial, formula (3.22) helps to evaluate series like

$$\sum_{k=0}^{\infty} c_k f(k)x^k \quad (3.23)$$

in a closed form. This idea was exploited by Schwatt [40] and more recently by the present author in [20]. For instance, when  $g(x) = e^x$ , (3.22) becomes

$$\sum_{k=0}^{\infty} f(k) \frac{x^k}{k!} = e^x \sum_{n=0}^{\infty} a_n \phi_n(x). \quad (3.24)$$

As shown in [20] this series transformation can be used for asymptotic series expansions of certain functions.

### *Leibniz Rule*

The higher-order Mellin derivative  $(xD)^n$  satisfies the Leibniz rule

$$(xD)^n (fg) = \sum_{k=0}^n \binom{n}{k} [(xD)^{n-k} f] [(xD)^k g]. \quad (3.25)$$

The proof is easy, by induction, and is left to the reader. We shall use this rule to prove the following proposition.

**Proposition 3.2.** *For all  $n, m = 0, 1, 2, \dots$*

$$\phi_{n+m}(x) = \sum_{k=0}^n \sum_{j=0}^m \binom{n}{k} \left\{ \begin{matrix} m \\ j \end{matrix} \right\} j^{n-k} x^j \phi_k(x). \quad (3.26)$$

*Proof.* One has

$$\phi_{n+m}(x) = (xD)^{n+m} e^x = (xD)^n (xD)^m e^x = (xD)^n (\phi_m(x) e^x), \quad (3.27)$$

which by the Leibniz rule (3.25) equals

$$\sum_{k=0}^n \binom{n}{k} [(xD)^{n-k} \phi_m(x)] [(xD)^k e^x]. \quad (3.28)$$

Using (3.3) and (3.16) we write

$$(xD)^{n-k} \phi_m(x) = \sum_{j=0}^m \left\{ \begin{matrix} m \\ j \end{matrix} \right\} j^{n-k} x^j, \tag{3.29}$$

and since also

$$(xD)^k e^x = \phi_k(x) e^x, \tag{3.30}$$

we obtain (3.26) from (3.27). The proof is completed. □

Setting  $x = 1$  in (3.25) yields an identity for the Bell numbers:

$$b_{n+m} = \sum_{k=0}^n \sum_{j=0}^m \binom{n}{k} \left\{ \begin{matrix} m \\ j \end{matrix} \right\} j^{n-k} b_k. \tag{3.31}$$

This identity was recently published by Spivey [32], who gave a combinatorial proof. After that Gould and Quaintance [16] obtained the generalization (3.26) together with two equivalent versions. The proof in [16] is different from the one above.

Using the Leibniz rule for  $xD$  we can prove also the following extension of property (2.24).

**Proposition 3.3.** *For any two integers  $n, m \geq 0$*

$$(xD)^n \phi_m(x) = \sum_{k=0}^m \left\{ \begin{matrix} m \\ k \end{matrix} \right\} k^n x^k = \sum_{k=0}^n \binom{n}{k} \phi_{m+k}(x) \phi_{n-k}(-x). \tag{3.32}$$

The proof is simple. Just compute

$$\begin{aligned} (xD)^n \phi_m(x) &= (xD)^n [(e^{-x})(\phi_m(x)e^x)] \\ &= \sum_{k=0}^n \binom{n}{k} [(xD)^{n-k} e^{-x}] [(xD)^k (\phi_m(x)e^x)] \end{aligned} \tag{3.33}$$

and (3.32) follows from (1.3).

For completeness we mention also the following three properties involving the operator  $Dx$ . Proofs and details are left to the reader:

$$\begin{aligned} (Dx)^n e^{ax} &= \frac{\phi_{n+1}(ax)}{ax} e^{ax}, \\ (Dx)^n f(x) &= \sum_{k=0}^n \binom{n+1}{k+1} x^k D^k f(x), \\ f(Dx)g(x) &= \sum_{k=0}^{\infty} c_k f(k+1)x^k, \end{aligned} \tag{3.34}$$

analogous to (1.3), (3.9), and (3.22) correspondingly.

For a comprehensive study of the Mellin derivative we refer to [41–43].

### More Stirling Numbers

The polynomials  $\phi_n$ ,  $n = 0, 1, \dots$ , form a basis in the linear space of all polynomials. Formula (3.4) shows how this basis is expressed in terms of the standard basis  $1, x, x^2, \dots, x^n, \dots$ . We can solve for  $x^k$  in (3.4) and express the standard basis in terms of the exponential polynomials

$$\begin{aligned} 1 &= \phi_0, \\ x &= \phi_1, \\ x^2 &= -\phi_1 + \phi_2, \\ x^3 &= 2\phi_1 - 3\phi_2 + \phi_3, \\ x^4 &= -6\phi_1 + 11\phi_2 - 6\phi_3 + \phi_4, \end{aligned} \tag{3.35}$$

and so forth. The coefficients here are also special numbers. If we write

$$x^n = \sum_{k=0}^n (-1)^{n-k} \begin{bmatrix} n \\ k \end{bmatrix} \phi_k, \tag{3.36}$$

then  $\begin{bmatrix} n \\ k \end{bmatrix}$  are the (absolute) Stirling numbers of first kind, as defined in [38]. (The numbers  $\begin{bmatrix} n \\ k \end{bmatrix}$  are nonnegative. The symbol  $s(n, k) = (-1)^{n-k} \begin{bmatrix} n \\ k \end{bmatrix}$  is used for Stirling numbers of the first kind with changing sign—see [28, 33, 39] for more details.)  $\begin{bmatrix} n \\ k \end{bmatrix}$  is the number of ways to arrange  $n$  objects into  $k$  cycles. According to this interpretation,

$$\begin{bmatrix} n \\ k \end{bmatrix} = (n-1) \begin{bmatrix} n-1 \\ k \end{bmatrix} + \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}, \quad n \geq 1. \tag{3.37}$$

### 4. Semiorthogonality of $\phi_n$

**Proposition 4.1.** *For every  $n, m = 1, 2, \dots$ , one has*

$$\int_0^\infty \phi_n(-x)\phi_m(-x)e^{-2x} \frac{dx}{x} = (-1)^{n-1} \frac{2^{n+m} - 1}{n + m} B_{n+m}. \tag{4.1}$$

Here  $B_k$  are the Bernoulli numbers. Note that the right-hand side is zero when  $k + m$  is odd, as all Bernoulli numbers with odd indices  $> 1$  are zeros.

Using the representation (3.4) in (4.1) and integrating termwise one obtains an equivalent form of (4.1):

$$\sum_{k=0}^n \sum_{j=0}^m (-1)^{k+j} \begin{Bmatrix} n \\ k \end{Bmatrix} \begin{Bmatrix} m \\ j \end{Bmatrix} \frac{(k + j - 1)!}{2^{k+j}} = (-1)^{n-1} \frac{2^{n+m} - 1}{n + m} B_{n+m}. \tag{4.2}$$

This (double sum) identity extends the known identity [38, page 317, Problem 6.76]

$$\sum_{j=0}^m (-1)^{j+1} \begin{Bmatrix} m \\ j \end{Bmatrix} \frac{j!}{2^{j+1}} = \frac{2^{m+1} - 1}{m + 1} B_{m+1}. \tag{4.3}$$

Namely, (4.3) results from (4.2) for  $n = 1$ . The presence of  $(-1)^{n-1}$  at the right-hand side in (4.1) is not a “break of symmetry,” because when  $n + m$  is even, then  $n$  and  $m$  are both even or both odd.

*Proof of the proposition.* Starting from

$$\Gamma(z) = \int_0^\infty x^{z-1} e^{-x} dx, \tag{4.4}$$

we set  $x = e^\lambda$ ,  $z = a + it$ , to obtain the representation

$$\Gamma(a + it) = \int_{-\infty}^{+\infty} e^{i\lambda t} e^{a\lambda} e^{-e^\lambda} d\lambda, \tag{4.5}$$

which is a Fourier transform integral. The inverse transform is

$$e^{a\lambda} e^{-e^\lambda} = \frac{1}{2\pi} \int_{\mathbb{R}} e^{-i\lambda t} \Gamma(a + it) dt. \tag{4.6}$$

When  $a = 1$ , this is

$$-e^\lambda e^{-e^\lambda} = \frac{d}{d\lambda} e^{-e^\lambda} = \frac{-1}{2\pi} \int_{\mathbb{R}} e^{-i\lambda t} \Gamma(1 + it) dt. \tag{4.7}$$

Differentiating (4.7)  $n - 1$  times for  $\lambda$  we find

$$\left(\frac{d}{d\lambda}\right)^n e^{-e^\lambda} = \phi_n(-e^\lambda)e^{-e^\lambda} = \frac{-1}{2\pi} \int_{\mathbb{R}} e^{-i\lambda t} (-it)^{n-1} \Gamma(1+it) dt, \quad (4.8)$$

and Parseval's formula yields the equation

$$\int_{\mathbb{R}} \phi_n(-e^\lambda) \phi_m(-e^\lambda) e^{-2e^\lambda} d\lambda = \frac{1}{2\pi} \int_{\mathbb{R}} (-it)^{n-1} (it)^{m-1} |\Gamma(1+it)|^2 dt, \quad (4.9)$$

or, with  $x = e^\lambda$

$$\int_0^\infty \phi_n(-x) \phi_m(-x) e^{-2x} \frac{dx}{x} = \frac{(-1)^n i^{n+m}}{2\pi} \int_{\mathbb{R}} t^{n+m-2} \frac{\pi t}{\sinh(\pi t)} dt. \quad (4.10)$$

The right-hand side is 0 when  $n + m$  is odd. When  $n + m$  is even, we use the integral [1, page 351]

$$\int_0^\infty \frac{t^{2p-1}}{\sinh(\pi t)} dt = \frac{2^{2p} - 1}{2p} (-1)^{p-1} B_{2p} \quad (4.11)$$

to finish the proof. □

Property (4.1) resembles the semiorthogonal property of the Bernoulli polynomials

$$\int_0^1 B_n(x) B_m(x) dx = (-1)^{n-1} \frac{n! m!}{(n+m)!} B_{n+m}, \quad (4.12)$$

see, for instance, [25, page 530].

## 5. Gamma Integrals

We use the technique in the previous section to compute certain Fourier integrals and evaluate the moments of  $\Gamma(a+it)$  and  $\Gamma(a+it)\Gamma(b-it)$ .

**Proposition 5.1.** For every  $n = 0, 1, \dots$  and  $a, b > 0$  one has

$$\int_{\mathbb{R}} e^{-i\mu t} t^n \Gamma(a + it) \Gamma(b - it) dt = i^n 2\pi e^{-b\mu} \sum_{k=0}^n \sum_{m=0}^k \binom{n}{k} \left\{ \begin{matrix} k \\ m \end{matrix} \right\} (-1)^m a^{n-k} \frac{\Gamma(a + b + m)}{(1 + e^{-\mu})^{a+b+m}}, \tag{5.1}$$

$$\int_{\mathbb{R}} e^{-i\lambda t} t^n \Gamma(a + it) dt = i^n 2\pi e^{a\lambda} e^{-e^\lambda} \sum_{k=0}^n \binom{n}{k} a^{n-k} \sum_{m=0}^k \left\{ \begin{matrix} k \\ m \end{matrix} \right\} (-1)^m e^{\lambda m}. \tag{5.2}$$

In particular, when  $\lambda = \mu = 0$ , one obtains the moments

$$G_n(a, b) \equiv \int_{\mathbb{R}} t^n \Gamma(a + it) \Gamma(b - it) dt = i^n \pi \sum_{k=0}^n \sum_{m=0}^k \binom{n}{k} \left\{ \begin{matrix} k \\ m \end{matrix} \right\} (-1)^m a^{n-k} \frac{\Gamma(a + b + m)}{2^{a+b+m-1}}, \tag{5.3}$$

$$G_n(a) \equiv \int_{\mathbb{R}} t^n \Gamma(a + it) dt = \frac{2\pi i^n}{e} \sum_{k=0}^n \sum_{m=0}^k \binom{n}{k} \left\{ \begin{matrix} k \\ m \end{matrix} \right\} (-1)^m a^{n-k}. \tag{5.4}$$

When  $n = 0$  in (5.1) one has the known integral

$$\int_{\mathbb{R}} e^{-i\mu t} \Gamma(a + it) \Gamma(b - it) dt = 2\pi \Gamma(a + b) e^{-b\mu} (1 + e^{-\mu})^{-a-b}, \tag{5.5}$$

which can be found in the form of an inverse Mellin transform in [44].

*Proof.* Using again (4.6)

$$e^{a\lambda} e^{-e^\lambda} = \frac{1}{2\pi} \int_{\mathbb{R}} e^{-i\lambda t} \Gamma(a + it) dt, \tag{5.6}$$

we differentiate both side  $n$  times

$$\left( \frac{d}{d\lambda} \right)^n [e^{a\lambda} e^{-e^\lambda}] = \frac{1}{2\pi} \int_{\mathbb{R}} e^{-i\lambda t} (-it)^n \Gamma(a + it) dt, \tag{5.7}$$

and then, according to the Leibniz rule and (1.2) the left-hand side becomes

$$\left( \frac{d}{d\lambda} \right)^n [e^{a\lambda} e^{-e^\lambda}] = e^{a\lambda} e^{-e^\lambda} \sum_{k=0}^n \binom{n}{k} \phi_k(-e^\lambda) a^{n-k}. \tag{5.8}$$

Therefore,

$$e^{a\lambda} e^{-e^\lambda} \sum_{k=0}^n \binom{n}{k} \phi_k(-e^\lambda) a^{n-k} = \frac{1}{2\pi} \int_{\mathbb{R}} e^{-i\lambda t} (-it)^n \Gamma(a+it) dt, \quad (5.9)$$

and (5.2) follows from here.

Replacing  $\lambda$  by  $\lambda - \mu$  we write (5.6) in the form

$$e^{b\lambda} e^{-b\mu} e^{-e^\lambda e^{-\mu}} = \frac{1}{2\pi} \int_{\mathbb{R}} e^{-i\lambda t} e^{i\mu t} \Gamma(b+it) dt, \quad (5.10)$$

and then Parseval's formula for Fourier integrals applied to (5.9) and (5.10) yields

$$\begin{aligned} e^{-b\mu} \sum_{k=0}^n \binom{n}{k} a^{n-k} \int_{\mathbb{R}} e^{(a+b)\lambda} e^{-e^\lambda(1+e^{-\mu})} \phi_k(-e^\lambda) d\lambda \\ = \frac{(-i)^n}{2\pi} \int_{\mathbb{R}} e^{-i\mu t} t^n \Gamma(a+it) \Gamma(b-it) dt. \end{aligned} \quad (5.11)$$

Returning to the variable  $x = e^\lambda$  we write this in the form

$$\begin{aligned} \frac{1}{2\pi} \int_{\mathbb{R}} e^{-i\mu t} t^n \Gamma(a+it) \Gamma(b-it) dt \\ = i^n e^{-b\mu} \sum_{k=0}^n \binom{n}{k} a^{n-k} \int_0^\infty \phi_k(-x) x^{a+b-1} e^{-x(1+e^{-\mu})} dx \\ = i^n e^{-b\mu} \sum_{k=0}^n \sum_{j=0}^k \binom{n}{k} \left\{ \begin{matrix} k \\ j \end{matrix} \right\} a^{n-k} (-1)^j \int_0^\infty x^{a+b+j-1} e^{-x(1+e^{-\mu})} dx \\ = i^n e^{-b\mu} \sum_{k=0}^n \sum_{j=0}^k \binom{n}{k} \left\{ \begin{matrix} k \\ j \end{matrix} \right\} a^{n-k} (-1)^j \frac{\Gamma(a+b+j)}{(1+e^{-\mu})^{a+b+j}}, \end{aligned} \quad (5.12)$$

which is (5.1). The proof is complete.  $\square$

Next, we observe that for any polynomial

$$p(t) = \sum_{n=0}^m a_n t^n \quad (5.13)$$

one can use (5.4) to write the following evaluation:

$$\int_{\mathbb{R}} p(t) \Gamma(a+it) dt = \sum_{n=0}^m a_n G_n(a). \quad (5.14)$$



In particular, when  $a = 1$  we have

$$G_n(1) = 2\pi i^n e^{-1} \phi_{n+1}(-1), \quad (5.15)$$

and therefore,

$$\int_{\mathbb{R}} p(t) \Gamma(1 + it) dt = \frac{2\pi}{e} \sum_{n=0}^m a_n i^n \phi_{n+1}(-1). \quad (5.16)$$

More applications can be found in the recent papers [9, 20, 21].

## References

- [1] A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev, *Integrals and Series. Vol. 1. Elementary Functions*, Gordon & Breach Science, New York, NY, USA, 1986.
- [2] H. W. Gould, *Topics in Combinatorics*, H. W. Gould, 2nd edition, 2000.
- [3] G. Dobinski, "Summerung der Reihe  $\sum n^m/n!$ , für  $m = 1, 2, 3, 4, 5, \dots$ ," *Archiv der Mathematik und Physik*, vol. 61, pp. 333–336, 1877.
- [4] W. Ligowski, "Zur summerung der Reihe...", *Archiv der Mathematik und Physik*, vol. 62, pp. 334–335, 1878.
- [5] J. A. Grunert, "Über die Summerung der Reihen...", *Journal für die reine und angewandte Mathematik*, vol. 25, pp. 240–279, 1843.
- [6] L. H. Harper, "Stirling behavior is asymptotically normal," *Annals of Mathematical Statistics*, vol. 38, pp. 410–414, 1967.
- [7] B. C. Berndt, *Ramanujan's Notebooks. Part I*, Springer, New York, NY, USA, 1985.
- [8] B. C. Berndt, *Ramanujan's Notebooks. Part II*, Springer, New York, NY, USA, 1989.
- [9] K. N. Boyadzhiev, "Polyexponentials," <http://www.arxiv.org/pdf/0710.1332>.
- [10] E. T. Bell, "Exponential polynomials," *Annals of Mathematics*, vol. 35, no. 2, pp. 258–277, 1934.
- [11] J. Touchard, "Nombres exponentiels et nombres de Bernoulli," *Canadian Journal of Mathematics*, vol. 8, pp. 305–320, 1956.
- [12] J. Touchard, "Proprietes arithmetiques de certains nombres recurrents," *Annales de la Société Scientifique de Bruxelles A*, vol. 53, pp. 21–31, 1933.
- [13] G.-C. Rota, *Finite Operator Calculus*, Academic Press, New York, NY, USA, 1975.
- [14] L. Carlitz, "Single variable Bell polynomials," *Collectanea Mathematica*, vol. 14, pp. 13–25, 1962.
- [15] H. W. Gould and J. Quaintance, "A linear binomial recurrence and the Bell numbers and polynomials," *Applicable Analysis and Discrete Mathematics*, vol. 1, no. 2, pp. 371–385, 2007.
- [16] H. W. Gould and J. Quaintance, "Implications of Spivey's Bell number formula," *Journal of Integer Sequences*, vol. 11, no. 3, article 08.3.7, 2008.
- [17] Y.-Q. Zhao, "A uniform asymptotic expansion of the single variable Bell polynomials," *Journal of Computational and Applied Mathematics*, vol. 150, no. 2, pp. 329–355, 2003.
- [18] L. Toscano, "Una classe di polinomi della matematica attuariale," *Rivista di Matematica della Università di Parma*, vol. 1, pp. 459–470, 1950.
- [19] R. P. Boas Jr. and R. C. Buck, *Polynomial Expansions of Analytic Functions*, Academic Press, New York, NY, USA, 1964.
- [20] K. N. Boyadzhiev, "A series transformation formula and related polynomials," *International Journal of Mathematics and Mathematical Sciences*, vol. 2005, no. 23, pp. 3849–3866, 2005.
- [21] K. N. Boyadzhiev, "On Taylor's coefficients of the Hurwitz zeta function," *JP Journal of Algebra, Number Theory and Applications*, vol. 12, no. 1, pp. 103–112, 2008.
- [22] A. Papoulis, *Probability, Random Variables, and Stochastic Processes*, McGraw-Hill, New York, NY, USA, 1991.
- [23] J. Riordan, *Combinatorial Identities*, John Wiley & Sons, New York, NY, USA, 1968.
- [24] J. Riordan, *An Introduction to Combinatorial Analysis*, John Wiley & Sons, New York, NY, USA, 1967.
- [25] J. Sandor and B. Crstici, *Handbook of Number Theory, Part II*, Springer/Kluwer Academic Publishers, Dordrecht, The Netherlands, 2004.

- [26] E. T. Bell, "Exponential numbers," *The American Mathematical Monthly*, vol. 41, no. 7, pp. 411–419, 1934.
- [27] N. G. de Bruijn, *Asymptotic Methods in Analysis*, Dover, New York, NY, USA, 3rd edition, 1981.
- [28] M. E. Dasef and S. M. Kautz, "Some sums of some significance," *The College Mathematics Journal*, vol. 28, pp. 52–55, 1997.
- [29] L. F. Epstein, "A function related to the series for  $e^{e^x}$ ," *Journal of Mathematics and Physics*, vol. 18, pp. 153–173, 1939.
- [30] M. Klazar, "Bell numbers, their relatives, and algebraic differential equations," *Journal of Combinatorial Theory. Series A*, vol. 102, no. 1, pp. 63–87, 2003.
- [31] J. Pitman, "Some probabilistic aspects of set partitions," *The American Mathematical Monthly*, vol. 104, no. 3, pp. 201–209, 1997.
- [32] M. Z. Spivey, "A generalized recurrence for Bell numbers," *Journal of Integer Sequences*, vol. 11, pp. 1–3, 2008.
- [33] H. W. Gould, *Catalan and Bell Numbers: Research Bibliography of Two Special Number Sequences*, H. W. Gould, 5th edition, 1979.
- [34] P. L. Butzer, M. Hauss, and M. Schmidt, "Factorial functions and Stirling numbers of fractional orders," *Results in Mathematics*, vol. 16, no. 1–2, pp. 16–48, 1989.
- [35] P. L. Butzer and M. Hauss, "On Stirling functions of the second kind," *Studies in Applied Mathematics*, vol. 84, no. 1, pp. 71–91, 1991.
- [36] L. Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, D. Reidel, Dordrecht, The Netherlands, 1974.
- [37] H. W. Gould, "Euler's formula for  $n$ th differences of powers," *The American Mathematical Monthly*, vol. 85, no. 6, pp. 450–467, 1978.
- [38] R. L. Graham, D. E. Knuth, and O. Patashnik, *Concrete Mathematics: A Foundation for Computer Science*, Addison-Wesley, Reading, Mass, USA, 2nd edition, 1994.
- [39] D. E. Knuth, "Two notes on notation," *The American Mathematical Monthly*, vol. 99, no. 5, pp. 403–422, 1992.
- [40] I. J. Schwatt, *An Introduction to the Operations with Series*, Chelsea, New York, NY, USA, 1962.
- [41] P. L. Butzer, A. A. Kilbas, and J. J. Trujillo, "Fractional calculus in the Mellin setting and Hadamard-type fractional integrals," *Journal of Mathematical Analysis and Applications*, vol. 269, no. 1, pp. 1–27, 2002.
- [42] P. L. Butzer, A. A. Kilbas, and J. J. Trujillo, "Compositions of Hadamard-type fractional integration operators and the semigroup property," *Journal of Mathematical Analysis and Applications*, vol. 269, no. 2, pp. 387–400, 2002.
- [43] P. L. Butzer, A. A. Kilbas, and J. J. Trujillo, "Mellin transform analysis and integration by parts for Hadamard-type fractional integrals," *Journal of Mathematical Analysis and Applications*, vol. 270, no. 1, pp. 1–15, 2002.
- [44] F. Oberhettinger, *Tables of Mellin Transforms*, Springer, New York, NY, USA, 1974.