

3-1-2001

Expression of PTHrP and PTHR (PTH/PTHrP-r) mRNAs and polypeptides in bovine ovary and stimulation of bovine blastocyst development in vitro following PTHrP treatment during oocyte maturation.

P H Watson

M E Westhusin

A J Watson

Follow this and additional works at: <https://ir.lib.uwo.ca/obs gynpub>

 Part of the [Obstetrics and Gynecology Commons](#)

Citation of this paper:

Watson, P H; Westhusin, M E; and Watson, A J, "Expression of PTHrP and PTHR (PTH/PTHrP-r) mRNAs and polypeptides in bovine ovary and stimulation of bovine blastocyst development in vitro following PTHrP treatment during oocyte maturation." (2001). *Obstetrics & Gynaecology Publications*. 61.
<https://ir.lib.uwo.ca/obs gynpub/61>

1 Accepted to Anatomy and Embryology

2 **Expression of PTHrp and PTHR (PTH/PTHrP-r) mRNAs and Polypeptides in Bovine Ovary**
3 **and Stimulation of Bovine Blastocyst Development *in vitro* Following PTHrP Treatment**
4 **during Oocyte Maturation ***

5

6 P.H. Watson¹, Mark E Westhusin² and Andrew J. Watson^{5,4},

7 Department of Medicine, The University of Western Ontario and the Lawson Research Institute¹,

8 Department of Veterinary Physiology and Pharmacology², College of Veterinary Medicine, Texas

9 A&M University, College Station, Texas, USA; Depts of Obstetrics and Gynaecology³ and

10 Physiology⁴, The University of Western Ontario, London, Ontario, Canada;

11

12 Short Title: Bovine Ovarian PTHrP

13 Key Words: blastocyst, *in vitro* fertilization, growth factors

14

15 ⁵All correspondence to: Dr. A.J. Watson, Departments of Obstetrics and Gynaecology and

16 Physiology, The University of Western Ontario, London, Ontario, Canada, N6A 5C1

17 phone:(519)661-3467; fax (519)661-3827; email Andrew.Watson@med.uwo.ca

18

19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

Abstract

Parathyroid hormone related protein (PTHrP) and its receptor have well established roles in the development and regulation of many tissues, including bone and mammary gland. The objectives of this study were: 1) to characterize the distribution of mRNAs encoding parathyroid hormone (PTH)-related protein (PTHrP) and receptor (PTHR) in bovine ovary; 2) to characterize the distribution of PTHrP and PTHR polypeptides in bovine ovary; and 3) to examine the influences of PTHrP (1-141) treatment during bovine oocyte maturation *in vitro* on blastocyst development. mRNAs encoding PTHrP and PTHR were detected by *in situ* hybridization methods in oocytes, and granulosa cells in all follicles from primordial to large antral. PTHrP and PTHR polypeptides displayed distinct distribution patterns with PTHrP polypeptides primarily confined to oocytes from primordial to large antral follicles. PTHrP polypeptides were detectable but at a reduced level in ovarian stroma and in granulosa and thecal layers. PTHR polypeptides were detected in oocytes of all follicular stages but were predominantly found in ovarian stroma, granulosa and theca follicular layers. Supplementation of serum-free cSOFMaa oocyte maturation medium with PTHrP (1-141) resulted in a concentration-dependent increase in development to the blastocyst stage *in vitro*. The results suggest that granulosa cells may be a primary site of PTHrP production and release. Oocytes from all follicular stages stained strongly for PTHrP polypeptides and PTHrP enhanced development to the blastocyst stage *in vitro*.

Introduction

42
43 With the characterization of effective, defined and protein-free media to support the
44 maturation of mammalian oocytes *in vitro* it is now possible to examine the roles played by specific
45 hormone and growth factor modulators during oocyte maturation and upon subsequent early
46 embryonic development (Saeki et al., 1991; Gardner, 1994; Eckert et al., 1995; Keskinetepe and
47 Brackett, 1996; Hill et al., 1997; Avery et al., 1998; Krishner and Bavister, 1999; Watson et al.,
48 2000). Parathyroid hormone related protein (PTHrP) was originally isolated from patients suffering
49 from humoral hypercalcemia of malignancy (Spiegel et. al., 1983; Strewler et. al., 1987; Stewart et.
50 al., 1987). As its name suggests, this molecule shares structural homology with parathyroid
51 hormone (PTH) (Stewart et. al., 1987; Horiuchi et. al. 1987), which enables PTHrP and PTH to
52 signal through a common PTH/PTHrP receptor (Rodan, et. al., 1988; Abou-Samra, et. al., 1992).
53 While PTH is predominantly (if not exclusively) produced by the parathyroid glands, PTHrP is
54 synthesized by many tissues including skin, brain, pancreas, adrenal glands, smooth muscle, heart,
55 lung, lactating breast, uterus, ovary and placenta (Urena et. al., 1993; Tian et. al., 1993; Lee et al.,
56 1995; Weaver et. al., 1995; Downey et. al., 1997; Li et. al., 1995; Vasavada, et. al., 1998; Curtis et.
57 al., 1998; Ferguson et. al., 1998; Moseley and Gillespie, 1995; Gutmann et. al., 1993). PTHrP is
58 predominantly a paracrine/autocrine regulator of cellular growth and differentiation (Moseley and
59 Gillespie, 1995). The expression and distribution of gene products encoding PTHrP and the PTHR
60 have not been investigated within the bovine ovary. The objectives of the present study were: 1) to
61 characterize the distribution of mRNAs encoding parathyroid hormone (PTH)-related protein
62 (PTHrP) and receptor (PTHR) in the bovine ovary; 2) to characterize the distribution of PTHrP and
63 PTHR polypeptides in the bovine ovary; and 3) to examine the influences of PTHrP (1-141)

64 treatment during bovine oocyte maturation *in vitro* on blastocyst development.

65 **Materials and methods**

66 Detection of PTHrP and PTHR mRNAs in Bovine Ovary by *in situ* Hybridization

67 Bovine ovaries were collected at a local slaughterhouse and were quartered and placed
68 into 4% paraformaldehyde in phosphate buffered saline (PBS) pH 7.2 fixative for overnight
69 fixation. Ovarian pieces were then processed for routine paraffin embedding. *In situ*
70 hybridization (ISH) was carried out using biotin-labeled sense and antisense riboprobes and the
71 Genpoint CSA kit (Dako Diagnostics, Mississauga, ON, Canada) as described (Watson, et. al.,
72 2000a; Watson, et. al., 2000b). Labeled riboprobes were prepared from plasmids containing
73 either a 2Kb cDNA corresponding to rat PTHR mRNA (Pausova, et.al., 1994; Amizuka, et.al.,
74 1997) or a 0.7 Kb cDNA corresponding to rat PTHrP mRNA (Yasuda et.al., 1989) using T3 and
75 T7 RNA polymerases (Gibco, Burlington, ON, Canada) and a nucleotide labeling mix containing
76 biotin-16-UTP (Boehringer-Mannheim, Burlington, ON, Canada). Tissue sections were
77 deparaffinized and rehydrated in a standard xylene and alcohol series and ISH performed
78 according to the manufacturer's directions. Briefly, slides were heated in Target Retrieval
79 Solution and proteinase K to undo protein cross-links caused by fixation. Endogenous peroxidase
80 activity was quenched in 0.3% hydrogen peroxide in methanol for 50 min at room temperature.
81 Sections were hybridized for 2 hours at 42°C with 5 ng/ml biotin-labeled riboprobe in the
82 supplied RNA hybridization buffer. After stringent washing at 55°C, sections were treated with
83 successive incubations in primary streptavidin-horse radish peroxidase, biotinyl-tyramide
84 solution and secondary streptavidin-horse radish peroxidase (Dako) for 15 minutes each at room
85 temperature. Color (golden-brown) was developed with the supplied diaminobenzidine diluted as

86 directed (Dako) and sections counterstained with Carazzi's haematoxylin (blue) prior to
87 dehydration and mounting. We have employed this procedure using the same cDNAs to generate
88 riboprobes to localize PTHrP and PTHR mRNAs in rat tissues (including ovary) in previous
89 studies (Watson, et. al., 2000a; Watson, et. al., 2000b).

90

91 Immunocytochemical (ICC) Detection of PTHrP and PTHR polypeptides in Bovine Ovary

92 ICC was performed as described (Watson et. al., 1995; Watson et. al., 2000a; Watson et. al.,
93 2000b). Briefly, tissue sections (as described above) were deparaffinized and rehydrated in a standard
94 xylene-ethanol series, post-fixed in 4% paraformaldehyde for 15 minutes and incubated for 10
95 minutes with 10 µg/ml proteinase K (Gibco, Burlington, ON, Canada) at room temperature to
96 retrieve fixation-concealed antigens. Sections were stabilized in 4% paraformaldehyde for a further
97 15 minutes before proceeding with the ICC procedure using the Vectastain ABC kit (Vector Labs,
98 Burlington, ON, Canada) following the manufacturer's directions. A polyclonal rabbit anti-rat PTHR
99 antibody was used: antibody PTH-IV raised against a portion of the first extracellular loop
100 (TLDEARLTEEELH; aa 249-262) (Babco, Berkeley, CA, USA) (Largo, Gomez-Garre, et. al., 1999).
101 The PTHrP antibody used was a mouse monoclonal raised against amino acids 34-53 of human
102 PTHrP (Oncogene Research Products, Cambridge, MA, USA). Both antisera are fully characterized
103 and their specificity for PTHrP and PTHR polypeptides is well-established (Watson et. al, 1999a;
104 Watson et. al., 1999b; Ferguson et. al., 1998, Largo et. al., 1999). Primary antibodies were diluted to
105 100 µg/ml in 1% BSA-PBS. As a control, some sections were treated with PTHR primary antibody
106 that had been pre-absorbed overnight at 4°C with the appropriate peptide. For the PTHrP antiserum,
107 control sections were treated with normal mouse serum instead of primary antibody. All sections

108 were counterstained with Carazzi's hematoxylin prior to dehydration and mounting.

109

110 Oocyte Collection, Insemination and Embryo Culture:

111 Cumulus oocyte complexes (COCs), were collected by razor blade slashing of slaughterhouse
112 ovaries within 4 h of removal from the animal (Wiemer et al., 1991; Watson et al., 1994). The COCs
113 were collected in oocyte collection medium (HEPES-buffered TCM-199 + 2% (v/v) NCS; Gibco
114 BRL, Burlington, ON, Canada) and then were washed 4 times in serum-free medium prior to
115 placement in oocyte maturation medium. Only denuded oocytes were discarded and a COC selection
116 strategy was not employed in this study. Following oocyte maturation (see below for specific
117 experimental conditions), oocytes were inseminated *in vitro* with frozen thawed bovine semen
118 (Semex Canada Inc., Guelph, ON, Canada) prepared using a "swim-up" method in Sperm TL
119 medium (HEPES-buffered modified Tyrodes solution as described in (Parrish et al., 1986). Matured
120 COCs were washed in sperm TL and placed in equilibrated fertilization drops (50 COCs/ 300 μ l
121 drop) composed of bicarbonate-buffered modified Tyrodes solution under light paraffin oil (Parrish
122 et al., 1986) BDH Inc., Toronto, ON, Canada). COCs and sperm (2.25×10^5 motile spermatozoa/drop)
123 were incubated for 18h at 39°C under 5% CO₂ in air atmosphere before removal with a fine bore
124 glass pipette of the cumulus investment including all corona cells. Inseminated oocytes (40-50) were
125 placed into embryo culture consisting initially of 20 μ l microdrops of citrate (0.5mM) and
126 polyvinylalcohol (PVA; 3 mg/ml) supplemented synthetic oviduct fluid medium (cSOFMaa)
127 (Keskinetepe et al., 1995; Kestinkepe and Brackett, 1996; Watson et al., 2000) + 1X non-essential
128 amino acids (NEA, Sigma-Aldrich Canada Ltd, Oakville, Ontario) and 1X essential amino acids
129 (MEM, Gibco, BRL) under paraffin oil in a humidified 5% CO₂ / 7% O₂ / 88% N₂ culture

130 atmosphere. Two days following initiation of culture, the microdrops were increased in volume by
131 addition of 20 μ l of cSOFMaa medium. On days 5 and 7 of culture 20 μ l of medium was removed
132 from each and replaced with 20 μ l of fresh medium, thus keeping the microdrop volume constant for
133 the remainder of the eight day culture interval. Cleavage and blastocyst frequencies were assessed on
134 day 3 and 8 post insemination respectively.

135

136 Experimental Design:

137 A PTHrP concentration experiment (0, 10^{-10} M, 1.59 ng/ml; 10^{-8} M, 15.9 ng/ml; and 10^{-6}
138 M, 159 ng/ml PTHrP) was conducted employing a randomized design that allocated equivalent
139 numbers of non-selected COCs (total number of 442 COCs representing 4 experimental
140 replicates with 25-30 oocytes allocated to each treatment group per replicate) to gonadotrophin
141 and estradiol 17- β -free cSOFMaa oocyte maturation medium employing a 5% CO₂ in air
142 atmosphere at 39°C for 22 h. Following maturation, oocyte pools were inseminated and zygotes
143 were placed into culture for assessment of frequency of development to the blastocyst stage as
144 described.

145

146 Statistical Analysis:

147 Culture data were analyzed using the SigmaStat (Jandel Scientific) software package.
148 One-way analysis of variance (ANOVA), followed by pairwise multiple comparisons
149 (Bonferroni Method), were used for analysis of differences in the means for two or more
150 populations. Differences of P# 0.05 were significant differences.

Results

Detection of PTHrP and PTHR mRNAs in Bovine Ovary by *in situ* Hybridization

PTHrP and PTHR mRNAs were localized in bovine ovarian tissue with biotinylated antisense riboprobes. Experiments were applied to tissue sections obtained from three bovine ovaries collected at slaughter and all *in situ* experiments were conducted a minimum of three times. In total over 100 follicles including stages from primordial to large antral follicles were examined. mRNAs encoding PTHrP were detected by *in situ* hybridization methods in oocytes, and granulosa cells in all follicles from primordial to large antral (Fig. 1 B, D, F, E). The granulosa cells of antral follicles displayed more intense signals for PTHrP mRNA (Fig 1E). A positive signal for PTHrP mRNAs was observed in thecal layers of larger follicles and in ovarian stroma in general (Figure 1 B,D,F, E). The signal observed for PTHrP sense controls (Fig. 1 A, C) was very low and indicated the specificity of the antisense PTHrP riboprobe. mRNAs encoding PTHR were detected in oocytes and granulosa cells in all follicles from primordial to large antral (Fig2 B,D,F, E). A positive signal for PTHR mRNA was observed in thecal layers of larger follicles and in ovarian stroma (Fig. 2 B,D, E,F). Sense PTHR mRNA controls displayed a very low background signal in granulosa and thecal layers (Fig. 2 A,C).

Immunocytochemical Detection of PTHrP and PTHR polypeptides in Bovine Ovary

PTHrP and PTHR immunoreactivity was studied in intact bovine ovarian tissue sections to assess the distribution of these polypeptides within the bovine ovary. PTHrP and PTHR polypeptides displayed distinct distribution patterns with PTHrP polypeptides primarily confined to oocytes from primordial to large antral follicles (Fig. 3 B,D,F). PTHrP polypeptides were also

173 localized to a lesser extent in ovarian stroma and reduced signals were obtained in granulosa
174 cell and thecal layers (Fig. 3 B,D, E, F). PTHrP immuno-specificity was determined by
175 demonstrating a very low background signal in normal mouse serum-treated tissue sections (Fig.
176 3 A,C). PTHR polypeptides were detected in oocytes of all follicular stages (Fig 4, B, D, F).
177 However, the most intense signals for PTHR polypeptides were localized to ovarian stroma,
178 granulosa cell and thecal follicular layers (Fig 4 B,D,E, F). Pre-absorbed antibody controls
179 indicated a high level of specificity for the PTHR immunolocalization signal (Fig 4A, C).

180

181 Influence of PTHrP during Oocyte Maturation on Development of Bovine *Zygotes In vitro*

182 cSOFMaa culture medium employed for bovine oocyte maturation was supplemented

183 with PTHrP and influences on development to the blastocyst stage were examined. No significant

184 differences in cleavage were observed among the four oocyte maturation treatment groups ($p <$

185 0.05; Fig. 5 and 6). Likewise, development to the 6-8 cell stage was not significantly influenced

186 by PTHrP treatment during oocyte maturation (Fig. 5 and 6). However, development to the

187 blastocyst stage as assessed by the proportion of total oocytes inseminated (Figure 5) or as a

188 proportion of cleaved zygotes progressing to the blastocyst stage was significantly enhanced

189 ($p < 0.05$) by the addition of PTHrP to the oocyte maturation medium. The beneficial influence of

190 PTHrP during oocyte maturation on blastocyst formation arose following the 6-8 cell stage since

191 PTHrP treatment during oocyte maturation lead to a significant increase ($p < 0.05$) in the

192 proportion of 6-8 cell zygotes progressing to the blastocyst stage (Figure 6).

193

194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215

DISCUSSION

This study characterized the distribution of mRNAs and polypeptides encoding PTHrP and the PTHR within bovine ovary and investigated the influence of PTHrP treatment during *in vitro* maturation of bovine oocytes on development to the blastocyst stage. The results indicate that the bovine ovary expresses both the mRNAs encoding PTHrP and PTHR and their polypeptides. The granulosa layer may be a primary site of PTHrP production (high mRNA signals) but the majority of this PTHrP does not appear to be stored within the granulosa layer (low polypeptide signal). Why the granulosa highly expresses PTHrP mRNA but contains little peptide is unclear. It may be that granulosa PTHrP mRNA is not translated well (or at all) or even that the PTHrP peptide is secreted to the antral cavity. The answer to these possibilities awaits further experimentation. Oocytes from all follicular stages stained intensely for PTHrP polypeptides. The cognate receptor for PTHrP, the PTHR, was expressed in both oocytes and granulosa cells of the bovine ovary suggesting that PTHrP may act in both an autocrine and paracrine fashion to regulate bovine follicle development. PTHrP may also be an important regulator of oocyte maturation since addition of PTHrP to bovine oocyte maturation medium enhanced the development of fertilized bovine eggs to the blastocyst stage *in vitro*.

PTHrP and its receptor, the PTHR, are widely expressed in both skeletal and extraskeletal tissues (Lee et al., 1995; Urena et. al., 1993; Tian et. al., 1993; Watson et. al, 2000a). Reports suggest that the distribution of PTHrP and its receptor are nearly ubiquitous since this receptor ligand duo are described in tissues as diverse as skin (Kaiser et. al., 1992), smooth muscle (Moseley and Gillespie, 1995; Williams et. al., 1994; Yamamoto et. al., 1992; Watson, et. al., 2000a), bone (Lee et. al., 1993; Kronenberg et. al., 1998), kidney (Lee et. al., 1996; Amizuka et.

216 al., 1997; Yang et. al., 1997; Watson, et. al., 2000a), uterus-placenta (Williams et. al., 1994;
217 Tucci and Beck, 1998; Watson, et. al., 2000b), brain (Weaver et. al., 1995), breast (Downey et.
218 al., 1997), intestine (Li et. al., 1995; Watson, et. al, 2000a), pancreas and cardiovascular system
219 (Vasavada, et. al., 1998). Few studies to date have investigated the expression or role of PTHrP
220 and the PTHR in the mammalian ovary. Ovarian expression of PTHrP was first demonstrated by
221 the humoral hypercalcemia of malignancy associated with ovarian small cell carcinomas and
222 PTHrP is now widely used as an ovarian tumour expression marker (Matias-Guiu et. al., 1994;
223 Inoue et. al., 1995). Northern blots were used to identify PTHR transcripts in rat ovary (Urena et.
224 al., 1993; Joun et. al., 1997) and PTHrP expression has been demonstrated in the ovary of the
225 developing frog (Danks et. al., 1997) and most recently in the porcine ovary (Garmey et al.,
226 2000). Both the PTHR and its ligand, PTHrP, have been identified in rat ovary (Watson, et. al,
227 2000a). Only one study has identified PTHrP as a component of human ovarian follicular fluid
228 and demonstrated that the granulosa-luteal cells were the source of the PTHrP (Gutmann et. al.,
229 1993). None of these studies have attempted to localize mRNAs encoding PTHrP and the PTHR
230 or their polypeptides as we have in the present study. In a recent study reporting nuclear
231 localization of the PTHR, we observed that transcripts encoding both the PTHR and PTHrP and
232 their polypeptides were present in rat ovary (Watson et. al., 2000a).

233 PTHrP is implicated as a regulator of embryo development (Behrendtsen et al.,1995; Lee
234 et al., 1995; Nowak et al., 1999). PTHrP and the PTHR are expressed during differentiation of
235 embryonal carcinoma or stem cells to primitive and parietal endoderm (van de Stolpe et. al.,
236 1993) and PTHrP is essential to the outgrowth of parietal endoderm from isolated mouse embryo
237 inner cell mass (ICM) regardless of the substratum used for outgrowth (Behrendtsen et al., 1995).

238 In the developing mouse embryo, expression of PTHR transcripts was detected in parietal
239 endoderm from day 5.5 p.c. onwards. In the embryo proper, PTHR transcripts were highly
240 expressed at sites of epithelial-mesenchyme interaction in developing intestine, lung, kidney and
241 dermis starting at day 9.5 p.c. (Karperien, et. al., 1994). A comprehensive survey of PTHrP and
242 PTHR expression during rat fetal development found that, in extraskeletal tissues, PTHrP mRNA
243 was largely expressed in surface epithelia while PTHR mRNA was mainly localized to the
244 adjacent mesenchyme (Lee et al., 1995). This was true for tissues as diverse as lung, choroid
245 plexus, teeth, heart and skin. These results are strongly suggestive of a paracrine role for PTHrP
246 during development. It has been proposed that PTHrP is a local, autocrine/paracrine regulator of
247 cell growth and differentiation (Tucci and Beck, 1998; Moseley and Gillespie, 1995). A
248 physiological role for PTHrP during mammalian endochondral bone formation and modeling is
249 well documented (Lee, et.al., 1996; Lanske et. al., 1996; Lanske and Kronenberg, 1998). In other
250 tissues PTHrP regulates smooth muscle relaxation, placental calcium transfer, breast
251 development and lactation, vascular smooth muscle tension and pancreatic β -cell growth (Mosely
252 and Gillespie, 1995; Vasavada et. al., 1998; Porter et. al., 1998). Very recently Nowak, et al.,
253 (1999) reported that PTHrP and transforming growth factor β (TGF- β) both interact to promote
254 murine blastocyst outgrowth demonstrating that PTHrP may be an important regulator of early
255 developmental events.

256 The characterization of optimized culture conditions for mammalian oocytes and early
257 embryos is an important priority. Progress in the development of effective defined media for
258 embryo culture has occurred rapidly in the last few years. Media supplements such as serum
259 have, however, only recently been removed from standard culture protocols and little research has

260 investigated the role of ovarian factors on oocyte maturation and early development *in vitro* under
261 defined serum-free culture conditions (Saeki et al., 1991; Gardner, 1994; Eckert et al., 1995;
262 Keskinetepe and Brackett, 1996; Hill et al., 1997; Avery et al., 1998; Krishner and Bavister, 1999;
263 Watson et al., 2000). Research from our laboratories has characterized cSOFMaa medium for
264 bovine oocyte maturation (Watson et al., 2000). Our results indicate that supplementation of
265 cSOFMaa oocyte maturation medium with PTHrP may be of benefit for supporting enhanced
266 numbers of bovine zygotes through to the blastocyst stage. Future studies will explore the
267 mechanism underlying this positive influence and will determine whether PTHrP should become
268 a routine supplement to mammalian oocyte maturation and embryo culture media.

269 **ACKNOWLEDGMENTS**

270 We thank John Looye, Dave Natale, Lisa Barcroft, and the ABEL laboratories (University of
271 Guelph) for assisting with ovary and oocyte collections, Dr. Shaffat Rabbani for the PTHrP (1-
272 141) peptide, Dr. Geoffrey N. Hendy for the PTHrP and PTHR probes and Drs John Eppig,
273 Randy Prather and Barry Bavister (members of the NIH Cooperative Program see below) for
274 critically reviewing this manuscript. This work was supported by the National Institute of Child
275 Health and Human Development, National Cooperative Program on Non-Human In Vitro
276 Fertilization and Embryo Development, USA.

277

278 **References**

- 279 Abou-Samra AB, Juppner H, Force T, Freeman MW, Kong XF, Schipani E, Urena P, Richards J
280 Bonventre JV, Potts Jr JT, Kronenberg HM and Segre GV (1992) Expression cloning of a
281 common receptor for parathyroid hormone and parathyroid hormone-related peptide from rat
282 osteoblast-like cells: a single receptor stimulates intracellular accumulation of both cAMP and
283 inositol triphosphates and increases intracellular free calcium. Proc Natl Acad Sci USA 89:2732-
284 2736
- 285
- 286 Amizuka N, Lee HS, Kwan MY, Arazani A, Warshawsky H, Hendy GN, Ozawa H, White JH and
287 Goltzman D (1997) Cell-specific expression of the parathyroid hormone (PTH)/PTH-related
288 peptide receptor gene in kidney from kidney-specific and ubiquitous promoters. Endocrinology
289 138:469-481
- 290
- 291 Avery B, Bavister BD and Greve T (1998) Development of bovine oocytes, in vitro matured in a
292 chemically defined protein-free medium, supplemented with different amino acid
293 formulations. Theriogenology 49:306.
- 294
- 295 Behrendtsen O, Alexander CM and Werb Z (1995) Cooperative interactions between extracellular
296 matrix, integrins and parathyroid hormone-related peptide regulate parietal endoderm
297 differentiation in mouse embryos. Development 121:4137-4148
- 298
- 299

300

301 Curtis NE, Thomas RJ, Gillespie MT, King RG, Rice GE and Wlodek ME (1998) Parathyroid
302 hormone-related protein (PTHrP) mRNA splicing and parathyroid hormone/PTHrP receptor
303 mRNA expression in human placenta and fetal membranes. *J Mol Endo* 21:225-234

304

305 Danks JA, McHale JC, Martin TJ and Ingleton PM (1997) Parathyroid hormone-related protein in
306 tissues of the emerging frog (*Rana temporaria*): immunohistochemistry and in situ hybridisation.
307 *J Anat* 190 (Pt 2): 229-238

308

309 Downey SE, Hoyland J, Freemont AJ, Knox F, Walls J and Bundred NJ (1997) Expression of the
310 receptor for parathyroid hormone-related protein in normal and malignant breast tissue. *J Pathol*
311 183:212-217

312

313 Eckert J and Niemann H (1995) In vitro maturation, fertilization and culture to blastocyst of
314 bovine oocytes in protein-free media. *Theriogenology* 43:1211-1225.

315

316 Ferguson II JE, Seaner RM, Bruns DE, Iezzoni JC and Bruns ME (1998) Expression and specific
317 immunolocalization of the human parathyroid hormone/parathyroid hormone-related protein
318 receptor in the uteroplacental unit. *Amer J Ob Gyn* 179: 321-329

319

320 Gardner DK (1994) Mammalian embryo culture in the absence of serum or somatic cell support.
321 *Cell Biology Inter* 18:1163-1179.

322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343

Garmey JC, Schnorr JA, Bruns ME, Bruns DE, Seaner RM, Ferguson JE, Jayes FCL, Aquirre C, and Veldhuis JD (2000) Expression of parathyroid hormone-related peptide (PTH-rp) and its receptor in the procine ovary:regulation by transforming growth factor- β and possible paracrine effects of granulose cell PTH-rp secretion on theca cells. Biol Reprod 62:334-339.

Gutmann JN, Burtis WJ, Dreyer BE, Andrade-Gordon P, Penzias AS, Polan ML and Insogna KL (1993) Human granulosa-luteal cells secrete parathyroid hormone-related protein in vivo and in vitro. J Clin Endocrinol & Metab 76:1314-1318

Hill JL, Wade MG, Nancarrow CD, Kelleher DL and Boland MP (1997) Influence of ovine oviductal amino acid concentrations and an ovine oestrus associated glycoprotein on development and viability of bovine embryos. Mol Reprod Dev 47:164-169.

Horiuchi N, Caulfield MP, Fisher JE, Goldman ME, McKee RL, Reagan JE, Levy JJ, Nutt RF, Rodan SB, Scholfield TL, Clemens TL and Rosenblatt M (1987) Similarity of synthetic peptide from human tumor to parathyroid hormone *in vivo* and *in vitro*. Science 238:1566-1568

Inoue H, Kikuchi Y, Hirata J, Wada S, Seki K and Nagata I (1995) Dysgerminoma of the ovary with hypercalcemia associated with elevated parathyroid hormone-related protein. Jap J Clin Oncol 25:113-117

344

345

346

347 Joun H, Lanske B, Karperien M, Qian F, Defize L and Abou-Samra A (1997) Tissue-specific
348 transcription start sites and alternative splicing of the parathyroid hormone (PTH)/PTH-related
349 peptide (PTHrP) receptor gene: a new PTH/PTHrP receptor splice variant that lacks the signal
350 peptide. *Endocrinology* 138:1742-1749

351

352 Kaiser SM, Laneuville P, Bernier SM, Rhim JS, Kremer R and Goltzman D (1992) Enhanced growth
353 of a human keratinocyte cell line induced by antisense RNA for parathyroid hormone-related
354 peptide. *J Biol Chem* 267:13623-13628

355

356 Karperien M, Lanser P, de Laat SW, Boonstra J, and Defize LH (1996) Parathyroid hormone
357 related peptide mRNA expression during murine postimplantation development:evidence
358 for involvement in multiple differentiation processes. *Int J Dev Biol* 40:599-608.

359

360 Keskinetepe L and Brackett BG (1996) *In vitro* developmental competence of *in vitro*-matured
361 bovine oocytes fertilized and cultured in completely defined media. *Biol Reprod* 55:333-
362 339.

363

364 Keskinetepe L, Burnley CA and Brackett BG (1995) Production of viable bovine blastocyst in
365 defined *in vitro* conditions. *Biol Reprod* 52:1410-1417.

366

367 Krisher RL and Bavister BD (1999) Enhanced glycolysis after maturation of bovine oocytes in
368 vitro is associated with increased developmental competence. *Mol Reprod Dev* 53:19-26.

369

370 Kronenberg HM, Lanske B, Kovacs CS, Chung U-I, Lee K, Segre GV, Schipani E and Juppner H
371 (1998) Functional analysis of the PTH/PTHrP network of ligands and receptors. *Rec Prog Hor*
372 *Res* 53:283-303

373

374 Lanske B, Karaplis AC, Lee K, Luz A, Vortkamp A, Pirro A, Karperien M, Defize LHK, Ho C,
375 Mulligan RC, Abou-Samra AB, Juppner H, Segre GV and Kronenberg HM (1996) PTH/PTHrP
376 receptor in early development and indian hedgehog-regulated bone growth. *Science* 273: 663-666

377

378 Lanske B and Kronenberg HM (1998) Parathyroid hormone-related peptide (PTHrP) and parathyroid
379 hormone (PTH)/PTHrP receptor. *Crit Rev Eukaryotic Gene Express* 8:297-320

380

381 Largo R, Gomez-Garre D, Santos S, Penaranda C, Blanco J, Esbrit P and Egido J (1999) Renal
382 expression of parathyroid hormone-related protein (PTHrP) and PTH/PTHrP receptor in a rat
383 model of tubulointerstitial damage. *Kid Inter* 55:82-90

384

385 Lee K, Brown D, Urena P, Ardaillou N, Ardaillou R, Deeds J and Segre GV (1996) Localization of
386 parathyroid hormone-parathyroid hormone-related peptide receptor mRNA in kidney. *Amer J*
387 *Physiol* 270:F186-F191

388

389 Lee K, Deeds JD, Bond AT, Juppner H, Abou-Samra A-B and Segre GV (1993) In situ localization of
390 PTH/PTHrP receptor mRNA in the bone of fetal and young rats. *Bone* 14:341-345

391

392

393 Lee K, Deeds JD and Segre GV (1995) Expression of parathyroid hormone-related peptide and its
394 receptor messenger ribonucleic acids during fetal development of rats. *Endocrinology* 136:453-
395 463

396

397 Lee K, Lanske B, Karaplis AC, Deeds JD, Kohno H, Nissenson RA, Kronenberg HM and Segre GV
398 (1996) Parathyroid hormone-related peptide delays terminal differentiation of chondrocytes
399 during endochondral bone formation. *Endocrinology* 137:5109-5118

400

401 Li H, Seitz PK, Thomas ML, Selvanayagam P, Rajaraman S and Cooper CW (1995) Widespread
402 expression of the parathyroid hormone-related peptide and PTH/PTHrP receptor genes in
403 intestinal epithelial cells. *Lab Invest* 73:864-870

404

405 Matia-Guiu X, Prat J, Young RH, Capen CC, Rosol TJ, Delellis DA and Scully RE (1994) Human
406 parathyroid hormone-related protein in ovarian small cell carcinoma. An immunohistochemical
407 study. *Cancer* 73:1878-1881

408

409 Moseley JM and Gillespie MT (1995) Parathyroid hormone-related protein. *Crit Rev Clin Lab Sci*

410 32:299-343

411

412 Nowak RA, Haimovici F, Biggers JD, and Erbach GT (1999) Transforming growth factor- β

413 stimulates mouse blastocyst outgrowth through a mechanism involving parathyroid

414 hormone-related protein. *Biol Reprod* 60: 85-93.

415

416 Parrish JJ, Susko-Parrish JL, Leibfried-Rutledge ML, Crister ES, Eyestone WH and First NL

417 (1986) Bovine *in vitro* fertilization with frozen thawed sperm. *Theriogenology* 25: 591-600.

418

419 Pausova Z, Bourdon J, Clayton D, Mattei M-G, Seldin MF, Janicic N, Riviere M, Szpirer J, Levan G,

420 Szpirer C, Goltzman D and Hendy GN (1994) Cloning of a parathyroid hormone/parathyroid

421 hormone-related peptide receptor (PTHrP) cDNA from a rat osteosarcoma (UMR106) cell line:

422 chromosomal assignment of the gene in the human, mouse and rat genomes. *Genomics* 20:20-26.

423

424 Porter SE, Sorenson RL, Dann P, Garcia-Ocana A, Stewart AF and Vasavada RC (1998) Progressive

425 pancreatic islet hyperplasia in the islet-targeted, parathyroid hormone-related protein-

426 overexpressing mouse. *Endocrinology* 139:3743-3751

427

428 Rodan SB, Noda M, Wesolowski G, Rosenblatt M and Rodan GA (1988) Comparison of postreceptor

429 effects of 1-34 human hypercalcemia factor and 1-34 human parathyroid hormone in rat

430 osteosarcoma cells. *J Clin Invest* 81:924-927

431

432 Saeki K, Hoshi M, Leibfried-Rutledge ML and First NL (1991) In vitro fertilization and
433 development of bovine oocytes matured in serum-free medium. *Biol Reprod* 44:256-260.
434

435 Spiegel AM, Saxe AW, Deftos LJ and Brennan MF (1983) Purification of peptides with parathyroid
436 hormone-like bioactivity from human and rat malignancies associated with hypercalcemia. *Hor*
437 *Metab Res* 15: 299-304

438 Stewart AF, Wu T, Goumas D, Burtis WJ and Broadus AE (1987) N-terminal amino acid sequence of
439 two novel tumor-derived adenylate cyclase-stimulating proteins: identification of parathyroid
440 hormone-like and parathyroid hormone-unlike domains. *Biochem Biophys Res Comm* 146:672-
441 678
442

443 Strewler GJ, Stern PH, Jacobs JW, Eveloff J, Klein RF, Leung SC, Rosenblatt M and Nissenson RA
444 (1987) Parathyroid hormone-like protein from human renal carcinoma cells: structural and
445 functional homology with parathyroid hormone. *J Clin Invest* 80:1803-1807
446

447 Tian J, Smogorzewski M, Kedes L and Massry SG (1993) Parathyroid hormone-parathyroid hormone-
448 related protein receptor messenger RNA is present in many tissues besides the kidney. *Amer J*
449 *Nephrol* 13:210-213
450

451 Tucci J and Beck F (1998) Expression of parathyroid hormone-related protein (PTHrP) and the
452 PTH/PTHrP receptor in the rat uterus during early pregnancy and following artificial decidualoma
453 induction. *J Reprod Fertil* 112:1-10

454

455 Urena P, Kong S-F, Abou-Samra A-B, Juppner H, Kronenberg HM, Potts Jr. JT and Segre GV (1993)

456 Parathyroid hormone (PTH)/PTH-related peptide receptor messenger ribonucleic acids are widely

457 distributed in rat tissues. *Endocrinology* 133:617-623

458

459

460

461 van de Stolpe A, Karperien M, Lowik CW, Juppner H, Segre GV, Abou-Samra AB, de Laat SW and

462 Defize LH (1993) Parathyroid hormone-related peptide as an endogenous inducer of parietal

463 endoderm differentiation. *J Cell Biol* 120: 235-243

464

465 Vasavada RC, Garcia-Ocana A, Massfelder T, Dann P and Stewart **AF** (1998) Parathyroid hormone-

466 related protein in the pancreatic islet and the cardiovascular system. *Rec Prog Hor Res* 53:305-

467 340

468

469 Watson AJ, Watson PH, Warnes D, Walker SK, Armstrong DT and Seamark **RF** (1994)

470 Preimplantation development of *in vitro*-matured and *in vitro*-fertilized ovine

471 zygotes: comparison between coculture on oviduct epithelial cell monolayers and culture

472 under low oxygen atmosphere. *Biol Reprod* 50:715-724.

473

474 Watson P, Lazowski D, Han V, Fraher L, Steer B and Hodsman A (1995) Parathyroid hormone restores

475 bone mass and enhances osteoblast insulin-like growth factor I gene expression in

476 ovariectomized rats. Bone 16:357-365.

477

478 Watson PH, Fraher LJ, Hendy GN, Chung U-I, Kisiel M, Natale BV and Hodsman **AB** (2000a) Nuclear
479 localization of the type 1 PTH/PTHrP receptor in rat tissues. J Bone Min Res 15 (6): 1033-1044.

480

481

482

483 Watson PH, Fraher LJ, Natale BV, Kisiel M, Hendy GN and Hodsman AB (2000b) Nuclear
484 localization of the type 1 PTH/PTHrP receptor in MC3T3-E1 cells: association with the cell
485 cycle. Bone 26 (3): 221-225.

486

487 Watson AJ, De Sousa P, Caveney A, Barcroft LC, Natale D, Urquhart J and Westhusin ME (2000)
488 Impact of bovine oocyte maturation media on oocyte transcripts levels, blastocyst development,
489 cell number and apoptosis. Biol Reprod 62:355-364.

490

491 Weaver DR, Deeds JD, Lee K and Segre GV (1995) Localization of parathyroid hormone-related
492 peptide (PTHrP) and PTH/PTHrP receptor mRNAs in rat brain. Mol Brain Res 28:296-310.

493

494 Wiemer KE, Watson AJ, Polanski V, McKenna AI, Fick GH and Schultz GA (1991) Effects of
495 maturation and co-culture treatments on the developmental capacity of early bovine embryos.
496 Mol Reprod Dev 30:330-338.

497

498 Williams ED, Leaver DD, Danks JA, Moseley JM and Martin TJ (1994) Effect of parathyroid
499 hormone-related protein (PTHrP) on the contractility of the myometrium and localisation of
500 PTHrP in the uterus of pregnant rats. *J Reprod Fertil* 102: 209-214.
501

502 Yamamoto M, Harm SC, Grasser WA and Thied MA (1992) Parathyroid hormone-related protein in
503 the rat urinary bladder: a smooth muscle relaxant produced locally in response to mechanical
504 stretch. *Proc Natl Acad Sci (USA)* 89:5326-5330.
505

506 Yang T, Hassan S, Huang YG, Smart AM, Briggs JP and Schnermann JB (1997) Expression of PTHrP,
507 PTH/PTHrP receptor, and Ca(2+)-sensing receptor mRNAs along the rat nephron. *Amer J*
508 *Physiol* 272:F751-F758
509

510 Yasuda T, Banville D, Rabbani SA, Hendy GN and Goltzman D (1989) Rat parathyroid hormone-like
511 peptide: comparison with the human homologue and expression in malignant and normal tissue.
512 *Mol Endocrinol* 3:518-525
513

514 **Figure 1: Detection of PTHrP mRNA in bovine ovary.** Representative photomicrographs
515 display A) sense control, B) detection of PTHrP mRNAs in primordial follicles included oocytes
516 and pre-granulosa cells (arrowheads), C) sense control, D) PTHrP mRNAs in oocyte (o) and
517 granulosa(g) layer of a secondary follicle and stromal (s) tissue, E) PTHrP mRNAs in granulosa
518 and theca (t) cell layers of antral follicles (a), F) PTHrP mRNAs in oocyte, granulosa cells, and
519 theca of antral follicles. Bar = 50 μ M.

520

521 **Figure 2: Detection of PTHR mRNAs in bovine ovary.** Representative photomicrographs
522 display A) sense control, B) detection of PTHR mRNAs in primordial follicles included oocytes
523 and pre-granulosa cells (arrowheads), C) sense control, D) PTHR mRNAs in oocyte (o) and
524 granulosa (g) layer of primordial and secondary follicles and stromal (s) tissue, E) PTHR mRNAs
525 in granulosa and theca (t) layers of antral (a) follicles, F) PTHR mRNAs in oocyte, granulosa
526 cells, and theca of early antral follicle. Bar = 50 μ M

527

528 **Figure 3: Detection of PTHrP polypeptides in bovine ovary** The photomicrographs represent
529 A) mouse normal serum control, B) detection of PTHrP polypeptides in primordial follicles
530 included oocytes and pre-granulosa cells (arrowheads), C) control, D) PTHrP polypeptides in
531 oocyte (o) of secondary follicle, E) PTHrP polypeptides in theca (t) layers of antral follicles, F)
532 PTHrP polypeptides in oocyte of early antral (a) follicle and in ovarian stroma (s). g, granulosa
533 Bar = 50 μ M.

534

535

536

537
538 **Figure 4: Detection of PTHR polypeptide in bovine ovary** The photomicrographs represent A)
539 pre-absorbed control, B) detection of PTHR polypeptides in primordial follicles included oocytes
540 and pre-granulosa cells (arrowheads) and ovarian stroma (s), C) control, D) PTHR polypeptides
541 in oocyte (o) and granulosa (g) cells of secondary follicle and stromal tissue, E) PTHR
542 polypeptides in granulosa and theca (t) layers of antral (a) follicles and stroma, F) PTHR
543 polypeptides in oocyte, granulosa cells and theca of antral follicle. Bar = 50 μ M


544
545

546 **Figure 5: Influence of PTHrP during oocyte maturation on blastocyst development.**

547 Proportion of inseminated oocytes cleaving (cleavage), or reaching the 6-8 cell stage, 6-8/insem,
548 or morula and blastocyst stage of development, mor/bl/insem and blastocysts blast/insem
549 displayed by oocytes matured in 0, 1.59, 15.9, and 159 ng/ml (0, 0.1, 1 and 10 nM, respectively)
550 of human PTHrP (1-141)–supplemented cSOFMaa. Cleavage, development to the 6-8 cell and
551 morulae /blastocyst stages did not vary significantly among the treatment groups. The proportion
552 of zygotes that progressed to the blastocyst stage varied significantly among the treatments. Bars
553 represent the mean \pm SEM of n=4 replicates. Values with different superscript letters are
554 significantly different ($p < 0.05$).


555

Figure 5: Influence of PTHrP During Bovine Oocyte Maturation on Development to the Blastocyst Stage



559 **Figure 6: Influence of PTHrP during oocyte maturation on progression of 6-8 cell embryos**
560 **to the blastocyst stage.** Proportion of 6-8 cell stage embryos (over cleaved embryos, 6-8/clvd),
561 morulae –blastocysts (over cleaved embryos, mor-blast/cleaved) blastocysts (over cleaved
562 embryos, blast/clvd) and 6-8 cell embryos progressing to the blastocyst stage, (blast/6-8)
563 displayed by oocytes matured in 0, 1.59, 15.9, and 159 ng/ml (0, 0.1, 1 and 10 nM, respectively)
564 of murine PTHrP (1-141)–supplemented cSOFMaa. Development to the 6-8 cell and morula-
565 blastocyst stages did not vary significantly among the treatment groups. However the proportion
566 of blastocysts and 6-8 cell embryos that progressed to the blastocyst stage varied significantly.
567 Bars represent the mean \pm SEM of n=4 replicates. Values with different superscript letters are
568 significantly different (p<0.05).
569

Figure 6: Influence of PTHrP During Oocyte Maturation on Development to the Blastocyst Stage



570