

Expression of the *MRP2* Gene-Encoded Conjugate Export Pump in Human Kidney Proximal Tubules and in Renal Cell Carcinoma

THOMAS P. SCHAUB,* JÜRGEN KARTENBECK,† JÖRG KÖNIG,*
HERBERT SPRING,† JOACHIM DÖRSAM,‡ GERD STAEHLER,‡
STEPHAN STÖRKEL,§ WALTER F. THON,|| and DIETRICH KEPPLER*

*Division of Tumor Biochemistry and †Division of Cell Biology, Deutsches Krebsforschungszentrum, Heidelberg; ‡Department of Urology, University of Heidelberg, Germany; §Institute of Pathology, Witten/Herdecke University, Wuppertal Clinic, Wuppertal; and ||Department of Urology, Siloah Clinic, Hannover, Germany.

Abstract. Human kidney proximal tubule epithelia express the ATP-dependent export pump for anionic conjugates encoded by the *MRP2* (*cMRP/cMOAT*) gene (symbol *ABCC2*). *MRP2*, the apical isoform of the multidrug resistance protein, is an integral membrane glycoprotein with a molecular mass of approximately 190 kD that was originally cloned from liver and localized to the canalicular (apical) membrane domain of hepatocytes. In this study, *MRP2* was detected in human kidney cortex by reverse transcription-PCR followed by sequencing of a 826-bp cDNA fragment and by immunoblotting using two different antibodies. Human *MRP2* was localized to the apical brush-border membrane domain of proximal tubules by double and triple immunofluorescence microscopy including laser scanning microscopy.

The expression of *MRP2* in renal cell carcinoma was studied

by reverse transcription-PCR and immunoblotting in samples from patients undergoing tumor-nephrectomy without prior chemotherapy. Clear-cell carcinomas, originating from the proximal tubule epithelium, expressed *MRP2* in 95% (18 of 19) of cases. Immunofluorescence microscopy of *MRP2* in clear-cell carcinoma showed a lack of a distinct apical-to-basolateral tumor cell polarity and an additional localization of *MRP2* on intracellular membranes. *MRP2*, the first cloned ATP-dependent export pump for anionic conjugates detected in human kidney, may be involved in renal excretion of various anionic endogenous substances, xenobiotics, and cytotoxic drugs. This conjugate-transporting ATPase encoded by the *MRP2* gene has a similar substrate specificity as the multidrug resistance protein *MRP1*, and may contribute to the multidrug resistance of renal clear-cell carcinomas.

Active secretion of endogenous and xenobiotic organic anions from blood into urine is a major physiologic function of kidney proximal tubule epithelia (1–6). Many different synthetic and biologic amphiphilic anions have long been known as substrates for basolateral uptake and luminal excretion, including *para*-aminohippurate (PAH), phenol red, carboxyfluorescein, bimane glutathione conjugates, and leukotriene C₄ (2–6). In particular, environmentally occurring cytotoxic, potentially mutagenic, and carcinogenic organic anions, like polychlorinated aromatic compounds and cytostatic drugs such as methotrexate, are actively eliminated from the blood circulation via the proximal tubules of the kidney (2–6). An uptake transporter accepting a large number of structurally diverse organic

anions, termed OAT1, has been cloned from rat kidney and localized to the basolateral membrane domain of proximal tubule epithelia (5,6).

Recently, we demonstrated that *MRP2*, a novel ATP-dependent transporter for amphiphilic organic anions, is expressed in the apical membrane domain of rat proximal tubule epithelia where it may function in the export of amphiphilic anionic compounds across the luminal membrane (7). The multidrug resistance proteins *MRP1* and *MRP2*, two integral membrane glycoproteins belonging to the ATP-binding cassette superfamily of transporters (8–11), have been identified as conjugate-transporting ATPases (12–18). *MRP2*, alternatively termed canalicular *MRP* (*cMRP*) (9) or canalicular multispecific organic anion transporter (*cMOAT*) (10,11), has been cloned from rat (9,10) and human liver (9,18) and tumor cells (11). These primary active transport proteins were functionally characterized as unidirectional export pumps for conjugates of lipophilic compounds with glutathione, glucuronate, or sulfate (12–18). The substrate specificities of *MRP1* and *MRP2* are similar, despite the fact that both proteins share only 49% amino acid identity (reviewed in reference (18)). The *MRP2*

Received February 3, 1998. Accepted December 21, 1998.

Correspondence to Dr. Dietrich Keppler, Deutsches Krebsforschungszentrum, Division of Tumor Biochemistry, Im Neuenheimer Feld 280, D-69120 Heidelberg, Germany. Phone: 49 6221 422400; Fax: 49 6221 422402; E-mail: d.keppler@dkfz-heidelberg.de

1046-6673/1006-1159

Journal of the American Society of Nephrology

Copyright © 1999 by the American Society of Nephrology

(*cMRP/cMOAT*) gene has been localized to chromosome 10q24 (11). On the mRNA level, human *MRP2* is expressed predominantly in the liver, in addition to kidney and small intestine (11,18–20).

In the present study, we extended our work on expression and localization of *MRP2* in the rat kidney (7) to human kidney obtained from patients undergoing tumor-nephrectomy. Furthermore, *MRP2* expression was studied in renal cell carcinomas, predominantly of the most frequent subtype, the clear-cell carcinoma (21–23). Primary and metastasizing renal cell carcinomas often exhibit an intrinsic resistance to different chemotherapeutic agents (23). Clinical trials with the cytostatic drug vinblastine, including dexverapamil as a modulator of MDR1 P-glycoprotein-mediated multidrug resistance, showed no significant improvement of survival rates of patients with metastasizing renal cell carcinoma (24).

Materials and Methods

Human Tissue Samples

Renal cell carcinoma and corresponding nontumorous kidney cortex samples were obtained from 21 patients (mean age 63.5 yr; range, 47 to 89). None of the patients had received chemotherapy. After tumor-nephrectomy, the neoplastic and the normal kidney tissues were separated, snap-frozen in isopentane precooled in liquid nitrogen, and further processed for reverse transcription (RT)-PCR, immunoblotting, or immunofluorescence microscopy. The warm ischemia period of the tissues ranged from 15 to 30 min until freezing. Informed consent for tissue delivery was obtained from patients before tumor-nephrectomy.

Materials

The protease inhibitors phenylmethylsulfonyl fluoride, aprotinin, leupeptin, pepstatin, and protein standard mixture (M_r 26,600 to 180,000) were purchased from Sigma Chemicals (Deisenhofen, Germany). RNase inhibitor (RNasin), StrataScriptTM Moloney murine leukemia virus reverse transcriptase, *Taq* DNA polymerase, and β -actin primers were from Stratagene (Heidelberg, Germany). Other chemicals were of analytical grade and delivered by Merck (Darmstadt, Germany).

Antibodies

The EAG5 polyclonal antibody was raised in rabbits against a synthetic peptide containing the amino acid sequence EAGIENVN-STKF at the carboxy terminus of the human *MRP2* protein, as described previously (9,25). The polyclonal antibody MLE was obtained by immunization of rabbits against the amino-terminal synthetic peptide of the human *MRP2* (MLEKFCNSTFWNSSFLD-SPEADLPLC) (9,18,25). The antibodies EAG5 and MLE did not cross-react with *MRP1*. The monoclonal mouse-anti-CD26 antibody directed against human dipeptidyl-peptidase IV was purchased from Dianova (Hamburg, Germany). Goat anti-rabbit secondary antibodies coupled to Texas red were from Dianova. The monoclonal mouse anti-calbindin-D (28 kD) antibody was purchased from Sigma (St. Louis, MO), and the polyclonal sheep antibody to Tamm-Horsfall glycoprotein was from Biogenesis (Poole, United Kingdom). Donkey anti-sheep secondary antibodies coupled to Rhodamine Red-X and amino-methylcoumarin acetate-conjugated goat anti-mouse antibodies were from Jackson ImmunoResearch Laboratories (West Grove, PA). Goat anti-rabbit secondary antibodies coupled to Texas red were

from Dianova. Goat anti-mouse secondary antibody coupled to FITC or cyanin 2-conjugate (Cy2) was from Biotrend (Cologne, Germany).

RNA Isolation, RT-PCR, and Subcloning

Total RNA was isolated from nontumorous kidney cortex and corresponding renal cell carcinomas by a guanidinium thiocyanate lysis procedure with subsequent purification by centrifugation on cesium chloride (26). To prevent amplification on genomic DNA, total RNA was treated, before reverse transcription, with 10 U of DNase I in 50 μ l of digestion buffer (100 mM sodium acetate, pH 5.0, 5 mM $MgSO_4$, and 40 U of the RNase inhibitor RNasin) at 37°C for 1 h. Then, total RNA (5 μ g) was reverse-transcribed using an oligo(dT)₁₈ primer for *MRP1* as described (7), or a sequence-specific reverse primer for *MRP2* (bases 4284 to 4267). The primer pairs used for PCR detection of the respective cDNA were based on the sequences from the GenBank/EMBL/DataBank with the accession nos. L05628 (*MRP1*; reference (8)) and X96395 (*MRP2*; reference (9)). For *MRP1*, the sense primer was 5'-ATCAAGACCGCTGTCAT-TGG-3' (nucleotides 1183–1202), and the antisense primer was 5'-GAGCAAGGATGACTTGCAGG-3' (nucleotides 1363–1344). For *MRP2*, the sense primer was 5'-CCACAGGCCGGATTGTG-3' (nucleotides 3227–3243), and the antisense primer was 5'-AAGATTCT-GAAGAGGCAG-3' (nucleotides 4052–4035). The commercial β -actin control primer pair was obtained from Stratagene. For each PCR, the cycling conditions were as follows: 94°C for 45 s; 60°C for 60 s; 72°C for 90 s (35 cycles). Amplified cDNA fragments were subcloned into the pCR2.1 vector (Invitrogen, Leek, The Netherlands). Positive clones were sequenced by the dideoxynucleotide chain termination method of Sanger using [α -³⁵S]dATP and the sequencing kit from Pharmacia Biotech (Freiburg, Germany). Dried gels were exposed to Kodak BioMax MR-1 film obtained from Sigma (Deisenhofen, Germany).

Immunoblotting

Human kidney cortex and inner medulla or renal cell carcinoma tissue (about 1.0 g each) was homogenized during thawing in 10 ml of lysis buffer (10 mM KCl, 1.5 mM $MgCl_2$, 10 mM Tris/HCl, pH 7.4) and 0.5% (wt/vol) sodium dodecyl sulfate supplemented with four proteinase inhibitors at the following final concentrations: 1 mM phenylmethylsulfonyl fluoride, 0.3 μ M aprotinin, 1 μ M pepstatin, and 1 μ M leupeptin at 4°C (7). The resulting suspension was further processed for crude membrane preparation as recently described (7), and 25 or 50 μ g of protein was loaded onto a 7.5% (wt/vol) sodium dodecyl sulfate-polyacrylamide gel, without boiling, and subjected to electrophoresis (27). After electrotransfer onto nitrocellulose membranes (Schleicher & Schuell, Dassel, Germany), the blots were blocked with Tris-buffered saline containing 0.05% Tween 20 and 10% (wt/vol) low-fat dry milk (Glücksklee, Frankfurt, Germany) for 1 h at room temperature and probed overnight with the polyclonal *MRP2* antibodies EAG5 and MLE at a dilution of 1:20,000 and 1:10,000, respectively. Antibody binding was visualized with a horse-radish peroxidase-conjugated goat anti-rabbit antibody (Bio-Rad, Munich, Germany) diluted 1:1000, followed by enhanced chemiluminescence detection with exposure on HyperfilmTM-MP (Amersham-Buchler, Braunschweig, Germany). A semiquantitative score was used for the signal intensity of the 190-kD band corresponding to *MRP2* after application of 50 μ g of protein: +++, strong; ++, positive; +, weak; –, no reactivity for *MRP2* using the EAG5 antibody. Canalicular membrane from human liver (9,18) served as an *MRP2*-rich positive control.

Immunofluorescence Microscopy

Samples from frozen kidney cortex and corresponding renal cell carcinoma were used for single and double-label immunofluorescence microscopy. Unfixed tissue sections (4 to 5 μ m thick) were prepared with a cryotome (Leica, FrigoCut 2800E; Nussloch, Germany) and air-dried for 2 to 4 h at room temperature. Before the incubation with the antibodies, the sections were fixed in 100% acetone precooled at -20°C for 10 min. For double- or triple-label immunofluorescence microscopy, the primary (rabbit EAG5, 1:100; mouse anti-CD26, 1:100 or 1:200; mouse anti-calbindin-D, 1:50 or 1:100; sheep anti-Tamm-Horsfall glycoprotein, 1:80 to 1:120) as well as the secondary antibodies (goat anti-mouse or anti-rabbit, or donkey anti-sheep) were applied simultaneously. Incubation with the primary and secondary antibodies was for 30 to 40 min each at room temperature. Unbound antibodies were removed by several washes with phosphate-buffered saline after each incubation step. As negative controls for the specific reactivity of the EAG5 and MLE antibodies, both preimmune sera were used. After a final wash with distilled water, the air-dried sections were mounted with Elvanol (CTI, Idstein/Taunus, Germany). As a further control, peptide competition with the EAG5 antibody was performed. For preabsorption of the EAG5 antibody, the synthetic peptide used to create this antibody (amino acids 1534–1545 of MRP2) was solubilized at concentrations of 100 and 10 μM in the antibody solution and allowed to bind for 30 min at room temperature before its application to the tissue sections. Fluorescence imaging micrographs were taken with a Zeiss-Axiophot microscope (Carl-Zeiss, Jena, Germany). FITC or Cy2 and Texas red were excited at 450 to 490 nm and visualized with a 515 to 565 and 610 nm bandpass filter, respectively. Photographs were taken on Kodak T-MAX 400

film. Confocal images in double or triple fluorescence were obtained with a Carl Zeiss LSM 510 ultraviolet laser scanning microscope. For our photographs, we used an argon ion laser (488 nm) and a HeNe laser (543 nm) with the corresponding barrier filters in the double track mode of the instrument (*i.e.*, each scan line was alternatively illuminated only with one wavelength to avoid bleed-through of the fluorescence dyes) for double fluorescence. For simultaneous triple fluorescence, we used an additional argon ion laser (364 nm) for the excitation of amino-methylcoumarin acetate.

Histopathology

Tissue samples for diagnostic histopathology were formaldehyde-fixed, paraffin-embedded, sectioned, and stained with hematoxylin and eosin. All tumor specimens were reviewed independently by a reference pathologist (S.S.). Tumor progression was estimated according to the postoperative tumor staging classification (28). Grading of tumor tissue was recorded as follows: well differentiated (G1), moderately differentiated (G2), and poorly differentiated (G3).

Table 1. MRP2 immunoreactivity in patients with renal-cell carcinoma^a

Patient	Stage (TNM) ^b	Grade (G) ^c	Histopathology	MRP2 Expression ^d	
				RCC	Kidney Cortex
1	pT2N0M1	G1	Clear-cell	+++	+++
2	pT3N0M0	G1	Clear-cell	+++	+++
3	pT2N0M0	G1	Clear-cell	+++	++
4	pT2N0M0	G1	Clear-cell	+++	++
5	pT2N0M0	G2	Clear-cell	+++	— ^e
6	pT2N0M0	G1	Clear-cell	++	— ^e
7	pT2N0M0	G1	Clear-cell	++	+++
8	pT2N0M1	G2	Clear-cell	++	+++
9	pT3N0M0	G2	Clear-cell	++	+++
10	pT2N0M2	G2	Clear-cell	++	+++
11	pT3N0M0	G2	Clear-cell	++	+++
12	pT2N0M0	G2	Clear-cell	++	+++
13	pT2N0M1	G2	Clear-cell	++	+++
14	pT2N0M0	G2	Clear-cell	++	++
15	pT3N2M1	G3	Clear-cell	++	+++
16	pT2N0M0	G3	Clear-cell	+	+++
17	pT2N0M1	G3	Clear-cell	+	+++
18	pT2N0M0	G3	Clear-cell	+	+++
19	pT4N0M0	G3	Clear-cell	—	+++

^a Patients underwent tumor surgery without preoperative chemotherapy or irradiation treatment. The mean age of all patients undergoing tumor-nephrectomy was 63.5 yr (range, 47 to 89). Immunoblotting was performed with the EAG5 antibody as shown in the examples given in Figures 2 and 3.

^b pTNM, postoperative tumor staging (28).
^c Pathohistologic tumor grading.

^d Immunoreactivity for MRP2: +++, strong; ++, positive; +, weak; —, negative.

^e No kidney tissue available because of nephron-sparing surgery.




Figure 1. Reverse transcription-PCR analysis of *MRP2* and *MRP1* mRNA expression in human kidney cortex (Cortex) and in renal cell carcinomas (RCC). Cortex 12 and RCC 12 were obtained from patient 12 diagnosed with clear-cell carcinoma (see Table 1); patient 20 suffered from chromophilic renal cell carcinoma. For PCR analysis, two pairs of primers were derived from the human *MRP2* and *MRP1* cDNA sequences yielding an 826-bp and a 181-bp fragment, respectively. A 661-bp fragment for β -actin generated with specific primers fitting the human sequence demonstrates the integrity of the isolated RNA. *MRP1* and *MRP2* amplification products were obtained for normal kidney cortex (patients 12 and 20) and from a corresponding clear-cell carcinoma sample (RCC 12). An amplification product for the *MRP2* cDNA fragment was not obtained with the chromophilic renal cell carcinoma (RCC 20).

Results

Detection of MRP2 mRNA in Human Kidney and in Renal Cell Carcinomas

The expression of *MRP2* (symbol *ABCC2*) and *MRP1* (symbol *ABCC1*) genes in kidney cortex and renal cell carcinoma from tumor-nephrectomized patients (cortex 12 and 20; renal cell carcinoma 12 and 20) was analyzed by reverse transcription and PCR amplification of cDNA fragments generated from isolated total RNA (Figure 1). The amplification products showed the expected size of 826 bp for *MRP2* and 181 bp for *MRP1* (Figure 1). As an internal control, the β -actin 661-bp cDNA fragment demonstrated the integrity of the isolated total RNA from the tissue specimens (Figure 1). Amplification products corresponding to *MRP2* and *MRP1* cDNA fragments were obtained from normal kidney cortex (cortex 12 and 20). The *MRP2* and *MRP1* cDNA fragments were further identified by subcloning and sequencing of 826 nucleotides for *MRP2* and of 181 nucleotides for *MRP1*. Both the *MRP2* and *MRP1* cDNA fragments showed a nucleotide sequence identity of 99.5% compared with the cloned sequences from human liver (*MRP2*; references (9) and (18)) and H69AR cells (*MRP1*; reference (8)). We have analyzed the mRNA from a total of six well preserved renal clear-cell carcinomas and consistently detected the amplification product of the *MRP2* cDNA. *MRP2* mRNA was below detectability in the two chromophilic renal cell carcinomas studied, but we did obtain an amplification product for *MRP1* mRNA (Figure 1).

Immunoblot Analysis of MRP2 in Human Kidney and Renal Cell Carcinomas

Immunoblots on crude membrane preparations from kidney cortex and from the corresponding renal cell carcinoma samples (Table 1) were probed with polyclonal antibodies directed

against human MRP2. The antibody EAG5 indicated a strong expression of the 190-kD glycoprotein in all kidney cortex samples (Figures 2 and 3; Table 1). In addition, the MLE antibody, directed against the amino terminus of MRP2, detected the protein at a molecular mass of approximately 190 kD in kidney cortex (Figures 2 and 3c).

Eighteen of 19 clear-cell carcinoma samples (95%) were positive for MRP2 expression using the antibody EAG5. This result was confirmed by the MLE antibody. Clear-cell carcinomas with G1 grading semiquantitatively showed a more intense immunoreactivity of MRP2 than the less differentiated tumors graded G3 (Table 1).




Figure 3. Immunoblot of MRP2 expression in crude membrane fractions from human kidney cortex, inner medulla, and renal cell carcinoma from patients with clear-cell carcinoma (patients 3 and 12). MRP2 was detected with antibodies EAG5 and MLE as a protein of approximately 190 kD in all kidney cortex samples and in clear-cell carcinoma (RCC 12). No reactivity for MRP2 was seen in kidney medulla regions (A) and in the two chromophilic renal cell carcinomas studied (RCC 20 in B and C).




Figure 2. Immunoblot showing MRP2 expression in crude membranes from human kidney cortex and a corresponding clear-cell renal carcinoma (RCC 4) (patient 4; Table 1). Detection of MRP2 with protein A-Sepharose-purified antibodies MLE and EAG5 shows the protein of approximately 190 kD.




Figure 4. Single-label (a) and double-label (b through g) immunofluorescence microscopy of MRP2 and dipeptidyl-peptidase IV in human adult kidney cortex. Tissue sections (from samples 1 and 5; Table 1) show fluorescent signals with the antibody EAG5 (EAG) directed against human MRP2 (a, b, d, and f). As a marker protein for the apical (luminal) membrane domain of renal proximal tubules, the monoclonal antibody anti-CD26 directed against human DPP-IV (DPP) was applied (29). Proximal tubule epithelia near the glomerulus (G) showed staining at the luminal brush-border membrane with the EAG5 (filled triangles in a, b, d, and f) and anti-DPP-IV antibodies (filled triangles in c, e, and g). Both antibodies did not react with tubular segments distal from the proximal tubules (open triangles). Poor immunoreactivity in some areas is a result of the period of warm ischemia during surgery and prior to freezing of the tissue (filled circles in d and e). Occasionally, autofluorescent particles can be detected in some tissue slices (arrowheads in a). Bars: 50 μ m in a through e; 100 μ m in f and g.

Localization of MRP2 and Dipeptidyl-Peptidase IV in Human Kidney and Renal Cell Carcinomas

MRP2 was localized in human kidney cortex by immunofluorescence microscopy. The EAG5 antibody indicated a fluorescence signal only in proximal tubule epithelia. The reaction product was predominantly localized in the luminal (apical) brush-border membrane domain (Figure 4, a, b, d, and f). The nephron segments following the proximal tubule, *i.e.*, thin limbs of Henle's loop, thick ascending limb of Henle's loop including the macula densa, distal collecting tubule, and the cortical and medullary collecting ducts, were negative for MRP2 (Figure 4). Dipeptidyl-peptidase IV (CD 26; EC 3.4.14.5) (DPPIV), a marker for the apical membrane domain (29), clearly colocalized with MRP2 on proximal tubule brush-border membranes (Figure 4, b through g).

MRP2 localization in different specimens of clear-cell carcinomas indicated a loss of the regular apical-to-basolateral polarity observed in normal proximal tubule epithelial cells (Figure 5, a and c). An apparent localization of MRP2 on intracellular membranes was detected in many of the clear-cell carcinoma cells. MRP2-positive cells also expressed DPPIV as shown by double-label immunofluorescence (some are indi-

cated by arrowheads in Figure 5). In a high number of clear-cells, reaction products with DPPIV were localized to intracellular membranes. A localization of MRP2 together with DPPIV to the plasma membrane of the tumor cells was observed only occasionally (*e.g.*, small arrows in Figure 5d). In many clear-cell carcinomas, a dissociation of MRP2 and DPPIV expression was observed, particularly with cells that were DPPIV-positive but MRP2-negative (Figure 5).

The immunofluorescence specificity of the EAG5 antibody was tested by two different experiments. Either the secondary antibody was used without prior incubation with primary EAG5 antibody, or the EAG5 antibody was preabsorbed with the synthetic peptide that had been used to raise this antibody. In both experiments, the normal staining of the apical membrane domain of proximal tubules (Figure 4, a, b, and d, and Figure 6a) did not appear (Figures 6, b and c). This proved that the secondary antibody used did not create an unspecific staining pattern and that the specific antibody reaction could be blocked by preincubation with the proper peptide. The specificity of this reaction was further confirmed by the fact that preincubation with EAG5 peptide did not influence the reac-




Figure 5. Double-label immunofluorescence microscopy of MRP2 and DPPIV in a compact-growing clear-cell renal carcinoma. Tissue sections (from tumor sample 15; Table 1) were stained with the antibodies EAG5 (EAG; a and c) and anti-DPPIV (DPP; b and d). Although only a subpopulation of the carcinoma cells expressed MRP2 (some corresponding cells in Panels a and b and Panels c and d that show colocalization with both antibodies are indicated by arrowheads in a and c), most of the clear cells were positive for DPPIV (b and d). Small arrow in Panel a points to cellular extensions of MRP2-positive cells. Bar, 50 μ m.




Figure 6. Single-label (a and b) and double-label (c and d) immunofluorescence microscopy of frozen sections of adult human kidney cortex for control of the specificity of the EAG5 antibody. Fluorescent signals at luminal membrane domains of renal proximal tubules can be seen after sequential application of the primary (EAG5) antibody and cyanin 2-coupled secondary (Cy) antibody (a). No staining appears when the secondary (Cy) antibody is used alone (b). The immunofluorescence reaction of the EAG5 antibody is completely lost when the antibody was preincubated with a 10 μ M solution of the synthetic peptide that had been used to raise this antibody (c; EAG & Pep), whereas the antibody reaction against DPPIV remained unchanged (d). G, glomerulus. Filled triangles in Panels c and d indicate some corresponding proximal tubules. Bar, 100 μ m.

tivity of the DPPIV antibody (Figure 6d) as shown by double-label immunofluorescence (Figure 6, c and d).

To further support the finding that MRP2 is restricted to proximal tubules, we performed double- and triple-label experiments using antibodies to MRP2 in combination with antibodies against Tamm-Horsfall glycoprotein, marking the thick ascending limb and the early distal convoluted tubule (30,31). In addition, we used antibodies to calbindin, reacting with the distal convoluted and connecting tubule (32,33). All three antibodies stained different nephron segments in the kidney cortex (Figure 7, a and b, and Figure 8, a through d). In medullary regions, however, only antibodies to Tamm-Horsfall glycoprotein reacted positively (Figure 7, c and d). These results demonstrate that human MRP2 is detectable only in the apical membrane domain of proximal tubules.

Discussion

In the present study, we demonstrate the expression of MRP2 in the apical membrane of human kidney proximal tubule epithelia. Moreover, we describe the expression of MRP2 in clear-cell renal carcinoma. Renal cell carcinoma is the most common malignancy of human adult kidney and accounts for 2 to 3% of all cancers (23). Approximately 80% of all renal cell carcinomas belong to the clear-cell subtype originating from proximal tubular epithelia (21–23). The intrinsic multidrug resistance phenotype of renal cell carcinomas

is still a major obstacle for chemotherapy of patients with primary and metastasizing adult kidney cancer (24,34).

MRP2 was detected in all nontumorous, human kidney cortex samples (Table 1), and its localization was confined to the apical membrane domain of proximal tubule epithelia (Figures 4 and 8). MRP2, also termed the hepatocyte canalicular multidrug resistance protein, cMRP (9,35), or canalicular multispecific organic anion transporter, cMOAT (10,11), has a broad substrate specificity for endogenous and xenobiotic amphiphilic anions similar to the substrate specificity established for the MRP1 transporter (12–17,36). This substrate specificity may account at least for some of the hitherto described active transport processes for organic anions from renal proximal tubules into the luminal space (1–4). Conjugation or complex formation of endogenous compounds and cytotoxic agents with glutathione and other anionic groups is often a prerequisite for transport by MRP1 and MRP2 (12–18,36). Transfection studies with *MRP1* cDNA in HeLa cells (reviewed in reference (16)) and work by Koike *et al.* (20) suggest that not only MRP1 but also MRP2 expression confers multidrug resistance to cancer cell lines. In the present study, we observed that MRP2 is expressed in 95% of the renal cell carcinomas with a clear-cell pathohistology (Table 1). The localization of MRP2 in clear-cells significantly differs from nontumorous kidney proximal tubule epithelia because of the predominant intracellular localization of this transport protein. It is presently not known




Figure 7. Double-label immunofluorescence of frozen sections of adult human kidney cortex (Co; a and b) and medulla (Me; c and d) after reaction with EAG5 for MRP2 (E; a and c) and Tamm-Horsfall (T; b and d) antibodies. In cortex areas, the antibodies clearly reacted with different segments of the nephron (a and b). Some structures including the glomeruli (G) were negative for both antibodies (arrowheads in a and b). In medullary regions (Me; c and d), the EAG5 (E) antibody to MRP2 was negative (c), while the antibody to Tamm-Horsfall protein (T) strongly reacted with various tubular segments (d). Bar (shown in b), 100 μ m; same magnification for all panels.

to what extent the MRP2 in intracellular membranes can be sorted to the plasma membrane of tumor cells when exposed to cytostatic drugs. However, recent studies by Cui *et al.* (37) indicate that MRP2 expression in cells confers multidrug resistance. Our semiquantitative analyses shown in Table 1 suggest that MRP2 expression might also correlate with the differentiation status of renal clear-cell carcinomas. MRP2 may be below detectability in dedifferentiated clear-cell tumors with G3 grading. Further investigations are required to clarify whether MRP2 expression is negatively correlated with the dedifferentiation of renal clear-cell carcinoma or with chromosomal aberrations during tumor progression.

In conclusion, we propose that MRP2 under physiologic conditions functions in renal elimination of endogenous and xenobiotic amphiphilic anions from blood into urine. MRP2 has the capacity to confer multidrug resistance (37) and, when

sorted to the plasma membrane, may contribute to the resistance of renal clear-cell carcinoma to a wide variety of cytotoxic chemotherapeutic agents. Future studies should define whether MRP2 is the only ATP-dependent export pump secreting amphiphilic organic anions into the lumen of kidney proximal tubules or whether additional members of the MRP family of transporters, which were identified recently (19), contribute to this process.

Acknowledgments

We thank the following colleagues for their contributions to this study: Dr. H. R. Rackwitz (Division of Cell Biology, Deutsches Krebsforschungszentrum, Heidelberg) for synthesizing peptides and conjugates for immunization; Dr. W. Hagmann (Division of Tumor Biochemistry, Deutsches Krebsforschungszentrum, Heidelberg) for his help in the generation of the MLE antibody; J. Arlt and K. Goetzke




Figure 8. Confocal laser scanning micrographs of frozen sections of human kidney after double (a through c) and triple (d) immunofluorescence labeling. Positive reactions with the EAG5 antibody to MRP2 (E; a and c) are seen on proximal tubules (some are marked with filled triangles in a), while other tubular segments are unstained (open triangles in a). After application of calbindin antibodies (C; b and c), positive reactions are seen on distal convoluted and connecting tubules (open triangles in b), leaving other tubular segments unstained (closed triangles in b). The merged image of a and b (c) reveals that the staining patterns of the EAG5 antibody for MRP2 (E, green fluorescence) and the calbindin antibody (C, red fluorescence) do not overlap. Triple-label fluorescence, shown in Panel d, with antibodies to calbindin (C, blue fluorescence), Tamm-Horsfall protein (T, red/orange fluorescence), and MRP2 (E, green fluorescence), demonstrates that the three different antigens are located at different cortical tubular segments of the nephron. G, glomerulus. Bars (shown in b and d), 100 μ m; same magnification for panels a through c.

(Division of Cell Biology, Deutsches Krebsforschungszentrum, Heidelberg) for their help in the immunofluorescence microscopy studies; and Dr. H. Ostertag (Institute of Pathology, Nordstadt Clinic, Hannover, Germany) and Dr. W. Hofmann (Institute of Pathology, Uni-

versity of Heidelberg) for the histologic examination of renal cell carcinomas. We thank the Deutsche Forschungsgemeinschaft, Bonn, for supporting parts of this work through the "Graduiertenkolleg Experimentelle Nieren- und Kreislaufforschung," Heidelberg, through

SFB 601, and through SFB 352/B3. Additional parts of this work were supported by the Forschungsschwerpunkt Transplantation, Heidelberg.

References

- Pritchard JB, Miller DS: Renal secretion of organic anions and cations. *Kidney Int* 49: 1649–1654, 1996
- Miller DS, Letcher S, Barnes DM: Fluorescence imaging study of organic anion transport from renal proximal tubule cell to lumen. *Am J Physiol* 40: F508–F520, 1996
- Masereeuw R, Russel FGM, Miller DS: Multiple pathways of organic anion secretion in renal proximal tubule revealed by confocal microscopy. *Am J Physiol* 271: F1173–F1182, 1996
- Ullrich KJ, Rumrich G: Luminal transport step of *para*-aminohippurate (PAH): Transport from PAH-loaded proximal tubular cells into the tubular lumen of the rat kidney in vivo. *Pflügers Arch* 433: 735–743, 1997
- Sekine T, Watanabe N, Hosoyamada M, Kanai Y, Endou H: Expression cloning and characterization of a novel multispecific organic anion transporter. *J Biol Chem* 272: 18526–18529, 1997
- Sweet DH, Wolff NA, Pritchard JB: Expression cloning and characterization of ROAT1: The basolateral organic anion transporter in rat kidney. *J Biol Chem* 272: 30088–30095, 1997
- Schaub TP, Kartenbeck J, König J, Vogel O, Witzgall R, Kriz W, Keppler D: Expression of the conjugate export pump encoded by the *mrp2* gene in the apical membrane of kidney proximal tubules. *J Am Soc Nephrol* 8: 1213–1221, 1997
- Cole SPC, Bhardwaj G, Gerlach JH, Mackie JE, Grant CE, Almquist KC, Stewart AJ, Kurz EU, Duncan AMV, Deeley RG: Overexpression of a transporter gene in a multidrug-resistant human lung cancer cell line. *Science* 258: 1650–1654, 1992
- Büchler M, König J, Brom M, Kartenbeck J, Spring H, Horie T, Keppler D: cDNA cloning of the hepatocyte canalicular isoform of the multidrug resistance protein, cMRP, reveals a novel conjugate export pump deficient in hyperbilirubinaemic mutant rats. *J Biol Chem* 271: 15091–15098, 1996
- Paulusma CC, Bosma PJ, Zaman GJR, Bakker CTM, Otter M, Scheffer GL, Scheper RJ, Borst P, Oude Elferink RPJ: Congenital jaundice in rats with a mutation in a multidrug resistance-associated protein gene. *Science* 271: 1126–1128, 1996
- Taniguchi K, Wada M, Kohno K, Nakamura T, Kawabe T, Kawakami M, Kagotani K, Okumura K, Akiyama S-I, Kuwano M: A human canalicular multispecific organic anion transporter (cMOAT) gene is overexpressed in cisplatin-resistant human cancer cell lines with decreased drug accumulation. *Cancer Res* 54: 4124–4129, 1996
- Jedlitschky G, Leier I, Buchholz U, Center M, Keppler D: ATP-dependent transport of glutathione S-conjugates by the multidrug resistance-associated protein. *Cancer Res* 54: 4833–4836, 1994
- Leier I, Jedlitschky G, Buchholz U, Cole SPC, Deeley RG, Keppler D: The *MRP* gene encodes an ATP-dependent export pump for leukotriene C₄ and structurally related conjugates. *J Biol Chem* 269: 27807–27810, 1994
- Müller M, Meijer C, Zaman GJR, Borst P, Scheper N, Mulder H, de Vries EGE, Jansen PLM: Overexpression of the gene encoding the multidrug resistance-associated protein results in increased ATP-dependent glutathione S-conjugate transport. *Proc Natl Acad Sci USA* 91: 13033–13037, 1994
- Jedlitschky G, Leier I, Buchholz U, Barnouin K, Kurz G, Keppler D: Transport of glutathione, glucuronate, and sulfate conjugates by the *MRP* gene-encoded conjugate export pump. *Cancer Res* 56: 988–994, 1996
- Loe DW, Deeley RG, Cole SPC: Biology of the multidrug resistance-associated protein, MRP. *Eur J Cancer* 32A: 945–957, 1996
- Chang X-B, Hou Y-X, Riordan JR: ATPase activity of purified multidrug resistance-associated protein. *J Biol Chem* 272: 30962–30968, 1997
- Keppler D, König J: Expression and localization of the conjugate export pump encoded by the *MRP2* (cMRP/cMOAT) gene in liver. *FASEB J* 11: 509–516, 1997
- Kool M, de Haas M, Scheffer GL, Scheper RJ, van Eijk MJT, Juijn JA, Baas F, Borst P: Analysis of expression of *cMOAT* (*MRP2*), *MRP3*, *MRP4*, and *MRP5*, homologues of the multidrug resistance-associated protein gene (*MRP1*), in human cancer cell lines. *Cancer Res* 57: 3537–3547, 1997
- Koike K, Kawabe T, Tanaka T, Toh S, Uchiumi T, Wada M, Akiyama S-I, Ono M, Kuwano M: A canalicular multispecific organic anion transporter (cMOAT) antisense cDNA enhances drug sensitivity in human hepatic cancer cells. *Cancer Res* 57: 5475–5479, 1997
- Thoenes W, Störkel S, Rumpelt HJ: Histopathology and classification of renal cell tumors (adenomas, oncocytomas and carcinomas): The basic cytological and histopathological elements and their use for diagnostics. *Pathol Res Pract* 181: 125–143, 1986
- Störkel S, van den Berg E: Morphological classification of renal cancer. *World J Urol* 13: 153–158, 1995
- Motzer RJ, Bander NH, Nanus DM: Renal-cell carcinoma. *N Engl J Med* 335: 865–875, 1996
- Motzer RJ, Lyn P, Fischer P, Lianes P, Ngo RL, Cordon-Cardo C, O'Brien JP: Phase I/II trial of dexverapamil plus vinblastine for patients with advanced renal cell carcinoma. *J Clin Oncol* 13: 1958–1965, 1995
- Schnölzer M, Alewood P, Jones A, Alewood D, Kent SBH: In situ neutralization in Boc-chemistry solid phase peptide synthesis: Rapid, high yield assembly of difficult sequences. *Int J Pept Protein Res* 40: 180–193, 1992
- Chomczynski P, Sacchi N: Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162: 156–159, 1987
- Towbin H, Staehelin T, Gordon J: Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: Procedure and some applications. *Proc Natl Acad Sci USA* 76: 4350–4354, 1979
- Sobin LH, Hermanek P, Hutter RV: TNM classification of malignant tumors: A comparison between the new (1987) and the old editions. *Cancer* 61: 2310–2314, 1988
- Kugler P, Wolf G, Scherberich J: Histochemical demonstration of peptidases in the human kidney. *Histochemistry* 83: 337–341, 1985
- Sikri KL, Alexander DP, Foster CL: Localization of Tamm-Horsfall glycoprotein in the normal rat kidney and the effect of adrenalectomy on its localization in the hamster and rat kidney. *J Anat* 135: 29–45, 1982
- Malagolini N, Cavallone D, Serafini-Cessi F: Intracellular transport, cell-surface exposure and release of recombinant Tamm-Horsfall glycoprotein. *Kidney Int* 52: 1340–1350, 1997
- Kumar R, Schaefer J, Grande JP, Roche PC: Immunolocalization of calcitriol receptor, 24-hydroxylase cytochrome P-450, and calbindin D28k in human kidney. *Am J Physiol* 266: F477–F485, 1994

33. Plotkin MD, Kaplan MR, Verlander JW, Lee WS, Brown D, Poch E, Gullans SR, Hebert SC: Localization of the thiazide sensitive Na-Cl cotransporter, rTSC1 in the rat kidney. *Kidney Int* 50: 174–183, 1996
34. Kim W-J, Kakehi Y, Kinoshita H, Arao S, Fukumoto M, Yoshida O: Expression patterns of multidrug-resistance (MDR1), multidrug resistance-associated protein (MRP), glutathione S-transferase-Pi (GST-Pi) and DNA topoisomerase II (Topo II) genes in renal cell carcinomas and normal kidney. *J Urol* 156: 506–511, 1996
35. Keppler D, Kartenbeck J: The canalicular conjugate export pump encoded by the *cmrp/cmrat* gene. In: *Progress in Liver Diseases*, Vol. 14, edited by Boyer JL, Ockner RK, Philadelphia, Saunders, 1996, pp 55–67
36. Jedlitschky G, Leier I, Buchholz U, Hummel-Eisenbeiss J, Burchell B, Keppler D: ATP-dependent transport of bilirubin glucuronides by the multidrug resistance protein MRP1 and its hepatocyte canalicular isoform MRP2. *Biochem J* 327: 305–310, 1997
37. Cui Y, König J, Buchholz U, Spring H, Leier I, Keppler D: Drug resistance and ATP-dependent conjugate transport mediated by the apical multidrug resistance protein, MRP2, permanently expressed in human and canine cells. *Mol Pharmacol* 55: 929–937, 1999