

 Open access • Journal Article • DOI:10.1088/1751-8121/AA5E8F

Extension of the $\mathfrak{o} \sim \mathfrak{s} \sim \mathfrak{p}(m|n)$ correspondence to the infinite-dimensional chiral spinors and self dual tensors — [Source link](#)

N. I. Stoilova, J. Thierry-Mieg, J. Van der Jeugt

Institutions: National Institutes of Health, Ghent University

Published on: 13 Mar 2017 - Journal of Physics A (IOP Publishing)

Topics: Representation theory of SU, Spinor, Irreducible representation, Irreducible component and Tensor

Related papers:

- [Spinors in Physics](#)
- [Spinor representations on a sphere](#)
- [On the representation theory of negative spin](#)
- [Finite- and infinite-dimensional representations of the orthosymplectic superalgebra \$OSP\(3,2\)\$](#)
- [On the quasi-simple irreducible representations of the Lorentz groups](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/extension-of-the-mathfrak-o-mathfrak-s-mathfrak-p-m-n-sim-3xu4eism42>

Extension of the $\mathfrak{osp}(m|n) \sim \mathfrak{so}(m-n)$ correspondence to the infinite-dimensional chiral spinors and self dual tensors

N.I. Stoilova¹, J. Thierry-Mieg² and J. Van der Jeugt³

¹Institute for Nuclear Research and Nuclear Energy, Boul. Tsarigradsko Chaussee 72, 1784 Sofia, Bulgaria

²NCBI, National Library of Medicine, National Institute of Health, 8600 Rockville Pike, Bethesda MD20894, USA

³Department of Applied Mathematics, Computer Science and Statistics, Ghent University, Krijgslaan 281-S9, B-9000 Gent, Belgium

E-mail: stoilova@inrne.bas.bg, mieg@ncbi.nlm.nih.gov and Joris.VanderJeugt@UGent.be

Short title: The $\mathfrak{osp}(m|n) \sim \mathfrak{so}(m-n)$ correspondence

MSC numbers: 17B10, 17B81, 81R05

Abstract

The spinor representations of the orthosymplectic Lie superalgebras $\mathfrak{osp}(m|n)$ are considered and constructed. These are infinite-dimensional irreducible representations, of which the superdimension coincides with the dimension of the spinor representation of $\mathfrak{so}(m-n)$. Next, we consider the self dual tensor representations of $\mathfrak{osp}(m|n)$ and their generalizations: these are also infinite-dimensional and correspond to the highest irreducible component of the p^{th} power of the spinor representation. We determine the character of these representations, and deduce a superdimension formula. From this, it follows that also for these representations the $\mathfrak{osp}(m|n) \sim \mathfrak{so}(m-n)$ correspondence holds.

1 Introduction

In space-time, space and time are not equivalent. The metric of $\mathfrak{so}(3,1)$, $ds^2 = dx^2 + dy^2 + dz^2 - dt^2$, counts time as negative distance. From the point of view of the photon travelling at speed of light between two distant galaxies, the proper time and distance are zero. This explains why the photon only has two degrees of freedom. Although it is described classically by a 4-component vector of $\mathfrak{so}(3,1)$, its two polarization states form an $\mathfrak{so}(2)$ vector, defined in the two-dimensional space transverse to the direction of propagation.

In order to avoid the mathematical difficulties in the rigorous formulation of quantum field theory in Minkowski space-time, one may use the formulation in Euclidean space $ds^2 = dx^2 + dy^2 + dz^2 + d\tau^2$, obtained by performing a so-called Wick rotation $\tau = it$, which changes the signature of the metric and effectively replaces $\mathfrak{so}(3,1)$ by $\mathfrak{so}(4)$. But this raises a problem, since the photon would acquire 4 degrees of freedom. The solution, inspired by Feynman [15], De Witt [12], Faddeev and Popov [14], and by so called BRST symmetry [2,6] (see also [5] and the references therein), is to add two anticommuting directions $(\bar{\theta}, \theta)$ and use a Euclidean metric in (x, y, z, τ) space and antisymmetric symplectic metric in $(\bar{\theta}, \theta)$ space: $ds^2 = dx^2 + dy^2 + dz^2 + d\tau^2 + d\bar{\theta}d\theta - d\theta d\bar{\theta}$, effectively replacing $\mathfrak{so}(3,1)$ by $\mathfrak{osp}(4|2)$ thus, replacing Minkowski space-time $\mathbb{R}^{3,1}$ by the superspace $\mathbb{R}^{4|2}$ [13].

Further, one uses the fact that the two Grassmann variables $\theta, \bar{\theta}$ are compensating the extra dimensions created by the Wick rotation, thus, recovering the $\mathfrak{so}(2)$ invariant transverse physical

space. In terms of representation theory, one could say that the 6-dimensional defining representation of $\mathfrak{osp}(4|2)$, with 4 ordinary variables and 2 Grassmann variables, has superdimension $4 - 2$, coinciding with the dimension of the defining representation of $\mathfrak{so}(2)$. This is a manifestation of what is called the $\mathfrak{osp}(4|2) \sim \mathfrak{so}(2)$ correspondence. More generally, the coincidence of superdimensions of $\mathfrak{osp}(m|n)$ representations with dimensions of $\mathfrak{so}(m - n)$ representations is called the $\mathfrak{osp}(m|n) \sim \mathfrak{so}(m - n)$ correspondence.

To quote other examples of this phenomenon for mass zero gauge fields, the $g_{\mu\nu}$ graviton and the anti-symmetric 3-tensors of $\mathfrak{so}(10, 1)$ supergravity follow the same pattern [9]. The number of degrees of freedom are respectively 44 and 84, equal to the respective dimensions of the metric and the 3-tensor of $\mathfrak{so}(9)$. Their so-called ghost spectra are given by the corresponding tensors of $\mathfrak{osp}(11|2)$ [4].

In the early days of Lie superalgebra representations, the $\mathfrak{osp}(m|n) \sim \mathfrak{so}(m - n) \sim \mathfrak{sp}(n - m)$ correspondence was established for various classes of representations [1, 18, 25]. For finite-dimensional tensors of $\mathfrak{osp}(m|n)$ it was observed that they follow the same rule: their superdimension is equal to the dimension of the corresponding tensor of $\mathfrak{so}(m - n)$. For example, the superdimension of the skew 3-tensor of $\mathfrak{osp}(10|4)$ is $6 \cdot 5 \cdot 4 / 6 = 20$, matching the skew 3-tensor of $\mathfrak{so}(6)$. For a precise statement of when this correspondence holds, see [22, Theorem 3.3]. Note that also for affine Lie superalgebras this correspondence was examined [23].

But the important case of chiral fermions and self dual tensors, which occurs for example in $\mathfrak{osp}(9|1)$ supergravity [19] remained unsolved. These representations have no simple covariant quantization scheme, probably because they have no finite-dimensional $\mathfrak{osp}(m|n)$ counter-part. The aim of the present study is to address this problem.

We show that the chiral spinors and self dual tensors of $\mathfrak{osp}(m|n)$ exist, that they are infinite-dimensional, and that they also follow the $\mathfrak{osp}(m|n) \sim \mathfrak{so}(m - n)$ correspondence, possibly paving the way for a new covariant quantization scheme.

In terms of (the distinguished) Dynkin diagrams of $\mathfrak{osp}(m|n)$, the spinor representation has Dynkin labels $[0, 0, \dots, 0, 1]$ and the self dual tensor $[0, 0, \dots, 0, 2]$. In this paper, we shall treat the irreducible representations (irreps) with Dynkin labels $[0, 0, \dots, 0, p]$, where p is a positive integer. Such representations do not appear in Kac's list of finite-dimensional irreps [21], so they are infinite-dimensional highest weight representations (see also [39]). For infinite-dimensional irreps, one has to be more careful with notions such as dimension and superdimension. It will be useful to view (super)dimension as some infinite formal power series in a variable t , such that the coefficient of t^k keeps track of the finite (super)dimension "at level k " from the highest weight (according to some gradation).

The technique used in this paper to compute superdimensions is as follows. We consider the branching to the $\mathfrak{gl}(m|n)$ subalgebra (i.e. $\mathfrak{osp}(2m + 1|2n) \rightarrow \mathfrak{gl}(m|n)$ and $\mathfrak{osp}(2m|2n) \rightarrow \mathfrak{gl}(m|n)$). The decomposition of the infinite-dimensional irrep $[0, 0, \dots, 0, p]$ according to this branching rule is a (infinite) direct sum of covariant tensor representations of $\mathfrak{gl}(m|n)$. For these tensor representations of $\mathfrak{gl}(m|n)$, labeled by a partition λ , the superdimension is a very simple expression: it actually reduces to the dimension of a $\mathfrak{gl}(m - n)$ irrep. This explains why it is so useful to determine the decomposition with respect to $\mathfrak{gl}(m|n)$. Note that this is equivalent to expressing the character of the $\mathfrak{osp}(2m(+1)|2n)$ irrep $[0, 0, \dots, 0, p]$ as an infinite sum of supersymmetric Schur functions (since the characters of the covariant tensor representations of $\mathfrak{gl}(m|n)$ are given by these Schur functions).

So the main problem is then the determination of the character of the $\mathfrak{osp}(2m(+1)|2n)$ irrep $[0, 0, \dots, 0, p]$ in an appropriate form. For this part of the problem, we are lucky. For

$\mathfrak{osp}(2m+1|2n)$, the character of $[0, 0, \dots, 0, p]$ has been determined in [35]. In fact, the irrep $[0, 0, \dots, 0, p]$ is (equivalent to) the “irreducible parastatistics Fock space of order p ”. A combined system of m parafermions and n parabosons generate the algebra $\mathfrak{osp}(2m+1|2n)$, and the Fock space $V(p)$ of order p (constructed in [35]) corresponds exactly to the irrep $[0, 0, \dots, 0, p]$. For the case of $\mathfrak{osp}(2m|2n)$, the character of $[0, 0, \dots, 0, p]$ can be obtained by considering the decomposition $\mathfrak{osp}(2m+1|2n) \supset \mathfrak{osp}(2m|2n)$.

The structure of the paper is as follows. In section 2 we describe, as an introductory example, the construction of $\mathfrak{osp}(m|2n)$ spinors. The ideas of this construction appeared already in [36], but are reviewed here because they open the way to the more general problem discussed in this paper. In the following sections, we describe some of the mathematical notions needed here. In section 3, we recall the basic notation and terminology related to partitions, remind the reader of symmetric and supersymmetric Schur functions and their relation to $\mathfrak{gl}(n)$ and $\mathfrak{gl}(m|n)$ irreps, and recall some useful (super)dimension formulas for the \mathfrak{gl} case. Section 4 deals with (super)dimensions of some infinite-dimensional representations of $\mathfrak{osp}(1|2n)$, which is a special case among the orthosymplectic Lie superalgebras. We shall see that the technique and results of this section will be crucial for the remainder of the paper. The core of the paper is in sections 5 ($\mathfrak{osp}(2m+1|2n)$) and 6 ($\mathfrak{osp}(2m|2n)$). The main results are as follows: the superdimension of the $\mathfrak{osp}(2m+1|2n)$ irrep $[0, 0, \dots, 0, p]$ is equal to the dimension of the $\mathfrak{so}(2m+1-2n)$ irrep $[0, 0, \dots, 0, p]$. Herein, if $2m+1-2n$ is negative, the algebra $\mathfrak{so}(2m+1-2n)$ should be interpreted as $\mathfrak{osp}(1|2n-2m)$. Secondly, the superdimension of the $\mathfrak{osp}(2m|2n)$ irrep $[0, 0, \dots, 0, p]$ is equal to the dimension of the $\mathfrak{so}(2m-2n)$ irrep $[0, 0, \dots, 0, p]$ (or $[0, 0, \dots, p, 0]$). In this case, if $2m-2n$ is negative, the algebra $\mathfrak{so}(2m-2n)$ should be interpreted as $\mathfrak{sp}(2n-2m)$. At the end of the paper, we also include a section on $D(2, 1; \alpha)$, in order to cover all type II orthosymplectic Lie superalgebras, and we give a summary of the main results in terms of Dynkin diagrams.

2 Construction of $\mathfrak{osp}(m|2n)$ spinors

In his famous lectures on the theory of spinors [8], Cartan has shown that there is no finite-dimensional spinor in curved space. But they exist as infinite-dimensional representations, analysed much later by Ne’eman [32]. In the same way, Kac [21] has shown that the orthosymplectic Lie superalgebras have no finite-dimensional spinors. Their infinite-dimensional counterpart, however, can be constructed very simply as follows [36].

Consider the vector space P^n of polynomials in n formal variables x^1, \dots, x^n with complex coefficients. This space is naturally graded by the total degree in the variables

$$P^n = \bigoplus_{k=0}^{\infty} P_k, \quad (1)$$

where each subspace P_k has dimension $\binom{n+k-1}{k}$. Its partition function, or t -dimension (see later for a formal definition), is

$$\dim_t(P^n) = \sum_{k=0}^{\infty} \dim(P_k) t^k = \frac{1}{(1-t)^n}. \quad (2)$$

The $2n$ operators of multiplication and partial derivation relative to x^i

$$\gamma_{2i-1} = x^i, \quad \gamma_{2i} = 2 \frac{\partial}{\partial x^i}, \quad i = 1, 2, \dots, n \quad (3)$$

form a Weyl algebra

$$\gamma_i \gamma_j - \gamma_j \gamma_i = 2\epsilon_{ij} I \quad (4)$$

where I is the identity and ϵ is a $2n$ dimensional block diagonal matrix where each 2×2 diagonal block is of the form

$$\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}.$$

In other words, ϵ is the canonical skew symmetric metric fixed by the $\mathfrak{sp}(2n)$ symplectic Lie algebra. And indeed, the $2n(2n+1)/2$ symmetric bilinear operators

$$J_{ij} = \frac{1}{4}(\gamma_i \gamma_j + \gamma_j \gamma_i) \quad i, j = 1, 2, \dots, 2n \quad (5)$$

satisfy the $\mathfrak{sp}(2n)$ commutation rules

$$[J_{ij}, J_{kl}] = J_{ij} J_{kl} - J_{kl} J_{ij} = \epsilon_{jk} J_{il} + \epsilon_{jl} J_{ik} + \epsilon_{il} J_{jk} + \epsilon_{ik} J_{jl}. \quad (6)$$

The Chevalley generators e_i, f_i, h_i , $i = 1, 2, \dots, n$ of $\mathfrak{sp}(2n)$, satisfying the standard Chevalley-Serre relations [33], are represented by the operators

$$e_i = x^{i+1} \frac{\partial}{\partial x^i}, \quad f_i = x^i \frac{\partial}{\partial x^{i+1}}, \quad h_i = x^{i+1} \frac{\partial}{\partial x^{i+1}} - x^i \frac{\partial}{\partial x^i}, \quad i = 1, \dots, n-1$$

$$e_n = \frac{1}{2} \left(\frac{\partial}{\partial x^n} \right)^2, \quad f_n = -\frac{1}{2} (x^n)^2, \quad h_n = -\frac{1}{2} (x^n \frac{\partial}{\partial x^n} + \frac{\partial}{\partial x^n} x^n).$$

Since these operators map polynomials of even (resp. odd) degree in x into polynomials of even (resp. odd) degree in x one obtains the two irreducible so-called metaplectic representations characterized by their highest weight 1 and x_n , the only monomial belonging to the kernel of all the e_i . By inspection, the Dynkin weights are:

$$\begin{array}{ccccccc} 0 & & 0 & & 0 & & -1/2 \\ \circ & \text{---} & \circ & \text{---} & \circ & \leftarrow & \circ \\ 1 & & 2 & & n-1 & & n \end{array}$$

$$\begin{array}{ccccccc} 0 & & 0 & & 1 & & -3/2 \\ \circ & \text{---} & \circ & \text{---} & \circ & \leftarrow & \circ \\ 1 & & 2 & & n-1 & & n \end{array}$$

In a similar way, the orthogonal Lie algebra $\mathfrak{so}(2m)$ can be realized by means of operators acting on polynomials of anti-commuting variables. For this purpose, consider the vector space Q^m of polynomials in m formal anti-commuting variables θ^α ($\alpha = 1, 2, \dots, m$) with complex coefficients. Since $(\theta^\alpha)^2 = 0$, this space is finite-dimensional

$$\dim(Q^m) = 2^m. \quad (7)$$

The following $2m$ operators

$$\gamma_{2\alpha-1} = \theta^\alpha + \frac{\partial}{\partial \theta^\alpha},$$

$$\gamma_{2\alpha} = i\left(\theta^\alpha - \frac{\partial}{\partial \theta^\alpha}\right), \quad \alpha = 1, 2, \dots, m$$

represent the $2m$ dimensional Clifford algebra

$$\gamma_\mu \gamma_\nu + \gamma_\nu \gamma_\mu = 2g_{\mu\nu} I, \quad \mu, \nu = 1, 2, \dots, 2m \quad (8)$$

where I is the identity and $g_{\mu\nu}$ is a $2m$ dimensional identity matrix. In other words, $g_{\mu\nu}$ is the canonical symmetric metric fixed by the $\mathfrak{so}(2m)$ orthogonal Lie algebra. And indeed, the $2m(2m-1)/2$ antisymmetric bilinear operators

$$J_{\mu\nu} = \frac{1}{4}(\gamma_\mu \gamma_\nu - \gamma_\nu \gamma_\mu), \quad \mu, \nu = 1, 2, \dots, 2m \quad (9)$$

satisfy the $\mathfrak{so}(2m)$ commutation rules

$$[J_{\mu\nu}, J_{\rho\sigma}] = J_{\mu\nu} J_{\rho\sigma} - J_{\rho\sigma} J_{\mu\nu} = g_{\nu\rho} J_{\mu\sigma} - g_{\nu\sigma} J_{\mu\rho} + g_{\mu\sigma} J_{\nu\rho} - g_{\mu\rho} J_{\nu\sigma}. \quad (10)$$

In other words, using polynomials in anti-commuting variables, one obtains a simple construction of the well known 2^m -dimensional Dirac matrices in $2m$ dimensions.

Since the $J_{\mu\nu}$ operators are even in θ , the representation space Q^m splits into the even and odd polynomials, which correspond to the left and right spinors, each of dimension 2^{m-1} . The $\mathfrak{so}(2m+1)$ algebra is obtained by defining the operator Γ_5 and adding the operators $J_{\mu 5} = -J_{5\mu}$:

$$\Gamma_5 = \gamma_1 \gamma_2 \cdots \gamma_{2m}, \quad J_{\mu 5} = \gamma_\mu \Gamma_5 = -\Gamma_5 \gamma_\mu = -J_{5\mu}, \quad \Gamma_5^2 = (-1)^m. \quad (11)$$

Since Γ_5 is even in the θ 's, the $J_{\mu 5}$ operators are odd and they connect the even and odd polynomials, so the space Q^m is irreducible and the dimension of the spinors of $\mathfrak{so}(2m+1)$ is 2^m .

The parallel between the \mathfrak{so} and \mathfrak{sp} construction is striking, but notice that the statistics is counter-intuitive: in spinor space $\mathfrak{sp}(2n)$ acts on ‘‘bosonic polynomials’’ and $\mathfrak{so}(2m+1)$ on Grassmann or ‘‘fermionic polynomials’’.

The generalization to the orthosymplectic Lie superalgebra $\mathfrak{osp}(2m|2n)$ is now immediate. One considers the space $P^{m|n}$ of superpolynomials in m anti-commuting and n commuting variables. As we will see later in general, this corresponds to the spinor representation of $\mathfrak{osp}(2m|2n)$. Using the same gradation by the total degree in x , the t -dimension is

$$\dim_t(P^{m|n}) = \frac{2^m}{(1-t)^n}. \quad (12)$$

The superdimension (difference between dimension of even and odd subspaces) is

$$\text{sdim}(P^{m|n}) = \dim_{-1}(P^{m|n}) = 2^{m-n}. \quad (13)$$

So for $m \geq n$, the superdimension of the (infinite-dimensional) spinor representation of $\mathfrak{osp}(2m|2n)$ coincides with the (finite) dimension (7) of the spinor representation of $\mathfrak{so}(2m-2n)$, fulfilling the correspondence being studied in this paper.

To construct the $\mathfrak{osp}(2m|2n)$ generators, one considers the $m(2m-1)$ antisymmetric $J_{\mu\nu}$, the $n(2n+1)$ symmetric J_{ij} and adds the $4mn$ odd operators

$$J_{\mu i} = \gamma_\mu \gamma_i, \quad \mu = 1, 2, \dots, 2m, \quad i = 1, 2, \dots, 2n. \quad (14)$$

Their anticommutators close on the previous even operators, completing the construction of the $\mathfrak{osp}(2m|2n)$ superspinor representation:

$$\{J_{\mu i}, J_{\nu j}\} = J_{\mu i} J_{\nu j} + J_{\nu j} J_{\mu i} = 4\epsilon_{ij} J_{\mu\nu} + 4g_{\mu\nu} J_{ij}. \quad (15)$$

The superspinor representation space $P^{m|n}$ naturally splits in four components

$$P^{m|n} = P_{0,0}^{m|n} \oplus P_{1,0}^{m|n} \oplus P_{0,1}^{m|n} \oplus P_{1,1}^{m|n}, \quad (16)$$

where $P_{i,j}^{m|n}$ denotes the subspace of polynomials of total even (resp. odd) degree in the variables θ for $i = 0$ (resp. $i = 1$) and of total even (resp. odd) degree in the variables x for $j = 0$ (resp. $j = 1$). Since the odd operators $J_{\mu i}$ are of odd degree in x and θ , they map these subspaces onto each other. On the other hand, these operators are of total even degree (in x and θ combined), so under the action of $\mathfrak{osp}(2m|2n)$ the superspinor representation space $P^{m|n}$ still splits into a “left sector” $P_{0,0}^{m|n} \oplus P_{1,1}^{m|n}$ and a “right sector” $P_{1,0}^{m|n} \oplus P_{0,1}^{m|n}$, each of superdimension 2^{m-n-1} .


The case $\mathfrak{osp}(2m+1|2n)$ follows by adding the $\mathfrak{so}(2m+1)$ operator Γ_5 as described above, and in this case the space $P^{m|n}$ is irreducible of superdimension 2^{m-n} .

In the following sections we shall be dealing with the self dual tensors of the orthosymplectic Lie superalgebras and their generalizations. These can be considered as the top irreducible component of the p^{th} symmetric tensor product of the superspinor. These self dual tensors are also infinite-dimensional and follow the $\mathfrak{osp}(m|n) \sim \mathfrak{so}(m-n)$ correspondence. Notice that we have given in this section a construction of the superspinor, with the explicit action of all the generators of $\mathfrak{osp}(m|n)$. As far as we know, there is no such simple construction for the self dual tensors. In the remainder of the paper, we only give the characters of the irreducible self dual tensors, and that is already sufficient to establish the correspondence.

3 Preliminaries


3.1 Partitions, symmetric functions and supersymmetric functions

We need in this paper some basic notions on partitions and symmetric functions, see [30] as a standard reference. A partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$ of weight $|\lambda|$ and length $\ell(\lambda) \leq n$ is a sequence of non-negative integers satisfying the condition $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$, such that their sum is $|\lambda|$, and $\lambda_i > 0$ if and only if $i \leq \ell(\lambda)$. To each such partition there corresponds a Young diagram F^λ consisting of $|\lambda|$ boxes arranged in $\ell(\lambda)$ left-adjusted rows of lengths λ_i for $i = 1, 2, \dots, \ell(\lambda)$. For example, the Young diagram of $\lambda = (5, 4, 4, 2)$ is given by



The conjugate partition λ' corresponds to the Young diagram of λ reflected about the main diagonal. In other words, λ'_j is the length of column j of F^λ . For the above example, $\lambda' = (4, 4, 3, 3, 1)$.

An important notion is the Frobenius notation [30] of a partition λ . If F^λ has $r = r(\lambda)$ boxes on the diagonal, r is said to be the rank of λ . In the above example, $r = 3$, denoted by crosses in the diagonal boxes:



The arm lengths $a_k = \lambda_k - k$ and leg lengths $b_k = \lambda'_k - k$ ($k = 1, \dots, r$) refer to the remaining boxes to the right or below the k th diagonal box, where $a_1 > a_2 > \dots > a_r \geq 0$ and $b_1 > b_2 > \dots > b_r \geq 0$. The Frobenius notation of λ is then

$$\lambda = \begin{pmatrix} a_1 & a_2 & \dots & a_r \\ b_1 & b_2 & \dots & b_r \end{pmatrix}.$$

Note that for our example we have $\lambda = \begin{pmatrix} 4 & 2 & 1 \\ 3 & 2 & 0 \end{pmatrix}$.

Denote by Λ_n the ring of symmetric polynomials with integer coefficients in the n independent variables $x = (x_1, x_2, \dots, x_n)$. The Schur functions [30] or S -functions $s_\lambda(x)$ form a \mathbb{Z} -basis of Λ_n , where λ runs over the set of all partitions of length at most n . There are various ways to define Schur functions. For a partition λ with $\ell(\lambda) \leq n$, one has

$$s_\lambda(x) = \frac{\det(x_i^{\lambda_j + n - j})_{1 \leq i, j \leq n}}{\det(x_i^{n - j})_{1 \leq i, j \leq n}}. \quad (17)$$

If $\ell(\lambda) > n$, one puts $s_\lambda(x) = 0$. Clearly, for the zero partition $\lambda = (0)$ (which is the only partition with Frobenius rank $r = 0$) one has $s_{(0)}(x) = 1$.

In terms of two sets of variables $x = (x_1, \dots, x_m)$ and $y = (y_1, \dots, y_n)$, one can define the ring $\Lambda_{m,n}$ of supersymmetric polynomials with integer coefficients [3]. This ring consists of all double symmetric polynomials in x and y (elements of $\Lambda_m \otimes \Lambda_n$) that satisfy the so-called cancellation property (i.e. when the substitution $x_1 = t, y_1 = -t$ is made in an element p of $\Lambda_m \otimes \Lambda_n$, the resulting polynomial is independent of t). For a partition λ , one can define supersymmetric Schur functions $s_\lambda(x|y)$ belonging to $\Lambda_{m,n}$. One possible definition is [3, 24]

$$s_\lambda(x|y) = \sum_{\mu, \nu} c_{\mu, \nu}^\lambda s_\mu(x) s_{\nu'}(y),$$

where the sum is over all partitions μ ($\ell(\mu) \leq m$) and ν ($\ell(\nu') \leq n$), and $c_{\mu, \nu}^\lambda$ are the Littlewood-Richardson coefficients [30]. The polynomials $s_\lambda(x|y)$ are identically zero when $\lambda_{m+1} > n$. Denote by $\mathcal{H}_{m,n}$ the set of all partitions with $\lambda_{m+1} \leq n$, i.e. the partitions (with their Young diagram) inside the (m, n) -hook. The set of $s_\lambda(x|y)$ with $\lambda \in \mathcal{H}_{m,n}$ form a \mathbb{Z} -basis of $\Lambda_{m,n}$.

3.2 Dimension formulas for $\mathfrak{gl}(n)$ and superdimensions for $\mathfrak{gl}(m|n)$

A finite-dimensional irreducible representation of the Lie algebra $\mathfrak{gl}(n)$ is characterized by a partition λ with $\ell(\lambda) \leq n$. In terms of the standard basis $\epsilon_1, \dots, \epsilon_n$ of the weight space of $\mathfrak{gl}(n)$, the highest weight of this representation is $\sum_{i=1}^n \lambda_i \epsilon_i$, and the representation space will be denoted by $V_{\mathfrak{gl}(n)}^\lambda$. Weyl's character formula for such representations corresponds to the right hand side of (17), and thus

$$\text{char } V_{\mathfrak{gl}(n)}^\lambda = s_\lambda(x), \quad (18)$$

where $x_i = e^{\epsilon_i}$.

There are two useful formulas for the dimension of $V_{\mathfrak{gl}(n)}^\lambda$, by specializing all $x_i = 1$ in the Schur function s_λ . One is Weyl's dimension formula:

$$\dim V_{\mathfrak{gl}(n)}^\lambda = \prod_{1 \leq i < j \leq n} \frac{(\lambda_i - \lambda_j + j - i)}{(j - i)}. \quad (19)$$

The other expression uses hook-lengths:

$$\dim V_{\mathfrak{gl}(n)}^\lambda = \prod_{(i,j) \in \lambda} \frac{n+j-i}{h_{ij}} = \prod_{(i,j) \in \lambda} \frac{n+j-i}{\lambda_i + \lambda'_j - i - j + 1}. \quad (20)$$

Herein, the product runs over all boxes (i, j) of the Young diagram F^λ , and $h_{ij} = \lambda_i + \lambda'_j - i - j + 1$ is the number of boxes of the hook with hook box (i, j) . For example, for the partition $(5, 4, 4, 2)$ and $n = 5$, the numbers $(n + j - i)$ and h_{ij} read, respectively,

5	6	7	8	9
4	5	6	7	
3	4	5	6	
2	3			

8	7	5	4	1
6	5	3	2	
5	4	2	1	
2	1			

Note, if λ is a partition with $\ell(\lambda) > n$, (20) gives automatically 0, whereas in (19) one has to impose that $\dim V_{\mathfrak{gl}(n)}^\lambda = 0$.

Just as the functions $s_\lambda(x)$ are characters of irreducible representations (or simple modules) of the Lie algebra $\mathfrak{gl}(n)$, the supersymmetric Schur functions are characters of a class of simple modules of the Lie superalgebra $\mathfrak{gl}(m|n)$, namely of the covariant representations [3]. For a partition $\lambda \in \mathcal{H}_{m,n}$, the corresponding covariant representation will be denoted by $V_{\mathfrak{gl}(m|n)}^\lambda$. In terms of the standard basis $\epsilon_1, \dots, \epsilon_m, \delta_1, \dots, \delta_n$ of the weight space of $\mathfrak{gl}(m|n)$, the highest weight of this representation is $\sum_{i=1}^m \lambda_i \epsilon_i + \sum_{j=1}^n \max(\lambda'_j - m, 0) \delta_j$. The main result of [3] is

$$\text{char } V_{\mathfrak{gl}(m|n)}^\lambda = s_\lambda(x|y), \quad (21)$$

where $x_i = e^{\epsilon_i}$ and $y_j = e^{\delta_j}$.

Any Lie superalgebra \mathfrak{g} is \mathbb{Z}_2 -graded: $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$. A Lie superalgebra module or representation V is also \mathbb{Z}_2 -graded: $V = V_0 \oplus V_1$. In our convention, the highest weight vector v of V will always be an even vector ($v \in V_0$). When V is finite-dimensional, one can speak of the dimension and superdimension of V :

$$\dim V = \dim V_0 + \dim V_1, \quad \text{sdim } V = \dim V_0 - \dim V_1.$$

Dimension formulas for covariant representations of $\mathfrak{gl}(m|n)$ are not particularly simple [31], but there is a great simplification when it comes to superdimensions [24]. The result depends on whether m is greater than, equal to, or less than n . It can be summarized as follows:

$$\text{sdim } V_{\mathfrak{gl}(n+k|n)}^\lambda = \dim V_{\mathfrak{gl}(k)}^\lambda, \quad \text{sdim } V_{\mathfrak{gl}(m|m+k)}^\lambda = (-1)^{|\lambda|} \dim V_{\mathfrak{gl}(k)}^{\lambda'}. \quad (22)$$

In particular, when $m = n$, $\text{sdim } V_{\mathfrak{gl}(n|n)}^\lambda = 0$ unless λ is the zero partition (0) (then $V_{\mathfrak{gl}(n|n)}^{(0)}$ is the trivial module with $\text{sdim } V_{\mathfrak{gl}(n|n)}^{(0)} = 1$). Note that (22) implies: when $\ell(\lambda) > k$ then $\text{sdim } V_{\mathfrak{gl}(n+k|n)}^\lambda = 0$; when $\lambda_1 > k$ then $\text{sdim } V_{\mathfrak{gl}(m|m+k)}^\lambda = 0$.

4 The Lie superalgebra $\mathfrak{osp}(1|2n)$

Although the main aim is to study superdimensions of certain infinite-dimensional representations of the Lie superalgebras $B(m, n) = \mathfrak{osp}(2m+1|2n)$ and $D(m, n) = \mathfrak{osp}(2m|2n)$, it will be useful to start with the special case $B(0, n) = \mathfrak{osp}(1|2n)$. Let us first fix some notation [16, 20, 21]. In the

common basis δ_j for the weight space of $\mathfrak{osp}(1|2n)$, the odd roots are given by $\pm\delta_j$ ($j = 1, \dots, n$), and the even roots are $\delta_i - \delta_j$ ($i \neq j$) and $\pm(\delta_i + \delta_j)$. The simple roots are

$$\delta_1 - \delta_2, \delta_2 - \delta_3, \dots, \delta_{n-1} - \delta_n, \delta_n. \quad (23)$$

It will be helpful to use the following \mathbb{Z} -gradation of $\mathfrak{g} = \mathfrak{osp}(1|2n)$: $\mathfrak{g} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_{+1} \oplus \mathfrak{g}_{+2}$, where each \mathfrak{g}_j is spanned by the root vectors corresponding to the following roots:

$$\begin{array}{ccccc} \mathfrak{g}_{-2} & \mathfrak{g}_{-1} & \mathfrak{g}_0 & \mathfrak{g}_{+1} & \mathfrak{g}_{+2} \\ -\delta_i - \delta_j & -\delta_i & \delta_i - \delta_j & \delta_i & \delta_i + \delta_j \end{array}$$

Note that $\mathfrak{g}_0 = \mathfrak{gl}(n)$.

We will consider some infinite-dimensional highest weight representations V of \mathfrak{g} , so at first sight it might be strange to speak of the dimension and superdimension of V . However, if the action of $\mathfrak{g}_0 = \mathfrak{gl}(n)$ on the highest weight vector v of V corresponds to a finite-dimensional \mathfrak{g}_0 module V_0 , then the \mathbb{Z} -gradation of \mathfrak{g} induces a \mathbb{Z} -gradation of V :

$$V = V_0 \oplus V_{-1} \oplus V_{-2} \oplus \dots$$

in terms of finite-dimensional \mathfrak{g}_0 modules. Then one defines:

$$\dim_t(V) = \sum_{i=0}^{\infty} \dim V_{-i} t^i, \quad \text{sdim}_t(V) = \sum_{i=0}^{\infty} \dim V_{-i} (-t)^i. \quad (24)$$

Clearly $\text{sdim}_t(V) = \dim_{-t}(V)$, and $\dim_t(V)$ is a formal power series in the variable t such that the coefficient of t^k counts the dimension ‘‘at level k ’’ according to the \mathbb{Z} -gradation.

For reasons that will become clear, we will consider the irreducible highest weight representation with highest weight given by $(-\frac{p}{2}, -\frac{p}{2}, \dots, -\frac{p}{2}) = \sum_{i=1}^n (-\frac{p}{2})\delta_i$, where p is a non-negative integer. This representation will be denoted by $V = V_{\mathfrak{osp}(1|2n)}^{(-p/2)^n}$. Its Dynkin labels according to the simple roots system (23) are $[0, 0, \dots, 0, -p]$. Clearly, with this choice of highest weight one has $\dim V_0 = 1$. Obviously, the structure of the irreducible highest weight representation with highest weight $(-\frac{p}{2}, -\frac{p}{2}, \dots, -\frac{p}{2})$ is completely the same as the irreducible lowest weight representation with lowest weight $(\frac{p}{2}, \frac{p}{2}, \dots, \frac{p}{2})$. This last representation is well known: it is the paraboson representation of order p . Its structure and character have been determined in [29]. Using the notation $x_i = e^{-\delta_i}$, the following character formula holds:

$$\text{char } V_{\mathfrak{osp}(1|2n)}^{(-p/2)^n} = (x_1 \cdots x_n)^{p/2} \sum_{\lambda, \ell(\lambda) \leq p} s_{\lambda}(x). \quad (25)$$

This is an infinite sum over all partitions of length at most p . Since $s_{\lambda}(x) = 0$ if $\ell(\lambda) > n$, the sum is actually over all partitions satisfying $\ell(\lambda) \leq \min(n, p)$.

In order to give a first expression of $\dim_t V_{\mathfrak{osp}(1|2n)}^{(-p/2)^n}$ one should (apart from the factor $(x_1 \cdots x_n)^{p/2}$) specify $x_i = t$ in (25). Thus one finds:

$$\dim_t V_{\mathfrak{osp}(1|2n)}^{(-p/2)^n} = \sum_{\lambda, \ell(\lambda) \leq \min(n, p)} \dim V_{\mathfrak{gl}(n)}^{\lambda} t^{|\lambda|}. \quad (26)$$

This is an infinite sum; it can be simplified using the alternative character formula also determined in [29]:

$$\text{char } V_{\mathfrak{osp}(1|2n)}^{(-p/2)^n} = (x_1 \cdots x_n)^{p/2} \frac{E_{n,p}(x)}{\prod_i (1 - x_i) \prod_{i < j} (1 - x_i x_j)}. \quad (27)$$

Herein,

$$E_{n,p}(x) = \sum_{\eta} (-1)^{c_{\eta}} s_{\eta}(x),$$

where the sum is over all partitions η of Frobenius form

$$\eta = \begin{pmatrix} a_1 & a_2 & \cdots & a_r \\ a_1 + p & a_2 + p & \cdots & a_r + p \end{pmatrix}$$

and $c_{\eta} = a_1 + \cdots + a_r + r = |a| + r$. So the sum in $E_{n,p}(x)$ is over all self-conjugate partitions to which r “legs” of p boxes are added. Note that $E_{n,p}(x) = 1$ if $p \geq n$. In the following, this trivial case will be left out and we shall deal only with $p < n$.

Specifying $x_i = t$ in (27) now yields

$$\dim_t V_{\mathfrak{osp}(1|2n)}^{(-p/2)^n} = \frac{E_{n,p}(t, \dots, t)}{(1-t)^n (1-t^2)^{n(n-1)/2}}. \quad (28)$$

So, let us therefore consider $E_{n,p}(x)$ in more detail, starting with $E_{n,0}(x)$. Since $s_{\lambda}(x) = 0$ when $\ell(\lambda) > n$, the expression $E_{n,0}(x)$ has only a finite number of terms:

$$E_{n,0}(x) = \sum_{r=0}^n \sum_{n-1 \geq a_1 > a_2 > \cdots > a_r \geq 0} (-1)^{|a|+r} s_{\begin{pmatrix} a_1 & a_2 & \cdots & a_r \\ a_1 & a_2 & \cdots & a_r \end{pmatrix}}(x). \quad (29)$$

It is easy to see that this expression has 2^n terms. Similarly, $E_{n,p}(x)$ has 2^{n-p} terms, and

$$E_{n,p}(x) = \sum_{r=0}^{n-p} \sum_{n-p-1 \geq a_1 > a_2 > \cdots > a_r \geq 0} (-1)^{|a|+r} s_{\begin{pmatrix} a_1 & a_2 & \cdots & a_r \\ a_1+p & a_2+p & \cdots & a_r+p \end{pmatrix}}(x). \quad (30)$$

Let us now turn to the dimension formula. In general, for a partition λ given in Frobenius notation

$$\lambda = \begin{pmatrix} a_1 & a_2 & \cdots & a_r \\ b_1 & b_2 & \cdots & b_r \end{pmatrix} \equiv \begin{pmatrix} a \\ b \end{pmatrix}$$

one can write

$$s_{\lambda}(t, \dots, t) = t^{r+|a|+|b|} s_{\lambda}(1, \dots, 1) = t^{r+|a|+|b|} \dim V_{\mathfrak{gl}(n)}^{\lambda}.$$

Using (20), one finds

$$\dim V_{\mathfrak{gl}(n)}^{\lambda} = \dim V_{\mathfrak{gl}(n)}^{\begin{pmatrix} a \\ b \end{pmatrix}} = \prod_{i=1}^r \frac{(n+a_i)!}{(n-b_i-1)!} \bigg/ \frac{\prod_{i=1}^r a_i! b_i! \prod_{i,j=1}^r (a_i + b_j + 1)}{\prod_{1 \leq i < j \leq r} (a_i - a_j)(b_i - b_j)}. \quad (31)$$

And hence

$$E_{n,p}(t, \dots, t) = \sum_{r=0}^{n-p} \sum_{n-p-1 \geq a_1 > a_2 > \cdots > a_r \geq 0} (-1)^{|a|+r} \dim V_{\mathfrak{gl}(n)}^{\begin{pmatrix} a \\ a+p \end{pmatrix}} t^{2|a|+(p+1)r}. \quad (32)$$

Let us complete this section by means of some examples:

$$\begin{aligned} E_{1,0}(t, \dots, t) &= 1 - t \\ E_{2,0}(t, \dots, t) &= 1 - 2t + 2t^3 - t^4 = (1-t)^2(1-t^2) \\ E_{3,0}(t, \dots, t) &= 1 - 3t + 8t^3 - 6t^4 - 6t^5 + 8t^6 - 3t^8 + t^9 = (1-t)^3(1-t^2)^3 \\ E_{3,1}(t, \dots, t) &= 1 - 3t^2 + 3t^4 - t^6 = (1-t^2)^3 \\ E_{3,2}(t, \dots, t) &= 1 - t^3 \\ E_{4,1}(t, \dots, t) &= 1 - 6t^2 + 15t^4 - 20t^6 + 15t^8 - 6t^{10} + t^{12} = (1-t^2)^6 \end{aligned}$$

Of course, due to the general factorization of $E_{n,0}(x)$ [26], one has $E_{n,0}(t, \dots, t) = (1-t)^2(1-t^2)^{n(n-1)/2}$. Some other general expressions following from (32) are:

$$\begin{aligned}
E_{n,1}(t, \dots, t) &= (1-t^2)^{n(n-1)/2} \\
E_{p+1,p}(t, \dots, t) &= 1-t^{p+1} \\
E_{p+2,p}(t, \dots, t) &= 1-(p+2)t^{p+1}+(p+2)t^{p+3}-t^{2p+4} \\
E_{p+3,p}(t, \dots, t) &= 1-\frac{(p+2)(p+3)}{2}t^{p+1}+(p+2)(p+4)t^{p+3}-\frac{(p+3)(p+4)}{2}t^{p+5} \\
&\quad -\frac{(p+3)(p+4)}{2}t^{2p+4}+(p+2)(p+4)t^{2p+6}-\frac{(p+2)(p+3)}{2}t^{2p+8}-t^{3p+9}
\end{aligned}$$

These were a few examples of $E_{n,p}(t, \dots, t)$. Including denominators according to (28) one finds the actual dimensions, e.g.:

$$\begin{aligned}
\dim_t V_{\mathfrak{osp}(1|6)}^{(0,0,0)} &= 1 \\
\dim_t V_{\mathfrak{osp}(1|6)}^{(-1/2,-1/2,-1/2)} &= \frac{1-3t^2+3t^4-t^6}{(1-t)^3(1-t^2)^3} = \frac{1}{(1-t)^3} = 1+3t+6t^2+10t^3+15t^4+\dots \\
\dim_t V_{\mathfrak{osp}(1|6)}^{(-1,-1,-1)} &= \frac{1-t^3}{(1-t)^3(1-t^2)^3} = 1+3t+9t^2+18t^3+36t^4+\dots \\
\dim_t V_{\mathfrak{osp}(1|6)}^{(-3/2,-3/2,-3/2)} &= \frac{1}{(1-t)^3(1-t^2)^3} = 1+3t+9t^2+19t^3+39t^4+\dots
\end{aligned}$$

5 The Lie superalgebra $\mathfrak{osp}(2m+1|2n)$

We will now treat one of the two main cases, the Lie superalgebra $B(m, n) = \mathfrak{osp}(2m+1|2n)$. The weight space of $\mathfrak{osp}(2m+1|2n)$ has basis $\epsilon_1, \dots, \epsilon_m, \delta_1, \dots, \delta_n$, and the distinguished set of simple roots is [16, 20]

$$\delta_1 - \delta_2, \dots, \delta_{n-1} - \delta_n, \delta_n - \epsilon_1, \epsilon_1 - \epsilon_2, \dots, \epsilon_{m-1} - \epsilon_m, \epsilon_m. \quad (33)$$

Also in this case there exists a useful \mathbb{Z} -gradation of $\mathfrak{g} = \mathfrak{osp}(2m+1|2n)$: $\mathfrak{g} = \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_{+1} \oplus \mathfrak{g}_{+2}$, where each \mathfrak{g}_j is spanned by the root vectors corresponding to the following roots:

\mathfrak{g}_{-2}	\mathfrak{g}_{-1}	\mathfrak{g}_0	\mathfrak{g}_{+1}	\mathfrak{g}_{+2}
$-\delta_i - \delta_j$	$-\delta_i$	$\delta_i - \delta_j$	δ_i	$\delta_i + \delta_j$
$-\epsilon_i - \epsilon_j \ (i \neq j)$	$-\epsilon_i$	$\epsilon_i - \epsilon_j$	ϵ_i	$\epsilon_i + \epsilon_j \ (i \neq j)$
$-\epsilon_i - \delta_j$		$\pm(\epsilon_i - \delta_j)$		$\epsilon_i + \delta_j$

So in this case $\mathfrak{g}_0 = \mathfrak{gl}(m|n)$. Note that this \mathbb{Z} -gradation is different from the more common \mathbb{Z} -gradation for which $\mathfrak{g}_0 = \mathfrak{g}_{\bar{0}}$ [16, 17].

Let us now consider the irreducible highest weight representation with highest weight given by $(\frac{p}{2}, \dots, \frac{p}{2}; -\frac{p}{2}, \dots, -\frac{p}{2}) = \sum_{i=1}^m \frac{p}{2}\epsilon_i + \sum_{i=1}^n (-\frac{p}{2})\delta_i$, where p is a non-negative integer. This representation will be denoted by $V = V_{\mathfrak{osp}(2m+1|2n)}^{(p/2)^m, (-p/2)^n}$. According to the simple roots system (33), its Dynkin labels are $[0, 0, \dots, 0, p]$. The action of $\mathfrak{g}_0 = \mathfrak{gl}(m|n)$ on the highest weight vector v is again trivial, so the \mathbb{Z} -gradation of \mathfrak{g} induces a \mathbb{Z} -gradation of V with $\dim V_0 = 1$, and one can again study $\dim_t(V)$ and $\text{sdim}_t(V)$. In the current situation, the structure of the irreducible highest weight representation with highest weight $(\frac{p}{2}, \dots, \frac{p}{2}; -\frac{p}{2}, \dots, -\frac{p}{2})$ is completely the same as the irreducible lowest weight representation with lowest weight $(-\frac{p}{2}, \dots, -\frac{p}{2}; \frac{p}{2}, \dots, \frac{p}{2})$. This

representation has been studied recently and its structure is well known: it is the parastatistics representation of order p [35]. Using the notation $x_i = e^{-\epsilon_i}$, $y_i = e^{-\delta_i}$, the following character formula holds:

$$\text{char } V_{\mathfrak{osp}(2m+1|2n)}^{(p/2)^m, (-p/2)^n} = (y_1 \cdots y_n / x_1 \cdots x_m)^{p/2} \sum_{\lambda, \lambda_1 \leq p} s_\lambda(x|y). \quad (34)$$

So here the sum is over all partitions λ inside the (m, n) -hook (otherwise $s_\lambda(x|y)$ is zero anyway) with $\lambda_1 \leq p$, or equivalently $\ell(\lambda') \leq p$. There is also an alternative character formula [35, Theorem 7], but we do not need it here.

In order to determine $\text{sdim}_t V_{\mathfrak{osp}(2m+1|2n)}^{(p/2)^m, (-p/2)^n}$, one should (apart from the factor in front of the above sum) specify $x_i = t$ and $y_j = -t$ in the above character, and so one finds

$$\begin{aligned} \text{sdim}_t V_{\mathfrak{osp}(2m+1|2n)}^{(p/2)^m, (-p/2)^n} &= \sum_{\lambda, \lambda_1 \leq p} s_\lambda(t, \dots, t | -t, \dots, -t) \\ &= \sum_{\lambda, \lambda_1 \leq p} s_\lambda(1, \dots, 1 | -1, \dots, -1) t^{|\lambda|} \\ &= \sum_{\lambda, \lambda_1 \leq p} \text{sdim } V_{\mathfrak{gl}(m|n)}^\lambda t^{|\lambda|}. \end{aligned} \quad (35)$$

We can now make use of the expressions and properties of $\mathfrak{gl}(m|n)$ superdimensions, given at the end of section 3, and thus specify the following three cases.

Case 1: $m = n$, $\mathfrak{osp}(2n+1|2n)$.

This is the simplest case, since all superdimensions of covariant representations of $\mathfrak{gl}(n|n)$ are zero, except when $\lambda = (0)$. Hence:

$$\text{sdim}_t V_{\mathfrak{osp}(2n+1|2n)}^{(p/2)^n, (-p/2)^n} = 1. \quad (36)$$

Case 2: $m = n + k$, $\mathfrak{osp}(2n+2k+1|2n)$.

Now it follows directly from (35) and (22) that

$$\text{sdim}_t V_{\mathfrak{osp}(2m+1|2n)}^{(p/2)^m, (-p/2)^n} = \sum_{\lambda, \lambda_1 \leq p} \dim V_{\mathfrak{gl}(k)}^\lambda t^{|\lambda|} = \sum_{\lambda, \lambda_1 \leq p, \ell(\lambda) \leq k} \dim V_{\mathfrak{gl}(k)}^\lambda t^{|\lambda|}. \quad (37)$$

So the original infinite sum reduces to a finite sum, running over all partitions λ with $\lambda_1 \leq p$ and $\ell(\lambda) \leq k$. In other words, the sum is over all partitions λ whose Young diagrams fit inside the $(k \times p)$ rectangle. However, such an expression is known, see [34] for a detailed investigation (although the origin goes back to work of Bracken and Green [7]). Indeed, in the branching $\mathfrak{so}(2k+1) \supset \mathfrak{gl}(k)$, the $\mathfrak{so}(2k+1)$ highest weight representation $V_{\mathfrak{so}(2k+1)}^{(p/2)^k}$ with highest weight $(\frac{p}{2}, \dots, \frac{p}{2})$ decomposes into the direct sum of covariant $\mathfrak{gl}(k)$ representation $V_{\mathfrak{gl}(k)}^\lambda$ with $\ell(\lambda') \leq p$. Thus we find:

$$\text{sdim}_t V_{\mathfrak{osp}(2m+1|2n)}^{(p/2)^m, (-p/2)^n} = \dim_t V_{\mathfrak{so}(2k+1)}^{(p/2)^k}, \quad (38)$$

or, in terms of Dynkin labels:

$$\text{sdim}_t [0, 0, \dots, 0, p]_{\mathfrak{osp}(2n+2k+1|2n)} = \dim_t [0, \dots, 0, p]_{\mathfrak{so}(2k+1)}. \quad (39)$$

Let us give a few examples. For $k = 3$ and $p = 1$, one finds

$$\text{sdim}_t [0, 0, \dots, 0, 1]_{\mathfrak{osp}(2n+7|2n)} = \dim_t [0, 0, 1]_{\mathfrak{so}(7)} = 1 + 3t + 3t^2 + t^3.$$

and thus $\text{sdim}[0, 0, \dots, 0, 1]_{\mathfrak{osp}(2n+7|2n)} = \dim[0, 0, 1]_{\mathfrak{so}(7)} = 8$. For $k = 3$ and $p = 2$, one gets

$$\text{sdim}_t[0, 0, \dots, 0, 2]_{\mathfrak{osp}(2n+7|2n)} = \dim_t[0, 0, 2]_{\mathfrak{so}(7)} = 1 + 3t + 9t^2 + 9t^3 + 9t^4 + 3t^5 + t^6.$$

and $\text{sdim}[0, 0, \dots, 0, 2]_{\mathfrak{osp}(2n+7|2n)} = \dim[0, 0, 2]_{\mathfrak{so}(7)} = 35$.

Case 3: $n = m + k$, $\mathfrak{osp}(2m + 1|2m + 2k)$.

Again following (35) and (22), one finds

$$\text{sdim}_t V_{\mathfrak{osp}(2m+1|2n)}^{(p/2)^m, (-p/2)^n} = \sum_{\lambda, \lambda_1 \leq p, \lambda_1 \leq k} (-1)^{|\lambda|} \dim V_{\mathfrak{gl}(k)}^{\lambda'} t^{|\lambda|} = \sum_{\mu, \ell(\mu) \leq \min(p, k)} \dim V_{\mathfrak{gl}(k)}^{\mu} (-t)^{|\mu|}. \quad (40)$$

The right hand side is the same expression as (26) (with $t \rightarrow -t$). Following (28) one can write

$$\text{sdim}_t V_{\mathfrak{osp}(2m+1|2n)}^{(p/2)^m, (-p/2)^n} = \dim_{-t} V_{\mathfrak{osp}(1|2k)}^{(-p/2)^k} = \frac{E_{k,p}(-t, \dots, -t)}{(1+t)^k (1-t^2)^{k(k-1)/2}}. \quad (41)$$

Herein, a finite expression for $E_{k,p}(-t, \dots, -t)$ can be found in (32). Note that in terms of Dynkin labels one could write:

$$\text{sdim}_t[0, 0, \dots, 0, p]_{\mathfrak{osp}(2m+1|2m+2k)} = \dim_{-t}[0, \dots, 0, -p]_{\mathfrak{osp}(1|2k)}. \quad (42)$$

Let us again consider some examples. For $k = 3$ and $p = 1$, one finds

$$\text{sdim}_t[0, 0, \dots, 0, 1]_{\mathfrak{osp}(2m+1|2m+6)} = \dim_{-t}[0, 0, -1]_{\mathfrak{osp}(1|6)} = \frac{1}{(1+t)^3} = 1 - 3t + 6t^2 - 10t^3 + 15t^4 - \dots.$$

One could say that the actual superdimension (for $t = 1$) is $1/8$. More general, for arbitrary k ,

$$\text{sdim}_t[0, 0, \dots, 0, 1]_{\mathfrak{osp}(2m+1|2m+2k)} = \dim_{-t}[0, \dots, 0, -1]_{\mathfrak{osp}(1|2k)} = \frac{1}{(1+t)^k}.$$

For $k = 3$ and $p = 2$, one gets

$$\begin{aligned} \text{sdim}_t[0, 0, \dots, 0, 2]_{\mathfrak{osp}(2m+1|2m+6)} &= \dim_{-t}[0, 0, -2]_{\mathfrak{osp}(1|6)} \\ &= \frac{1+t^3}{(1+t)^3(1-t^2)^3} = 1 - 3t + 9t^2 - 18t^3 + 36t^4 - \dots \end{aligned}$$

6 The Lie superalgebra $\mathfrak{osp}(2m|2n)$

The second main case treated here is the Lie superalgebra $D(m, n) = \mathfrak{osp}(2m|2n)$. The weight space of $\mathfrak{osp}(2m|2n)$ has the same basis $\epsilon_1, \dots, \epsilon_m, \delta_1, \dots, \delta_n$, and the distinguished set of simple roots is now

$$\delta_1 - \delta_2, \dots, \delta_{n-1} - \delta_n, \delta_n - \epsilon_1, \epsilon_1 - \epsilon_2, \dots, \epsilon_{m-2} - \epsilon_{m-1}, \epsilon_{m-1} - \epsilon_m, \epsilon_{m-1} + \epsilon_m. \quad (43)$$

It will be helpful to see $D(m, n)$ as a subalgebra of $B(m, n)$. In fact, using the \mathbb{Z} -gradation $\mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_{+1} \oplus \mathfrak{g}_{+2}$ of $\mathfrak{g} = \mathfrak{osp}(2m+1|2n)$ introduced in the previous section, it is easy to see that $\mathfrak{osp}(2m|2n) = \mathfrak{g}_{-2} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_{+2}$, with root structure

$$\begin{array}{ccc} \mathfrak{g}_{-2} & \mathfrak{g}_0 & \mathfrak{g}_{+2} \\ -\delta_i - \delta_j & \delta_i - \delta_j & \delta_i + \delta_j \\ -\epsilon_i - \epsilon_j \ (i \neq j) & \epsilon_i - \epsilon_j & \epsilon_i + \epsilon_j \ (i \neq j) \\ -\epsilon_i - \delta_j & \pm(\epsilon_i - \delta_j) & \epsilon_i + \delta_j \end{array}$$

So also in this case $\mathfrak{g}_0 = \mathfrak{gl}(m|n)$, and it will be useful to consider the representation of the previous section according to $\mathfrak{osp}(2m+1|2n) \supset \mathfrak{osp}(2m|2n) \supset \mathfrak{gl}(m|n)$.

The purpose is to study the irreducible highest weight representation of $\mathfrak{osp}(2m|2n)$ with highest weight given by $(\frac{p}{2}, \dots, \frac{p}{2}; -\frac{p}{2}, \dots, -\frac{p}{2}) = \sum_{i=1}^m \frac{p}{2} \epsilon_i + \sum_{i=1}^n (-\frac{p}{2}) \delta_i$, where p is a non-negative integer, and denoted by $V = V_{\mathfrak{osp}(2m|2n)}^{(p/2)^m, (-p/2)^n}$. According to the simple roots system (43), its Dynkin labels are $[0, 0, \dots, 0, p]$.

Contrary to $V_{\mathfrak{osp}(2m+1|2n)}^{(p/2)^m, (-p/2)^n}$, $V_{\mathfrak{osp}(2m|2n)}^{(p/2)^m, (-p/2)^n}$ has not (yet) been studied in the context of parastatistics. However, the techniques of [35] can also be used in the current case. In particular, let v be the vector of highest weight $(\frac{p}{2}, \dots, \frac{p}{2}; -\frac{p}{2}, \dots, -\frac{p}{2})$, then $\mathbb{C}v$ is a trivial one-dimensional module of $\mathfrak{g}_0 \oplus \mathfrak{g}_{+2}$, and we consider the induced module

$$\overline{V}_{\mathfrak{osp}(2m|2n)}^{(p/2)^m, (-p/2)^n} = \text{Ind}_{\mathfrak{g}_0 \oplus \mathfrak{g}_{+2}}^{\mathfrak{osp}(2m|2n)} \mathbb{C}v.$$

Using the notation $x_i = e^{-\epsilon_i}$, $y_i = e^{-\delta_i}$, and the above root structure, the character is given by:

$$\text{char } \overline{V}_{\mathfrak{osp}(2m|2n)}^{(p/2)^m, (-p/2)^n} = (y_1 \cdots y_n / x_1 \cdots x_m)^{p/2} \frac{\prod_{i,j} (1 + x_i y_j)}{\prod_{i < j} (1 - x_i x_j) \prod_{i \leq j} (1 - y_i y_j)}. \quad (44)$$

Cummins and King [10,11] obtained an expansion of the above product in terms of supersymmetric Schur functions:

$$\begin{aligned} \frac{\prod_{i,j} (1 + x_i y_j)}{\prod_{i < j} (1 - x_i x_j) \prod_{i \leq j} (1 - y_i y_j)} &= \sum_{\beta} s_{\beta}(x|y) \\ &= 1 + s_{1,1}(x|y) + s_{2,2}(x|y) + s_{1,1,1,1}(x|y) + s_{3,3}(x|y) + s_{2,2,1,1}(x|y) + s_{1,1,1,1,1,1}(x|y) + \cdots, \end{aligned} \quad (45)$$

i.e. the sum is over all partitions β such that each part of β appears twice. This can also be written as:

$$\frac{\prod_{i,j} (1 + x_i y_j)}{\prod_{i < j} (1 - x_i x_j) \prod_{i \leq j} (1 - y_i y_j)} = \sum_{\delta} s_{\delta'}(x|y), \quad (46)$$

where the sum is now over all partitions δ with even parts only. Let us denote by \mathcal{B} the set of partitions for which each part appears twice (including the zero partition), and by \mathcal{D} the set of partitions with even parts only. So

$$\text{char } \overline{V}_{\mathfrak{osp}(2m|2n)}^{(p/2)^m, (-p/2)^n} = (y_1 \cdots y_n / x_1 \cdots x_m)^{p/2} \sum_{\lambda \in \mathcal{B}} s_{\lambda}(x|y). \quad (47)$$

Now we return to the irreducible representation $V_{\mathfrak{osp}(2m|2n)}^{(p/2)^m, (-p/2)^n}$. On the one hand, $V_{\mathfrak{osp}(2m|2n)}^{(p/2)^m, (-p/2)^n}$ is a quotient module of $\overline{V}_{\mathfrak{osp}(2m|2n)}^{(p/2)^m, (-p/2)^n}$ with character (47). On the other hand, $V_{\mathfrak{osp}(2m|2n)}^{(p/2)^m, (-p/2)^n}$ is a submodule of $V_{\mathfrak{osp}(2m+1|2n)}^{(p/2)^m, (-p/2)^n}$ in the decomposition $\mathfrak{osp}(2m+1|2n) \supset \mathfrak{osp}(2m|2n)$, with character (34). From this observation, one can deduce:

$$\text{char } V_{\mathfrak{osp}(2m|2n)}^{(p/2)^m, (-p/2)^n} = (y_1 \cdots y_n / x_1 \cdots x_m)^{p/2} \sum_{\lambda \in \mathcal{B}, \lambda_1 \leq p} s_{\lambda}(x|y). \quad (48)$$

To determine $\text{sdim}_t V_{\mathfrak{osp}(2m|2n)}^{(p/2)^m, (-p/2)^n}$, one should (apart from the factor in front of the above sum) specify again $x_i = t$ and $y_j = -t$ in the above character, and so one finds

$$\begin{aligned} \text{sdim}_t V_{\mathfrak{osp}(2m|2n)}^{(p/2)^m, (-p/2)^n} &= \sum_{\lambda \in \mathcal{B}, \lambda_1 \leq p} s_\lambda(t, \dots, t | -t, \dots, -t) \\ &= \sum_{\lambda \in \mathcal{B}, \lambda_1 \leq p} s_\lambda(1, \dots, 1 | -1, \dots, -1) t^{|\lambda|} \\ &= \sum_{\lambda \in \mathcal{B}, \lambda_1 \leq p} \text{sdim} V_{\mathfrak{gl}(m|n)}^\lambda t^{|\lambda|}. \end{aligned} \quad (49)$$

As in the previous section, we now make use of properties of $\mathfrak{gl}(m|n)$ superdimensions.

Case 1: $m = n$, $\mathfrak{osp}(2n|2n)$.

$$\text{sdim}_t V_{\mathfrak{osp}(2n|2n)}^{(p/2)^n, (-p/2)^n} = 1. \quad (50)$$

Case 2: $m = n + k$, $\mathfrak{osp}(2n + 2k|2n)$.

It follows directly from (49) and (22) that

$$\text{sdim}_t V_{\mathfrak{osp}(2m|2n)}^{(p/2)^m, (-p/2)^n} = \sum_{\lambda \in \mathcal{B}, \lambda_1 \leq p} \dim V_{\mathfrak{gl}(k)}^\lambda t^{|\lambda|} = \sum_{\lambda \in \mathcal{B}, \lambda_1 \leq p, \ell(\lambda) \leq k} \dim V_{\mathfrak{gl}(k)}^\lambda t^{|\lambda|}. \quad (51)$$

This is a finite sum. The summation is over all partitions λ with each part appearing twice and whose Young diagrams fit inside the $(k \times p)$ rectangle. Using a character formula for $\mathfrak{so}(2k)$, it is easy to see that this corresponds to the character of an irreducible $\mathfrak{so}(2k)$ representation (see Appendix). There is a distinction between the cases k even and k odd. We find:

$$\text{sdim}_t V_{\mathfrak{osp}(2m|2n)}^{(p/2)^m, (-p/2)^n} = \begin{cases} \dim_t V_{\mathfrak{so}(2k)}^{(p/2)^k} & \text{for } k \text{ even,} \\ \dim_t V_{\mathfrak{so}(2k)}^{(p/2)^{k-1}, -p/2} & \text{for } k \text{ odd;} \end{cases} \quad (52)$$

or, in terms of Dynkin labels:

$$\text{sdim}_t [0, 0, \dots, 0, p]_{\mathfrak{osp}(2n+2k|2n)} = \begin{cases} \dim_t [0, \dots, 0, 0, p]_{\mathfrak{so}(2k)} & \text{for } k \text{ even.} \\ \dim_t [0, \dots, 0, p, 0]_{\mathfrak{so}(2k)} & \text{for } k \text{ odd.} \end{cases} \quad (53)$$

As an example, let $k = 5$ and $p = 1$:

$$\text{sdim}_t [0, 0, \dots, 0, 1]_{\mathfrak{osp}(2n+10|2n)} = \dim_t [0, 0, 0, 1, 0]_{\mathfrak{so}(10)} = 1 + 10t^2 + 5t^4,$$

and thus $\text{sdim}[0, 0, \dots, 0, 1]_{\mathfrak{osp}(2n+10|2n)} = \dim[0, 0, 0, 1, 0]_{\mathfrak{so}(10)} = 16$. For $k = 5$ and $p = 2$, one gets

$$\text{sdim}_t [0, 0, \dots, 0, 2]_{\mathfrak{osp}(2n+10|2n)} = \dim_t [0, 0, 0, 2, 0]_{\mathfrak{so}(10)} = 1 + 10t^2 + 55t^4 + 45t^6 + 15t^8,$$

and $\text{sdim}[0, 0, \dots, 0, 2]_{\mathfrak{osp}(2n+10|2n)} = \dim[0, 0, 0, 2, 0]_{\mathfrak{so}(10)} = 126$.

Case 3: $n = m + k$, $\mathfrak{osp}(2m|2m + 2k)$.

From (49) and (22) one obtains

$$\begin{aligned} \text{sdim}_t V_{\mathfrak{osp}(2m|2n)}^{(p/2)^m, (-p/2)^n} &= \sum_{\lambda \in \mathcal{B}, \lambda_1 \leq p} \dim V_{\mathfrak{gl}(k)}^{\lambda'} (-t)^{|\lambda|} = \sum_{\lambda \in \mathcal{B}, \lambda_1 \leq \min(p, k)} \dim V_{\mathfrak{gl}(k)}^{\lambda'} (-t)^{|\lambda|} \\ &= \sum_{\lambda \in \mathcal{D}, \ell(\lambda) \leq \min(p, k)} \dim V_{\mathfrak{gl}(k)}^\lambda (-t)^{|\lambda|}. \end{aligned} \quad (54)$$

This is an infinite sum, over all partitions λ with even parts only and with $\ell(\lambda) \leq \min(p, k)$. Interestingly, this expression is known and follows from the character of the infinite-dimensional irreducible $\mathfrak{sp}(2k)$ module with highest weight $(-\frac{p}{2}, \dots, -\frac{p}{2})$ [28]. Thus we find:

$$\text{sdim}_t V_{\mathfrak{osp}(2m|2n)}^{(p/2)^m, (-p/2)^n} = \dim_{-t} V_{\mathfrak{sp}(2k)}^{(-p/2)^k}, \quad (55)$$

or, in terms of Dynkin labels:

$$\text{sdim}_t [0, 0, \dots, 0, p]_{\mathfrak{osp}(2m|2m+2k)} = \dim_{-t} [0, \dots, 0, -p/2]_{\mathfrak{sp}(2k)}. \quad (56)$$

As an example, consider first $p = 1$. The series of partitions appearing in (54) is

$$(0) + (2) + (4) + (6) + (8) + \dots$$

So, for $k = 3$, one finds

$$\begin{aligned} \text{sdim}_t [0, 0, \dots, 0, 1]_{\mathfrak{osp}(2m|2m+6)} &= \dim_{-t} [0, 0, -1/2]_{\mathfrak{sp}(6)} \\ &= 1 + 6t^2 + 15t^4 + 28t^6 + 45t^8 + \dots = \frac{1}{2} \left(\frac{1}{(1+t)^3} + \frac{1}{(1-t)^3} \right). \end{aligned}$$

Also in general, one has

$$\text{sdim}_t [0, 0, \dots, 0, 1]_{\mathfrak{osp}(2m|2m+2k)} = \dim_{-t} [0, \dots, 0, -1/2]_{\mathfrak{sp}(2k)} = \frac{1}{2} \left(\frac{1}{(1+t)^k} + \frac{1}{(1-t)^k} \right).$$

For $p = 2$, the series of partitions (54) is

$$(0) + (2) + (4) + (2, 2) + (6) + (4, 2) + (8) + (6, 2) + (4, 4) + \dots$$

For $k = 3$, one finds

$$\text{sdim}_t [0, 0, \dots, 0, 2]_{\mathfrak{osp}(2m|2m+6)} = \dim_{-t} [0, 0, -1]_{\mathfrak{sp}(6)} = 1 + 6t^2 + 21t^4 + 55t^6 + 120t^8 + \dots$$

7 The Lie superalgebra $D(2, 1; \alpha)$

Among the families of simple orthosymplectic Lie superalgebras, there are besides $B(m, n) = \mathfrak{osp}(2m+1|2n)$ and $D(m, n) = \mathfrak{osp}(2m|2n)$, still $C(n+1) = \mathfrak{osp}(2|2n)$ and $D(2, 1; \alpha)$. Since $C(n+1)$ is a type 1 Lie superalgebra (in the sense of Kac [20]), there seem to be no representations that one could describe as “spinor” or “self dual tensors”. So we shall not include $C(n+1)$ in this paper. By the way, all finite-dimensional irreducible representations of $C(n+1)$ are known [38].

For the exceptional family $D(2, 1; \alpha)$ (where α is a real parameter that can be chosen positive), there are representations that one could consider as spinors or self dual tensors. The weight space of $D(2, 1; \alpha)$ is usually expressed in the basis $\epsilon_1, \epsilon_2, \epsilon_3$, with the distinguished set of simple roots given by

$$2\epsilon_2, \epsilon_1 - \epsilon_2 - \epsilon_3, 2\epsilon_3. \quad (57)$$

The purpose of this section is to consider irreducible representations $V(\Lambda)$ with highest weight Λ given by the Dynkin labels $[0, 0, p]$ in the above simple root system. These are infinite-dimensional representations. For $p = 1$ this is the “spinor” irrep of $D(2, 1; \alpha)$, for $p = 2$ the “self dual tensor”. In terms of the ϵ -basis, the highest weight is given by $\Lambda = (-\frac{\alpha p}{\alpha+1}, 0, p)$. Such representations are “doubly atypical”. All irreducible highest weight representations of $D(2, 1; \alpha)$ have been described

in [37]. For the irreps under consideration one can describe the structure by means of the decomposition with respect to the even subalgebra $\mathfrak{su}(1,1) \oplus \mathfrak{su}(2) \oplus \mathfrak{su}(2)$ (isomorphic to $\mathfrak{sp}(2) \oplus \mathfrak{so}(4)$). From [37], the following decompositions hold:

$$p = 1: \quad V(\Lambda) \rightarrow \left(-\frac{\alpha p}{\alpha + 1}; 0, 1\right) \oplus \left(-\frac{\alpha p}{\alpha + 1} - 1; 1, 0\right) \quad (58)$$

$$p \geq 2: \quad V(\Lambda) \rightarrow \left(-\frac{\alpha p}{\alpha + 1}; 0, p\right) \oplus \left(-\frac{\alpha p}{\alpha + 1} - 1; 1, p - 1\right) \oplus \left(-\frac{\alpha p}{\alpha + 1} - 2; 0, p - 2\right). \quad (59)$$

The weights of such an $\mathfrak{su}(1,1)$ irrep with highest weight μ ($\mu < 0$) are given by $(\mu, \mu - 2, \mu - 4, \dots)$: this is a negative discrete series representation. So for the case $p = 1$, with $\mu = -\frac{\alpha p}{\alpha + 1}$, one has:

- at level μ : the $\mathfrak{su}(2) \oplus \mathfrak{su}(2)$ irrep $(0, 1)$ of dimension 2
- at level $\mu - 1$: the $\mathfrak{su}(2) \oplus \mathfrak{su}(2)$ irrep $(1, 0)$ of dimension 2
- at level $\mu - 2$: the $\mathfrak{su}(2) \oplus \mathfrak{su}(2)$ irrep $(0, 1)$ of dimension 2
- at level $\mu - 3$: the $\mathfrak{su}(2) \oplus \mathfrak{su}(2)$ irrep $(1, 0)$ of dimension 2
- etc.

Thus one could say that the superdimension of $V(\Lambda)$ according to the above gradation (i.e. according to the $\mathfrak{su}(1,1)$ diagonal element) is given by

$$2 - 2t + 2t^2 - 2t^3 + 2t^4 - \dots = \frac{2}{1 + t}.$$

In the limit $t \rightarrow 1$, this gives superdimension 1.

In the previous section, we always considered a gradation with respect to the subalgebra $\mathfrak{gl}(m|n)$. For the irrep under consideration, the decomposition with respect to $D(2, 1; \alpha) \rightarrow \mathfrak{gl}(2|1)$ consists of

- a $\mathfrak{gl}(2|1)$ singlet with weight $(\mu, 0, 1)$
- a $\mathfrak{gl}(2|1)$ irrep with weights $(\mu, 0, -1), (\mu - 1, -1, 0), (\mu - 1, 1, 0), (\mu - 2, 0, 1)$, of dimension 4 and superdimension 0
- a $\mathfrak{gl}(2|1)$ irrep with weights $(\mu - 2, 0, -1), (\mu - 3, -1, 0), (\mu - 3, 1, 0), (\mu - 4, 0, 1)$, of dimension 4 and superdimension 0
- etc.

Hence also according to this gradation the superdimension is $1 + 0 + 0 + \dots = 1$, whereas the dimension is $1 + 4 + 4 + \dots$.

For the case $p \geq 2$, the decomposition is given by (59). The dimensions of the $\mathfrak{su}(2) \oplus \mathfrak{su}(2)$ irreps appearing here are, respectively, $p + 1$, $2p$ and $p - 1$. According to the gradation by the $\mathfrak{su}(1, 1)$ diagonal element, one finds as superdimension:

$$\begin{aligned} & (p + 1) - t(2p) + t^2((p + 1) + (p - 1)) - t^3(2p) + t^4((p + 1) + (p - 1)) + \dots \\ & = (p + 1) - 2pt + 2pt^2 - 2pt^3 + 2pt^4 + \dots = 1 - p + \frac{2p}{1 + t}. \end{aligned}$$

In the limit $t \rightarrow 1$, this gives superdimension 1. Also according to the decomposition with respect to $\mathfrak{gl}(2|1)$ one finds a singlet followed by an infinite series of irreps of dimension 4 and superdimension 0, thus the same result.

Conclusion: all $D(2, 1; \alpha)$ irreps $[0, 0, p]$ have superdimension 1, which is the same as the dimension of the $\mathfrak{so}(2)$ irrep $[p]$.

8 Summary: Dynkin diagrams

The main results of sections 5 and 6 can be best summarized and illustrated by means of Dynkin diagrams, carrying the Kac-Dynkin labels of the representations studied here.

For the case $B(m, n) = \mathfrak{osp}(2m + 1|2n)$, one has, in an obvious notation, from (39) and (42):

$$\begin{aligned} & \text{sdim} \left(\begin{array}{ccccccccc} 0 & 0 & & 0 & 0 & 0 & & 0 & p \\ \circ & \circ & \cdots & \circ & \otimes & \circ & \cdots & \circ & \circ \\ 1 & 2 & & n-1 & n & n+1 & & m+n-1 & m+n \end{array} \right) = \\ & \text{dim} \left(\begin{array}{ccccccc} 0 & 0 & & 0 & p \\ \circ & \circ & \cdots & \circ & \circ \\ 1 & 2 & & m-n-1 & m-n \end{array} \right) & \text{if } m > n \\ & \text{dim} \left(\begin{array}{ccccccc} 0 & 0 & & 0 & -p \\ \circ & \circ & \cdots & \circ & \bullet \\ 1 & 2 & & n-m-1 & n-m \end{array} \right) & \text{if } m < n \end{aligned}$$

For the case $D(m, n) = \mathfrak{osp}(2m|2n)$, the result is, following (53) and (56):

$$\begin{aligned}
& \text{sdim} \left(\begin{array}{c} \begin{array}{ccccccccc} \circ & \cdots & \circ & \circ & \circ & \cdots & \circ & \circ & \circ \\ 1 & & n-1 & n & n+1 & & m+n-2 & m+n-1 & m+n \end{array} \\ \begin{array}{c} \circ \\ \circ \end{array} \\ \begin{array}{c} 0 \\ p \end{array} \end{array} \right) = \\
& \text{dim} \left(\begin{array}{c} \begin{array}{ccccccc} \circ & \circ & \cdots & \circ & \circ & \circ & \circ \\ 1 & 2 & & m-n-2 & m-n-1 & m-n & \end{array} \\ \begin{array}{c} \circ \\ \circ \end{array} \\ \begin{array}{c} p \\ 0 \end{array} \end{array} \right) & \quad \text{if } m > n \text{ and } m - n \text{ odd} \\
& \text{dim} \left(\begin{array}{c} \begin{array}{ccccccc} \circ & \circ & \cdots & \circ & \circ & \circ & \circ \\ 1 & 2 & & m-n-2 & m-n-1 & m-n & \end{array} \\ \begin{array}{c} \circ \\ \circ \end{array} \\ \begin{array}{c} 0 \\ p \end{array} \end{array} \right) & \quad \text{if } m > n \text{ and } m - n \text{ even} \\
& \text{dim} \left(\begin{array}{c} \begin{array}{ccccccc} \circ & \circ & \cdots & \circ & \circ & \circ & \circ \\ 1 & 2 & & n-m-1 & n-m & & \end{array} \\ \begin{array}{c} \circ \\ \circ \end{array} \\ \begin{array}{c} 0 \\ -p/2 \end{array} \end{array} \right) & \quad \text{if } m < n
\end{aligned}$$

Acknowledgments

NIS is thankful to Professor V.K. Dobrev for constructive discussions. NIS and JVdJ were supported by the Joint Research Project “Lie superalgebras - applications in quantum theory” in the framework of an international collaboration programme between the Research Foundation – Flanders (FWO) and the Bulgarian Academy of Sciences. NIS was partially supported by Bulgarian National Science Fund, grant DFNI T02/6. This research (JT-M) was supported in part by the Intramural Research Program of the NIH, U.S. National Library of Medicine.

Appendix

In this appendix we give character formulas for the irreducible representations of $\mathfrak{so}(2k)$ which are relevant in the context of this paper. Following the notation of this paper, these are the representations with Dynkin labels $[0, 0, \dots, 0, 0, p]$ and $[0, 0, \dots, 0, p, 0]$ (with p a positive integer), or with highest weights $(\frac{p}{2}, \frac{p}{2}, \dots, \frac{p}{2}, \frac{p}{2})$ and $(\frac{p}{2}, \frac{p}{2}, \dots, \frac{p}{2}, -\frac{p}{2})$ respectively. The modules will be denoted as $V_{\mathfrak{so}(2k)}^{(p/2)^k}$ and $V_{\mathfrak{so}(2k)}^{(p/2)^{k-1}, -p/2}$. Of course, the characters of these representations could be deduced from Weyl’s formula. Here, however, it is more useful to give these characters as an expansion in Schur functions. This expression can actually be deduced from the $\mathfrak{so}(2k) \rightarrow \mathfrak{gl}(n)$ branching rule, given in [27, Section IV]. Working out these branchings, there is a distinction between k even and k odd. Explicitly, one finds:

Case k even:

$$\text{char}[0, \dots, 0, p]_{\text{so}(2k)} = \text{char } V_{\text{so}(2k)}^{(p/2)^k} = (x_1 \cdots x_k)^{-p/2} \sum_{\lambda \in \mathcal{B}: \lambda_1 \leq p, \ell(\lambda) \leq k} s_\lambda(x). \quad (60)$$

So the sum is over all partitions λ for which each part appears twice, and such that the Young diagram of λ fits inside the rectangle of width p and height k .

$$\text{char}[0, \dots, p, 0]_{\text{so}(2k)} = \text{char } V_{\text{so}(2k)}^{(p/2)^{k-1}, -p/2} = (x_1 \cdots x_k)^{-p/2} \sum_{\lambda \in \mathcal{B}: \lambda_1 \leq p, \ell(\lambda) \leq k-2} s_{(p, \lambda)}(x). \quad (61)$$

Here the sum is again over all partitions λ for which each part appears twice, but such that the Young diagram of $\mu = (p, \lambda)$ fits inside the rectangle of width p and height k .

Case k odd:

$$\text{char}[0, \dots, 0, p]_{\text{so}(2k)} = \text{char } V_{\text{so}(2k)}^{(p/2)^k} = (x_1 \cdots x_k)^{-p/2} \sum_{\lambda \in \mathcal{B}: \lambda_1 \leq p, \ell(\lambda) \leq k-1} s_{(p, \lambda)}(x). \quad (62)$$

$$\text{char}[0, \dots, p, 0]_{\text{so}(2k)} = \text{char } V_{\text{so}(2k)}^{(p/2)^{k-1}, -p/2} = (x_1 \cdots x_k)^{-p/2} \sum_{\lambda \in \mathcal{B}: \lambda_1 \leq p, \ell(\lambda) \leq k-1} s_\lambda(x). \quad (63)$$

Note that in (63), due to the fact that λ has an even number of parts (since $\lambda \in \mathcal{B}$), the condition $\ell(\lambda) \leq k-1$ could just as well be replaced by $\ell(\lambda) \leq k$ since k is odd.

References

- [1] Balantekin A B and Bars I 1981 Representations of supergroups *J. Math. Phys.* **22** 1810-1818
- [2] Becchi C, Rouet A and Stora A 1975 Renormalization of the abelian Higgs-Kibble model *Commun. Math. Phys.* **42** 127-162; 1976 Renormalization of gauge theories *Ann. Phys.* 287-321; Tyutin I V 1975, Lebedev Institute Preprint: FIAN-39 (unpublished)
- [3] Berele A and Regev A 1987 Hook Young diagrams with applications to combinatorics and to representations of Lie superalgebras *Adv. Math.* **64** 118-175
- [4] Baulieu L and Thierry-Mieg J 1983 Covariant quantization of non-Abelian antisymmetric tensor gauge theories *Nucl. Phys. B* **228** 259-284
- [5] Bonora L 2016 BRST and supermanifolds *Nucl. Phys. B* **912** 103-118
- [6] Bonora L and Tonin M 1981 Superfield formulation of extended BRS symmetry *Phys. Lett. B* **98** 48-50
- [7] Bracken A J and Green H S 1972 Algebraic Identities for Parafermi Statistics of Given Order *Nuovo Cim.* **9** 349-365
- [8] Cartan É 1938 *Leçons sur la Théorie des Spineurs I, II* Paris: Hermann; English translation 1966 *The Theory of Spinors* Paris: Hermann

- [9] Cremmer E, Julia B and Scherk J 1978 Supergravity Theory in Eleven-Dimensions *Phys. Lett.* **76B** 409-412
- [10] Cummins C 1986 Application of S-function techniques to representations of Lie superalgebras and symmetry breaking *PhD Thesis* University of Southampton
- [11] Cummins C J and King R C 1987 Some noteworthy S-function identities *Preprint Centre des Recherche Mathematiques* Université de Montréal, Canada
- [12] De Witt B S 1965 *Dynamical Theory of Groups and Fields* (New York: Gordon and Breach)
- [13] Delbourgo R and Jarvis P D 1982 Extended BRS invariance and $OSp(4/2)$ supersymmetry *J. Phys. A: Math. Gen.* **15** 611-625; Delbourgo R, Jarvis P D and Thompson G 1982 Extended Becchi-Rouet-Stora invariance for gravity via local $Osp(4/2)$ supersymmetry *Phys. Rev. D* **26** 775-786
- [14] Faddeev L D and Popov V N 1967 Feynman Diagrams for the Yang-Mills Field *Phys. Lett. B* **25** 29-30
- [15] Feynman R P 1963 Quantum Theory of Gravitation *Acta Phys. Polon.* **24** 697-722
- [16] Frappat L, Sciarrino A and Sorba P 2000 *Dictionary on Lie Algebras and Superalgebras* London: Academic Press
- [17] Gould M D and Isaac P S 2015 Reduced matrix elements of the orthosymplectic Lie superalgebra *J. Phys A: Math. Theor.* **48** 025201
- [18] Green H S and Jarvis P D 1983 Casimir invariants, characteristic identities, and Young diagrams for color algebras and superalgebras *J. Math. Phys.* **24** 1681-1687
- [19] Green M B and Schwarz J H 1983 Extended Supergravity in 10 Dimensions *Phys. Lett. B* **122** 143-147
- [20] Kac V G 1977 Lie superalgebras *Adv. Math.* **26** 8-96
- [21] Kac V G 1978 Representations of Classical Lie Superalgebras *Lect. Notes in Math.* **626** 597-626
- [22] Kac V G and Wakimoto M 1994 Integrable highest weight modules over affine superalgebras and number theory *Progress in Math.* **123** 415-456
- [23] Kac V G and Wakimoto M 2016 Representations of affine superalgebras and mock theta functions. III *Izv. Math* **80** 693-750
- [24] King R C 1983 Supersymmetric functions and the Lie supergroup $U(m/n)$ *Ars. Combin.* **16A** 269-287
- [25] King R C 1983 Generalised Young tableaux for Lie algebras and superalgebras *Lect. Notes in Phys.* **180** 41-47
- [26] King R C 1990 S-functions and characters of Lie algebras and superalgebras *IMA Volumes in Mathematics and its Applications* **19** 226-261
- [27] King R C and Wybourne B G 2000 Analogies between finite-dimensional irreducible representations of $SO(2n)$ and infinite-dimensional irreducible representations of $Sp(2n, R)$ *J. Math. Phys.* **41** 5002-5019

- [28] King R C 2013 *From Palev's Study of Wigner Quantum Systems to New Results on Sums of Schur Functions* Proceedings in Mathematics & Statistics **36** 61-75
- [29] Lievens S, Stoilova N I and Van der Jeugt J 2008 The paraboson Fock space and unitary irreducible representations of the Lie superalgebra $\mathfrak{osp}(1|2n)$ *Commun. Math. Phys.* **281** 805-826
- [30] Macdonald I G 1995 *Symmetric Functions and Hall Polynomials. 2nd edition* Oxford: Oxford University Press
- [31] Moens E M and Van der Jeugt J 2004 On dimension formulas for $gl(m|n)$ representations *J. Lie Theory* **14** 523-535
- [32] Ne'eman Y 1978 Spinor type fields with linear, affine and general coordinate transformations *Ann. Inst. Henri Poincare A* **28** 369-378
- [33] Serre J 1987 *Complex Semisimple Lie Algebras* New York: Springer-Verlag, pp. 48-49 and 52-55
- [34] Stoilova N I and Van der Jeugt J 2008 The parafermion Fock space and explicit $\mathfrak{so}(2n + 1)$ representations *J. Phys A: Math. Theor.* **41** 075202
- [35] Stoilova N I and Van der Jeugt J 2015 A class of infinite-dimensional representations of the Lie superalgebra $\mathfrak{osp}(2m + 1|2n)$ and the parastatistics Fock space *J. Phys A: Math. Theor.* **48** 155202
- [36] Thierry-Mieg J 1984 Superspin et dimension négative *C. R. Acad. Sc. Paris, Série II* **299** (19) 1309-1312
- [37] Van der Jeugt J 1985 Irreducible representations of the exceptional Lie superalgebras $D(2, 1; \alpha)$ *J. Math. Phys.* **26** 913-924
- [38] Van der Jeugt J 1991 Character formulae for the Lie superalgebra $C(n)$ *Commun. Algebra* **19** 199-222
- [39] Yamane H 2014 Exposition on classification of finite dimensional irreducible representations of the Lie superalgebra $B(m, n)$ *J. Phys.: Conference Series* **563** 012035