

EXTENSION THEOREMS FOR REDUCTIVE GROUP ACTIONS ON COMPACT KAEHLER MANIFOLDS

BY ANDREW J. SOMMESE

Communicated by Stephen S. Shatz, March 13, 1975

Let G be a connected complex reductive Lie group. Noting [3], [7], [8] that G has the structure of a linear algebraic group, let \bar{G} be any projective manifold in which G is Zariski open and which induces the above algebraic structure on G . The purpose of the present note is to announce

PROPOSITION I. *Let G be as above and act holomorphically on a compact Kaehler manifold X . Assume that the Lie algebra of holomorphic vector fields on X generated by G is annihilated by every holomorphic one form. Let $\Phi: Y \rightarrow X$ be a holomorphic map where Y is a normal reduced analytic space. Consider the equivariant map $\Phi': G \times Y \rightarrow X$; Φ' extends meromorphically (in the sense of Remmert) to $\bar{G} \times Y$.*

REMARKS. The condition on vector fields annihilated by one forms is automatically satisfied if (cf. [12]–[14]) $H^1(X, \mathbb{Q}) = 0$, or G is semisimple, or if every generator of the solvable radical of G has a fixed point, or if G is a linear algebraic group acting algebraically on a projective X . Taking Y to be a point, one gets the orbits of G to be Zariski open in their closures which are analytic sets. A simple corollary is the classical result that there is only one structure of a linear algebraic group on G (cf. [7]), and in fact any reductive connected subgroup of an algebraic group over \mathbb{C} is an algebraic subgroup.

As a further application of the techniques used, a new proof of an improved form of a fixed point theorem (cf. [12], [13], [14]) of the author is given:

PROPOSITION II. *Let S be a connected solvable Lie group acting holomorphically on a compact Kaehler manifold X . The following are equivalent:*
(a) *S has a fixed point on X .*

AMS (MOS) subject classifications (1970). Primary 32M05, 32J25, 53C55; Secondary 22E10.

Key words and phrases. Reductive group actions, Kaehler manifolds, linear algebraic groups.

Copyright © 1975, American Mathematical Society

(b) S leaves a compact set in a fibre of the Albanese map invariant.

(c) S has a fixed point within any compact set K on X that S leaves invariant.

(d) The Lie algebra of vector-fields that S generates on X is annihilated by every holomorphic one form on X .

REMARK. The assertion (c) where K is any compact set is new; the method of proof allows one to relax the compactness of X and show if, in addition, $H^1(X, \mathcal{O}_X) = 0$, then (c) is true.

The following is the fundamental observation on which everything rests.

LEMMA. Let X be a compact Kaehler manifold and $\rho: \mathbf{C}^* \rightarrow \text{Aut}(X)$ a holomorphic \mathbf{C}^* action that has at least one fixed point. Let $A: \mathbf{C}^* \rightarrow X$ be a holomorphic equivariant map onto an orbit: then A extends to a homomorphic equivariant map \tilde{A} of $\mathbf{C}P^1$ to X .

PROOF. Assume without loss of generality that $A(\mathbf{C}^*)$ is not a point. Let μ be a Kaehler metric on X and ω the associated Kaehler form. Assume that μ has been averaged with respect to the circle subgroup $S^1 \subseteq \mathbf{C}^*$. Let χ be the holomorphic vector-field on X associated to $\rho: \mathbf{C}^* \rightarrow \text{Aut}(X)$.

Because of equivariance, the Jacobian, dA , of A , maps some constant multiple of $z(\partial/\partial z)$ onto the restriction of the vector-field χ to $A(\mathbf{C}^*)$. Without loss of generality this constant is assumed to be one.

Let $A^*\mu = a(r) dz \otimes d\bar{z}$ where $a(r)$ is positive and depends only on r due to the S^1 averaging of μ . $A^*\omega = (i/2)a(r) dz \wedge d\bar{z}$.

$$\begin{aligned} \mu(\chi, \chi) &= \mu\left(dA\left(z\frac{\partial}{\partial z}\right), dA\left(z\frac{\partial}{\partial z}\right)\right) = A^*\mu\left(z\frac{\partial}{\partial z}, z\frac{\partial}{\partial z}\right) \\ &= a(r)|z|^2 \leq M < \infty \end{aligned}$$

where $\sup_X \mu(\chi, \chi) = M < \infty$.


Now by Lichnerowicz [5] there exists a C^∞ function ϕ on X such that $\bar{\partial}\phi = \omega(\chi)$. Pulling back and, without confusion, letting ϕ stand for $A^*\phi = \phi(A(z))$, one has

$$\frac{i}{2} z a(r) d\bar{z} = \frac{\partial\phi}{\partial\bar{z}} d\bar{z} \quad \text{or} \quad \frac{i}{2} z a(r) = \frac{\partial\phi}{\partial\bar{z}}.$$

Now fix one circle, say the unit circle $C_1 \subset \mathbf{C}^*$ and let $C_R = \{z \in \mathbf{C}^* \mid |z| = R\}$. Assume $R > 1$; C_1 and C_R bound an annulus \tilde{A} with $\partial\tilde{A} = C_R - C_1$. Now

$$\begin{aligned} \int_{\tilde{A}} A^* \mu &= \int_{\tilde{A}} \int \frac{i}{2} a(r) dz \wedge d\bar{z} = - \int_{\tilde{A}} \int \frac{\partial \phi}{\partial \bar{z}} \frac{d\bar{z} \wedge dz}{z} \\ &= - \int_{C_R} \phi \frac{dz}{z} + \int_{C_1} \phi \frac{dz}{z} = \frac{1}{i} \int_0^{2\pi} \phi(Re^{i\theta}) d\theta - C \end{aligned}$$

with C a constant. Now $|\int_0^{2\pi} \phi(Re^{i\theta}) d\theta| \leq M' < \infty$ since ϕ is the pullback of a bounded function on X .



Therefore $\int_{\tilde{A}} A^* \mu \leq M'' < \infty$ where M'' is a positive constant independent of R . Thus by Bishop's extension theorem (cf. [1], [2]), A extends holomorphically over ∞ . An identical argument gives extension at 0. Q.E.D.

Using the above Lemma and the Levi-Griffiths-Shiffman-Siu extension theorem (cf. [2], [9], [10], [11]) repeatedly, one proves the result for $SL(2, \mathbb{C})$ and groups of the form $(\mathbb{C}^*)^n$ that have a fixed point on X . Then one proves it for one parameter unipotent subgroup of G by using the above $SL(2, \mathbb{C})$ result on an $SL(2, \mathbb{C})$ in G containing the subgroup; this can be done by Jacobson-Morosow (cf. [4]). One now proves it for a Borel subgroup of G and uses an argument depending on the fact that one has a locally trivial fibring of G over G/B which is compact.

In the very interesting paper [6] of Lieberman, related matters are discussed.

BIBLIOGRAPHY

1. E. Bishop, *Conditions for the analyticity of certain sets*, Michigan Math. J. 11 (1964), 289–304. MR 29 #6057.
2. P. A. Griffiths, *Two theorems on extensions of holomorphic mappings*, Invent. Math. 14 (1971), 27–62. MR 45 #2202.
3. G. Hochschild and G. D. Mostow, *Automorphisms of affine algebraic groups*, J. Algebra 13 (1969), 535–543. MR 41 #315.

4. N. Jacobson, *Lie algebras*, Interscience Tracts in Pure and Appl. Math., no. 10, Interscience, New York and London, 1962. MR 26 #1345.
5. A. Lichnerowicz, *Variétés kähleriennes et première classe de Chern*, Differential Geometry 1 (1967), 195–223. MR 37 #2150.
6. D. Lieberman, *Holomorphic vector-fields and rationality* (unpublished manuscript).
7. G. D. Mostow, *Representative functions on a Lie group*, Some Recent Advances in the Basic Sciences, Vol. 2 (Proc. Annual Sci. Conf., Belfer Grad. School Sci., Yeshiva Univ., New York, 1965–1966), Belfer Graduate School of Science, Yeshiva University, New York, 1969, pp. 209–226. MR 42 #6156.
8. J.-P. Serre, *Algèbres de Lie semi-simples complexes*, Benjamin, New York and Amsterdam, 1966. MR 35 #6721.
9. B. Shiffman, *Extensions of positive line bundles and meromorphic maps*, Invent. Math. 15 (1972), 332–347.
10. Y.-T. Siu, *Analyticity of sets associated to Lelong numbers and the extension of meromorphic maps*, Bull. Amer. Math. Soc. 79 (1973), 1200–1205.
11. ———, *Analyticity of sets associated to Lelong numbers and the extension of closed positive currents*, Invent. Math. 27 (1974), 53–156.
12. A. J. Sommese, *Algebraic properties of the period mapping*, Thesis, Princeton Univ., 1973.
13. ———, *Borel's fixed point theorem for Kaehler manifolds and an application*, Proc. Amer. Math. Soc. 41 (1973), 51–54. MR 48 #579.
14. ———, *Holomorphic vector-fields on compact Kaehler manifolds*, Math. Ann. 210 (1974), 75–82.

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, NEW HAVEN, CONNECTICUT 06520