

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Faculty Publications in Food Science and  
Technology

Food Science and Technology Department

---

2019

## Extraction of astaxanthin from engineered *Camelina sativa* seed using ethanol-modified supercritical carbon dioxide

Liyang Xie

Edgar B. Cahoon

Yue Zhang

Ozan Ciftci

Follow this and additional works at: <https://digitalcommons.unl.edu/foodsciefacpub>



Part of the [Chemical Engineering Commons](#), and the [Food Science Commons](#)

---

This Article is brought to you for free and open access by the Food Science and Technology Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications in Food Science and Technology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

# Extraction of astaxanthin from engineered *Camelina sativa* seed using ethanol-modified supercritical carbon dioxide

Liyang Xie,<sup>1</sup> Edgar Cahoon,<sup>2</sup> Yue Zhang,<sup>1</sup> & Ozan N. Ciftci<sup>1</sup>

<sup>1</sup> Department of Food Science and Technology, University of Nebraska–Lincoln, Lincoln, NE 68588-6205, USA

<sup>2</sup> Center for Plant Science Innovation and Department of Biochemistry, University of Nebraska–Lincoln, Lincoln, NE 68588, USA

Corresponding author — O.N. Ciftci, email ciftci@unl.edu

## Abstract

Natural astaxanthin, a high-value carotenoid that is currently extracted mainly from marine organisms, was extracted from engineered camelina seed using ethanol-modified supercritical carbon dioxide (SC-CO<sub>2</sub>) for the first time, and compared with hexane and accelerated solvent extraction using hexane and ethanol. Response surface methodology (RSM) based on central composite rotatable design was employed to investigate the effect of pressure (30–45 MPa), temperature (40–60 °C), and ethanol concentration (10–35%, w/w). RSM-optimized conditions (41.6 MPa, 36.6 °C and 42.0% ethanol concentration) predicted the astaxanthin concentration as 437 µg/g oil, whereas the actual concentration was 421 ± 14 µg/g oil. Astaxanthin concentration in accelerated solvent extracted oil was significantly lower than that in ethanol-modified SC-CO<sub>2</sub>- and hexane-extracted oils ( $P < 0.05$ ). Oils extracted with ethanol-modified SC-CO<sub>2</sub> had the highest antioxidant activity. Results indicated that ethanol-modified SC-CO<sub>2</sub> extraction method can be successfully used as a green method to extract astaxanthin from high oil feedstocks.

**Keywords:** Astaxanthin, Supercritical carbon dioxide, Extraction, Camelina seed


---

Published in *The Journal of Supercritical Fluids* 143 (2019), pp 171–178.

doi: 10.1016/j.supflu.2018.08.013

Copyright © 2018 Elsevier B.V. Used by permission.

Submitted 27 June 2018; revised 17 August 2018; accepted 19 August 2018; published 20 August 2018.



## 1. Introduction

Astaxanthin (3,3-dihydroxy- $\beta$ -carotene-4,4-dione) is a red color fat-soluble pigment which belongs to the keto-carotenoid family. It has been reported that astaxanthin's antioxidant activity is ten times higher than other carotenoids such as beta-carotene and lutein, and over 500 hundred times higher than tocopherol [1]. Growing evidence also shows that astaxanthin provides beneficial effects on human health, including the enhancement of general well-being and immune system, protection against lipid membrane peroxidation and DNA damage, gastrointestinal cancers, degenerative ailments such as Parkinson's and Alzheimer's diseases; chronic inflammatory diseases; metabolic disorders such as diabetes; and cardiovascular diseases [2,3].

Astaxanthin is the highest value carotenoid that is currently used in the food, feed, nutraceutical and pharmaceutical industries. Currently, more than 95% of the astaxanthin used in aquaculture is synthesized artificially [2]. There is an increasing demand for the natural astaxanthin due to growing demand for natural ingredients. Currently, the most popular source of natural astaxanthin is microalgae *Haematococcus pluvialis*. Various microorganisms such as green algae *Haematococcus pluvialis* and *Chlorella zofingiensis*, red yeast *Phaffia rhodozyma* and marine bacterium *Agrobacterium aurantiacum* can produce astaxanthin [4]. Other natural sources of astaxanthin include wild Pacific sockeye salmon, lobster, arctic shrimp, crab, crawfish, red trout, algae, and krill. However, the potential of current sources is limited and is not sufficient to supply the global astaxanthin market [5].

In recent years, there have been some attempts to synthesize astaxanthin in carrot [6], tomato [7], and corn [5] through metabolic engineering. However, the engineered plants make the development of new extraction methods crucial as current methods are based on marine organisms, which are high water content materials. Camelina seed contains high amount of oil (€40%) and its water content is much lower than that of current natural astaxanthin sources, which in turn affect the selection of the solvent and extraction method and conditions for efficient recovery of astaxanthin from the source.

Astaxanthin is sensitive to light and oxygen; therefore, may degrade during extraction. Moreover, there is a growing demand for clean extraction methods by consumers and manufacturers for labeling and marketing purposes. Supercritical carbon dioxide (SC-CO<sub>2</sub>) has been used as an alternative clean extraction method for various oilseeds such as flaxseed [8], sunflower seed [9] and watermelon seed [10] and various lipid compounds such as lycopene from tomato seeds [11] and lutein from spinach [12]. Pure SC-CO<sub>2</sub> is nonpolar; therefore, it extracts nonpolar compounds. Because astaxanthin is slightly polar, the polarity of SC-CO<sub>2</sub> was improved by modifying ethanol, which is a food grade solvent, to extract astaxanthin from *Haematococcus pluvialis* [13], Brazilian redspotted shrimp waste [14] and tiger shrimp [15].

There is no reported study on the extraction of astaxanthin from an oilseed using SC-CO<sub>2</sub> or ethanol-modified SC-CO<sub>2</sub>. In this study, engineered camelina seed was used for the first time as an alternative nonmarine astaxanthin source. The main objective of this study was to investigate the feasibility of ethanol-modified SC-CO<sub>2</sub> extraction of astaxanthin from engineered camelina seed. The specific objectives were: (i) to study the effects of ethanol modified-SC-CO<sub>2</sub> extraction parameters, namely, pressure, temperature and ethanol concentration on the extraction of astaxanthin using response surface methodology (RSM); (ii) to optimize the ethanol-modified SC-CO<sub>2</sub> extraction conditions using RSM; (iii) to compare the SC-CO<sub>2</sub>,<sub>opt</sub> extraction with conventional hexane extraction and accelerated ethanol and hexane extractions in terms of oil yield and astaxanthin concentrations of the extracted oils; and iv) to evaluate the effect of the extraction method on the antioxidant activity of the astaxanthin-rich camelina seed oils.

## 2. Materials and methods

### 2.1. Materials

Astaxanthin-enriched camelina seeds were provided by the Plant Innovation Center at the University of Nebraska-Lincoln. Seeds were ground using an analytical mill (A11 basic, IKA Works, Inc., Wilmington, NC, USA) and sieved to obtain the particles smaller than 0.3 mm. CO<sub>2</sub> (99.99% purity) was purchased from Matheson (Lincoln, NE, USA). Astaxanthin from *Haematococcus pluvialis* (≥97% purity) and trans-β-apo-8'-carotenal (≥96% purity) standards were purchased from Sigma Aldrich (St Louis, MO, USA). Rac-5,7-Dimethyltocol was purchased from Matreya LLC. (State College, PA, USA). All other reagents and solvents were of analytical or chromatographic grade.

### 2.2. Ethanol-modified SC-CO<sub>2</sub> extraction

Ethanol-modified SC-CO<sub>2</sub> extractions were carried out in a laboratory scale SC-CO<sub>2</sub> extraction system (SFT 110, Supercritical Fluids, Inc., Newark, DE, USA). Schematic diagram of the system was reported previously [16]. For each experiment, the extraction vessel was loaded with 10 g of ground camelina seed. The air in the vessel was flushed out by opening the CO<sub>2</sub> cylinder before each run for 2 min. Then, the shutoff valve was closed, and the extraction vessel heated to the extraction temperature in the oven of the system. After reaching the set extraction temperature, CO<sub>2</sub> was pumped into the system using the high-pressure CO<sub>2</sub> pump, and ethanol was pumped into the system at an inlet point before the extraction vessel using the co-solvent pump at the predetermined flow rates to attain the set ethanol concentrations. Extraction pressure was monitored and maintained constant using the CO<sub>2</sub> pump. A static extraction time of 20 min was established by keeping the shut-off valve closed. Then, the shut-off valve was opened, and the extracted oil was collected continuously in an amber glass sample collection vial held in a cold trap at -10 °C. CO<sub>2</sub> flow rate was maintained at 1 L/min (measured at ambient conditions) with a heated micrometering valve, and measured by the gas flow meter placed after the sample collection vial. The ethanol in the extracted oils was evaporated under nitrogen flow at 40 °C. The amount of oil extracted was determined gravimetrically and the oil

yield (% w/w) was obtained by dividing the mass of oil extracted by the mass of camelina seed used for extraction. The headspace of the vials containing the extracted oil were filled with nitrogen and stored at  $-20\text{ }^{\circ}\text{C}$  until analyzed for astaxanthin concentration.

### 2.2.1. Experimental design

RSM based on a central composition rotatable design (CCRD) with three variables at three levels were used to investigate the effects of extraction variables (pressure, temperature, and ethanol concentration) on the astaxanthin concentration of the extracted oils. Extraction time was limited to 180 min. The three different levels of the three variables were represented in codes as  $-1$ ,  $0$ , and  $+1$ . Two extreme levels were coded as  $-1.68$  and  $+1.68$ . The actual levels of the coded and uncoded variables generated by the Design Expert software (Stat-Ease Inc., Minneapolis, MN, USA) were shown in **Table 1**. The total number of experiments was  $20 (2^k + 2k + 6)$ , where  $k$  is the number of independent variables. In order to determine the pure error, five replications were performed at the center point. The levels of the variables were determined based on the capabilities of the SC- $\text{CO}_2$  extraction system and the preliminary study.

### 2.3. Hexane (Soxhlet) extraction

Ground camelina seeds (12 g) were extracted with hexane (250 mL) in a Soxhlet apparatus for 6 h in dark to prevent photooxidation of astaxanthin. Hexane was separated from the oil using a rotary vacuum evaporator (Buchi Labortechnik AG, model B-490, Flawil, Switzerland) at  $22\text{ }^{\circ}\text{C}$  after each extraction. The resulting oil was weighed, and the total oil yield was reported as (weight of oil/weight of ground seed

**Table 1.** Independent variables and levels used for central composite rotatable design (CCRD).

Variable	Symbol coded	Levels				
		$-1.68$	$-1$	$0$	$+1$	$+1.68$
Pressure (MPa)	$X_1$	24.9	30.0	37.5	45.0	50.1
Temperature ( $^{\circ}\text{C}$ )	$X_2$	33.2	40.0	50.0	60.0	66.8
Ethanol concentration (%)	$X_3$	1.5	10.0	22.5	35.0	43.5

used for extraction)  $\times 100$ . The oil extracts were flushed with nitrogen and stored at  $-20\text{ }^{\circ}\text{C}$  until analyzed for astaxanthin concentration.

#### 2.4. Accelerated solvent extraction

Accelerated solvent extractions (ASE) using hexane and ethanol were performed using an accelerated solvent extractor (Dionex ASE 350, Sunnyvale, CA, USA) according to Zaghdoui et al. [17] with minor modifications. Ground camelina seed (3 g) was mixed with diatomaceous earth (1:3) to reduce the dead volume and then loaded into the 34mL-extraction cell. Extractions were performed at 1500 psi (10.3 MPa) and  $40\text{ }^{\circ}\text{C}$  for 5 static cycles of 5 min. The cell was rinsed with the extraction solvent and the solvent was purged from the cell with nitrogen for 60 s. The solvent in the extract was evaporated under the nitrogen, and the oils were stored at  $-20\text{ }^{\circ}\text{C}$  until analyzed for astaxanthin concentration.

#### 2.5. Analysis of astaxanthin

Astaxanthin concentration of the oil samples were determined by a reversed phase high performance liquid chromatography (RP-HPLC) according to Du et al. [18] with minor modifications. The samples were dissolved in acetone (70 mg/mL) and 15  $\mu\text{L}$  of trans- $\beta$ -Apo-8'-carotenal (internal standard) solution (1 mg/mL in acetone) was added to each sample. An aliquot (10  $\mu\text{L}$ ) was injected into an HPLC (Agilent 1100, Agilent Technologies, Waldbronn, Germany) equipped with a diode array detector (DAD). Samples were separated on a C18 column (150 $\times$ 4.6 mm; 5  $\mu\text{m}$  particle size; Phenomenex Inc., Torrance, CA, USA) using a mobile phase of methanol: acetonitrile (3:97, v/v) at a flow rate of 1 mL/min. The column temperature was set at  $30\text{ }^{\circ}\text{C}$  and the elution was detected at 474 nm.

#### 2.6. Tocopherol analysis

Tocopherols were analyzed according to the method of Belayneh et al. [19] with minor modification. Ten mg of each extract was dissolved in 1 mL methanol: dichloromethane (9:1, v/v) solvent mixture and 20  $\mu\text{L}$  of rac-5,7-Dimethyltocol (internal standard) solution (0.05 mg/mL) was added onto each sample. Then the samples were analyzed by a

high performance liquid chromatography (HPLC) (1200 Series, Agilent Technologies, Inc., Santa Clara, CA, USA) equipped with a fluorescence detector set at an excitation wavelength of 292 nm and an emission wavelength of 330 nm. An aliquot (70  $\mu$ L) of prepared solution was separated on a reversed-phase Eclipse XDB-C18 column (150 $\times$ 4.6 mm; 5  $\mu$ m particle size; Agilent Technologies, Inc., Santa Clara, CA, USA) using a mobile phase of methanol: water (95:5, v/v) at a flow rate of 1.5 mL/min. Retention time of tocopherol standards was used for identification. Tocopherol content was expressed as the sum of all tocopherols in mg tocopherol per kg oil.

### 2.7. Measurement of antioxidant activity


The antioxidant capacity of the oils obtained by different extraction methods was measured by ABTS<sup>•+</sup> radical cation. The ABTS<sup>•+</sup> radical cation was generated by reacting 7 mmol/L ABTS and 2.45 mmol/L potassium persulfate after incubation at room temperature for 16 h in the dark. The ABTS<sup>•+</sup> radical solution was diluted with ethanol to an absorbance of  $0.700 \pm 0.02$  at 734 nm. 6 mg of each extract was added to react with 2 mL of ABTS solution. The mixture was stored in the dark for 6 min and the absorbance at 734 nm was recorded using an Evolution 201 UV–vis Spectrophotometer (ThermoFisher, Waltham, MA). Ethanol was used as the control. Each measurement was conducted in duplicate. The scavenging of free radical was calculated according to the Eq. (1):

$$\text{ABTS Scavenging activity (\%)} = \frac{\text{Absorbance of control} - \text{Absorbance of sample}}{\text{Absorbance of control}} \quad (1)$$

### 2.8. Statistical analysis

Design Expert software 10.0.6 was used for regression and graphical analysis of the data. A quartic polynomial equation that correlates the response (astaxanthin concentration,  $\mu$ g/g oil) as a function of the independent variables and their interaction was developed. Analysis of variance (ANOVA) was used to determine the significance of the model through regression and mean square of residue error. The coefficient of determination ( $R_2$ ) was used to assess the quality of the






**Fig. 1.** Ethanol-modified SC-CO<sub>2</sub> extraction curves of camelina seed oil at 30 MPa and 50 °C at varying ethanol concentrations; n=2.

developed model. Analysis of the data to determine the statistical differences was performed by ANOVA and least-squares difference (LSD) using the SAS (version 9.4, SAS Institute Inc., Cary, NC, USA) at 95% confidence interval.

### 3. Results and discussion

Preliminary studies were performed to observe the effect of ethanol concentration on the astaxanthin extraction and to determine the range of ethanol concentrations to be used in the RSM design, because ethanol-modified SC-CO<sub>2</sub> extraction of astaxanthin was not performed on an oilseed before. Total oil yield increased with increasing ethanol content in the SC-CO<sub>2</sub> (**Fig. 1**). The highest oil yield of 25.6% was obtained at 15% ethanol concentration, whereas oil yield was only 18.1% with pure SC-CO<sub>2</sub>. Moreover, ethanol addition into SC-CO<sub>2</sub> increased the rate of the extraction, which is observed from the slope of the extraction lines in the first 120 min of the extraction (linear region) [20]. Astaxanthin content of the seed was 200 µg/g seed, including both free and esterified astaxanthin. Preliminary studies revealed that the astaxanthin content of the oils increased with increasing ethanol concentration (**Fig. 2**). Astaxanthin content of the oil extracted with pure SC-CO<sub>2</sub> was 190 µg/g oil, whereas it was 304 µg/g oil for the oil extracted with 15% ethanol in SC-CO<sub>2</sub>. In a study where astaxanthin was extracted from redspotted shrimp waste using SC-CO<sub>2</sub>, the highest yield (2.3% dry wt.) was obtained at 30 MPa



**Fig. 2.** Effect of the ethanol concentration in the SC-CO<sub>2</sub> on the astaxanthin concentration of the extracted oils obtained at 30 MPa and 50 °C. Different lowercase letters are significantly different for each extraction conditions ( $p < 0.05$ );  $n=2$ .

and 50 °C [14]. Camargo et al. [2] reported that the astaxanthin concentration increased from 26 to 35 µg/g dry residue when SC-CO<sub>2</sub> was modified with 15% ethanol at 30 MPa and 50 °C. SC-CO<sub>2</sub> is non-polar; therefore, it cannot extract polar compounds and has limited capacity to extract slightly polar compounds. Ethanol increases the polarity of SC-CO<sub>2</sub>; therefore, increases the solubility of the astaxanthin in the ethanol-modified SC-CO<sub>2</sub> due to the slightly polar structure of astaxanthin. In addition, the SC-CO<sub>2</sub>-expanded ethanol helps to swell the pores of the seeds, which in turn improves the interaction of the solvent with the inner parts of the seed matrix. In a study by Lopez et al. [21], maximum astaxanthin yield was obtained with 15% ethanol, whereas, any further increase in the ethanol amount in the SC-CO<sub>2</sub> caused a considerable decrease in the astaxanthin yield. Effect of ethanol content of the SC-CO<sub>2</sub> depends on the matrix, polarity of the target compound, extraction pressure and temperature; therefore, optimization of the extraction conditions is required for each case.

### 3.1. Model fitting

Multiple regression was used to determine the effect of pressure, temperature and ethanol concentration on the SC-CO<sub>2</sub> extraction of astaxanthin from camelina seed from the 20 runs that were generated by the RSM (**Table 2**). **Table 3** shows the statistical analysis of the quartic polynomial model based on ANOVA. After examining the lack of

**Table 2.** Experimental variables ( $X_1$ , pressure;  $X_2$ , temperature;  $X_3$ , ethanol concentration) and responses.

Run	$X_1$	$X_2$	$X_3$	Astaxanthin concentration ( $\mu\text{g/g}$ oil)	
				Actual	Predicted
1	-1	+1	-1	274	274
2	0	0	0	305	298
3	-1	-1	-1	261	262
4	-1	+1	+1	283	283
5	0	0	0	299	298
6	+1	-1	+1	364	364
7	-1.68	0	0	287	287
8	+1	+1	+1	276	276
9	0	-1.68	0	342	342
10	0	0	-1.68	208	215
11	0	0	0	298	298
12	+1	+1	-1	328	328
13	+1.68	0	0	396	396
14	0	0	0	310	298
15	-1	-1	+1	269	270
16	0	0	+1.68	375	381
17	0	+1.68	0	360	360
18	+1	-1	-1	270	270
19	0	0	0	287	298
20	0	0	0	300	298

fit, a quartic polynomial model was found to be adequate to explain the relationship between the astaxanthin yield and the extraction parameters. Backward-elimination was applied to refine the model by eliminating the insignificant terms. Model P-value < 0.0001, an insignificant lack of fit (P-value = 0.2351) and a higher coefficient of determination ( $R^2 = 0.989$ ) confirmed the suitability of the model to explain the relationship within the range of variables studied.

Regression coefficients were determined to predict the polynomial model for astaxanthin concentration and Eq. (2), expressed in coded variables, was obtained:

$$\begin{aligned}
 Y = & -20175.12 + 1161.47X_1 + 925.47X_2 - 111.28X_3 - 52.32X_1X_2 \\
 & + 5.61X_1X_3 + 0.60X_2X_3 - 16.28X_1^2 - 9.50X_2^2 - 0.02X_1X_2X_3 \\
 & + 0.73X_1^2X_2 - 0.06X_1^2X_3 + 0.53X_1X_2^2 - 0.007X_1^2X_2^2 \quad (2)
 \end{aligned}$$

where  $X_1$  is pressure,  $X_2$  is temperature,  $X_3$  is ethanol concentration. The coefficients in front of every term ( $X_1$ ,  $X_2$ , and  $X_3$ ) illustrate the effect of

**Table 3.** ANOVA for the fitted quartic polynomial model for optimization of extraction conditions.

Source	Sum of squares	Degrees of freedom	Mean of square	F-value	Prob > F	Significance
Model	38016.86	13	2924.37	42.99	<0.0001	***
X <sub>1</sub>	5955.77	1	5955.77	87.55	<0.0001	***
X <sub>2</sub>	177.47	1	177.47	2.61	0.1574	n.s.
X <sub>3</sub>	13804.57	1	13804.57	202.92	<0.0001	***
X <sub>1</sub> X <sub>2</sub>	388.37	1	388.37	5.71	0.0541	n.s.
X <sub>1</sub> X <sub>3</sub>	76.76	1	76.76	1.13	0.3290	n.s.
X <sub>2</sub> X <sub>3</sub>	2606.42	1	2606.42	38.31	0.0008	***
X <sub>1</sub> <sup>2</sup>	3067.75	1	3067.75	45.10	0.0005	***
X <sub>2</sub> <sup>2</sup>	4509.25	1	4509.25	66.29	0.0002	***
X <sub>1</sub> X <sub>2</sub> X <sub>3</sub>	2766.19	1	2766.19	40.66	0.0007	***
X <sub>1</sub> <sup>2</sup> X <sub>2</sub>	118.24	1	118.24	1.74	0.2355	n.s.
X <sub>1</sub> <sup>2</sup> X <sub>3</sub>	5848.17	1	5848.17	85.97	<0.0001	***
X <sub>1</sub> X <sub>2</sub> <sup>2</sup>	613.73	1	613.73	9.02	0.0239	n.s.
X <sub>1</sub> <sup>2</sup> X <sub>2</sub> <sup>2</sup>	6594.35	1	6594.35	96.94	<0.0001	***
Lack of Fit	108.97	1	108.97	1.82	0.2351	n.s.
Pure Error	299.20	5	59.84			
Cor Total	38425.03	19				


CV=2.71%, R<sup>2</sup>=0.9894

\*\*\* P &lt; 0.001; n.s., not significant.

a factor and the interaction among the factors, respectively. The positive sign in front of the terms indicates a synergistic effect, while the negative sign indicates an antagonistic effect. In the fitting model for this response variable, the pressure ( $p < 0.001$ ) and ethanol concentration ( $p < 0.001$ ) affected the astaxanthin concentration linearly. Besides, an interaction of temperature and co-solvent was found.

### 3.2. Effect of extraction parameters on the astaxanthin yield

**Fig. 3** presents the effects of extraction parameters on the astaxanthin concentration at -1, 0, and +1 levels of the three variables. Unlike quadratic or cubic polynomial model, the RSM plots of the quartic model were complex. At lower pressure (-1 level), ethanol concentration and temperature did not have a significant effect on the astaxanthin concentration (Fig. 3a). At moderate pressures (0 level), increasing ethanol concentration increased the astaxanthin yield at lower temperatures. At higher pressure (+1 level), increasing ethanol concentration at lower temperatures increased the astaxanthin yield; however,




**Fig. 3.** Response surface plots of astaxanthin concentration at -1, 0, and +1 levels of the independent variables.

increasing ethanol concentration resulted in lower astaxanthin yields at higher temperatures. Effect of ethanol concentration and pressure on the astaxanthin yield at all temperature levels were similar. Regardless of the temperature level, at high ethanol concentration level, moderate pressure resulted in the highest astaxanthin concentration, whereas moderate pressure led to the lowest astaxanthin concentration when ethanol concentration was low. At all ethanol levels, moderate pressures yielded higher astaxanthin at all temperatures. However, at lower and higher pressures, astaxanthin yield was lower at lower and higher temperatures at all ethanol levels. The temperature had two opposing effects; the density of CO<sub>2</sub> decreased with increasing temperature, which resulted in a reduced solvent power, on the other hand, increasing temperature led to an increase in the solute vapor pressure which affected the extraction positively. Furthermore, the temperature could affect the interaction between the solute and co-solvent molecules like hydrogen bond.

Extraction conditions were optimized to obtain the highest astaxanthin concentration using the RSM-developed model. The optimal extraction conditions were 41.6 MPa, 36.6 °C with 42.0% ethanol (w/w) at 1 L/min CO<sub>2</sub> flow rate (measured at ambient conditions). The predicted optimum conditions were verified by three additional independent extractions at those optimum conditions. Optimum predicted astaxanthin concentration was 437 µg/g oil, while the oil yield was 28.3%. The actual astaxanthin concentration was 421 ± 14 µg/g oil, while the oil yield was 29 ± 2%. These results indicated that the experimental values were in good agreement with the predicted one.


Once the optimum extraction conditions were obtained, an extraction curve of the camelina seed oil was generated (**Fig. 4**) to investigate the effect of extraction time. In the experimental design, extraction time was limited to 180 min to avoid long extraction times during optimization. However, it was found that an extraction time of 120 min at the optimized conditions is enough for the maximum recovery of the astaxanthin. It was found that the oil extracted in the first one hour of the extraction has the highest astaxanthin concentration, meaning astaxanthin was extracted at a higher rate compared to camelina seed oil. This information is useful to obtain high purity products and to lower the processing costs with shorter extraction times.



**Fig. 4.** Experimental extraction curves for oil yield and astaxanthin concentration at RSM-optimized conditions (41.6 MPa pressure, 36.6 °C temperature, and 42.0% ethanol concentration, w/w); n=2.

### 3.3. Comparison with other extraction methods

**Fig. 5** presents the comparison of the optimized ethanol-modified SC-CO<sub>2</sub> extraction (SC-CO<sub>2, opt</sub>) with conventional hexane extraction and accelerated solvent extraction using hexane and ethanol as solvents. There was no significant difference between the oil yield and



**Fig. 5.** Oil yield (black) and astaxanthin concentration (gray) obtained by SC-CO<sub>2, opt</sub>, ASE and Hexane (soxhlet) extraction. SC-CO<sub>2, opt</sub>: Optimized SC-CO<sub>2</sub> extraction; AEE: Accelerated ethanol extraction; AHE: Accelerated hexane extraction. Different capital letters mean significant differences ( $p < 0.05$ ) of the oil yield obtained from different extraction methods. Different lowercase letters mean significant differences ( $p < 0.05$ ) of the astaxanthin concentration obtained from different extraction methods; n=2.

the astaxanthin concentration in the oils extracted with SC-CO<sub>2, opt</sub> and hexane. Astaxanthin concentration of the oil extracted with SC-CO<sub>2, opt</sub> was 421 µg/g oil, whereas it was 418 µg/g oil in the hexane-extracted one. The lowest oil yield (23%) was obtained with accelerated hexane extraction (AHE), whereas the lowest astaxanthin concentration (75 µg/g oil) was obtained with accelerated ethanol extraction (AEE). For SCCO<sub>2, opt</sub> the ethanol consumption was 166 mL, whereas it was 250 mL for the hexane (Soxhlet) extraction. ASE required approximately 50 mL solvent (ethanol or hexane), which was much less than hexane and SCCO<sub>2, opt</sub> extractions. Lower solvent consumption is desired due to processing costs related to solvent storage, pumping, and separation. Although ASE required a shorter extraction time and consumed less solvent, the astaxanthin concentration was very low compared to SC-CO<sub>2, opt</sub> and hexane extraction methods. The lowest astaxanthin yield with AEE was due to high polarity of ethanol. Ethanol extracts polar compounds in the seed such as phospholipids and sugars but cannot extract slightly polar astaxanthin efficiently. A higher oil yield by AEE compared to AHE, but a lower β-carotene concentration in the oil was previously reported by Eller et al. [11]. Compared to the hexane extraction, a shorter extraction time of SC-CO<sub>2, opt</sub> allowed us to extract more astaxanthin, although two values were not significantly different. Reyes et al. [22] reported astaxanthin extraction recoveries from *Haematococcus pluvialis* up to 82.3% using ethanol-modified SC-CO<sub>2</sub>.

High ethanol concentrations in the optimized conditions suggested that better yields are obtained when a CO<sub>2</sub>-expanded ethanol was formed. Actually, the mechanism of extraction in a CO<sub>2</sub>-expanded ethanol may be different than ethanol-modified SC-CO<sub>2</sub> extraction where pressure and temperature play the critical role by changing the solubility of astaxanthin in the supercritical phase, whereas when the ethanol content in the SC-CO<sub>2</sub> approaches 50%, the mixture behaves like a pressurized liquid [23]. However, the low astaxanthin yields from AEE showed that the extraction mechanism of the AEE and the CO<sub>2</sub>-expanded ethanol extraction is different. Both higher pressures and high ethanol contents at the optimized ethanol-modified SC-CO<sub>2</sub> extraction played the role by improving the mass transfer by higher pressures and improving polarity by the presence of ethanol. It must be noted that, in this study, the matrix contained high amount of oil which acted



as co-solvent; therefore, the extraction mechanism was different than that of extracted from high water-content marine animals. Previously, Krichnavaruk et al. [24] used soybean oil, olive oil and ethanol as co-solvent to extract astaxanthin from microalgae *Haematococcus pluvisialis*. At the same extraction conditions, olive oil gave a comparable result to that with ethanol, however, soybean oil gave the lowest extraction efficiency. Those results suggested that the properties of the oil (e.g., fatty acid composition and viscosity) and the solubility of the oil in the SC-CO<sub>2</sub> could affect the astaxanthin extraction efficiency. In another study, transgenic maize was extracted with ethanol in combination with added commercial maize oil, and it was found that this combination was as effective as tetrahydrofuran and chloroform [5]. In the study of transgenic maize [5], the astaxanthin concentration of the final oil was 1 µg/g oil, which was considerably lower than the astaxanthin concentration obtained in this study.

In general, the polarity of the solvent used in the extraction should be the same as or similar with that of the target compounds, however, in some cases, using the mixture of polar and non-polar solvents may get higher recoveries [25]. For example, Yu et al. [26] used a mixture of hexane and methanol as solvents to extract amitraz and 2,4- dimethylaniline (2,4- DMA) from animal tissues. Because amitraz is less polar than 2,4- DMA, higher ratio of methanol in the mixture should result in a higher recovery for 2,4- DMA but not for amitraz. However, the highest recovery for both amitraz and 2,4- DMA was obtained when hexane/ methanol ratio was at a ratio of 1:9 (v/v) [26]. Solvents used in the extraction not only act as the carrier of target compound, but also have other functionalities. For example, the solvent can also swell the pores of the seeds or other materials. Thus, using the mixture of hexane and ethanol of ASE might result in higher recovery for oil, astaxanthin or both of them. However, mixing ethanol with hexane can be a concern for the manufacturers as the use of hexane cannot generate a product that complies with clean labeling.

#### *3.4. Tocopherol content of the extracts obtained from different extraction methods*

Tocopherols are important minor lipid components in camelina seed, and are best known for their antioxidant activities. **Table 4** presents

**Table 4.** Effect of extraction method on the tocopherol composition of the oils.

Extraction method	Tocopherol content (mg/kg oil)			
	$\delta$	$\gamma$ and $\beta$	$\alpha$	Total
AEE	20 ± 0 <sup>c</sup>	610 ± 8 <sup>a</sup>	5 ± 0 <sup>a</sup>	633 ± 8 <sup>a</sup>
AHE	17 ± 1 <sup>a</sup>	621 ± 5 <sup>a</sup>	12 ± 1 <sup>c</sup>	650 ± 6 <sup>a</sup>
SC-CO <sub>2, opt</sub>	17 ± 1 <sup>ab</sup>	613 ± 13 <sup>a</sup>	9 ± 0 <sup>b</sup>	640 ± 13 <sup>a</sup>
Hexane	18 ± 1 <sup>bc</sup>	732 ± 12 <sup>b</sup>	13 ± 0 <sup>c</sup>	763 ± 11 <sup>b</sup>

Values followed by the different letters indicate significant difference ( $p < 0.05$ );  $n=2$ .

AEE: Accelerated ethanol extraction; AHE: Accelerated hexane extraction; SC-CO<sub>2, opt</sub>: Optimized ethanol-modified SC-CO<sub>2</sub> extraction.

the tocopherol composition of the oils obtained by four different extraction methods. The oils extracted with hexane contained the highest amount of tocopherols (763 mg/kg oil). Tocopherol content of the oils extracted with AEE, AHE, and SC-CO<sub>2, opt</sub> ranged between 633 and 650 mg/kg oil, and there was no significant difference among them ( $p > 0.05$ ).  $\gamma$ - and  $\beta$ -Tocopherols were the dominant tocopherols in all oils and their level ranged from 610 mg/kg oil (AEE) to 732 mg/kg oil (hexane). Low concentration of  $\delta$ -tocopherol (17–20 mg/kg oil) followed by  $\alpha$ -tocopherol (5–13 mg/kg oil) was detected in all samples. No significant difference was observed between the tocopherol contents of the oils extracted by AEE, AHE, and SC-CO<sub>2, opt</sub>. Tocopherols are nonpolar compounds and they show good solubility both in hexane and pure SC-CO<sub>2</sub>; therefore, increasing the polarity of SC-CO<sub>2</sub> by ethanol decreased the extraction of tocopherols. Reports on the tocopherol content of camelina seed oil are scarce. Previously, tocopherol content of hexane- and SC-CO<sub>2</sub>-extracted non-engineered camelina seed oil was reported as 653 and 766 mg/kg, respectively [16]. It was shown that modification of SC-CO<sub>2</sub> with ethanol up to 10% ethanol at 35/45 MPa and 50/70 °C did not cause a significant difference in the extraction of tocopherols, but in this study it was shown that a drastic increase in the ethanol content of the SC-CO<sub>2</sub> decreases the extraction of tocopherols significantly compared to pure SC-CO<sub>2</sub>.

**Table 5.** Effect of extraction method on the ABTS radical scavenging activity of the oils.

Extraction method	ABTS scavenging activity (%)
AEE	32 ± 1 <sup>a</sup>
AHE	39 ± 1 <sup>b</sup>
SC-CO <sub>2, opt</sub>	65 ± 1 <sup>d</sup>
Hexane	48 ± 2 <sup>c</sup>

Values followed by the different letters indicate significant difference ( $p < 0.05$ );  $n=2$ .

AEE: Accelerated ethanol extraction; AHE: Accelerated hexane extraction; SC-CO<sub>2, opt</sub>: Optimized ethanol-modified SC-CO<sub>2</sub> extraction.

### 3.5. Antioxidant activity of the extracts obtained from different extraction methods

The ABTS radical scavenging activity of engineered camelina seed oils obtained from four extraction methods is shown in **Table 5**. The oil extracted from SC-CO<sub>2, opt</sub> showed the highest scavenging activity (65%), followed by hexane (48%), and AHE (39%). The lowest scavenging activity was obtained from AEE (32%). Results have shown that the ABTS scavenging activity increased with increasing astaxanthin content of the oils (Fig. 5). Tocopherols are natural antioxidants; however, it was found that astaxanthin content was the major factor determining the antioxidant activity of the oils. Even though the hexane-extracted oil had the highest tocopherol content, it did not have the highest antioxidant activity.

Previously, Reyes et al. [22] stated that the antioxidant activity increased as the astaxanthin concentration in the extract from *Haematococcus pluvialis* increased regardless of the extraction method. However, Jaime et al. [27] reported that the carotenoids in *Haematococcus pluvialis* extracted by AEE had higher antioxidant activity than those by AHE. The extraction condition and target compound could affect the selection of the appropriate solvent, resulting in different results. The oil extracted by SC-CO<sub>2, opt</sub> had higher antioxidant activity compared to that obtained by hexane extraction, even though their astaxanthin concentration in the oils were similar, showing the possible effect of other polar minor lipid compounds with antioxidant properties. Shao et al. [28] compared the antioxidant activity of essential oils extracted from *Anoectochilus roxburghii* by hexane extraction and

SC-CO<sub>2</sub> extraction and also reported that the oils extracted by SC-CO<sub>2</sub> showed higher antioxidant activity than those by hexane.

It should be noted that the material being investigated plays an important role in the antioxidant properties. The ABTS method measured the total antioxidant capacity of the extract, the specific chemical compositions of the extract could affect the antioxidant activity. Tocopherols are natural compounds that can have a synergistic effect on the antioxidant activity of the oils. Kang et al. [29] reported that the highest antioxidant activity of paprika leave was achieved by the highest amount of lutein and  $\gamma$ -tocopherol in the extract. Our results suggested that tocopherols did not have a significant contribution to the antioxidant activity of the samples because the antioxidant activity of astaxanthin is much higher than that of tocopherols.

#### 4. Conclusions

Engineered camelina seed was found to be a promising alternative source for astaxanthin. Ethanol-modified SC-CO<sub>2</sub> can be successfully used as a green extraction technique to extract astaxanthin. RSM was useful in optimizing the ethanol-modified SC-CO<sub>2</sub> extraction of astaxanthin from camelina oil seed. Ethanol-modified SC-CO<sub>2</sub> was more effective than AEE and AHE, and as effective as hexane to extract astaxanthin from camelina seed. Compared to hexane extraction, ethanol-modified SC-CO<sub>2</sub> extraction can provide protection to astaxanthin against oxidation and also eliminates the hazardous solvents from processing.

**Acknowledgments** — Authors thank Anji Reddy Konda for the assistance in tocopherol analysis, and Dr. Mark Wilkins from the Industrial Agricultural Products Center for giving us access to the accelerated solvent extraction system.

#### References

- [1] S. Dong, Y. Huang, R. Zhang, S. Wang, Y. Liu, Four different methods comparison for extraction of astaxanthin from green alga *Haematococcus pluvialis*, *Sci. World J.* 2014 (2014).

- [2] A.P. Sanchez-Camargo, M.A.A. Meireles, A.L.K. Ferreira, E. Saito, F.A. Cabral, Extraction of  $\omega$ -3 fatty acids and astaxanthin from Brazilian redspotted shrimp waste using supercritical CO<sub>2</sub> + ethanol mixtures, *J. Supercrit. Fluids* 61 (2012) 71–77.
- [3] J.P. Yuan, J. Peng, K. Yin, J.H. Wang, Potential health-promoting effects of astaxanthin: a high-value carotenoid mostly from microalgae, *Mol. Nutr. Food Res.* 55 (2011) 150–165.
- [4] P. Kittikaiwan, S. Powthongsook, P. Pavasant, A. Shotipruk, Encapsulation of *Haematococcus pluvialis* using chitosan for astaxanthin stability enhancement, *Carbohydr. Polym.* 70 (2007) 378–385.
- [5] J. Breitenbach, M. Nogueira, G. Farre, C. Zhu, T. Capell, P. Christou, G. Fleck, U. Focken, P.D. Fraser, G. Sandmann, Engineered maize as a source of astaxanthin: processing and application as fish feed, *Transgenic Res.* 25 (2016) 785–793.
- [6] J. Jayaraj, R. Devlin, Z. Punja, Metabolic engineering of novel ketocarotenoid production in carrot plants, *Transgenic Res.* 17 (2008) 489–501.
- [7] J.C. Huang, Y.J. Zhong, J. Liu, G. Sandmann, F. Chen, Metabolic engineering of tomato for high-yield production of astaxanthin, *Metab. Eng.* 17 (2013) 59–67.
- [8] R.Y. Khattab, M.A. Zeitoun, Quality evaluation of flaxseed oil obtained by different extraction techniques, *LWT – Food Sci. Technol.* 53 (2013) 338–345.
- [9] A. Rai, B. Mohanty, R. Bhargava, Supercritical extraction of sunflower oil: a central composite design for extraction variables, *Food Chem.* 192 (2016) 647–659.
- [10] A. Rai, B. Mohanty, R. Bhargava, Modeling and response surface analysis of supercritical extraction of watermelon seed oil using carbon dioxide, *Sep. Purif. Technol.* 141 (2015) 354–365.
- [11] F.J. Eller, J.K. Moser, J.A. Kenar, S.L. Taylor, Extraction and analysis of tomato seed oil, *J. Am. Oil Chem. Soc.* 87 (2010) 755–762.
- [12] L. Jaime, E. Vazquez, T. Fornari, M.C. Lopez-Hazas, M.R. Garcia-Risco, S. Santoyo, G. Reglero, Extraction of functional ingredients from spinach (*Spinacia oleracea* L.) using liquid solvent and supercritical CO<sub>2</sub> extraction, *J. Sci. Food Agric.* 95 (2015) 722–729.
- [13] S. Machmudah, A. Shotipruk, M. Goto, M. Sasaki, T. Hirose, Extraction of astaxanthin from *Haematococcus pluvialis* using supercritical CO<sub>2</sub> and ethanol as entrainer, *Ind. Eng. Chem. Res.* 45 (2006) 3652–3657.
- [14] A.P. Sanchez-Camargo, H.A. Martinez-Correa, L.C. Paviani, F.A. Cabral, Supercritical CO<sub>2</sub> extraction of lipids and astaxanthin from Brazilian redspotted shrimp waste (*Farfantepenaeus paulensis*), *J. Supercrit. Fluids* 56 (2011) 164–173.
- [15] S.A. Radzali, B.S. Baharin, R. Othman, M. Markom, R.A. Rahman, Co-solvent selection for supercritical fluid extraction of astaxanthin and other carotenoids from *Penaeus monodon* waste, *J. Oleo Sci.* 63 (2014) 769–777.
- [16] H.D. Belayneh, R.L. Wehling, A.K. Reddy, E.B. Cahoon, O.N. Ciftci, Ethanol-modified supercritical carbon dioxide extraction of the bioactive lipid components of *Camelina sativa* seed, *J. Am. Oil Chem. Soc.* (2017) 1–11.

- [17] K. Zaghdoudi, S. Pontvianne, X. Framboisier, M. Achard, R. Kudaibergenova, M. Ayadi-Trabelsi, J. Kalthoum-cherif, R. Vanderesse, C. Frochot, Y. Guiavarc'h, Accelerated solvent extraction of carotenoids from: Tunisian kaki (*Diospyros kaki* L.), peach (*Prunus persica* L.) and apricot (*Prunus armeniaca* L.), *Food Chem.* 184 (2015) 131–139.
- [18] P. Du, M. Jin, L. Yang, G. Chen, C. Zhang, F. Jin, H. Shao, M. Yang, X. Yang, Y. She, S. Wang, L. Zheng, J. Wang, Determination of astaxanthin in feeds using high performance liquid chromatography and an efficient extraction method, *J. Liq. Chromatogr. Relat. Technol.* 39 (2016) 35–43.
- [19] H.D. Belayneh, R.L. Wehling, E. Cahoon, O.N. Ciftci, Extraction of omega-3-rich oil from *Camelina sativa* seed using supercritical carbon dioxide, *J. Supercrit. Fluids* 104 (2015) 153–159.
- [20] O.N. Ciftci, J. Calderon, F. Temelli, Supercritical carbon dioxide extraction of corn distiller's dried grains with solubles: experiments and mathematical modeling, *J. Agric. Food Chem.* 60 (2012) 12482–12490.
- [21] M. Lopez, L. Arce, J. Garrido, A. Ríos, M. Valcarcel, Selective extraction of astaxanthin from crustaceans by use of supercritical carbon dioxide, *Talanta* 64 (2004) 726–731.
- [22] F.A. Reyes, J.A. Mendiola, E. Ibanez, J.M. del Valle, Astaxanthin extraction from *Haematococcus pluvialis* using CO<sub>2</sub>-expanded ethanol, *J. Supercrit. Fluids* 92 (2014) 75–83.
- [23] M.T. Golmakani, J.A. Mendiola, K. Rezaei, E. Ibanez, Expanded ethanol with CO<sub>2</sub> and pressurized ethyl lactate to obtain fractions enriched in  $\gamma$ -linolenic acid from *Arthrospira platensis* (Spirulina), *J. Supercrit. Fluids* 62 (2012) 109–115.
- [24] S. Krichnavaruk, A. Shotipruk, M. Goto, P. Pavasant, Supercritical carbon dioxide extraction of astaxanthin from *Haematococcus pluvialis* with vegetable oils as cosolvent, *Bioresour. Technol.* 99 (2008) 5556–5560.
- [25] H. Sun, X. Ge, Y. Lv, A. Wang, Application of accelerated solvent extraction in the analysis of organic contaminants, bioactive and nutritional compounds in food and feed, *J. Chromatogr. A* 1237 (2012) 1–23.
- [26] H. Yu, Y. Tao, T. Le, D. Chen, A. Ishsan, Y. Liu, Y. Wang, Z. Yuan, Simultaneous determination of amitraz and its metabolite residue in food animal tissues by gas chromatography-electron capture detector and gas chromatography–mass spectrometry with accelerated solvent extraction, *J. Chromatogr. B* 878 (2010) 1746–1752.
- [27] L. Jaime, I. Rodriguez-Meizoso, A. Cifuentes, S. Santoyo, S. Suarez, E. Ibanez, F.J. Senorans, Pressurized liquids as an alternative process to antioxidant carotenoids' extraction from *Haematococcus pluvialis* microalgae, *LWT – Food Sci. Technol.* 43 (2010) 105–112.
- [28] Q. Shao, Y. Deng, H. Liu, A. Zhang, Y. Huang, G. Xu, M. Li, Essential oils extraction from *Anoectochilus roxburghii* using supercritical carbon dioxide and their antioxidant activity, *Ind. Crops Prod.* 60 (2014) 104–112.
- [29] J.H. Kang, S. Kim, B. Moon, Optimization by response surface methodology of lutein recovery from paprika leaves using accelerated solvent extraction, *Food Chem.* 205 (2016) 140–145.