

Extremal Behaviour in Multiagent Contract Negotiation

Paul E. Dunne

PED@CSC.LIV.AC.UK

Dept. of Computer Science, University of Liverpool, Liverpool L69 7ZF, United Kingdom

Abstract

¹ We examine properties of a model of resource allocation in which several agents exchange resources in order to optimise their individual holdings. The schemes discussed relate to well-known negotiation protocols proposed in earlier work and we consider a number of alternative notions of “rationality” covering both quantitative measures, e.g. cooperative and individual rationality and more qualitative forms, e.g. Pigou-Dalton transfers. While it is known that imposing particular rationality and structural restrictions on the form of exchanges may render these unable to realise every reallocation of the resource set, in this paper we address the issue of the number of restricted rational exchanges that may be required to implement a particular reallocation when it *is* possible to do so. We construct examples showing that this number may be exponential (in the number of resources m), even when all of the agent utility functions are monotonic. We further show that k agents may achieve in a single exchange a reallocation requiring exponentially many rational exchanges if at most $k - 1$ agents can participate, this same reallocation being unrealisable by any sequences of rational exchanges in which at most $k - 2$ agents are involved.

1. Introduction

Mechanisms for negotiating allocation of resources within a group of agents form an important body of work within the study of multiagent systems. Typical abstract models derive from game-theoretic perspectives in economics and among the issues that have been addressed are strategies that agents use to obtain a particular subset of the resources available, e.g. [14, 18, 21], and protocols by which the process of settling upon some allocation of resources among the agents involved is agreed, e.g. [5, 6, 7, 15].

The setting we are concerned with is encapsulated in the following definition.

Definition 1 *A resource allocation setting is defined by a triple $\langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle$ where*

$$\mathcal{A} = \{A_1, A_2, \dots, A_n\} \quad ; \quad \mathcal{R} = \{r_1, r_2, \dots, r_m\}$$

are, respectively, a set of (at least two) agents and a collection of (non-shareable) resources. A utility function, u , is a mapping from subsets of \mathcal{R} to rational values. Each agent $A_i \in \mathcal{A}$ has associated with it a particular utility function u_i , so that \mathcal{U} is $\langle u_1, u_2, \dots, u_n \rangle$. An allocation P of \mathcal{R} to \mathcal{A} is a partition $\langle P_1, P_2, \dots, P_n \rangle$ of \mathcal{R} . The value $u_i(P_i)$ is called the utility of the resources assigned to A_i .

Two major applications in which the abstract view of Definition 1 has been exploited are e-commerce and distributed task realisation. In the first \mathcal{R} represents some collection of commodities offered for sale and individual agents seek to acquire a subset of these, the

1. This report is a revised and extended version of ULCS-04-001 which it subsumes.

“value” an agent attaches to a specific set being described by that agent’s utility function. In task planning, the “resource” set describes a collection of sub-tasks to be performed in order to realise some complex task, e.g. the “complex task” may be to transport goods from a central warehouse to some set of cities. In this example \mathcal{R} describes the locations to which goods must be dispatched and a given allocation defines those places to which an agent must arrange deliveries. The utility functions in such cases model the cost an agent associates with carrying out its allotted sub-tasks.

Within the very general context of Definition 1, a number of issues arise stemming from the observation that it is unlikely that some *initial allocation* will be seen as satisfactory either with respect to the views of all agents in the system or with respect to diverse global considerations. Thus, by proposing changes to the initial assignment individual agents seek to obtain a “better” allocation. This scenario raises two immediate questions: how to evaluate a given partition and thus have a basis for forming improved or optimal allocations; and, the issue underlying the main results of this paper, what restrictions should be imposed on the form that proposed exchanges may take.

We shall subsequently review some of the more widely studied approaches to defining conditions under which some allocations are seen as “better” than others. For the purposes of this introduction we simply observe that such criteria may be either *quantitative* or *qualitative* in nature. As an example of the former we have the approach wherein the “value” of an allocation P is simply the sum of the values given by the agents’ utility functions to the subsets of \mathcal{R} they have been apportioned within P , i.e. $\sum_{i=1}^n u_i(P_i)$: this is the so-called *utilitarian social welfare*, which to avoid repetition we will denote by $\sigma_u(P)$. A natural aim for agents within a commodity trading context is to seek an allocation under which σ_u is *maximised*. One example of a *qualitative* criterion is “*envy freeness*”: informally, an allocation, P , is envy-free if no agent assigns greater utility to the resource set (P_j) held by another agent than it does with respect to the resource set (P_i) it has actually been allocated, i.e. for each distinct pair $\langle i, j \rangle$, $u_i(P_i) \geq u_i(P_j)$.

In very general terms there are two approaches that have been considered in treating the question of how a finite collection of resources might be distributed among a set of agents in order to optimise some criterion of interest: “contract-net” based methods, e.g. [8, 10, 11, 20, 21] deriving from the work of [25]; and “combinatorial auctions”, e.g. [16, 17, 22, 24, 26] amongst others. In this article our concerns focus on the first of these, however, before discussing it in greater depth we compare the two approaches.

One may view the strategy underlying combinatorial auctions as investing the computational effort into a “pre-processing” stage following which a given allocation is determined. Thus a controlling agent (the “auctioneer”) is supplied with a set of *bids* – pairs $\langle S_j, p_j \rangle$ wherein S_j is some subset of the available resources and p_j the price agent A_j is prepared to pay in order to acquire S_j . The problem faced by the auctioneer is to decide which bids to accept in order to maximise the overall profit subject to the constraint that each item can be obtained by at most one agent. By imposing restrictions on the structure of bids some efficient algorithmic methods that can identify optimal or near-optimal² allocations have been discovered, e.g. [26]. In general, however, this optimisation problem is NP-hard, even if only a good approximate solution is sought, cf. [23]. In addition to these compu-

2. By “near-optimal” we mean in the sense of being “good” approximations.

tational problems, one further difficulty that can arise in implementations of combinatorial auction protocols is their potential vulnerability to manipulation by agents. For example, agents may assume multiple identities in order to submit several bids: so-called “false-name” bids concerning which the susceptibility of various protocols together with possible defence mechanisms have been studied in depth, e.g. [19, 28, 29].

What we shall refer to as “contract-net schemes” typically eschew the precomputation stage and subordination to a controlling arbiter employed in auction mechanisms, seeking instead to realise a suitable allocation by an agreed sequence of exchanges. The *contract-net* (in its most general instantiation) for scenarios of m resources distributed among n agents, is the complete directed graph with n^m vertices (each of which is associated with a distinct allocation). In this way a possible exchange $\langle P, Q \rangle$ is represented as an edge directed from the vertex labelled with P to that labelled Q . Viewed thus, identifying a sequence of exchanges can be interpreted as a search process which, in principle, individual agents may conduct in an autonomous fashion. We defer a discussion of why a *sequence* of exchanges may be needed, noting only that some mechanism for dealing with combinatorial problems may be used so that the number of “legal” exchanges described in the contract-net graph is reduced: for any given agent and allocation, P , the number of new allocations “reachable” in a single exchange from P becomes bounded by some polynomial function of the number of resources. As a result at any stage each agent has only a relatively small number of possibilities to consider, so in principle allowing better allocations to be identified by a sequence of *local* improvements.

While it is known that contract-net schemes have a number of computationally undesirable aspects some of which we review subsequently, we argue that the capability to move from an initial allocation to some notionally better one via a sequence of local adjustments offers significant benefits. In discussing these we stress it is not our intention to promote the superiority (or otherwise) of contract-net approaches over auction mechanisms, but rather to indicate that such schemes may provide a more suitable model in *some* arenas. It is, of course, equally true that in other environments the algorithmic advantages resulting from suitable auction mechanisms render these a more appropriate optimisation technique.

We have suggested one potential benefit in our earlier introductory description, namely, the capability for agents to negotiate independently of a central arbiter. Thus, whether some allocation of the resources is actually effected depends not on the bid ranking algorithm employed by an auctioneer but solely on the desires and intentions of the participating agents. This capability for independent action raises a number of interesting issues. Although we have outlined auction mechanisms as shifting the computational burden in choosing an allocation to a pre-processing stage handled by the controlling agent charged with resolving competing bids, it is, of course, the case that this does not obviate the requirement for (potentially significant) computational effort on the part of the bidding agents. It is well-known assuming a “pure” scenario in which each agent can only submit a limited number of bids – an environment which methods aiming to detect false-name bidding seek to ensure – that an agent is faced with a highly non-trivial strategic issue: deciding which set(s) of resources and prices are “most likely” to bring about a desired outcome. Notice that we are faced with two extremes in such cases: a fully competitive situation where bids are formulated independently of consultation with other agents; and a fully cooperative one whereby the agents *locally* agree a partition of the resources and bid accordingly. In this

latter case, assuming agents behave honestly and enter bids only for the resource subsets they have agreed, the auctioneer is presented with a *fait accompli*: accept every bid (or be prepared to make a sub-optimal profit). In total, by allowing collusion between agents prior to submitting bids, appropriate contract-net schemes provide a means of undermining the role of the auctioneer. One can interpret such collusion as a rather extreme form of “strategic manipulation” in which a coalition of all agents forms to bring about a partition of resources with which each is satisfied.

This example of how carrying out a sequence of “local negotiations” may be used externally to manipulate the outcome of an auction process raises the question of the extent to which regimes without a controlling auctioneer could be *internally* manipulated to the advantage of particular agents. While a detailed analysis of such possibilities is outside the scope of this article, we mention one very basic way in which a coalition of agents, \mathcal{C} could act collectively in their own interests against the interests of other agents. Thus it may be that for some subset, S say, of the available resources, were \mathcal{C} collectively to acquire S , two outcomes would result: every agent in \mathcal{C} could achieve an allocation that it viewed as optimal; and no agent outside \mathcal{C} would be able to realise a similarly regarded allocation from those resources remaining, i.e. $\mathcal{R} \setminus S$. A very preliminary study of such coalitional issues is presented in [9], where such S are termed *critical sets* for a coalition. One result of interest as regards such manipulative processes, is that without even considering the strategic complexities that arise for \mathcal{C} in *obtaining* a critical set the problem of *recognising* if such a resource set exists is already of significant complexity: [9] showing it to be Σ_2^P -complete even if \mathcal{C} comprises only two members out of a society of four agents. While this is only a very basic class of manipulative behaviour, the level of complexity involved suggests that identifying critical sets is unlikely to be a productive investment of computational effort.

The preceding discussion has identified general criteria that may apply in assessing the worth of a specific allocation and presented an overview of two approaches that have been studied in terms of optimising quantitative measures. Henceforward we consider contract-net schemes with particular emphasis on their properties when limits are imposed on the form that sequences of exchanges can assume in order to bring about better allocations.

Before proceeding with the precise matter of our results, however, we deal with a question raised earlier: why a *sequence* of exchanges should be required. Suppose, for example, the aims are to bring about an allocation, P_{opt} , that maximises utilitarian social welfare (σ_u) or to improve that of an initial allocation P_{init} . It is certainly the case that the exchange $\langle P_{init}, P_{opt} \rangle$ is a *single* exchange, i.e. described by a single directed edge in the contract-net graph, and thus, all that any agent has to do is to identify those items in its possession that are expendable and which agents should be given these, as well as those items it lacks and the agents from whom these must be acquired. Now, even if we make the assumption that every agent is aware of the allocation held by any other agent, such “single-shot” solutions are not feasible approaches.

One difficulty is that just as determining optimal outcomes for combinatorial auctions faces issues of computational intractability, so too, these arise in the context of identifying allocations with various properties. Thus deciding if a given allocation *could* be improved with respect to σ_u is NP-complete, as is the problem of deciding if a given lower bound on σ_u can be achieved, cf. [8]. We note that these results apply even in the case of scenarios involving only two agents both of whom employ *monotone* utility functions, i.e. with the

property that for any two sets, S and T , $S \subseteq T$ implies $u(S) \leq u(T)$. This situation is unchanged if one moves from quantitative measures such as σ_u to *qualitative* indicators. So, again with two agents and monotone utility functions, [8] establish that it is co-NP-complete to decide whether an allocation is “*Pareto optimal*”, i.e. one such that any allocation under which some agent’s utility improves does so at the expense of another agent’s decreasing. Similarly, [9] has recently shown that deciding if a resource allocation setting admits an *envy-free* distribution is NP-complete in a two agent environment³. In summary, one problem with single exchange solutions to obtain improved or optimal allocations is that from a given initial point in the contract-net graph it is computationally hard for the agents to agree if there are better allocations available at all. There are, however, other reasons why one might regard single exchange solutions as infeasible. For example, even if a new allocation P_{final} is agreed, this might differ quite radically from the initial state represented by P_{init} : it may not be possible to implement $\langle P_{init}, P_{final} \rangle$ in a *single* exchange even if only two agents are involved since the environment in which the trading process is implemented may not be suited to handling exchanges with large numbers of resources; similarly the protocol used for describing contracts may not allow arbitrarily large numbers of resources to be dealt with.

In total, in order to proceed by way of local improvements some means of managing the exponentially large search space is needed. To begin, we first formalise the concepts of *exchange* and *contract path*.

Definition 2 Let $\langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle$ be a resource allocation setting. An exchange is a pair $\langle P, Q \rangle$ where $P = \langle P_1, \dots, P_n \rangle$ and $Q = \langle Q_1, \dots, Q_n \rangle$ are distinct partitions of \mathcal{R} . The effect of implementing the exchange $\langle P, Q \rangle$ is that the allocation of resources specified by P is replaced with that specified by Q . Following the notation of [11] for an exchange $\delta = \langle P, Q \rangle$, we use \mathcal{A}^δ to indicate the subset of \mathcal{A} involved, i.e. $A_k \in \mathcal{A}^\delta$ if and only if $P_k \neq Q_k$.

Let $\delta = \langle P, Q \rangle$ be an exchange. A contract path realising δ is a sequence of allocations

$$\Delta = \langle P^{(1)}, P^{(2)}, \dots, P^{(t-1)}, P^{(t)} \rangle$$

in which $P = P^{(1)}$ and $P^{(t)} = Q$. The length of Δ , denoted $|\Delta|$ is $t - 1$, i.e. the number of exchanges in Δ .

There are two methods which we can use to reduce the number of exchanges that a single agent may have to consider in seeking to move from some allocation to another, thereby avoiding the need to choose from exponentially many alternatives: *structural* and *rationality* constraints. Structural constraints limit the permitted exchanges to those which bound the number of resources and/or the number of agents involved, but take no consideration of the view any agent may have as to whether its allocation has improved. In contrast, rationality constraints restrict exchanges, $\langle P, Q \rangle$ to those in which Q “improves” upon P according to particular criteria. In this article we consider two classes of structural constraint: *O*-contracts, defined and considered in [20], and what we shall refer to as *M(k)*-contracts.

Definition 3 Let $\delta = \langle P, Q \rangle$ be an exchange involving a reallocation of \mathcal{R} among \mathcal{A} .

3. The utility functions in this proof are not monotone, however, we conjecture that imposing such a restriction will not result in any reduction in complexity.

a. δ is a one contract (*O-contract*) if

O1. $\mathcal{A}^\delta = \{i, j\}$.

O2. There is a unique resource $r \in P_i \cup P_j$ for which $Q_i = P_i \cup \{r\}$ and $Q_j = P_j \setminus \{r\}$ (with $r \in P_j$) or $Q_j = P_j \cup \{r\}$ and $Q_i = P_i \setminus \{r\}$ (with $r \in P_i$)

b. For a value $k \geq 2$, the exchange $\delta = \langle P, Q \rangle$ is an $M(k)$ -contract if $2 \leq |\mathcal{A}^\delta| \leq k$ and $\cup_{i \in \mathcal{A}^\delta} Q_i = \cup_{i \in \mathcal{A}^\delta} P_i$.

Thus, *O*-contracts involve the transfer of *exactly one* resource from a particular agent to another, resulting in the number of exchanges compatible with any given allocation being exactly $(n - 1)m$: each of the m resources can be reassigned from its current owner to any of the other $n - 1$ agents.

Rationality constraints arise in a number of different ways. For example, from the standpoint of an individual agent A_i a given exchange $\langle P, Q \rangle$ may have three different outcomes: $u_i(P_i) < u_i(Q_i)$, i.e. A_i values the allocation Q_i as superior to P_i ; $u_i(P_i) = u_i(Q_i)$, i.e. A_i is indifferent between P_i and Q_i ; and $u_i(P_i) > u_i(Q_i)$, i.e. A_i is worse off after the exchange. When global optima such as utilitarian social welfare are to be maximised, there is the question of what incentive there is for any agent to accept an exchange $\langle P, Q \rangle$ under which they are left with a less valuable resource holding. The standard approach to this latter question is to introduce the notion of a *pay-off* function, i.e. in order for A_i to accept an exchange under which it suffers a reduction in utility, A_i receives some payment sufficient to compensate for its loss. Of course such compensation must be made by other agents in the system who in providing it do not wish to pay in excess of any gain. In defining notions of pay-off the interpretation is that in any transaction each agent A_i makes a payment, π_i : if $\pi_i < 0$ then A_i is given $-\pi_i$ in return for accepting an exchange; if $\pi_i > 0$ then A_i contributes π_i to the amount to be distributed among those agents whose pay-off is negative.

This notion of “sensible transfer” is captured by the concept of *individual rationality*, and is often defined in terms of an appropriate pay-off vector existing. It is not difficult, however, to show that such definitions are equivalent to the following.

Definition 4 An exchange $\langle P, Q \rangle$ is individually rational (*IR*) if and only if $\sigma_u(Q) > \sigma_u(P)$.

We shall consider alternative bases for rationality constraints later: these are primarily of interest within so-called *money free* settings (so that compensatory payment for a loss in utility is not an option).

The central issue of interest in this paper concerns the properties of the contract-net graph when the allowed exchanges must satisfy both a structural *and* a rationality constraint. Thus, if we consider arbitrary predicates Φ on exchanges $\langle P, Q \rangle$ – where the cases of interest are Φ combining a structural and rationality condition – we have,

Definition 5 For Φ a predicate over distinct pairs of allocations, a contract path

$$\langle P^{(1)}, P^{(2)}, \dots, P^{(t-1)}, P^{(t)} \rangle$$

realising $\langle P, Q \rangle$ is a Φ -path if for each $1 \leq i < t$, $\langle P^{(i)}, P^{(i+1)} \rangle$ is a Φ -exchange, i.e. $\Phi(P^{(i)}, P^{(i+1)})$ holds. We say that Φ is complete if any exchange δ may be realised by a Φ -path. We, further, say that Φ is complete with respect to Ψ -exchanges (where Ψ is a predicate over distinct pairs of allocations) if any exchange δ for which $\Psi(\delta)$ holds may be realised by a Φ -path.

The main interest in earlier studies of these ideas has been in areas such as identifying necessary and/or sufficient conditions on exchanges to be complete with respect to particular criteria, e.g. [20]; and in establishing “convergence” and termination properties, e.g. [10, 11] consider exchange types, Φ , such that every maximal⁴ Φ -path ends in a Pareto optimal allocation. In [20], Sandholm examines how restrictions e.g. with $\Phi(P, Q) = \top$ if and only if $\langle P, Q \rangle$ is an O -contract, may affect the existence of contract paths to realise exchanges. Of particular interest, from the viewpoint of heuristics for exploring the contract-net graph, are cases where $\Phi(P, Q) = \top$ if and only if the exchange $\langle P, Q \rangle$ is individually rational. For the case of O -contracts the following are known:

Theorem 1

- a. O -contracts are complete.
- b. IR O -contracts are not complete with respect to IR exchanges.

Of course one might question why paths combining both structural and rationality constraints should be of interest, particularly in view of Theorem 1: since it is possible to move between any pair of allocations via O -contracts alone why impose further limits? To illustrate the point that rationality constraints may also be needed, consider the example now outlined.

Suppose the agents within a particular setting are observing the following protocol:

A reallocation of resources is agreed over a sequence of stages, each of which involves communication between two agents, A_i and A_j . This communication consists of A_i issuing a proposal to A_j of the form (buy, r, p) , offering to purchase r from A_j for a payment of p ; or $(sell, r, p)$, offering to transfer r to A_j in return for a payment p . The response from A_j is simply *accept* (following which the exchange is implemented) or *reject*.

This, of course, is a very simple negotiation structure, however consider its operation within a two agent setting in which one agent, A_1 say, wishes to bring about an allocation P_{fin} (and thus can devise a plan – sequence of exchanges – to realise this from an initial allocation P_{init}) while the other agent, A_2 , does *not* know P_{fin} . In addition, assume that A_1 is the only agent that makes proposals and that a final allocation is fixed either when A_1 is “satisfied” or as soon as A_2 *rejects* any offer.

While A_2 *could* be better off if P_{fin} is realised, it may be the case that the only proposals A_2 will accept are those under which it does not lose, i.e. A_2 is not prepared to suffer a short-term loss even if it is suggested that a long-term gain will result. Thus if some

4. “Maximal” in the sense that if $\langle P^{(1)}, \dots, P^{(t)} \rangle$ is such a path, then for every allocation, Q , $\Phi(P^{(t)}, Q)$ does not hold.

agents are sceptical about the *bona fides* of others then they will be inclined to accept *only* exchanges from which they can perceive an *immediate* benefit, i.e. those which are individually rational. There are several reasons why an agent may embrace such attitudes within the schema outlined: once an exchange has been implemented A_2 may lose utility but no further proposals are made by A_1 so that the loss is “permanent”. We note that even if we enrich the basic protocol so that A_1 can describe P_{fin} to A_2 *before* any formal exchange of resources takes place, if the exchange is implemented by an O -contract path (via the sequence of stages outlined), A_2 may still reject offers under which it suffers a loss, since it is unwilling to rely on the subsequent exchanges that would ameliorate its loss actually being proposed⁵. Although the position taken by A_2 in the setting just described may appear unduly cautious, we would claim that it does reflect “real” behaviour in certain contexts. Outside the automated allocation and negotiation models in multiagent systems that we have been reviewing, there are many examples of actions by individuals where promised long-term gains are insufficient to engender the acceptance of short term loss. Consider “chain letter” schemes (or their more subtle manifestation as “pyramid selling” enterprises): such have a natural lifetime bounded by the size of the population in which they circulate, but may breakdown before this is reached. Faced with a request to “send \$10 to the five names at the head of the list and forward the letter to ten others after adding your name” despite the possibility of significant gain after a temporary loss of \$50, to ignore such blandishments is not seen as overly sceptical and cautious: there may be reluctance to accept that one will eventually receive sufficient recompense in return and suspicion that the name order has been manipulated.

In total for the scenario we have described, if A_1 wishes to bring about an allocation P_{fin} then faced with the view adopted by A_2 and the limitations imposed by the exchange protocol, the only “effective plan” that A_1 could adopt is to find a sequence of *rational*, i.e. IR, exchanges to propose to A_2 . In the consideration of algorithmic and complexity issues presented in [8] one difficulty with such plan formulation is already apparent, that is:

Theorem 2 *Even in the case $n = 2$ and with monotone utility functions the problem of deciding if an IR O -contract path exists to realise the IR exchange $\langle P, Q \rangle$ is NP-hard.*

Thus deciding if any rational plan is possible is already computationally hard. In this article we demonstrate that, even if an appropriate rational plan exists, in extreme cases, there may be significant problems: the number of exchanges required could be exponential in the number of resources, so affecting both the time it will take for the schema outlined to conclude and the space that an agent will have to dedicate to storing it. Thus in his proof of Theorem 1 (b), Sandholm observes that when an IR O -contract path exists for a given IR exchange, it may be the case that its length exceeds m , i.e. some agent passes a resource to another and then accepts the same resource at a later stage. The typical form of the results that we derive can be summarised as:

For Φ a structural constraint (O -contract or $M(k)$ -contract) and Ψ a rationality constraint, e.g. $\Psi(P, Q)$ holds if $\langle P, Q \rangle$ is individually rational, there are

5. We note that even if A_1 attempts to construct an ordering of exchanges under which any “irrational” exchange reduces the value of its own holding, there is one problem: A_2 may reject subsequent offers after the “irrational” exchanges so that A_1 is worse off.

resource allocation settings $\langle \mathcal{A}_n, \mathcal{R}_m, \mathcal{U} \rangle$ in which there is an exchange $\langle P, Q \rangle$ satisfying all of the following.

- a. $\langle P, Q \rangle$ is a Ψ -exchange.
- b. $\langle P, Q \rangle$ can be realised by a contract path on which every exchange satisfies the structural constraint Φ and the rationality constraint Ψ .
- c. Every such contract path has length at least $g(m)$.

For example, we show that there are instances for which the shortest IR O -contract path has length exponential in m .⁶

In the next section we will be interested in lower bounds on the values of the following functions: we introduce these in general terms to avoid unnecessary subsequent repetition.

Definition 6 *Let $\langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle$ be a resource allocation setting. Additionally let Φ and Ψ be two predicates on exchanges. For an exchange $\delta = \langle P, Q \rangle$ the partial function $L^{\text{opt}}(\delta, \langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle, \Phi)$ is the length of the shortest Φ -contract path realising $\langle P, Q \rangle$ if such a path exists (and is undefined if no such path is possible). The partial function $L^{\text{max}}(\langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle, \Phi, \Psi)$ is*

$$L^{\text{max}}(\langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle, \Phi, \Psi) = \max_{\Psi\text{-exchanges } \delta} L^{\text{opt}}(\delta, \langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle, \Phi)$$

Finally, the partial function $\rho^{\text{max}}(n, m, \Phi, \Psi)$ is

$$\rho^{\text{max}}(n, m, \Phi, \Psi) = \max_{\mathcal{U}=\langle u_1, u_2, \dots, u_n \rangle} L^{\text{max}}(\langle \mathcal{A}_n, \mathcal{R}_m, \mathcal{U} \rangle, \Phi, \Psi)$$

where consideration is restricted to those Ψ -exchanges $\delta = \langle P, Q \rangle$ for which a realising Φ -path exists.

The three measures, L^{opt} , L^{max} and ρ^{max} distinguish different aspects regarding the length of contract-paths. The function L^{opt} is concerned with Φ -paths realising a single exchange $\langle P, Q \rangle$ in a given resource allocation setting $\langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle$: the property of interest being the number of exchanges in the *shortest*, i.e. optimal length, Φ -path. We stress that L^{opt} is a *partial* function whose value is undefined in the event that $\langle P, Q \rangle$ cannot be realised by a Φ -path in the setting $\langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle$. The function L^{max} is defined in terms of L^{opt} , again in the context of a specific resource allocation setting. The behaviour of interest for L^{max} , however, is not simply the length of Φ -paths realising a specific $\langle P, Q \rangle$ but the “worst-case” value of L^{opt} for exchanges which are Ψ -exchanges. We note the qualification that L^{max} is defined only for Ψ -exchanges that *are* capable of being realised by Φ -paths, and thus do not consider cases for which no appropriate contract path exists. Thus, if it should be the case that no Ψ -exchange in the setting $\langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle$ can be realised by a Φ -path then the value $L^{\text{max}}(\langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle, \Phi, \Psi)$ is undefined, i.e. L^{max} is also a partial function. We may interpret any *upper* bound on L^{max} in the following terms: if $L^{\text{max}}(\langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle, \Phi, \Psi) \leq K$ then any Ψ -exchange *for which a Φ -path exists* can be realised by a Φ -path of length at most K .

Our main interest will centre on ρ^{max} which is concerned with the behaviour of L^{max} as a function of n and m and ranges over *all* n -tuples of utility functions $\langle u : 2^{\mathcal{R}} \rightarrow \mathbf{Q} \rangle^n$. Our

6. [20] gives an upper bound on the length of such paths which is also exponential in m , but does not explicitly state any lower bound other than that already referred to.

approach to obtaining lower bounds for this function is *constructive*, i.e. for each $\langle \Phi, \Psi \rangle$ that is considered, we show how the utility functions \mathcal{U} may be defined in a setting with m resources so as to yield a lower bound on $\rho^{\max}(n, m, \Phi, \Psi)$. In contrast to the measures L^{opt} and L^{\max} , the function ρ^{\max} is not described in terms of a single fixed resource allocation setting. It is, however, still a *partial* function: depending on $\langle n, m, \Phi, \Psi \rangle$ it may be the case that in *every* n agent, m resource allocation setting, regardless of which choice of utility functions is made, there is no Ψ -exchange, $\langle P, Q \rangle$ capable of being realised by Φ -path, and for such cases the value of $\rho^{\max}(n, m, \Phi, \Psi)$ will be undefined⁷.

It is noted, at this point, that the definition of ρ^{\max} allows *arbitrary* utility functions to be employed in constructing “worst-case” instances. While this is reasonable in terms of general lower bound results, as will be apparent from the given constructions the utility functions actually employed are highly artificial (and unlikely to feature in “real” application settings). We shall attempt to address this objection by further considering bounds on the following variant of ρ^{\max} :

$$\rho_{\text{mono}}^{\max}(n, m, \Phi, \Psi) = \max_{\mathcal{U}=\langle u_1, u_2, \dots, u_n \rangle : \text{each } u_i \text{ is monotone}} L^{\max}(\langle \mathcal{A}_n, \mathcal{R}_m, \mathcal{U} \rangle, \Phi, \Psi)$$

Thus, $\rho_{\text{mono}}^{\max}$ deals with resource allocation settings within which all of the utility functions must satisfy a monotonicity constraint.

The main results of this article are presented in the next section. We consider two general classes of contract path: O -contract paths under various rationality conditions; and, similarly, $M(k)$ -contract paths for arbitrary values of $k \geq 2$. Our results are concerned with the construction of resource allocation settings $\langle \mathcal{A}, \mathcal{R}_m, \mathcal{U} \rangle$ for which given some rationality requirement, e.g. that exchanges be individually rational, there is some exchange $\langle P, Q \rangle$ that satisfies the rationality condition, can be realised by an O -contract path (respectively, $M(k)$ -contract path), but with the number of exchanges required by such paths being exponential in m . We additionally obtain slightly weaker (but still exponential) lower bounds for O -contract paths within settings of monotone utility functions, i.e. for the measure $\rho_{\text{mono}}^{\max}$, outlining how similar results may be derived for $M(k)$ -contract paths.

In the resource allocation settings constructed for demonstrating these properties with $M(k)$ -contract paths, the constructed exchange $\langle P, Q \rangle$ is realisable with a *single* $M(k+1)$ -contract but unrealisable by any *rational* $M(k-1)$ -contract path. Conclusions and some directions for further work are presented in the final section.

7. In recognising the *possibility* that $\rho^{\max}(n, m, \Phi, \Psi)$ could be undefined, we are *not* claiming that such behaviour arises with any of the instantiations of $\langle \Phi, \Psi \rangle$ considered subsequently: in fact it will be clear from the constructions that, denoting by $\rho_{\Phi, \Psi}^{\max}(n, m)$ the function $\rho^{\max}(n, m, \Phi, \Psi)$ for a fixed instantiation of $\langle \Phi, \Psi \rangle$, with the restricted exchange types and rationality conditions examined, the function $\rho_{\Phi, \Psi}^{\max}(n, m)$ is a *total* function. Whether it is possible to formulate “sensible” choices of $\langle \Phi, \Psi \rangle$ with which $\rho_{\Phi, \Psi}^{\max}(n, m)$ is undefined for some values of $\langle n, m \rangle$ (and, if so, demonstrating examples of such) is, primarily, only a question of combinatorial interest, whose development is not central to the concerns of the current article.

2. Lower Bounds on Path Length

2.1 Overview

The strategy employed in proving our results involves two parts: for a given class of restricted contract paths we proceed as follows in obtaining lower bounds on $\rho^{\max}(n, m, \Phi, \Psi)$.

- a. For the contract-net graph partitioning m resources among n agents, construct a path, $\Delta_m = \langle P^{(1)}, P^{(2)}, \dots, P^{(t)} \rangle$ realising an exchange $\langle P^{(1)}, P^{(t)} \rangle$. For the *structural* constraint, Φ' influencing Φ it is then proved that:
 - a1. The contract path Δ_m is a Φ' -path, i.e. for each $1 \leq i < t$, the exchange $\langle P^{(i)}, P^{(i+1)} \rangle$ satisfies the structural constraint Φ' .
 - a2. For any pair of allocations $P^{(i)}$ and $P^{(i+j)}$ occurring in Δ_m , if $j \geq 2$ then the exchange $\langle P^{(i)}, P^{(i+j)} \rangle$ is *not* a Φ' -exchange.

Thus (a1) ensures that Δ_m is a suitable contract path, while (a2) will guarantee that there is exactly one allocation, $P^{(i+1)}$, that can be reached *within* Δ_m from any given allocation $P^{(i)}$ in Δ_m by means of a Φ' -exchange.

- b. Define utility functions $\mathcal{U}_n = \langle u_1, \dots, u_n \rangle$ with the following properties
 - b1. The exchange $\langle P^{(1)}, P^{(t)} \rangle$ is a Ψ -exchange.
 - b2. For the *rationality* constraint, Φ'' influencing Φ , every exchange $\langle P^{(i)}, P^{(i+1)} \rangle$ is a Φ'' -exchange.
 - b3. For every allocation $P^{(i)}$ in the contract path Δ and every allocation Q other than $P^{(i+1)}$ the exchange $\langle P^{(i)}, Q \rangle$ is *not* a Φ -exchange, i.e. it violates either the structural constraint Φ' or the rationality constraint Φ'' .

Thus, (a1) and (b2) ensure that $\langle P^{(1)}, P^{(t)} \rangle$ has a defined value with respect to the function L^{opt} for the Ψ -exchange $\langle P^{(1)}, P^{(t)} \rangle$, i.e. a Φ -path realising the exchange is possible. The properties given by (a2) and (b3) indicate that (within the constructed resource allocation setting) the path Δ_m is the *unique* Φ -path realising $\langle P^{(1)}, P^{(t)} \rangle$. It follows that $t - 1$, the length of this path, gives a *lower bound* on the value of L^{\max} and hence a lower bound on $\rho^{\max}(n, m, \Phi, \Psi)$.

Before continuing it will be useful to fix some notational details.

We use \mathcal{H}_m to denote the m -dimensional hypercube. Interpreted as a directed graph, \mathcal{H}_m has 2^m vertices each of which is identified with a distinct m -bit label. Using $\alpha = a_1 a_2 \dots a_m$ to denote an arbitrary such label, the edges of \mathcal{H}_m are formed by

$$\{ \langle \alpha, \beta \rangle : \alpha \text{ and } \beta \text{ differ in exactly one bit position} \}$$

We identify m -bit labels $\alpha = a_1 a_2 \dots a_m$ with subsets S^α of \mathcal{R}_m , via $r_i \in S^\alpha$ if and only if $a_i = 1$. Similarly, any subset S of \mathcal{R} can be described by a binary word, $\beta(S)$, of length m , i.e. $\beta(S) = b_1 b_2 \dots b_m$ with $b_i = 1$ if and only if $r_i \in S$. For a label α we use $|\alpha|$ to denote the number of bits with value 1, so that $|\alpha|$ is the size of the subset S^α . If α and β are m -bit labels, then $\alpha\beta$ is a $2m$ -bit label, so that if \mathcal{R}_m and \mathcal{T}_m are disjoint sets, then $\alpha\beta$ describes

the union of the subset S^α of \mathcal{R}_m with the subset S^β of \mathcal{T}_m . Finally if $\alpha = a_1 a_2 \dots a_m$ is an m -bit label then $\bar{\alpha}$ denotes the label formed by changing all 0 values in α to 1 and *vice versa*. In this way, if S^α is the subset of \mathcal{R}_m described by α then $\bar{\alpha}$ describes the set $\mathcal{R}_m \setminus S^\alpha$. To avoid an excess of superscripts we will, where no ambiguity arises, use α both to denote the m -bit label and the subset of \mathcal{R}_m described by it, e.g. we write $\alpha \subset \beta$ rather than $S^\alpha \subset S^\beta$.

For $n = 2$ the contract-net graph induced by O -contracts can be viewed as the m -dimensional hypercube \mathcal{H}_m : the m -bit label, α associated with a vertex of \mathcal{H}_m describing the allocation $\langle \alpha, \bar{\alpha} \rangle$ to $\langle A_1, A_2 \rangle$. In this way the set of IR O -contracts define a subgraph, \mathcal{G}_m of \mathcal{H}_m with any directed path from $\beta(P)$ to $\beta(Q)$ in \mathcal{G}_m corresponding to a possible IR O -contract path from the allocation $\langle P, \mathcal{R} \setminus P \rangle$ to the allocation $\langle Q, \mathcal{R} \setminus Q \rangle$.

2.2 O -contract paths

Our first result clarifies one issue in the presentation of [20, Proposition 2].

Theorem 3 *Let $\Phi(P, Q)$ be the predicate which holds whenever $\langle P, Q \rangle$ is an IR O -contract and $\Psi(P, Q)$ that which holds whenever $\langle P, Q \rangle$ is IR. For $m \geq 7$*

$$\rho^{\max}(2, m, \Phi, \Psi) \geq \left(\frac{77}{256} \right) 2^m - 2$$

Proof. Consider a path $\mathcal{C} = \langle \alpha_1, \alpha_2, \dots, \alpha_t \rangle$ in \mathcal{H}_m , with the following property⁸

$$\forall 1 \leq i < j \leq t \ (j \geq i + 2 \Rightarrow (\alpha_i \text{ and } \alpha_j \text{ differ in at least 2 positions})) \quad (\text{SC})$$

e.g. if $m = 4$ then

$$\emptyset, \{r_1\}, \{r_1, r_3\}, \{r_1, r_2, r_3\}, \{r_2, r_3\}, \{r_2, r_3, r_4\}, \{r_2, r_4\}, \{r_1, r_2, r_4\}$$

is such a path as it corresponds to the sequence $\langle 0000, 1000, 1010, 1110, 0110, 0111, 0101, 1101 \rangle$.

Choose $\mathcal{C}^{(m)}$ to be a *longest* such path with this property that could be formed in \mathcal{H}_m , letting $\Delta_m = \langle P^{(1)}, P^{(2)}, \dots, P^{(t)} \rangle$ be the sequence of allocations with $P^{(i)} = \langle \alpha_i, \bar{\alpha}_i \rangle$. We now define the utility functions u_1 and u_2 so that for $\gamma \subseteq \mathcal{R}_m$,

$$u_1(\gamma) + u_2(\bar{\gamma}) = \begin{cases} k & \text{if } \gamma = \alpha_k \\ 0 & \text{if } \gamma \notin \{\alpha_1, \alpha_2, \dots, \alpha_t\} \end{cases}$$

With this choice, the contract path Δ_m describes the *unique* IR O -contract path realising the IR exchange $\langle P^{(1)}, P^{(t)} \rangle$: that Δ_m is an IR O -contract path is immediate, since

$$\sigma_u(P^{(i+1)}) = i + 1 > i = \sigma_u(P^{(i)})$$

That it is unique follows from the fact that for all $1 \leq i \leq t$ and $i + 2 \leq j \leq t$, the exchange $\langle P^{(i)}, P^{(j)} \rangle$ is not an O -contract (hence there are no “short-cuts” possible), and for each $P^{(i)}$ there is exactly one IR O -contract that can follow it, i.e. $P^{(i+1)}$.⁹

8. This defines the so-called “snake-in-the-box” codes introduced in [12].

9. In our example with $m = 4$, the sequence $\langle 0000, 1000, 1001, 1101 \rangle$, although defining an O -contract path gives rise to an exchange which is not IR, namely that corresponding to $\langle 1000, 1001 \rangle$.

From the preceding argument it follows that any lower bound on the length of $\mathcal{C}^{(m)}$, i.e. a sequence satisfying the condition (SC), is a lower bound on $\rho^{\max}(2, m, \Phi, \Psi)$. These paths in \mathcal{H}_m were originally studied by Kautz [12] in the context of coding theory and the lower bound on their length of $(77/256)2^m - 2$ established by Abbott and Katchalski [1]. \square

There are a number of alternative formulations of “rationality” which can also be considered. For example

Definition 7 *Let $\delta = \langle P, Q \rangle$ be an exchange.*

- a. δ is cooperatively rational if for every agent, A_i , $u_i(Q_i) \geq u_i(P_i)$ and there is at least one agent, A_j , for whom $u_j(Q_j) > u_j(P_j)$.
- b. δ is equitable if $\min_{i \in \mathcal{A}^\delta} u_i(Q_i) > \min_{i \in \mathcal{A}^\delta} u_i(P_i)$.
- c. δ is a Pigou-Dalton exchange if $\mathcal{A}^\delta = \{i, j\}$, $u_i(P_i) + u_j(P_j) = u_i(Q_i) + u_j(Q_j)$ and $|u_i(Q_i) - u_j(Q_j)| < |u_i(P_i) - u_j(P_j)|$ (where $|\dots|$ is absolute value).

There are a number of views we can take concerning the rationality conditions given in Definition 7. One shared feature is that, unlike the concept of individual rationality for which some provision to compensate agents who suffer a loss in utility is needed, i.e. individual rationality presumes a “money-based” system, the forms defined in Definition 7 allow concepts of “rationality” to be given in “money-free” environments. Thus, in a cooperatively rational exchange, no agent involved suffers a loss in utility and *at least one* is better off. It may be noted that given the characterisation of Definition 4 it is immediate that any cooperatively rational exchange is perforce also individually rational; the converse, however, clearly does not hold in general. In some settings, an equitable exchange may be neither cooperatively nor individually rational. One may interpret such exchanges as one method of reducing inequality between the values agents place on their allocations: for those involved in an equitable exchange, it is ensured that the agent who places least value on their current allocation will obtain a resource set which is valued more highly. It may, of course, be the case that some agents suffer a *loss* of utility: the condition for an exchange to be equitable limits how great such a loss could be. Finally the concept of Pigou-Dalton exchange originates from and has been studied in depth within the theory of exchange economies. This is one of many approaches that have been proposed, again in order to describe exchanges which reduce inequality between members of an agent society. In terms of the definition given, such exchanges encapsulate the so-called Pigou-Dalton principle in economic theory: that any transfer of income from a wealthy individual to a poorer one should reduce the disparity between them. We note that, in principle, we could define related rationality concepts based on several extensions of this principle that have been suggested, e.g. [3, 4, 13].

Using the same O -contract path constructed in Theorem 3, we need only vary the definitions of the utility functions employed in order to obtain,

Corollary 1 *For each of the cases below,*

- a. $\Phi(\delta)$ holds if and only if δ is a cooperatively rational O -contract.
 $\Psi(\delta)$ holds if and only if δ is cooperatively rational.
- b. $\Phi(\delta)$ holds if and only if δ is an equitable O -contract.
 $\Psi(\delta)$ holds if and only if δ is equitable.

- c. $\Phi(\delta)$ holds if and only if δ is a Pigou-Dalton O -contract.
 $\Psi(\delta)$ holds if and only if δ Pigou-Dalton exchange.

$$\rho^{\max}(2, m, \Phi, \Psi) \geq \left(\frac{77}{256}\right)2^m - 2$$

Proof. We employ exactly the same sequence of allocations, Δ_m described in the proof of Theorem 3 but modify the utility functions $\langle u_1, u_2 \rangle$ for each case.

- a. Choose $\langle u_1, u_2 \rangle$ with $u_2(\gamma) = 0$ for all $\gamma \subseteq \mathcal{R}$ and

$$u_1(\gamma) = \begin{cases} k & \text{if } \gamma = \alpha_k \\ 0 & \text{if } \gamma \notin \{\alpha_1, \dots, \alpha_t\} \end{cases}$$

The resulting O -contract path is cooperatively rational: the utility enjoyed by A_2 remains constant while that enjoyed by A_1 increases by 1 with each exchange. Any deviation from this contract path (employing an alternative O -contract) will result in a loss of utility for A_1 .

- b. Choose $\langle u_1, u_2 \rangle$ with $u_2(\gamma) = u_1(\bar{\gamma})$ and

$$u_1(\gamma) = \begin{cases} k & \text{if } \gamma = \alpha_k \\ 0 & \text{if } \gamma \notin \{\alpha_1, \dots, \alpha_t\} \end{cases}$$

The O -contract path is equitable: both A_1 and A_2 increase their respective utility values by 1 with each exchange. Again, any O -contract deviating from this will result in both agents losing some utility.

- c. Choose $\langle u_1, u_2 \rangle$ as

$$u_1(\gamma) = \begin{cases} k & \text{if } \gamma = \alpha_k \\ 0 & \text{if } \gamma \notin \{\alpha_1, \dots, \alpha_t\} \end{cases} \quad ; \quad u_2(\gamma) = \begin{cases} 2^m - k & \text{if } \bar{\gamma} = \alpha_k \\ 2^m & \text{if } \bar{\gamma} \notin \{\alpha_1, \dots, \alpha_t\} \end{cases}$$

To see that the O -contract path consists of Pigou-Dalton exchanges, it suffices to note that $u_1(\alpha_i) + u_2(\bar{\alpha}_i) = 2^m$ for each $1 \leq i \leq t$. In addition, $|u_2(\bar{\alpha}_{i+1}) - u_1(\alpha_{i+1})| = 2^m - 2i - 2$ which is strictly less than $|u_2(\bar{\alpha}_i) - u_1(\alpha_i)| = 2^m - 2i$. Finally, any O -contract $\langle P, Q \rangle$ which deviates from this sequence will not be a Pigou-Dalton exchange since

$$|u_2(Q_2) - u_1(Q_1)| = 2^m > |u_2(P_2) - u_1(P_1)|$$

which violates one of the conditions required of Pigou-Dalton exchanges. \square

The construction for two agent settings, easily extends to larger numbers.

Corollary 2 For each of the choices of $\langle \Phi, \Psi \rangle$ considered in Theorem 3 and Corollary 1, and all $n \geq 2$,

$$\rho^{\max}(n, m, \Phi, \Psi) \geq \left(\frac{77}{256}\right)2^m - 2$$

Proof. Fix allocations in which A_1 is given α_1 , A_2 allocated $\bar{\alpha}_1$, and A_j assigned \emptyset for each $3 \leq j \leq n$. Using identical utility functions $\langle u_1, u_2 \rangle$ as in each of the previous cases, we employ for u_j : $u_j(\emptyset) = 1$, $u_j(S) = 0$ whenever $S \neq \emptyset$ ($\langle \Phi, \Psi \rangle$ as in Theorem 3); $u_j(S) = 0$ for all S (Corollary 1(a)); $u_j(\emptyset) = 2^m$, $u_j(S) = 0$ whenever $S \neq \emptyset$ (Corollary 1(b)); and, finally, $u_j(S) = 2^m$ for all S , (Corollary 1(c)). Considering a realisation of the Ψ exchange $\langle P^{(1)}, P^{(t)} \rangle$ the only Φ -contract path admissible is the path Δ_m defined in the related proofs. This gives the lower bound stated. \square

We note, at this point, some other consequences of Corollary 1 with respect to [11, Theorems 1, 3], which state

Fact 1 *We recall that a Φ -path, $\langle P^{(1)}, \dots, P^{(t)} \rangle$ is maximal if for each allocation Q , $\langle P^{(t)}, Q \rangle$ is not a Φ -exchange.*

- a. *If $\langle P^{(1)}, \dots, P^{(t)} \rangle$ is any maximal path of cooperatively rational exchanges then $P^{(t)}$ is Pareto optimal.*
- b. *If $\langle P^{(1)}, \dots, P^{(t)} \rangle$ is any maximal path of equitable exchanges then $P^{(t)}$ maximises the value $\sigma_e(P) = \min_{1 \leq i \leq n} u_i(P_i)$, i.e. the so-called egalitarian social welfare.*

The sequence of cooperatively rational exchanges in Corollary 1(a) terminates in the Pareto optimal allocation $P^{(t)}$: the allocation for A_2 always has utility 0 and there is no allocation to A_1 whose utility can exceed t . Similarly, the sequence of equitable exchanges in Corollary 1(b) terminates in the allocation $P^{(t)}$, for which $\sigma_e(P^{(t)}) = t$ the maximum that can be attained for the instance defined. In both cases, however, the optima are reached by sequences of exponentially many (in m) exchanges: thus, although Fact 1 guarantees convergence of particular exchange sequences to optimal states, it may be the case, as illustrated in Corollary 1(a–b) that the process of convergence takes considerable time.

We conclude our results concerning O -contracts by presenting a lower bound on $\rho_{\text{mono}}^{\max}$, i.e. the length of paths when the utility functions are required to be *monotone*.

In principle one could attempt to construct appropriate monotone utility functions that would have the desired properties with respect to the path used in Theorem 3. It is, however, far from clear whether such a construction is possible. We do not attempt to resolve this question here. Whether an exact translation could be accomplished is, ultimately, a question of purely combinatorial interest: since our aim is to demonstrate that exponential length contract paths are needed with monotone utility functions we are not, primarily, concerned with obtaining an optimal bound.

Theorem 4 *With $\Phi(P, Q)$ and $\Psi(P, Q)$ be defined as in Theorem 3 and $m \geq 14$*

$$\rho_{\text{mono}}^{\max}(2, m, \Phi, \Psi) \geq \begin{cases} \left(\frac{77}{128}\right) 2^{m/2} - 3 & \text{if } m \text{ is even} \\ \left(\frac{77}{128}\right) 2^{(m-1)/2} - 3 & \text{if } m \text{ is odd} \end{cases}$$

Proof. We describe the details only for the case of m being even: the result when m is odd is obtained by a simple modification which we shall merely provide in outline.

Let $m = 2s$ with $s \geq 7$. For any path

$$\Delta_s = \langle \alpha_1, \alpha_2, \dots, \alpha_t \rangle$$

in \mathcal{H}_s (where α_i describes a subset of \mathcal{R}_s by an s -bit label), the path $double(\Delta_s)$ in \mathcal{H}_{2s} is defined by

$$\begin{aligned} double(\Delta_s) &= \langle \alpha_1 \overline{\alpha_1}, \alpha_2 \overline{\alpha_2}, \dots, \alpha_i \overline{\alpha_i}, \alpha_{i+1} \overline{\alpha_{i+1}}, \dots, \alpha_t \overline{\alpha_t} \rangle \\ &= \langle \beta_1, \beta_3, \dots, \beta_{2i-1}, \beta_{2i+1}, \dots, \beta_{2t-1} \rangle \end{aligned}$$

(The reason for successive indices of β increasing by 2 will become clear subsequently)

Of course, $double(\Delta_s)$ does not describe an O -contract path¹⁰: it is, however, not difficult to interpolate appropriate allocations, β_{2i} , in order to convert it to such a path. Consider the subsets, β_{2i} (with $1 \leq i < t$) defined as follows:

$$\beta_{2i} = \begin{cases} \alpha_{i+1} \overline{\alpha_i} & \text{if } \alpha_i \subset \alpha_{i+1} \\ \alpha_i \overline{\alpha_{i+1}} & \text{if } \alpha_i \supset \alpha_{i+1} \end{cases}$$

If we now consider the path, $ext(\Delta_s)$, within \mathcal{H}_{2s} given by

$$ext(\Delta_s) = \langle \beta_1, \beta_2, \beta_3, \dots, \beta_{2(t-1)}, \beta_{2t-1} \rangle$$

then this satisfies,

- a. If Δ_s has property (SC) of Theorem 3 in \mathcal{H}_s then $ext(\Delta_s)$ has property (SC) in \mathcal{H}_{2s} .
- b. If j is odd then $|\beta_j| = s$.
- c. If j is even then $|\beta_j| = s + 1$.

From (a) and the bounds proved in [1] we deduce that $ext(\Delta_s)$ can be chosen so that with $P^{(i)}$ denoting the allocation $\langle \beta_i, \overline{\beta_i} \rangle$

- d. $ext(\Delta_s)$ describes an O -contract path from $P^{(1)}$ to $P^{(2t-1)}$.
- e. For each pair $\langle i, j \rangle$ with $j \geq i + 2$, the exchange $\langle P^{(i)}, P^{(j)} \rangle$ is *not* an O -contract.
- f. If Δ_s is chosen as in the proof of Theorem 3 then the number of exchanges in $ext(\Delta_s)$ is as given in the statement of the present theorem.

We therefore fix Δ_s as the path from Theorem 3 so that in order to complete the proof we need to construct utility functions $\langle u_1, u_2 \rangle$ that are monotone and with which $ext(\Delta_s)$ defines the unique IR O -contract path realising the IR exchange $\langle P^{(1)}, P^{(2t-1)} \rangle$.

The choice for u_2 is relatively simple. Given $S \subseteq \mathcal{R}_{2s}$,

$$u_2(S) = \begin{cases} 0 & \text{if } |S| \leq s - 2 \\ 2t + 1 & \text{if } |S| = s - 1 \\ 2t + 2 & \text{if } |S| \geq s \end{cases}$$

10. In terms of the classification described in [20], it contains only *swap* exchanges (S -contracts): each exchange swaps exactly one item in β_{2i-1} with an item in $\overline{\beta_{2i-1}}$ in order to give β_{2i+1} .

In this t is the number of allocations in Δ_s . The behaviour of u_2 is clearly monotone.

The construction for u_1 is rather more complicated. Its main idea is to make use of the fact that the size of each set β_i occurring in $\text{ext}(\Delta_s)$ is very tightly constrained: $|\beta_i|$ is either s or $s + 1$ according to whether i is odd or even. We first demonstrate that each set of size $s + 1$ can have at most two strict subsets (of size s) occurring within $\text{ext}(\Delta_s)$: thus, every S of size $s + 1$ has exactly 2 or 1 or 0 subsets of size s on $\text{ext}(\Delta_s)$. To see this suppose the contrary. Let γ , β_{2i-1} , β_{2j-1} , and β_{2k-1} be such that $|\gamma| = s + 1$ with

$$\beta_{2i-1} \subset \gamma ; \beta_{2j-1} \subset \gamma ; \beta_{2k-1} \subset \gamma$$

Noting that $\beta_{2i-1} = \alpha_i \bar{\alpha}_i$ and that Δ_s has the property (SC) it must be the case that (at least) two of the s -bit labels from $\{\alpha_i, \alpha_j, \alpha_k\}$ differ in at least two positions. Without loss of generality suppose this is true of α_i and α_k . As a result we deduce that the sets β_{2i-1} and β_{2k-1} have at most $s - 2$ elements in common, i.e. $|\beta_{2i-1} \cap \beta_{2k-1}| \leq s - 2$: $\beta_{2i-1} = \alpha_i \bar{\alpha}_i$ and $\beta_{2k-1} = \alpha_k \bar{\alpha}_k$ so in any position at which α_i differs from α_k , $\bar{\alpha}_i$ differs from $\bar{\alpha}_k$ at exactly the same position. In total $|\beta_{2i-1} \setminus \beta_{2k-1}| \geq 2$, i.e. there are (at least) two elements of β_{2i-1} that do not occur in β_{2k-1} ; and in the same way $|\beta_{2k-1} \setminus \beta_{2i-1}| \geq 2$, i.e. there are (at least) two elements β_{2k-1} that do not occur in β_{2i-1} . The set γ , however, has only $s + 1$ members and so cannot have *both* β_{2i-1} and β_{2k-1} as subsets: this would require

$$\beta_{2i-1} \cap \beta_{2k-1} \cup \beta_{2i-1} \setminus \beta_{2k-1} \cup \beta_{2k-1} \setminus \beta_{2i-1} \subseteq \gamma$$

but, as we have just seen,

$$|\beta_{2i-1} \cap \beta_{2k-1} \cup \beta_{2i-1} \setminus \beta_{2k-1} \cup \beta_{2k-1} \setminus \beta_{2i-1}| \geq s + 2$$

One immediate consequence of the argument just given is that for any set γ of size $s + 1$ there are exactly two strict subsets of γ occurring on $\text{ext}(\Delta_s)$ if and only if $\gamma = \beta_{2i-1} \cup \beta_{2i+1} = \beta_{2i}$ for some value of i with $1 \leq i < t$. We can now characterise each subset of \mathcal{R}_{2s} of size $s + 1$ as falling into one of three categories.

C1. *Good* sets, given by $\{\gamma : \gamma = \beta_{2i}\}$.

C2. *Digressions*, consisting of

$$\{\gamma : \beta_{2i-1} \subset \gamma \text{ and } \gamma \neq \beta_{2i}\}$$

C3. *Inaccessible* sets, consisting of

$$\{\gamma : \gamma \text{ is neither } \textit{Good} \text{ nor a } \textit{Digression}\}$$

Good sets are those describing allocations to A_1 within the path defined by $\text{ext}(\Delta_s)$; *Digressions* are the allocations that could be reached using an O -contract from a set of size s on $\text{ext}(\Delta_s)$, i.e. β_{2i-1} , but differ from the set that actually occurs in $\text{ext}(\Delta_s)$, i.e. β_{2i} . Finally, *Inaccessible* sets are those that do not occur on $\text{ext}(\Delta_s)$ and cannot be reached via an O -contract from any set on $\text{ext}(\Delta_s)$.

We now define u_1 as

$$u_1(\gamma) = \begin{cases} 2i - 1 & \text{if } \gamma = \beta_{2i-1} \\ 2i + 1 & \text{if } \gamma = \beta_{2i} \\ 2i & \text{if } |\gamma| = s + 1 \text{ and } \gamma \text{ is a } Digression \\ 0 & \text{if } |\gamma| \leq s - 1 \\ 0 & \text{if } |\gamma| = s \text{ and } \gamma \notin ext(\Delta_s) \\ 2t - 1 & \text{if } \gamma \text{ is } Inaccessible \text{ or } |\gamma| \geq s + 2 \end{cases}$$

It remains only to prove for these choices of $\langle u_1, u_2 \rangle$ that the O -contract path $\langle P^{(1)}, \dots, P^{(2t-1)} \rangle$ defined from $ext(\Delta_s)$ is the unique IR O -contract path realising the IR exchange $\langle P^{(1)}, P^{(2t-1)} \rangle$ and that u_1 is monotone.

To show that $\langle P^{(1)}, \dots, P^{(2t-1)} \rangle$ is IR we need to demonstrate

$$\forall 1 \leq j < 2t - 1 \quad u_1(\beta_j) + u_2(\overline{\beta_j}) < u_1(\beta_{j+1}) + u_2(\overline{\beta_{j+1}})$$

We have via the definition of $\langle u_1, u_2 \rangle$

$$\begin{aligned} u_1(\beta_{2i-1}) + u_2(\overline{\beta_{2i-1}}) &= 2(t + i) + 1 \\ &< u_1(\beta_{2i}) + u_2(\overline{\beta_{2i}}) \\ &= 2(t + i) + 2 \\ &< u_1(\beta_{2i+1}) + u_2(\overline{\beta_{2i+1}}) \\ &= 2(t + i) + 3 \end{aligned}$$

Thus, via Definition 4, it follows that $ext(\Delta_s)$ gives rise to an IR O -contract path.

To see that this path is the unique IR O -contract path implementing $\langle P^{(1)}, P^{(2t-1)} \rangle$, consider any position $P^{(j)} = \langle \beta_j, \overline{\beta_j} \rangle$ and allocation Q other than $P^{(j+1)}$ or $P^{(j-1)}$. It may be assumed that the exchange $\langle P^{(j)}, Q \rangle$ is an O -contract. If $j = 2i - 1$ then $\sigma_u(P^{(2i-1)}) = 2(t + i) + 1$ and $|\beta_j| = s$. Hence $|Q_1| \in \{s - 1, s + 1\}$. In the former case, $u_1(Q_1) = 0$ and $u_2(Q_2) = 2t + 2$ from which $\sigma_u(Q) = 2t + 2$ and thus $\langle P^{(j)}, Q \rangle$ is not IR. In the latter case $u_1(Q_1) = 2i$ since Q_1 is a *Digression* and $u_2(Q_2) = 2t + 1$ giving $\sigma_u(Q) = 2(t + i) + 1$. Again $\langle P^{(j)}, Q \rangle$ fails to be IR since Q fails to give any increase in the value of σ_u . We are left with the case $j = 2i$ so that $\sigma_u(P^{(2i)}) = 2(t + i) + 2$ and $|\beta_j| = s + 1$. Since $\langle P^{(j)}, Q \rangle$ is assumed to be an O -contract this gives $|Q_1| \in \{s, s + 2\}$. For the first possibility Q_1 could not be a set on $ext(\Delta_s)$: β_{2i-1} and β_{2i+1} are both subsets of β_{2i} and there can be at most two such subsets occurring on $ext(\Delta_s)$. It follows, therefore, that $u_1(Q_1) = 0$ giving $\sigma_u(Q) = 2t + 2$ so that $\langle P^{(j)}, Q \rangle$ is not IR. In the second possibility, $u_1(Q_1) = 2t - 1$ but $u_2(Q_2) = 0$ as $|Q_2| = s - 2$ so the exchange would result in an overall loss. We deduce that for each $P^{(j)}$ the only IR O -contract consistent with it is the exchange $\langle P^{(j)}, P^{(j+1)} \rangle$.

The final stage is to prove that the utility function u_1 is indeed a *monotone* function. Suppose S and T are subsets of \mathcal{R}_{2s} with $S \subset T$. We need to show that $u_1(S) \leq u_1(T)$. We may assume that $|S| = s$, that S occurs as some set within $ext(\Delta_s)$, and that $|T| = s + 1$. If $|S| < s$ or $|S| = s$ but does not occur on $ext(\Delta_s)$ we have $u_1(S) = 0$ and the required inequality holds; if $|S| \geq s + 1$ then in order for $S \subset T$ to be possible we would need $|T| \geq s + 2$, which would give $u_1(T) = 2t - 1$ and this is the maximum value that any subset is assigned by u_1 . We are left with only $|S| = s$, $|T| = s + 1$ and S on $ext(\Delta_s)$ to consider. It has already been shown that there are at most two subsets of T that can occur on $ext(\Delta_s)$. Consider the different possibilities:

- a. $T = \beta_{2i}$ so that exactly two subsets of T occur in $\text{ext}(\Delta_s)$: β_{2i-1} and β_{2i+1} . Since $u_1(\beta_{2i}) = 2i + 1$ and this is at least $\max\{u_1(\beta_{2i-1}), u_1(\beta_{2i+1})\}$, should S be either of β_{2i-1} or β_{2i+1} then $u_1(S) \leq u_1(T)$ as required.
- b. T is a *Digression* from $S = \beta_{2i-1}$, so that $u_1(T) = 2i$ and $u_1(S) = 2i - 1$ and, again, $u_1(S) \leq u_1(T)$.

We deduce that u_1 is monotone completing our lower bound proof for $\rho_{\text{mono}}^{\max}$ for even values of m .

We conclude by observing that a similar construction can be used if $m = 2s + 1$ is odd: use the path $\text{ext}(\Delta_s)$ described above but modifying it so that one resource (r_m) is always held by A_2 . Only minor modifications to the utility function definitions are needed. \square

Example 1 For $s = 3$, we can choose $\Delta_3 = \langle 000, 001, 101, 111, 110 \rangle$ so that $t = 5$. This gives $\text{double}(\Delta_3)$ as

$$\langle 000111, 001110, 101010, 111000, 110001 \rangle$$

with the O -contract path being defined from $\text{ext}(\Delta_3)$ which is

$$\langle 000111, 001111, 001110, 101110, 101010, 111010, 111000, 111001, 110001 \rangle$$

Considering the 15 subsets of size $s + 1 = 4$, gives

$$\begin{aligned} \text{Good} &= \{001111, 101110, 111010, 111001\} \\ \text{Digression} &= \{010111, 100111, 101011, 110011, 110101, 011110, 11110\} \\ \text{Inaccessible} &= \{011011, 011101, 101101, 110110\} \end{aligned}$$

Notice that we choose to view both of the sets in $\{110011, 110101\}$ as a *Digression*: in principle we could continue from $\beta_9 = 110001$ using either, however, in order to simplify the construction the path is halted at β_9 .

The monotone utility functions, $\langle u_1, u_2 \rangle$, employed in proving Theorem 4 are defined so that the path arising from $\text{ext}(\Delta_s)$ is IR: in the event of either agent suffering a loss of utility the gain made by the other is sufficient to provide a compensatory payment. A natural question that now arises is whether the bound obtained in Theorem 4 can be shown to apply when the rationality conditions preclude any monetary payment, e.g. for cases where the concept of rationality is one of those given in Definition 7. Our next result shows that if we set the rationality condition to enforce cooperatively rational or equitable exchanges then the bound of Theorem 4 still holds.

Theorem 5 For each of the cases below and $m \geq 14$

- a. $\Phi(\delta)$ holds if and only if δ is a cooperatively rational O -contract.
 $\Psi(\delta)$ holds if and only if δ is cooperatively rational.
- b. $\Phi(\delta)$ holds if and only if δ is an equitable O -contract.
 $\Psi(\delta)$ holds if and only if δ is equitable.

$$\rho_{\text{mono}}^{\max}(2, m, \Phi, \Psi) \geq \begin{cases} \left(\frac{77}{128}\right) 2^{m/2} - 3 & \text{if } m \text{ is even} \\ \left(\frac{77}{128}\right) 2^{(m-1)/2} - 3 & \text{if } m \text{ is odd} \end{cases}$$

Proof. We again illustrate the constructions only for the case of m being even, noting the modification to deal with odd values of m outlined at the end of the proof of Theorem 4. The path $\text{ext}(\Delta_s)$ is used for both cases.

For (a), we require $\langle u_1, u_2 \rangle$ to be defined as monotone functions with which $\text{ext}(\Delta_s)$ will be the unique cooperatively rational O -contract path to realise the cooperatively rational exchange $\langle P^{(1)}, P^{(2t-1)} \rangle$ where $P^{(j)} = \langle \beta_j, \bar{\beta}_j \rangle$. In this case we set $\langle u_1, u_2 \rangle$ to be,

$$\langle u_1(\gamma), u_2(\bar{\gamma}) \rangle = \begin{cases} \langle i, i \rangle & \text{if } \gamma = \beta_{2i-1} \\ \langle i+1, i \rangle & \text{if } \gamma = \beta_{2i} \\ \langle i, i-1 \rangle & \text{if } |\gamma| = s+1 \text{ and } \gamma \text{ is a } Digression \\ \langle 0, 2t-1 \rangle & \text{if } |\gamma| \leq s-1 \\ \langle 0, 2t-1 \rangle & \text{if } |\gamma| = s \text{ and } \gamma \notin \text{ext}(\Delta_s) \\ \langle 2t-1, 0 \rangle & \text{if } \gamma \text{ is } Inaccessible \text{ or } |\gamma| \geq s+2 \end{cases}$$

Since,

$$\begin{aligned} \langle u_1(\beta_{2i-1}), u_2(\bar{\beta}_{2i-1}) \rangle &= \langle i, i \rangle \\ \langle u_1(\beta_{2i}), u_2(\bar{\beta}_{2i}) \rangle &= \langle i+1, i \rangle \\ \langle u_1(\beta_{2i+1}), u_2(\bar{\beta}_{2i+1}) \rangle &= \langle i+1, i+1 \rangle \end{aligned}$$

it is certainly the case that $\langle P^{(1)}, P^{(2t-1)} \rangle$ and all exchanges on the O -contract path defined by $\text{ext}(\Delta_s)$ are cooperatively rational. Furthermore if $Q = \langle \gamma, \bar{\gamma} \rangle$ is any allocation other than $P^{(j+1)}$ then the exchange $\langle P^{(j)}, Q \rangle$ will fail to be a cooperatively rational O -contract. For suppose the contrary letting $\langle P^{(j)}, Q \rangle$ without loss of generality be an O -contract, with $Q \notin \{P^{(j-1)}, P^{(j+1)}\}$ – we can rule out the former case since we have already shown such an exchange is not cooperatively rational. If $j = 2i - 1$ so that $\langle u_1(\beta_j), u_2(\bar{\beta}_j) \rangle = \langle i, i \rangle$ then $|\gamma| \in \{s-1, s+1\}$: the former case leads to a loss in utility for A_1 ; the latter, (since γ is a *Digression*) a loss in utility for A_2 . Similarly, if $j = 2i$ so that $\langle u_1(\beta_j), u_2(\bar{\beta}_j) \rangle = \langle i+1, i \rangle$ then $|\gamma| \in \{s, s+2\}$: for the first $\gamma \notin \text{ext}(\Delta_s)$ leading to a loss of utility for A_1 ; the second results in a loss of utility for A_2 . It follows that the path defined by $\text{ext}(\Delta_s)$ is the unique cooperatively rational O -contract path that realises $\langle P^{(1)}, P^{(2t-1)} \rangle$.

It remains only to show that these choices for $\langle u_1, u_2 \rangle$ define monotone utility functions.

Consider u_1 and suppose S and T are subsets of \mathcal{R}_{2s} with $S \subset T$. If $|S| \leq s-1$, or S does not occur on $\text{ext}(\Delta_s)$ then $u_1(S) = 0$. If $|T| \geq s+2$ or is *Inaccessible* then $u_1(T) = 2t-1$ which is the maximum value attainable by u_1 . So we may assume that $|S| = s$, occurs on $\text{ext}(\Delta_s)$, i.e. $S = \beta_{2i-1}$, for some i , and that $|T| = s+1$ and is either a *Good* set or a *Digression*. From the definition of u_1 , $u_1(S) = i$: if $T \in \{\beta_{2i}, \beta_{2i-2}\}$ then $u_1(T) \geq i = u_1(S)$; if T is a *Digression* $u_1(T) = i = u_1(S)$. We deduce that if $S \subseteq T$ then $u_1(S) \leq u_1(T)$, i.e. the utility function is monotone.

Now consider u_2 with S and T subsets of \mathcal{R}_{2s} having $S \subset T$. If $|T| \geq s+1$ or $\mathcal{R}_{2s} \setminus T$ does not occur in $\text{ext}(\Delta_s)$ then $u_2(T) = 2t-1$ its maximal value. If $|S| \leq s-2$

or $\mathcal{R}_{2s} \setminus S$ is *Inaccessible* then $u_2(S) = 0$. Thus we may assume that $T = \overline{\beta_{2i-1}}$ giving $u_2(T) = i$ and $|S| = s - 1$, so that $\mathcal{R}_{2s} \setminus S$ is either a *Digression* or one of the *Good* sets $\{\beta_{2i}, \beta_{2i-2}\}$. If $\mathcal{R}_{2s} \setminus S$ is a *Digression* then $u_2(S) = i - 1$; if it is the *Good* set β_{2i-2} then $u_2(S) = i - 1 < u_2(T)$; if it is the *Good* set β_{2i} then $u_2(S) = i = u_2(T)$. It follows that u_2 is monotone completing the proof of part (a).

For (b) we use,

$$\langle u_1(\gamma), u_2(\bar{\gamma}) \rangle = \begin{cases} \langle 2i - 1, 2i \rangle & \text{if } \gamma = \beta_{2i-1} \\ \langle 2i + 1, 2i \rangle & \text{if } \gamma = \beta_{2i} \\ \langle 2i, 2i - 1 \rangle & \text{if } |\gamma| = s + 1 \text{ and } \gamma \text{ is a } \textit{Digression} \\ \langle 0, 2t - 1 \rangle & \text{if } |\gamma| \leq s - 1 \\ \langle 0, 2t - 1 \rangle & \text{if } |\gamma| = s \text{ and } \gamma \notin \textit{ext}(\Delta_s) \\ \langle 2t - 1, 0 \rangle & \text{if } \gamma \text{ is } \textit{Inaccessible} \text{ or } |\gamma| \geq s + 2 \end{cases}$$

These choices give $\textit{ext}(\Delta_s)$ as the unique equitable O -contract path to realise the equitable exchange $\langle P^{(1)}, P^{(2t-1)} \rangle$, since

$$\begin{aligned} \min\{u_1(\beta_{2i-1}), u_2(\overline{\beta_{2i-1}})\} &= 2i - 1 \\ \min\{u_1(\beta_{2i}), u_2(\overline{\beta_{2i}})\} &= 2i \\ \min\{u_1(\beta_{2i+1}), u_2(\overline{\beta_{2i+1}})\} &= 2i + 1 \end{aligned}$$

each exchange $\langle P^{(j)}, P^{(j+1)} \rangle$ is equitable. If $Q = \langle \gamma, \bar{\gamma} \rangle$ is any allocation other than $P^{(j+1)}$ then the exchange $\langle P^{(j)}, Q \rangle$ is not an equitable O -contract. Assume that $\langle P^{(j)}, Q \rangle$ is an O -contract, and that $Q \notin \{P^{(j-1)}, P^{(j+1)}\}$. If $j = 2i - 1$, so that $P^{(j)} = \langle \beta_{2i-1}, \overline{\beta_{2i-1}} \rangle$ and $\min\{u_1(\beta_{2i-1}), u_2(\overline{\beta_{2i-1}})\} = 2i - 1$ then $|\gamma| \in \{s - 1, s + 1\}$. In the first of these $\min\{u_1(\gamma), u_2(\bar{\gamma})\} = 0$; in the second $\min\{u_1(\gamma), u_2(\bar{\gamma})\} = 2i - 1$ since γ must be a *Digression*. This leaves only $j = 2i$ with $P^{(j)} = \langle \beta_{2i}, \overline{\beta_{2i}} \rangle$ and $\min\{u_1(\beta_{2i}), u_2(\overline{\beta_{2i}})\} = 2i$. For this, $|\gamma| \in \{s, s + 2\}$: if $|\gamma| = s$ then $\min\{u_1(\gamma), u_2(\bar{\gamma})\} \leq 2i - 1$ (with equality when $\gamma = \beta_{2i-1}$); if $|\gamma| = s + 2$ then $\min\{u_1(\gamma), u_2(\bar{\gamma})\} = 0$. In total these establish that $\textit{ext}(\Delta_s)$ is the unique equitable O -contract path realising the equitable exchange $\langle P^{(1)}, P^{(2t-1)} \rangle$.

That the choices for $\langle u_1, u_2 \rangle$ describe monotone utility functions can be shown by a similar argument to that of part (a). \square

Discussion

That we can demonstrate similar extremal behaviours for contract path length with rationality constraints in both money-based (individual rationality) and money-free (cooperative rationality, equitable) settings irrespective of whether monotonicity properties are assumed, has some interesting parallels with other contexts in which monotonicity is relevant. In particular we can observe that in common with the complexity results already noted from [8] – deciding if an allocation is Pareto optimal, or if an allocation maximises σ_u , or if an IR O -contract path exists – requiring utility functions to be monotone does not result in a setting which is computationally more tractable. In a related, but somewhat different context, we have further indications that monotonicity may not suffice to ensure reasonable computational properties. Thus, [27] present a multiagent coalitional model, in which the feasibility of a particular set of outcomes with respect to a given set of agents is described

by a propositional logic function.¹¹ In this context a natural requirement to impose is that if a set of outcomes is feasible for some coalition \mathcal{C} then it is also feasible for any superset of \mathcal{C} : in informal terms, such “coalition monotonic” environments, require coalitions to be capable of realising any set of goals that could be brought about by any of their subsets. In [27] the computational complexity of a number of decision questions is studied, e.g. does a given coalition have *any* feasible set of outcomes; does *every* coalition have a feasible set of outcomes, etc. In the general setting, where no monotonicity requirement is imposed, the problem complexity ranges from NP–complete and CO-NP–complete to completeness for a class thought strictly to contain $\Sigma_2^P \cup \Pi_2^P$. It turns out, however, that even when consideration is limited to coalition monotonic cases, such restrictions tend not to affect the problem complexity: of the ten cases for which both general and coalition monotonic examples are examined, only two yield a reduction in complexity for the restricted form.

2.3 $M(k)$ -contract paths

We now turn to similar issues with respect to $M(k)$ -contracts, recalling that in one respect these offer a form of exchange that does not fit into the classification of [20] since $M(k)$ -contracts permit *two* agents to exchange resources. In one regard, however, $M(k)$ -contracts are not as general as the multiagent contracts (M -contracts) of [20] since a preset bound (k) is specified for the number of agents involved.

Our main result on $M(k)$ -contract paths is the following development of Theorem 3.

Theorem 6 *Let $\Phi_k(P, Q)$ be the predicate which holds whenever $\langle P, Q \rangle$ is an IR $M(k)$ -contract. For all $k \geq 3$, $n \geq k$ and $m \geq \binom{k}{2}$, there is a resource allocation setting $\langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle$ and an IR exchange $\delta = \langle P, Q \rangle$ for which,*

$$\begin{aligned} L^{\text{opt}}(\delta, \langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle, \Phi_k) &= 1 & (a) \\ L^{\text{opt}}(\delta, \langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle, \Phi_{k-1}) &\geq 2^{\lfloor 2m/k(k-1) \rfloor} - 1 & (b) \\ L^{\text{opt}}(\delta, \langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle, \Phi_{k-2}) &\text{is undefined} & (c) \end{aligned}$$

Before presenting the proof, we comment about the formulation of the theorem statement and give an overview of the proof structure.

We first note that the lower bounds (where defined) have been phrased in terms of the function L^{opt} as opposed to ρ^{max} used in the various results on O -contract paths in Section 2.2. It is, of course, the case that the bound claimed for $L^{\text{opt}}(\delta, \langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle, \Phi_{k-1})$ will also be a lower bound on $\rho^{\text{max}}(n, m, \Phi_{k-1}, \Psi)$ when $n \geq k$ and $\Psi(P, Q)$ holds whenever the exchange $\langle P, Q \rangle$ is IR. The statement of Theorem 6, however, claims rather more than this, namely that a *specific* resource allocation setting $\langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle$ can be defined for each $n \geq k$ and each m , together with an IR exchange $\langle P, Q \rangle$ in such a way that: $\langle P, Q \rangle$ can be achieved by a *single* $M(k)$ -contract and *cannot* be realised by an IR $M(k-2)$ -contract path. Recalling that L^{opt} is a *partial* function, the latter property is equivalent to the claim made in part (c) for the exchange $\langle P, Q \rangle$ of the theorem statement. Furthermore, this same exchange although achievable by an IR $M(k-1)$ -contract path can be so realised only by one whose length is as given in part (b) of the theorem statement.

11. These are termed “qualitative coalitional games” (QCGs) in [27]: the approach has some features in common with, but differs from, the concept of “effectivity function” as discussed in [2].

Regarding the proof itself, there are a number of notational complexities which we have attempted to ameliorate by making some simplifying assumptions concerning the relationship between m – the size of the resource set \mathcal{R} – and k – the number of agents which are needed to realise $\langle P, Q \rangle$ in a *single* IR exchange. In particular, we shall assume that m is an exact multiple of $\binom{k}{2}$. We observe that by employing a similar device to that used in the proof of Theorem 4 we can deal with cases for which m does not have this property: if $m = s \binom{k}{2} + q$ for integer values $s \geq 1$ and $1 \leq q < \binom{k}{2}$, we simply employ exactly the same construction using $m - q$ resources with the “missing” q resources from \mathcal{R}_m being allocated to A_1 and never being reallocated within the $M(k - 1)$ -contract path. This approach accounts for the rounding operation ($\lfloor \dots \rfloor$) in the exponent term of the lower bound. We shall also assume that the number of agents in \mathcal{A} is exactly k . Within the proof we use a running example for which $k = 4$ and $m = 18 = 3 \times 6$ to illustrate specific features.

We first give an outline of its structure.

Given $\langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle$ a resource allocation setting involving k agents and m resources, our aim is to define an IR $M(k - 1)$ -contract path

$$\Delta = \langle P^{(1)}, P^{(2)}, \dots, P^{(t)} \rangle$$

that realises the IR $M(k)$ exchange $\langle P^{(1)}, P^{(t)} \rangle$. We will use d to index particular allocations within Δ , so that $1 \leq d \leq t$.

In order to simplify the presentation we employ a setting in which the k agents are $\mathcal{A} = \{A_0, A_1, \dots, A_{k-1}\}$. Recalling that $m = s \binom{k}{2}$, the resource set \mathcal{R}_m is formed by the union of $\binom{k}{2}$ pairwise disjoint sets of size s . Given distinct values i and j with $0 \leq i < j \leq k - 1$, we use $\mathcal{R}^{i,j}$ to denote one of these subsets with $\{r_1^{\{i,j\}}, r_2^{\{i,j\}}, \dots, r_s^{\{i,j\}}\}$ the s resources that form $\mathcal{R}^{\{i,j\}}$.

There are two main ideas underpinning the structure of each $M(k - 1)$ -contract in Δ .

Firstly, in the initial and subsequent allocations, the resource set $\mathcal{R}^{\{i,j\}}$ is partitioned between A_i and A_j and any reallocation of resources between A_i and A_j that takes place within the exchange $\langle P^{(d)}, P^{(d+1)} \rangle$ will involve *only* resources in this set. Thus, for every allocation $P^{(d)}$ and each pair $\{i, j\}$, if $h \notin \{i, j\}$ then $P_h^{(d)} \cap \mathcal{R}^{\{i,j\}} = \emptyset$. Furthermore, for $\delta = \langle P^{(d)}, P^{(d+1)} \rangle$ should *both* A_i and A_j be involved, i.e. $\{A_i, A_j\} \subseteq \mathcal{A}^\delta$, then this reallocation of $\mathcal{R}^{\{i,j\}}$ between A_i and A_j will be an O -contract. That is, either exactly one element of $\mathcal{R}^{\{i,j\}}$ will be moved from $P_i^{(d)}$ to become a member of the allocation $P_j^{(d+1)}$ or exactly one element of $\mathcal{R}^{\{i,j\}}$ will be moved from $P_j^{(d)}$ to become a member of the allocation $P_i^{(d+1)}$. In total, every $M(k - 1)$ -contract δ in Δ consists of a *simultaneous* implementation of $\binom{k-1}{2}$ O -contracts: a single O -contract for each of the distinct pairs $\{A_i, A_j\}$ of agents from the $k - 1$ agents in \mathcal{A}^δ .

The second key idea is to exploit one well-known property of the s -dimensional hypercube network: for every $s \geq 2$, \mathcal{H}_s contains a *Hamiltonian cycle*, i.e. a simple directed cycle

formed using only the edges of \mathcal{H}_s and containing all 2^s vertices¹². Now, suppose

$$\mathcal{S}^{(v)} = \underline{v}^{(0)}, \underline{v}^{(1)}, \dots, \underline{v}^{(i)}, \dots, \underline{v}^{(2^s-1)}, \underline{v}^{(0)}$$

is a Hamiltonian cycle in the hypercube \mathcal{H}_s and

$$\mathcal{S}^{(w)} = \underline{w}^{(0)}, \underline{w}^{(1)}, \dots, \underline{w}^{(i)}, \dots, \underline{w}^{(2^s-1)}, \underline{w}^{(0)}$$

the Hamiltonian cycle in which $\underline{w}^{(i)}$ is obtained by complementing each bit in $\underline{v}^{(i)}$. As we have described in the overview of Section 2.1 we can interpret the s -bit label $\underline{v} = v_1 v_2 \dots v_s$ as describing a particular subset of $\mathcal{R}^{\{i,j\}}$, i.e. that subset in which $r_k^{\{i,j\}}$ occurs if and only if $v_k = 1$. Similarly from any subset of $\mathcal{R}^{\{i,j\}}$ we may define a unique s -bit word. Now suppose that $P_i^{(d)}$ is the allocation held by A_i in the allocation $P^{(d)}$ of Δ . The exchange $\delta = \langle P^{(d)}, P^{(d+1)} \rangle$ will affect $P_i^{(d)} \cap \mathcal{R}^{\{i,j\}}$ in the following way: if $i \notin \mathcal{A}^\delta$ or $j \notin \mathcal{A}^\delta$ then $P_i^{(d+1)} \cap \mathcal{R}^{\{i,j\}} = P_i^{(d)} \cap \mathcal{R}^{\{i,j\}}$ and $P_j^{(d+1)} \cap \mathcal{R}^{\{i,j\}} = P_j^{(d)} \cap \mathcal{R}^{\{i,j\}}$. Otherwise we have $\{i, j\} \subseteq \mathcal{A}^\delta$ and the (complementary) holdings $P_i^{(d)} \cap \mathcal{R}^{\{i,j\}}$ and $P_j^{(d)} \cap \mathcal{R}^{\{i,j\}}$ define (complementary) s -bit labels of vertices in \mathcal{H}_s : if these correspond to places $\langle \underline{v}^{(h)}, \underline{w}^{(h)} \rangle$ in the Hamiltonian cycle, then in $P_i^{(d+1)}$ and $P_j^{(d+1)}$ the s -bit labels defined from $P_i^{(d+1)} \cap \mathcal{R}^{\{i,j\}}$ and $P_j^{(d+1)} \cap \mathcal{R}^{\{i,j\}}$ produce the s -bit labels $\underline{v}^{(h+1)}$ and $\underline{w}^{(h+1)}$, i.e. the vertices that succeed $\underline{v}^{(h)}$ and $\underline{w}^{(h)}$ in the Hamiltonian cycles. In total, for each j , A_i initially holds either the subset of $\mathcal{R}^{\{i,j\}}$ that maps to $\underline{v}^{(0)}$ or that maps to $\underline{w}^{(0)}$ and, at the conclusion of the $M(k-1)$ -path, holds the subset that maps to $\underline{v}^{(2^s-1)}$ (or $\underline{w}^{(2^s-1)}$). The final detail is that the progression through the Hamiltonian cycles is conducted over a series of *rounds* each round comprising k $M(k-1)$ -exchanges.

We have noted that each $M(k-1)$ -contract, $\langle P^{(d)}, P^{(d+1)} \rangle$ that occurs in this path Δ can be interpreted as a set of $\binom{k-1}{2}$ distinct O -contracts. An important property of the utility functions employed is that unless $p \geq k-1$ there will be *no individually rational* $M(p)$ -contract path that realises the exchange $\langle P^{(d)}, P^{(d+1)} \rangle$, i.e. the $\binom{k-1}{2}$ O -contract exchanges must occur *simultaneously* in order for the progression from $P^{(d)}$ to $P^{(d+1)}$ to be IR. Although the required exchange could be realised by a sequence of O -contracts (or, more generally, any suitable $M(k-2)$ -contract path), such realisations will *not* describe an IR contract path. The construction of utility functions to guarantee such behaviour provides the principal component in showing that the IR exchange $\langle P^{(1)}, P^{(t)} \rangle$ cannot be realised with an IR $M(k-2)$ -contract path: if Q is *any* allocation for which $\langle P^{(1)}, Q \rangle$ is an $M(k-2)$ -contract then $\langle P^{(1)}, Q \rangle$ is not IR.

We now proceed with the proof of Theorem 6.

Proof. (of Theorem 6) Fix $\mathcal{A} = \{A_0, A_1, \dots, A_{k-1}\}$. \mathcal{R} consists of $\binom{k}{2}$ pairwise disjoint sets of s resources

$$\mathcal{R}^{\{i,j\}} = \{r_1^{\{i,j\}}, r_2^{\{i,j\}}, \dots, r_s^{\{i,j\}}\}$$

12. This can be shown by an easy inductive argument. For $s = 2$, the sequence $\langle 00, 01, 11, 10, 00 \rangle$ defines a Hamiltonian cycle in \mathcal{H}_2 . Inductively assume that $\langle \alpha_1, \alpha_2, \dots, \alpha_p, \alpha_1 \rangle$ (with $p = 2^s$) is such a cycle in \mathcal{H}_s then $\langle 0\alpha_1, 1\alpha_1, 1\alpha_p, 1\alpha_{p-1}, \dots, 1\alpha_2, 0\alpha_2, \dots, 0\alpha_p, 0\alpha_1 \rangle$ defines a Hamiltonian cycle in \mathcal{H}_{s+1} .

For $k = 4$ and $s = 3$ these yield $\mathcal{A} = \{A_0, A_1, A_2, A_3\}$ and

$$\begin{aligned}\mathcal{R}^{\{0,1\}} &= \{r_1^{\{0,1\}}, r_2^{\{0,1\}}, r_3^{\{0,1\}}\} \\ \mathcal{R}^{\{0,2\}} &= \{r_1^{\{0,2\}}, r_2^{\{0,2\}}, r_3^{\{0,2\}}\} \\ \mathcal{R}^{\{0,3\}} &= \{r_1^{\{0,3\}}, r_2^{\{0,3\}}, r_3^{\{0,3\}}\} \\ \mathcal{R}^{\{1,2\}} &= \{r_1^{\{1,2\}}, r_2^{\{1,2\}}, r_3^{\{1,2\}}\} \\ \mathcal{R}^{\{1,3\}} &= \{r_1^{\{1,3\}}, r_2^{\{1,3\}}, r_3^{\{1,3\}}\} \\ \mathcal{R}^{\{2,3\}} &= \{r_1^{\{2,3\}}, r_2^{\{2,3\}}, r_3^{\{2,3\}}\}\end{aligned}$$

We use two ordering structures in defining the $M(k-1)$ -contract path.

a.

$$\mathcal{S}^{(v)} = \underline{v}^{(0)}, \underline{v}^{(1)}, \dots, \underline{v}^{(i)}, \dots, \underline{v}^{(2^s-1)}, \underline{v}^{(0)}$$

a Hamiltonian cycle in \mathcal{H}_s , where without loss of generality, $\underline{v}^{(0)} = 111 \dots 11$.

b.

$$\mathcal{S}^{(w)} = \underline{w}^{(0)}, \underline{w}^{(1)}, \dots, \underline{w}^{(i)}, \dots, \underline{w}^{(2^s-1)}, \underline{w}^{(0)}$$

the complementary Hamiltonian cycle to this, so that $\underline{w}^{(0)} = 000 \dots 00$.

Thus for $k = 4$ and $s = 3$ we obtain

$$\begin{aligned}\text{a. } \mathcal{S}^{(v)} &= \langle 111, 110, 010, 011, 001, 000, 100, 101 \rangle \\ \text{b. } \mathcal{S}^{(w)} &= \langle 000, 001, 101, 100, 110, 111, 011, 010 \rangle\end{aligned}$$

We can now describe the $M(k-1)$ -contract path.

$$\Delta = \langle P^{(1)}, P^{(2)}, \dots, P^{(t)} \rangle$$

Initial Allocation: $P^{(1)}$.

Define the $k \times k$ Boolean matrix, $B = [b_{i,j}]$ (with $0 \leq i, j \leq k-1$) by

$$b_{i,j} = \begin{cases} \perp & \text{if } i = j \\ \neg b_{j,i} & \text{if } i > j \\ \neg b_{i,j-1} & \text{if } i < j \end{cases}$$

We then have for each $1 \leq i \leq k$,

$$P_i^{(1)} = \bigcup_{j=0}^{i-1} \{R^{\{j,i\}} : b_{i,j} = \top\} \cup \bigcup_{j=i+1}^{k-1} \{R^{\{i,j\}} : b_{i,j} = \top\}$$

Thus, in our example,

$$B = \begin{bmatrix} \perp & \top & \perp & \top \\ \perp & \perp & \top & \perp \\ \top & \perp & \perp & \top \\ \perp & \top & \perp & \top \end{bmatrix}$$

Yielding the starting allocation

$$\begin{aligned}
P_0^{(1)} &= \mathcal{R}^{\{0,1\}} \cup \mathcal{R}^{\{0,3\}} &= \langle 111, 000, 111 \rangle &\subseteq \mathcal{R}^{\{0,1\}} \cup \mathcal{R}^{\{0,2\}} \cup \mathcal{R}^{\{0,3\}} \\
P_1^{(1)} &= \mathcal{R}^{\{1,2\}} &= \langle 000, 111, 000 \rangle &\subseteq \mathcal{R}^{\{0,1\}} \cup \mathcal{R}^{\{1,2\}} \cup \mathcal{R}^{\{1,3\}} \\
P_2^{(1)} &= \mathcal{R}^{\{0,2\}} \cup \mathcal{R}^{\{2,3\}} &= \langle 111, 000, 111 \rangle &\subseteq \mathcal{R}^{\{0,2\}} \cup \mathcal{R}^{\{1,2\}} \cup \mathcal{R}^{\{2,3\}} \\
P_3^{(1)} &= \mathcal{R}^{\{1,3\}} &= \langle 000, 111, 000 \rangle &\subseteq \mathcal{R}^{\{0,3\}} \cup \mathcal{R}^{\{1,3\}} \cup \mathcal{R}^{\{2,3\}}
\end{aligned}$$

The third column in $P_i^{(1)}$ indicating the 3-bit labels characterising each of the subsets of $\mathcal{R}^{\{i,j\}}$ for the three values that j can assume.

Rounds: The initial allocation is changed over a series of *rounds*

$$Q^1, Q^2, \dots, Q^z$$

each of which involves exactly k distinct $M(k-1)$ -contracts. We use $Q^{x,p}$ to indicate the allocation resulting after stage p in round x where $0 \leq p \leq k-1$. We note the following:

- The initial allocation, $P^{(1)}$ will be denoted by $Q^{0,k-1}$.
- $Q^{x,0}$ is obtained using a single $M(k-1)$ -contract from $Q^{x-1,k-1}$ (when $x \geq 1$).
- $Q^{x,p}$ is obtained from using a single $M(k-1)$ -contract $Q^{x,p-1}$ (when $0 < p \leq k-1$).

Our final item of notation is that of the *cube position of i with respect to j in an allocation P* , denoted $\chi(i, j, P)$. Letting \underline{u} be the s -bit string describing $P_i \cap \mathcal{R}^{\{i,j\}}$ in some allocation P , $\chi(i, j, P)$ is the *index* of \underline{u} in the Hamiltonian cycle $S^{(v)}$ (when $\mathcal{R}^{\{i,j\}} \subseteq P_i^{(1)}$) or the Hamiltonian cycle $S^{(w)}$ (when $\mathcal{R}^{\{i,j\}} \subseteq P_j^{(1)}$). When $P = Q^{x,p}$ for some allocation in the sequence under construction we employ the notation $\chi(i, j, x, p)$, noting that one invariant of our path will be $\chi(i, j, x, p) = \chi(j, i, x, p)$, a property that certainly holds true of $P^{(1)} = Q^{0,k-1}$ since $\chi(i, j, 0, k-1) = \chi(j, i, 0, k-1) = 0$.

The sequence of allocations in Δ is built as follows. Since $Q^{1,0}$ is the immediate successor of the initial allocation $Q^{0,k-1}$, it suffices to describe how $Q^{x,p}$ is formed from $Q^{x,p-1}$ (when $p > 0$) and $Q^{x+1,0}$ from $Q^{x,k-1}$. Let $Q^{y,q}$ be the allocation to be formed from $Q^{x,p}$. The exchange $\delta = \langle Q^{x,p}, Q^{y,q} \rangle$ will be an $M(k-1)$ contract in which $\mathcal{A}^\delta = \mathcal{A} \setminus \{A_q\}$. For each pair $\{i, j\} \subseteq \mathcal{A}^\delta$ we have $\chi(i, j, x, p) = \chi(j, i, x, p)$ in the allocation $Q^{x,p}$. In moving to $Q^{y,q}$ exactly one element of $\mathcal{R}^{\{i,j\}}$ is reallocated between A_i and A_j in such a way that in $Q^{y,q}$, $\chi(i, j, y, q) = \chi(i, j, x, p) + 1$, since A_i and A_j are tracing complementary Hamiltonian cycles with respect to $\mathcal{R}^{\{i,j\}}$ this ensures that $\chi(j, i, y, q) = \chi(j, i, x, p) + 1$, thereby maintaining the invariant property.

Noting that for each distinct pair $\langle i, j \rangle$, we either have $\mathcal{R}^{\{i,j\}}$ allocated to A_i in $P^{(1)}$ or $\mathcal{R}^{\{i,j\}}$ allocated to A_j in $P^{(1)}$, the description just outlined indicates that the allocation $P^{(d)} = Q^{x,p}$ is completely specified as follows.

The cube position, $\chi(i, j, x, p)$, satisfies,

$$\chi(i, j, x, p) = \begin{cases} 0 & \text{if } x = 0 \text{ and } p = k-1 \\ 1 + \chi(i, j, x-1, k-1) & \text{if } x \geq 1, p = 0, \text{ and } p \notin \{i, j\} \\ \chi(i, j, x-1, k-1) & \text{if } x \geq 1, p = 0, \text{ and } p \in \{i, j\} \\ 1 + \chi(i, j, x, p-1) & \text{if } 1 \leq p \leq k-1, \text{ and } p \notin \{i, j\} \\ \chi(i, j, x, p-1) & \text{if } 1 \leq p \leq k-1, \text{ and } p \in \{i, j\} \end{cases}$$

For each i , the subset of $\mathcal{R}^{\{i,j\}}$ that is held by A_i in the allocation $Q^{x,p}$ is,

$$\begin{aligned} \underline{v}(\chi(i,j,x,p)) & \text{ if } \mathcal{R}^{\{i,j\}} \subseteq P_i^{(1)} \\ \underline{w}(\chi(i,j,x,p)) & \text{ if } \mathcal{R}^{\{i,j\}} \subseteq P_j^{(1)} \end{aligned}$$

(where we recall that s -bit labels in the hypercube \mathcal{H}_s are identified with subsets of $\mathcal{R}^{\{i,j\}}$.)

The tables below illustrates this process for our example.

d	x	p	A_0			A_1			A_2			A_3			$\mathcal{A}^{(P^{(d-1)}, P^{(d)})}$
			$i j$	$i j$	$i j$	$i j$	$i j$	$i j$	$i j$	$i j$	$i j$	$i j$	$i j$	$i j$	
1	0	3	111	000	111	000	111	000	111	000	111	000	111	000	–
2	1	0	111	000	111	000	110	001	111	001	110	000	110	001	$\{A_1, A_2, A_3\}$
3	1	1	111	001	110	000	110	001	110	001	010	001	110	101	$\{A_0, A_2, A_3\}$
4	1	2	110	001	010	001	110	101	110	001	010	101	010	101	$\{A_0, A_1, A_3\}$
5	1	3	010	101	010	101	010	101	010	101	010	101	010	101	$\{A_0, A_1, A_2\}$
6	2	0	010	101	011	101	011	100	010	100	011	101	011	100	$\{A_1, A_2, A_3\}$
7	2	1	010	100	001	101	011	100	011	100	001	100	011	110	$\{A_0, A_2, A_3\}$
8	2	2	011	100	001	100	011	110	011	100	001	110	001	110	$\{A_0, A_1, A_3\}$
9	2	3	001	110	001	110	001	110	001	110	001	110	001	110	$\{A_0, A_1, A_2\}$
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots

Subsets of $\mathcal{R}^{\{i,j\}}$ held by A_i in $Q^{x,p}$ ($k = 4, s = 3$)

d	x	p	A_0			A_1			A_2			A_3			$\mathcal{A}^{(P^{(d-1)}, P^{(d)})}$
			$i j$	$i j$	$i j$	$i j$	$i j$	$i j$	$i j$	$i j$	$i j$	$i j$	$i j$		
1	0	3	0	0	0	0	0	0	0	0	0	0	0	0	–
2	1	0	0	0	0	0	1	1	0	1	1	0	1	1	$\{A_1, A_2, A_3\}$
3	1	1	0	1	1	0	1	1	1	1	2	1	1	2	$\{A_0, A_2, A_3\}$
4	1	2	1	1	2	1	1	2	1	1	2	2	2	2	$\{A_0, A_1, A_3\}$
5	1	3	2	2	2	2	2	2	2	2	2	2	2	2	$\{A_0, A_1, A_2\}$
6	2	0	2	2	2	2	3	3	2	3	3	2	3	3	$\{A_1, A_2, A_3\}$
7	2	1	2	3	3	2	3	3	3	3	4	3	3	4	$\{A_0, A_2, A_3\}$
8	2	2	3	3	4	3	3	4	3	3	4	4	4	4	$\{A_0, A_1, A_3\}$
9	2	3	4	4	4	4	4	4	4	4	4	4	4	4	$\{A_0, A_1, A_2\}$
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\vdots

Cube Positions $\chi(i, j, x, p)$ ($k = 4, s = 3$)

It is certainly the case that this process of applying successive rounds of k exchanges could be continued, however, we wish to do this only so long as it is not possible to go from some allocation $P^{(d)}$ in the sequence to another $P^{(d+r)}$ for some $r \geq 2$ via an $M(k-1)$ -contract.

Now if $Q^{x,p}$ and $Q^{y,q}$ are distinct allocations generated by the process above then the exchange $\delta = \langle Q^{x,p}, Q^{y,q} \rangle$ is an $M(k-1)$ -contract if and only if for some A_i , $Q_i^{x,p} = Q_i^{y,q}$. It follows that if $\langle P^{(d)}, P^{(d+r)} \rangle$ is an $M(k-1)$ -contract for some $r > 1$, then for some i and all $j \neq i$, $P_i^{(d+r)} \cap \mathcal{R}^{\{i,j\}} = P_i^{(d)} \cap \mathcal{R}^{\{i,j\}}$.

To determine the minimum value of $r > 1$ with which $P_i^{(d+r)} = P_i^{(d)}$, we observe that without loss of generality we need consider only the case $d = i = 0$, i.e. we determine the minimum number of exchanges before $P_0^{(1)}$ reappears. First note that in each round, Q^x , if $\chi(0, j, x-1, k-1) = p$ then $\chi(0, j, x, k-1) = p+k-2$, i.e. each round advances the cube position $k-2$ places: $\chi(0, j, x-1, k-1) = \chi(0, j, x, 0)$ and $\chi(0, j, x, j) = \chi(0, j, x, j-1)$. We can also observe that $P_0^{(1)} = Q_0^{0, k-1} \neq Q_0^{x, p}$ for any p with $0 < p < k-1$, since

$$\chi(0, 1, x, p) = \chi(0, 2, x, p) = \dots = \chi(0, k-1, x, p)$$

only in the cases $p = 0$ and $p = k-1$. It follows that our value $r > 1$ must be of the form qk where q must be such that $q(k-2)$ is an *exact multiple* of 2^s . From this observation we see that,

$$\min\{ r > 1 : P_0^{(1)} = P_0^{(1+r)} \} = \min\{ qk : q(k-2) \text{ is a multiple of } 2^s \}$$

Now, if k is *odd* then $q = 2^s$ is the minimal such value, so that $r = k2^s$. If k is even then it may be uniquely written in the form $z2^l + 2$ where z is *odd* so giving q as 1 (if $l \geq s$) or 2^{s-l} (if $l \leq s$), so that these give $r = k$ and $r = z2^s + 2^{s-l+1}$, e.g. for $k = 4$ and $s = 3$, we get $k = 1 \times 2^1 + 2$ so that $r = 2^3 + 2^{3-1+1} = 16$ and in our example $P_0^{(1)} = P_0^{(17)}$ may be easily verified. In total,

$$r \geq \begin{cases} k2^s & \text{if } k \text{ is odd} \\ k & \text{if } k = z2^l + 2, z \text{ is odd, and } l \geq s \\ 2^s & \text{if } k = z2^l + 2, z \text{ is odd and } l \leq s \end{cases}$$

All of which immediately give $r \geq 2^s$ (in the second case $k \geq 2^s$, so the inequality holds trivially), and thus we can continue the chain of $M(k-1)$ contracts for at least 2^s moves. Recalling that $m = s \binom{k}{2}$, this gives the length of the $M(k-1)$ -contract path

$$\Delta = \langle P^{(1)}, P^{(2)}, \dots, P^{(t)} \rangle$$

written in terms of m and k as at least¹³

$$2^{m / \binom{k}{2}} - 1 = 2^{\frac{2m}{k(k-1)}} - 1$$

It remains to define appropriate utility functions $\mathcal{U} = \langle u_0, \dots, u_{k-1} \rangle$ in order to ensure that Δ is the unique IR $M(k-1)$ -contract path realising the IR $M(k)$ -exchange $\langle P^{(1)}, P^{(t)} \rangle$. In defining \mathcal{U} it will be convenient to denote Δ as the path

$$\Delta = \langle Q^{0, k-1}, Q^{1, 0}, Q^{1, 1}, \dots, Q^{1, k-1}, \dots, Q^{x, p}, \dots, Q^{r, k-1} \rangle$$

and, since $rk \geq 2^s$, we may without loss of generality, focus on the first 2^s allocations in this contract path.

Recalling that $\chi(i, j, x, p)$ is the index of the s -bit label \underline{u} corresponding to $Q_i^{x, p} \cap \mathcal{R}^{\{i, j\}}$ in the relevant Hamiltonian cycle – i.e. $\mathcal{S}^{(v)}$ if $\mathcal{R}^{\{i, j\}} \subseteq Q_i^{0, k}$, $\mathcal{S}^{(w)}$ if $\mathcal{R}^{\{i, j\}} \subseteq Q_j^{0, k-1}$ – we note the following properties of the sequence of allocations defined by Δ that hold for each distinct i and j .

13. We omit the rounding operation $\lceil \dots \rceil$ in the exponent, which is significant only if m is not an exact multiple of $\binom{k}{2}$, in which event the device described in our overview of the proof is applied.

P1. $\forall x, p \chi(i, j, x, p) = \chi(j, i, x, p)$

P2. If $Q^{y,q}$ is the immediate successor of $Q^{x,p}$ in Δ then $\chi(i, j, y, q) \leq \chi(i, j, x, p) + 1$ with equality if and only if $q \notin \{i, j\}$.

P3. $\forall i', j'$ with $0 \leq i', j' \leq k - 1$, $\chi(i, j, x, k - 1) = \chi(i', j', x, k - 1)$.

The first two properties have already been established in our description of Δ . The third follows from the observation that within each round Q^x , each cube position is advanced by exactly $k - 2$ in progressing from $Q^{x-1,0}$ to $Q^{x,k-1}$.

The utility function u_i is now given, for $S \subseteq \mathcal{R}_m$, by

$$u_i(S) = \begin{cases} \sum_{j \neq i} \chi(i, j, x, p) & \text{if } S = Q_i^{x,p} \text{ for some } 0 \leq x \leq r, 0 \leq p \leq k - 1 \\ -2^{km} & \text{otherwise} \end{cases}$$

We claim that, with these choices,

$$\Delta = \langle Q^{0,k-1}, Q^{1,0}, Q^{1,1}, \dots, Q^{1,k-1}, \dots, Q^{x,p}, \dots, Q^{r,k-1} \rangle$$

is the unique IR $M(k - 1)$ -contract path realising the IR $M(k)$ -exchange $\langle Q^{0,k-1}, Q^{r,k-1} \rangle$. Certainly, Δ is an IR $M(k - 1)$ -contract path: each exchange $\delta = \langle Q^{x,p}, Q^{y,q} \rangle$ on this path has $|\mathcal{A}^\delta| = k - 1$ and since for each agent A_i in $\mathcal{A}^\delta = \mathcal{A} \setminus \{A_q\}$ the utility of $Q_i^{y,q}$ has increased by exactly $k - 2$, i.e. each cube position of i with respect to j whenever $q \notin \{i, j\}$ has increased, it follows that $\sigma_u(Q^{y,q}) > \sigma_u(Q^{x,p})$ and hence $\langle Q^{x,p}, Q^{y,q} \rangle$ is IR.

We now show that Δ is the *unique* IR $M(k - 1)$ -contract path continuation of $Q^{0,k-1}$. Suppose $\delta = \langle Q^{x,p}, P \rangle$ is an exchange that deviates from the contract path Δ (having followed it through to the allocation $Q^{x,p}$). Certainly both of the following must hold of P : for each i , $P_i \subseteq \cup_{j \neq i} \mathcal{R}^{\{i,j\}}$; and there is a k -tuple of pairs $\langle (x_0, p_0), \dots, (x_{k-1}, p_{k-1}) \rangle$ with which $P_i = Q_i^{x_i, p_i}$, for if either fail to be the case for some i , then $u_i(P_i) = -2^{km}$ with the consequent effect that $\sigma_u(P) < 0$ and thence not IR. Now, if $Q^{y,q}$ is the allocation that would succeed $Q^{x,p}$ in Δ then $P \neq Q^{y,q}$, and thus for at least one agent, $Q_i^{x_i, p_i} \neq Q_i^{y,q}$. It cannot be the case that $Q_i^{x_i, p_i}$ corresponds to an allocation occurring *strictly later* than $Q_i^{y,q}$ in Δ since such allocations could not be realised by an $M(k - 1)$ -contract. In addition, since $P_i = Q_i^{x_i, p_i}$ it must be the case that $|\mathcal{A}^\delta| = k - 1$ since exactly $k - 1$ cube positions in the holding of A_i must change. It follows that there are only two possibilities for (y_i, p_i) : P_i reverts to the allocation immediately preceding $Q_i^{x_i, p_i}$ or advances to the holding $Q_i^{y,q}$. It now suffices to observe that an exchange in which some agents satisfy the first of these while the remainder proceed in accordance with the second either does not give rise to a valid allocation or cannot be realised by an $M(k - 1)$ -contract. On the other hand if P corresponds to the allocation preceding $Q^{x,p}$ then δ is not IR. We deduce, therefore, that the *only* IR $M(k - 1)$ exchange that is consistent with $Q^{x,p}$ is that prescribed by $Q^{y,q}$.

This completes the analysis needed for the proof of part (b) of the theorem. It is clear that since the system contains only k agents, any exchange $\langle P, Q \rangle$ can be effected with a single $M(k)$ -contract, thereby establishing part (a). For part (c) – that the IR exchange $\langle P^{(1)}, P^{(t)} \rangle$ cannot be realised using an *individually rational* $M(k - 2)$ -contract path, it suffices to observe that since the class of IR $M(k - 2)$ -contracts are a subset of the class of IR $M(k - 1)$ -contracts, were it the case that an IR $M(k - 2)$ -contract path existed to

implement $\langle P^{(1)}, P^{(t)} \rangle$, this would imply that Δ was not the *unique* IR $M(k-1)$ -contract path. We have, however, proved that Δ is unique, and part (c) of the theorem follows. \square

We obtain a similar development of Corollary 1 in

Corollary 3 For all $k \geq 3$, $n \geq k$, $m \geq \binom{k}{2}$ and each of the cases below,

- a. $\Phi_k(\delta)$ holds if and only if δ is a cooperatively rational $M(k)$ -contract.
 $\Psi(\delta)$ holds if and only if δ is cooperatively rational.
- b. $\Phi_k(\delta)$ holds if and only if δ is an equitable $M(k)$ -contract.
 $\Psi(\delta)$ holds if and only if δ is equitable.

there is a resource allocation setting $\langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle$ and a Ψ -exchange $\delta = \langle P, Q \rangle$ for which

$$\begin{aligned} L^{\text{opt}}(\delta, \langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle, \Phi_k) &= 1 & (a) \\ L^{\text{opt}}(\delta, \langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle, \Phi_{k-1}) &\geq 2^{\lfloor 2m/k(k-1) \rfloor} - 1 & (b) \\ L^{\text{opt}}(\delta, \langle \mathcal{A}, \mathcal{R}, \mathcal{U} \rangle, \Phi_{k-2}) &\text{is undefined} & (c) \end{aligned}$$

Proof. As with the proof of Corollary 1 in relation to Theorem 3, in each case we employ the contract path from the proof of Theorem 6, varying the definition of $\mathcal{U} = \langle u_1, u_2, \dots, u_k \rangle$ in order to establish each result. Thus let

$$\begin{aligned} \Delta_m &= \langle P^{(1)}, P^{(2)}, \dots, P^{(r)}, \dots, P^{(t)} \rangle \\ &= \langle Q^{0,k-1}, Q^{1,0}, \dots, Q^{x,p}, \dots, Q^{z,r} \rangle \end{aligned}$$

be the $M(k-1)$ -contract path realising the $M(k)$ -exchange $\langle P^{(1)}, P^{(t)} \rangle$ described in the proof of Theorem 6, this path having length $t \geq 2^{\lfloor 2m/k(k-1) \rfloor} - 1$.

- a. The utility functions $\mathcal{U} = \langle u_0, \dots, u_{k-1} \rangle$ of Theorem 6 ensure that $\langle P^{(1)}, P^{(t)} \rangle$ is cooperatively rational and that Δ_m is a cooperatively rational $M(k-1)$ -contract path realising $\langle P^{(1)}, P^{(t)} \rangle$: the utility held by A_i never decreases in value and there is at least one agent (in fact exactly $k-1$) whose utility increases in value. Furthermore Δ_m is the unique cooperatively rational $M(k-1)$ -contract path realising $\langle P^{(1)}, P^{(t)} \rangle$ since, by the same argument used in Theorem 6, any deviation will result in some agent suffering a loss of utility.
- b. Set the utility functions $\mathcal{U} = \langle u_0, \dots, u_{k-1} \rangle$ as,

$$u_i(S) = \begin{cases} -1 & \text{if } S \neq Q_i^{x,p} \text{ for any } Q^{x,p} \in \Delta_m \\ xk^2 + k - i & \text{if } S = Q_i^{x,k-1} \\ (x-1)k^2 + k + p & \text{if } S = Q_0^{x,p}, p < k-1 \text{ and } i = 0 \\ (x-1)k^2 + k - i + p + 1 & \text{if } S = Q_i^{x,p}, p < i-1 \text{ and } i \neq 0. \\ xk^2 + 1 & \text{if } S = Q_i^{x,i-1} = Q_i^{x,i} \text{ and } i \neq 0. \\ xk^2 + 1 + p - i & \text{if } S = Q_i^{x,p}, p > i \text{ and } i \neq 0 \end{cases}$$

To see that these choices admit Δ_m as an equitable $M(k-1)$ -contract path realising the equitable exchange $\langle Q^{0,k-1}, Q^{z,r} \rangle$, we first note that

$$\min_{0 \leq i \leq k-1} \{u_i(Q_i^{z,r})\} > 1 = \min_{0 \leq i \leq k-1} \{u_i(Q_i^{0,k-1})\}$$

thus, $\langle Q^{0,k-1}, Q^{z,r} \rangle$ is indeed equitable. Consider any exchange $\delta = \langle Q^{x,p}, Q^{y,q} \rangle$ occurring within Δ_m . It suffices to show that

$$\min_{0 \leq i \leq k-1} \{u_i(Q_i^{x,p})\} \neq u_q(Q_q^{x,p})$$

since $A_q \notin \mathcal{A}^\delta$, and for all other agents $u_i(Q_i^{y,q}) > u_i(Q_i^{x,p})$. We have two possibilities: $q = 0$ (in which case $p = k - 1$ and $y = x + 1$); $q > 0$ (in which case $p = q - 1$). Consider the first of these: $u_0(Q_0^{x,k-1}) = xk^2 + k$, however,

$$\min\{u_i(Q_i^{x,k-1})\} = xk^2 + 1 = u_{k-1}(Q_{k-1}^{x,k-1})$$

and hence every exchange $\langle Q^{x,k-1}, Q^{x+1,0} \rangle$ forming part of Δ_m is equitable.

In the remaining case, $u_q(Q_q^{x,q-1}) = xk^2 + 1$ and

$$\begin{aligned} \min\{u_i(Q_i^{x,q-1})\} &\leq u_0(Q_0^{x,q-1}) \\ &= (x-1)k^2 + k + q - 1 \\ &< xk^2 - (k^2 - 2k + 1) \\ &= xk^2 - (k-1)^2 \\ &< xk^2 + 1 \\ &= u_q(Q_q^{x,q-1}) \end{aligned}$$

and thus the remaining exchanges $\langle Q^{x,q-1}, Q^{x,q} \rangle$ within Δ_m are equitable. By a similar argument to that employed in Theorem 6 it follows that Δ_m is the unique equitable $M(k-1)$ -contract path realising $\langle Q^{0,k-1}, Q^{z,r} \rangle$.

□

Monotone Utility Functions and $M(k)$ -contract paths

The device used to develop Theorem 3 to obtain the path of Theorem 4 can be applied to the rather more intricate construction of Theorem 6, thereby allowing exponential lower bounds on $\rho_{\text{mono}}^{\max}(n, m, \Phi_k, \Psi)$ to be derived. We will merely outline the approach rather than present a detailed technical exposition. We recall that it became relatively straightforward to define suitable monotone utility functions once it was ensured that the subset sizes of interest – i.e. those for allocations arising in the O -contract path – were forced to fall into a quite restricted range. The main difficulty that arises in applying similar methods to the path Δ of Theorem 6 is the following: in the proof of Theorem 4 we consider two agents so that converting Δ_s from a setting with s resources in Theorem 3 to $\text{ext}(\Delta_s)$ with $2s$ resources in Theorem 4 is achieved by combining “complementary” allocations, i.e. $\alpha \subseteq \mathcal{R}_s$ with $\bar{\alpha} \subseteq \mathcal{T}_s$. We can exploit two facts, however, to develop a path $\text{multi}(\Delta)$ for which monotone utility functions could be defined: the resource set \mathcal{R}_m in Theorem 6 consists of $\binom{k}{2}$ disjoint sets of size s ; and any exchange δ on the path Δ involves a reallocation of $\mathcal{R}^{\{i,j\}}$ between A_i and A_j when $\{i,j\} \subseteq \mathcal{A}^\delta$. Thus letting \mathcal{T}_m be formed by $\binom{k}{2}$ disjoint sets, $\mathcal{T}^{\{i,j\}}$ each of size s . Suppose that $P_i^{(d)}$ is described by

$$\alpha_{i,0}^{(d)} \alpha_{i,1}^{(d)} \cdots \alpha_{i,i-1}^{(d)} \alpha_{i,i+1}^{(d)} \cdots \alpha_{i,k-1}^{(d)}$$

with $\alpha_{i,j}^{(d)}$ the s -bit label corresponding to the subset of $R^{\{i,j\}}$ that is held by A_i in $P^{(d)}$. Consider the sequence of allocations,

$$multi(\Delta) = \langle C^{(1)}, C^{(2)}, \dots, C^{(t)} \rangle$$

in a resource allocation setting have k agents and $2m$ resources – $\mathcal{R}_m \cup \mathcal{T}_m$ for which $C_i^{(d)}$ is characterised by

$$\beta_{i,0}^{(d)} \beta_{i,1}^{(d)} \cdots \beta_{i,i-1}^{(d)} \beta_{i,i+1}^{(d)} \cdots \beta_{i,k-1}^{(d)}$$

In this, $\beta_{i,j}^{(d)}$, indicates the subset of $\mathcal{R}^{\{i,j\}} \cup \mathcal{T}^{\{i,j\}}$ described by the $2s$ -bit label,

$$\beta_{i,j}^{(d)} = \alpha_{i,j}^{(d)} \overline{\alpha_{i,j}^{(d)}}$$

i.e. $\alpha_{i,j}^{(d)}$ selects a subset of $\mathcal{R}^{\{i,j\}}$ while $\overline{\alpha_{i,j}^{(d)}}$ a subset of $\mathcal{T}^{\{i,j\}}$.

It is immediate from this construction that for each allocation $C^{(d)}$ in $multi(\Delta)$ and each A_i , it is always the case that $|C_i^{(d)}| = (k-1)s$. It follows, therefore, that the only subsets that are relevant to the definition of *monotone* utility functions with which an analogous result to Theorem 6 for the path $multi(\Delta)$ are those of size $(k-1)s$: if $S \subseteq \mathcal{R}_m \cup \mathcal{T}_m$ has $|S| < (k-1)s$, we can fix $u_i(S)$ as a small enough negative value; similarly if $|S| > (k-1)s$ then $u_i(S)$ can be set to a large enough positive value¹⁴.

Our description in the preceding paragraphs is merely intended to outline how Theorem 6 is developed to apply to monotone utility functions. Extending this outline to a formal lower bound proof, is largely a technical exercise employing much of the analysis already introduced: since nothing significantly new is required for such an analysis we shall not give a detailed presentation of it.

3. Conclusions and Further Work

Our aim in this article has been to develop the earlier studies of Sandholm [20] concerning the scope and limits of particular “practical” contract forms. While [20] has established that insisting on individual rationality in addition to the structural restriction prescribed by O -contracts leads to scenarios which are incomplete (in the sense that there are individually rational exchanges that cannot be realised by individually rational O -contracts) our focus has been with respect to exchanges which can be realised by restricted contract paths, with the intention of determining to what extent the combination of structural and rationality conditions increases the number of exchanges required. We have shown that, using a number of natural definitions of rationality, for settings involving m resources, *rational* O -contract paths of length $\Omega(2^m)$ are needed, whereas without the rationality restriction on individual exchanges, at most m O -contracts suffice to realise any exchange. We have also considered a class of exchanges – $M(k)$ -contracts – that were not examined in [20], establishing for these cases that, when particular rationality conditions are imposed, $M(k-1)$ -contract

14. It is worth noting that the “interpolation” stage used in Theorem 4 is not needed in forming $multi(\Delta)$: the exchange $\langle C^{(d)}, C^{(d+1)} \rangle$ is an $M(k-1)$ -contract. We recall that in going from Δ_s of Theorem 3 to $ext(\Delta_s)$ the intermediate stage – $double(\Delta_s)$ – was not an O -contract path.

paths of length $\Omega(2^{2m/k^2})$ are needed to realise an exchange that can be achieved by a single $M(k)$ -contract.

We note that our analyses have primarily been focused on worst-case lower bounds on path length when appropriate paths exist, and as such there are several questions of practical interest that merit further discussion. It may be noted that the path structures and associated utility functions are rather artificial, being directed to attaining a path of a specific length meeting a given rationality criterion. We have seen, however, in Theorems 4 and 5 as outlined in our discussion concluding Section 2.3 that the issue of exponential length contract paths continues to arise even when we require the utility functions to satisfy a monotonicity condition. We can identify two classes of open question that arise from these results.

Firstly, focusing on IR O -contract paths, it would be of interest to identify “natural” restrictions on utility functions which would ensure that, *if* an exchange $\langle P, Q \rangle$ can be implemented by an IR O -contract path, then it can be realised by one whose length is polynomially bounded in m . We can interpret Theorem 4, as indicating that monotonicity does *not* guarantee “short” IR contract paths. We note, however, that there *are* some restrictions that suffice. To use a rather trivial example, if the number of *distinct* values that σ_u can assume is at most m^p for some *constant* p then no IR O -contract path can have length exceeding m^p : successive exchanges must strictly increase σ_u and if this can take at most K different values then no IR contract path can have length exceeding K . As well as being of practical interest, classes of utility function with the property being considered would also be of some interest regarding one complexity issue. The result proved in [8] establishing that deciding if an IR O -contract path exists is NP-hard, gives a *lower bound* on the computational complexity of this problem. At present, no (non-trivial) *upper bound* on this problem’s complexity has been demonstrated. Our results in Theorems 3 and 4 indicate that *if* this decision problem is in NP (thus its complexity would be NP-complete rather than NP-hard) then the required polynomial length existence certificate must be something other than the path itself. We note that the proof of NP-hardness in [8] constructs an instance in which σ_u can take at most $O(m)$ distinct values: thus, from our example of a restriction ensuring that if such are present then IR O -contract paths are “short”, this result of [8] indicates that the question of *deciding* their existence might remain computationally hard.

Considering restrictions on the form of utility functions is one approach that could be taken regarding finding “tractable” cases. An alternative, would be to gain some insight into what the “average” path length is likely to be. In attempting to address this question, however, a number of challenging issues arise. The most immediate of these concerns, of course, the notion of modeling a distribution on utility function given our definitions of rationality in terms of the value agents attach to their resource holdings. In principle an average-case analysis of scenarios involving exactly *two* agents could be carried out in purely graph-theoretic terms, i.e. without the complication of considering utility functions directly. It is unclear, however, whether such a graph-theoretic analysis obviating the need for consideration of literal utility functions, can be extended beyond settings involving exactly two agents. One difficulty arising with three or more agents is that our utility

functions are *context independent*, i.e. given an allocation $\langle X, Y, Z \rangle$ to three agents, $u_1(X)$ is unchanged should $Y \cup Z$ be redistributed among A_2 and A_3 ¹⁵.

As one final set of issues that may merit further study we raise the following. In our constructions, the individual exchanges on a contract path must satisfy both a *structural* condition (be an O -contract or involve at most k agents), and a *rationality* constraint. Focusing on O -contracts we have the following extremes: from [20], at most m O -contracts suffice to realise any rational exchange; from our results above, $\Omega(2^m)$ *rational* O -contracts are needed to realise some rational exchanges. There are a number of mechanisms we can employ to relax the condition that every single exchange be an O -contract and be rational. For example, allow a path to contain a some number of exchanges which are not O -contracts (but must still be IR) or insist that all exchanges are O -contracts but allow some to be irrational. Thus, in the latter case, if we go to the extent of allowing up to m irrational O -contracts, then any rational exchange can be realised efficiently. It would be of some interest to examine issues such as the effect of allowing a *constant* number, t , of irrational exchanges and questions such as whether there are situations in which t irrational contracts yield a ‘short’ contract path but $t-1$ force one of exponential length. Of particular interest, from an application viewpoint, is the following: define a $(\gamma(m), O)$ -path as an O -contract path containing at most $\gamma(m)$ O -contracts which are not individually rational. We know that if $\gamma(m) = 0$ then individually rational $(0, O)$ -paths are not complete with respect to individually rational exchanges; similarly if $\gamma(m) = m$ then (m, O) -paths are complete with respect to individually rational exchanges. A question of some interest would be to establish if there is some $\gamma(m) = o(m)$ for which $(\gamma(m), O)$ -paths are complete with respect to individually rational exchanges *and* with the maximum length of such a contract path bounded by a polynomial function of m .

References

- [1] H. L. Abbott and M. Katchalski. On the construction of snake in the box codes. *Utilitas Mathematica*, 40:97–116, 1991.
- [2] J. Abdou and H. Keiding. *Effectivity Functions in Social Choice Theory*. Kluwer Academic Publishers: Dordrecht, The Netherlands, 1991.
- [3] A. Atkinson. On the measurement of inequality. *Jnl. Econ. Theory*, 2:244–263, 1970.
- [4] A. Chateauneaf, T. Gajdos, and P.-H. Wilthien. The principle of strong diminishing transfer. *Jnl. Econ. Theory*, 103:311–333, 2002.
- [5] F. Dignum and M. Greaves (editors). *Issues in Agent Communication*. Springer-Verlag, 2000.
- [6] P.E. Dunne. Prevarication in dispute protocols. *Proc. Ninth International Conf. on A.I. and Law*, (ICAIL’ 03), Edinburgh, June 2003, ACM Press, pages 12–21

15. The term “context dependent” is that used in [9]: while such utility functions appear to have been neglected in the computational and algorithmic analysis of resource allocation problems, the idea is well-known to game-theoretic models in economics, where the term “allocative externality” is employed.

- [7] P.E. Dunne and P.J. McBurney. Optimal utterances in dialogue protocols. *Proc. Second International Joint Conf. on Autonomous Agents and Multiagent Systems (AAMAS'03)*, July 2003, ACM Press, pages 608-615
- [8] P.E. Dunne, M.J. Wooldridge, and M. Laurence. The complexity of contract negotiation. Technical Report, ULCS-03-002 Dept. of Computer Science, Univ. of Liverpool, March 2003 (submitted)
- [9] P.E. Dunne. Context Dependence in Multiagent Resource Allocation. Technical Report, Dept. of Computer Science, Univ. of Liverpool, 2004 (short version to appear *Proc. ECAI'04*), August 2004
- [10] U. Endriss, N. Maudet, F. Sadri, and F. Toni. On optimal outcomes of negotiations over resources. *Proc. Second International Joint Conf. on Autonomous Agents and Multiagent Systems (AAMAS'03)*, July 2003, ACM Press, pages 177 - 184
- [11] U. Endriss and N. Maudet. Welfare engineering in multiagent systems. In A. Omicini, P. Petta, and J. Pitt, (editors), *Proc. Fourth International Workshop on Engineering Societies in the Agents World (ESAW-2003)*, October 2003. Postproceedings to be published by Springer-Verlag
- [12] W. H. Kautz. Unit distance error checking codes. *IRE Trans. on Electronic Computers*, 7:179–180, 1958
- [13] S.-Ch. Kolm. Unequal inequalities. *Jnl. Econ. Theory*, 13:82–111, 1976.
- [14] S. Kraus. *Strategic negotiation in multiagent environments*. MIT Press, 2001.
- [15] P. McBurney, S. Parsons, and M. Wooldridge. Desiderata for argumentation protocols. In *Proc. First International Joint Conf. on Autonomous Agents and Multiagent Systems (AAMAS'02)*, pages 402–409. ACM Press, 2002.
- [16] D. C. Parkes and L. H. Ungar. Iterative combinatorial auctions: theory and practice. In *Proc. 17th National Conf. on Artificial Intelligence (AAAI-00)*, pages 74–81, 2000
- [17] D. C. Parkes and L. H. Ungar. Preventing strategic manipulation in iterative auctions: proxy agents and price adjustment. In *Proc. 17th National Conf. on Artificial Intelligence (AAAI-00)*, pages 82–89, 2000
- [18] J. S. Rosenschein and G. Zlotkin. *Rules of Encounter*. MIT Press, 1994.
- [19] Y. Sakurai, M. Yokoo and S. Matsubara. A limitation of the generalized Vickrey auction in electronic commerce: robustness against false-name bids. In *Proc. of the Sixteenth National Conference on Artificial Intelligence, AAAI-99*, pages 86–92, 1999
- [20] T. W. Sandholm. Contract types for satisficing task allocation: I theoretical results. In *AAAI Spring Symposium: Satisficing Models*, 1998.
- [21] T. W. Sandholm. Distributed rational decision making. In *Multiagent Systems* (Editor: G. Weiß) pages 201–258, MIT Press, 1999.

- [22] T. W. Sandholm, S. Suri, A. Gilpin, and D. Levine. CABOB: A fast optimal algorithm for combinatorial auctions. In *Proc. IJCAI-01*, pages 1102-1108, 2001
- [23] T. W. Sandholm. Algorithm for optimal winner determination in combinatorial auctions. *Artificial Intelligence*, 135:1-54, 2002
- [24] T. W. Sandholm and S. Suri. BOB: Improved winner determination in combinatorial auctions and generalizations. *Artificial Intelligence*, 145:33-58, 2003
- [25] R. G. Smith. The contract net protocol: high-level communication and control in a distributed problem solver. *IEEE Trans. on Computers*, C-29(12):1104-1113, 1980.
- [26] M. Tennenholz. Some tractable combinatorial auctions. In *Proc. 17th National Conf. on Artificial Intelligence (AAAI-00)*, 2000
- [27] M. Wooldridge and P. E. Dunne. On the computational complexity of qualitative coalitional games. to appear *Artificial Intelligence*, (in press, 2004)
- [28] M. Yokoo, Y. Sakurai and S. Matsubara. The effect of false-name declarations in mechanism design: towards collective decision making on the Internet. In *Proc. of the Twentieth International Conf. on Distributed Computing Systems, ICDCS-2000*, pages 146-153, 2000
- [29] M. Yokoo, Y. Sakurai and S. Matsubara. The effect of false-name bids in combinatorial auctions: new fraud in internet auctions. *Games and Economic Behavior*, 46(1):174-188, 2004