

EXTREMAL LYAPUNOV EXPONENTS: AN INVARIANCE PRINCIPLE AND APPLICATIONS

ARTUR AVILA ^{1,2} AND MARCELO VIANA ²

ABSTRACT. We propose a new approach to analyzing dynamical systems that combine hyperbolic and non-hyperbolic (“center”) behavior, e.g. partially hyperbolic diffeomorphisms. A number of applications illustrate its power.

We find that any ergodic automorphism of the 4-torus with two eigenvalues in the unit circle is stably Bernoulli among symplectic maps. Indeed, any nearby symplectic map has no zero Lyapunov exponents, unless it is volume preserving conjugate to the automorphism itself. Another main application is to accessible skew-product maps preserving area on the fibers. We prove, in particular, that if the genus of the fiber is at least 2 then the Lyapunov exponents must be different from zero and vary continuously with the map.

These, and other dynamical conclusions, originate from a general Invariance Principle we prove in here. It is formulated in terms of smooth cocycles, that is, fiber bundle morphisms acting by diffeomorphisms on the fibers. The extremal Lyapunov exponents measure the smallest and largest exponential rates of growth of the derivative along the fibers. The Invariance Principle states that if these two numbers coincide then the fibers carry some amount of structure which is transversely invariant, that is, invariant under certain canonical families of homeomorphisms between fibers.

1. INTRODUCTION

A core issue in the theory of partially hyperbolic dynamical systems is the possible vanishing of center Lyapunov exponents. Recall that a diffeomorphism $f : M \rightarrow M$ is called partially hyperbolic if the tangent bundle admits a continuous Df -invariant splitting

$$T_x M = E_x^s \oplus E_x^c \oplus E_x^u, \quad x \in M$$

such that $Df|_{E_x^s}$ is a uniform contraction, $Df|_{E_x^u}$ is a uniform expansion, and the behavior of $Df|_{E_x^c}$ lies in between those two (not quite as contracting nor as expanding, respectively). When the vectors in the center bundle E^c also have non-zero Lyapunov exponents, that is, when

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \|Df^n(x)v\| \neq 0 \quad \forall v \in E_x^c$$

at typical points $x \in M$, one can build on (non-uniform) hyperbolicity theory to derive important geometric and statistical information on the dynamics. See Pesin [38, 39], Ledrappier, Young [32, 34, 35], Katok [29], Barreira, Pesin, Schmeling [8], Young [45, 46], and Alves, Bonatti, Viana [2, 15].

Date: March 15, 2010.

¹ CNRS UMR 7599, Laboratoire de Probabilités et Modèles Aléatoires, Université Pierre et Marie Curie, Boîte Postale 188, 75252 Paris Cedex 05, France.

² IMPA – Estrada D. Castorina 110, Jardim Botânico, 22460-320 Rio de Janeiro – Brazil.

So, one would like to know: *When is it possible to remove vanishing center Lyapunov exponents by a small perturbation of the map? Supposing center Lyapunov exponents do vanish (stably), what can be said about the dynamical behavior?*

Substantial progress on this kind of questions has been achieved recently in the simpler, but closely related setting of linear cocycles. A linear cocycle over a transformation $g : X \rightarrow X$ is a map $F : \mathcal{E} \rightarrow \mathcal{E}$ on a vector bundle $\mathcal{E} \rightarrow X$ such that

$$\begin{array}{ccc} F : \mathcal{E} & \rightarrow & \mathcal{E} \\ & \downarrow & \downarrow \\ g : X & \rightarrow & X \end{array}$$

commutes and the action $F_x : \mathcal{E}_x \rightarrow \mathcal{E}_{g(x)}$ on every fiber is by a linear isomorphism. As before, the exponential rates of growth or decay of iterates of vectors

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \|F_x^n v\|, \quad v \in \mathcal{E}_x$$

are called Lyapunov exponents. By Oseledec's [37], they are well defined at almost every point $x \in X$, relative to any g -invariant probability measure μ .

Assuming the base dynamics (g, μ) is fairly "chaotic" (hyperbolic, possibly in a non-uniform fashion), there is now a good understanding of such issues as the existence of non-zero Lyapunov exponents (see Bonatti, Gomez-Mont, Viana [14, 43]) or the simplicity of the Lyapunov spectrum (see Avila, Bonatti, Viana [5, 6, 16]), in line with the classical theory of random matrices developed by Furstenberg [23], Ledrappier [33], Guivarc'h, Raugi [25], Gol'dsheid, Margulis [24], and other authors. In a nutshell, for generic linear cocycles the Lyapunov exponents are not all zero. Even more, at least in the so-called fiber bunched case, all Lyapunov exponents are generically distinct.

In this paper we propose to extend that family of ideas to a general non-linear context, suitable, in particular, to addressing such questions as we stated before in the realm of partially hyperbolic dynamics. The power of this kind of analysis is made evident by the following surprising rigidity phenomenon we have discovered, and which motivates our discussion below.

Let $A : \mathbb{T}^4 \rightarrow \mathbb{T}^4$ be a linear automorphism with exactly two eigenvalues on the unit circle and assume that no eigenvalue is a root of unity. The latter condition means that A is ergodic relative to the Haar measure on the 4-torus. Rodriguez-Hertz [26] proves that A is even stably ergodic, that is to say, every nearby volume preserving map $f : \mathbb{T}^4 \rightarrow \mathbb{T}^4$ is ergodic. Notice that A is a partially hyperbolic diffeomorphism, with 2-dimensional center bundle E^c that corresponds to the eigenvalues of norm 1. Clearly, the center Lyapunov exponents are identically zero.

The hypotheses imply that A preserves some symplectic form on \mathbb{T}^4 . Let this form be fixed once and for all.

Theorem A. *There exists a neighborhood \mathcal{U} of A in the space of smooth symplectic diffeomorphisms on \mathbb{T}^4 such that for every $f \in \mathcal{U}$,*

- *either f is non-uniformly hyperbolic, that is, all Lyapunov exponents are non-zero almost everywhere,*
- *or f is conjugate to A by some volume preserving homeomorphism.*

In particular, every $f \in \mathcal{U}$ is equivalent to a Bernoulli shift.

In other words, the center Lyapunov exponents must become non-zero whenever A is perturbed, unless the perturbation leaves the dynamics unchanged up

to volume preserving conjugacy. Moreover, A is stably Bernoulli, a much stronger property than ergodicity.

The proof of Theorem A has many ingredients, but the main thread is that in order to prove conjugacy to the original automorphism one must recover a commutative group structure in the torus compatible with the dynamics. In the hardest case to analyze, such structure is obtained as the completion of an *almost periodic translation structure* on the center leaves, which is itself a refinement of a *conformal structure*. Thus, at the basis of the entire proof, one needs to connect vanishing of center Lyapunov exponents with existence of invariant conformal structures.

With these motivations in mind, we now describe the abstract Invariance Principle that is the foundation of this paper. It is formulated in terms of smooth cocycles $F : \mathcal{E} \rightarrow \mathcal{E}$, a natural extension of the notion of linear cocycles where \mathcal{E} is now taken to be a fiber bundle whose fibers are manifolds, and F is taken to act by diffeomorphisms on the fibers. Lyapunov exponents are defined in this setting by

$$\lim \frac{1}{n} \log \|DF_x(\xi)v\|, \quad v \in T_\xi \mathcal{E}_x, \quad \xi \in \mathcal{E}_x.$$

In a few words, the Invariance Principle states that if the Lyapunov exponents vanish then the fibers carry some amount of structure which is transversely invariant, that is, invariant under certain canonical homeomorphisms between the fibers. The precise statement will be given in the next section, once we have introduced all the necessary notions. In the applications, we exploit such transversely invariant structure, together with information on the fiber (e.g. its topology), to deduce some very precise information on the dynamics.

Besides Theorem A, we apply this approach to certain area preserving cocycles. Namely, suppose the fibers of \mathcal{E} are modeled on some compact surface N and the cocycle $F : \mathcal{E} \rightarrow \mathcal{E}$ is such that all $F_x : \mathcal{E}_x \rightarrow \mathcal{E}_g(x)$ preserve some given area form. *Assuming the genus of N is at least 2*, together with a few additional conditions, we can prove that *the Lyapunov exponents of F are non-zero at almost every point*. Moreover, the Lyapunov exponents of every nearby cocycle are close to the Lyapunov exponents of F .

The idea of extending Furstenberg's theory to a non-linear set-up is, of course, not new. Let us mention, in particular, Carverhill [19] and Baxendale [9], who both dealt with i.i.d. situations. For instance, Baxendale's (discrete time) stochastic flows of diffeomorphisms correspond to the particular case of smooth cocycles where the base dynamics is a Bernoulli shift and the cocycle depends on only one coordinate in shift space. For the sharpest results Baxendale [9] also assumes the stationary measure to be absolutely continuous on the fibers.

However, the formulation we propose here seems particularly suited for applications to various situations in Dynamics. Indeed, several applications of the present methods have been found in the meantime, some of which had not been foreseen. In our joint paper with Santamaria [3] the Invariance Principle is refined and combined with other techniques to handle cocycles over partially hyperbolic systems. This is also a main ingredient in our joint paper with Wilkinson [7], where new connections between the dynamics of partially hyperbolic systems and the measure-theoretical properties of their invariant foliations are unveiled.

In a setting of dissipative systems, Viana, Yang [44] combine methods in the present paper with other ideas to prove existence and finiteness of physical (or Sinai-Ruelle-Bowen) measures for partially hyperbolic maps with 1-dimensional center.

The Invariance Principle allows them to treat the case of vanishing center Lyapunov exponent, which was not covered by [2, 15]. Also very recently, F. and J. Rodriguez-Hertz, R. Ures, A. Tahzibi announced further applications of these methods, to entropy maximizing measures of partially hyperbolic maps. Again, the Invariance Principle allows them to deal with the case when the maximizing measures have zero center exponents.

Acknowledgments. We are grateful to Jairo Bochi, Carlos Bocker, Jimmy Santamaría, Amie Wilkinson, and Jiagang Yang for several useful discussions. This work was started while the authors were visiting the Collège de France. It was partly conducted during the period A. A. served as a Clay Research Fellow. M. V. was partially supported by CNPq, FAPERJ, and PRONEX-Dynamical Systems.

2. STATEMENT OF RESULTS

In the sequel we give the precise definitions and statements that lead to the conclusions outlined in the Introduction.

2.1. Smooth cocycles. Let $(\hat{M}, \hat{\mathcal{B}}, \hat{\mu})$ be a probability space and $\hat{f} : \hat{M} \rightarrow \hat{M}$ be a measurable transformation preserving $\hat{\mu}$. Let N be a Riemannian manifold, not necessarily complete, and let $\text{Diff}^1(N)$ be endowed with a uniform C^1 norm. Let $\hat{P} : \hat{\mathcal{E}} \rightarrow \hat{M}$ be a measurable fiber bundle with fibers modeled on N . By this we mean $\hat{\mathcal{E}}$ comes with a countable system of bijections

$$(1) \quad \hat{P}^{-1}(U_n) \rightarrow U_n \times N$$

that map each fiber $\hat{\mathcal{E}}_{\hat{x}} = \hat{P}^{-1}(\hat{x})$ onto $\{\hat{x}\} \times N$, and all coordinate changes are measurable maps of the form

$$(2) \quad (U_m \cap U_n) \times N \rightarrow (U_m \cap U_n) \times N, \quad (\hat{x}, \hat{\xi}) \mapsto (\hat{x}, g_{\hat{x}}(\hat{\xi}))$$

where $g_{\hat{x}} : N \rightarrow N$ is a diffeomorphism depending measurably on the base point \hat{x} and such that both the derivative $Dg_{\hat{x}}(\hat{\xi})$ and its inverse are uniformly continuous and uniformly bounded. Then one may consider a Riemannian metric on the fibers, varying measurably with the base point, transported from N via these coordinates. This metric depends on the choice of the coordinates, but only up to a uniformly bounded factor, which does not affect the notions that follow.

A *smooth cocycle* over \hat{f} is a measurable map $\hat{F} : \hat{\mathcal{E}} \rightarrow \hat{\mathcal{E}}$ such that $\hat{P} \circ \hat{F} = \hat{f} \circ \hat{P}$, every

$$\hat{F}_{\hat{x}} : \hat{\mathcal{E}}_{\hat{x}} \rightarrow \hat{\mathcal{E}}_{\hat{f}(\hat{x})}$$

is a diffeomorphism depending measurably on \hat{x} , and the derivative $D\hat{F}_{\hat{x}}(\hat{\xi})$ and its inverse are uniformly bounded in norm. Then the functions

$$(\hat{x}, \hat{\xi}) \mapsto \log \|D\hat{F}_{\hat{x}}(\hat{\xi})\| \quad \text{and} \quad (\hat{x}, \hat{\xi}) \mapsto \log \|D\hat{F}_{\hat{x}}(\hat{\xi})^{-1}\|$$

are integrable, relative to any probability measure \hat{m} on $\hat{\mathcal{E}}$. The *extremal Lyapunov exponents* of \hat{F} at a point $(\hat{x}, \hat{\xi}) \in \hat{\mathcal{E}}$ are

$$\begin{aligned} \lambda_+(\hat{F}, \hat{x}, \hat{\xi}) &= \lim_{n \rightarrow \infty} \frac{1}{n} \log \|D\hat{F}_{\hat{x}}^n(\hat{\xi})\| . \\ \lambda_-(\hat{F}, \hat{x}, \hat{\xi}) &= \lim_{n \rightarrow \infty} \frac{1}{n} \log \|D\hat{F}_{\hat{x}}^n(\hat{\xi})^{-1}\|^{-1} . \end{aligned}$$

The limits exist \hat{m} -almost everywhere if \hat{m} is invariant under \hat{F} , by the subadditive ergodic theorem (Kingman [31]). Notice that

$$\lambda_{-}(\hat{F}, \hat{x}, \hat{\xi}) \leq \lambda_{+}(\hat{F}, \hat{x}, \hat{\xi}),$$

because $\|D\hat{F}_{\hat{x}}^n(\hat{\xi})\| \|D\hat{F}_{\hat{x}}^n(\hat{\xi})^{-1}\| \geq 1$. Denote

$$\lambda_{\pm} = \lambda_{\pm}(\hat{F}, \hat{m}) = \int \lambda_{\pm}(\hat{F}, \hat{x}, \hat{\xi}) d\hat{m}(\hat{x}, \hat{\xi}).$$

If (\hat{F}, \hat{m}) is ergodic then $\lambda_{\pm}(\hat{F}, \hat{x}, \hat{\xi}) = \lambda_{\pm}$ for \hat{m} -almost every $(\hat{x}, \hat{\xi})$. Throughout, we shall only be interested in measures \hat{m} that project down to μ under \hat{P} .

2.2. Invariance Principle - measurable. The main technical tool developed in this paper is a measurability criterion for the disintegration along the fibers of probability measures invariant under a cocycle. This is inspired by the main result in Ledrappier [33]: while Ledrappier's original formulation was for linear cocycles, ours applies to any deformation of a smooth cocycle, a notion that we also introduce in here.

Take $(\hat{M}, \hat{\mathcal{B}}, \mu)$ to be a Lebesgue space, that is, a separable probability space which is complete mod 0. See Rokhlin [41, §2-§3]. Then any probability \hat{m} on $\hat{\mathcal{E}}$ such that $\hat{P}_{*}\hat{m} = \hat{\mu}$ admits a family $\{\hat{m}_{\hat{x}} : \hat{x} \in \hat{M}\}$ of probabilities such that $\hat{x} \mapsto \hat{m}_{\hat{x}}$ is $\hat{\mathcal{B}}$ -measurable, every $\hat{m}_{\hat{x}}$ is supported inside the fiber $\hat{\mathcal{E}}_{\hat{x}}$ and

$$\hat{m}(E) = \int \hat{m}_{\hat{x}}(E) d\hat{\mu}(\hat{x})$$

for any measurable set $E \subset \hat{\mathcal{E}}$. Moreover, such a family is essentially unique. We call it the *disintegration* of \hat{m} and refer to the $\hat{m}_{\hat{x}}$ as its *conditional probabilities* along the fibers.

Assume that \hat{f} is invertible. A σ -algebra $\mathcal{B}_0 \subset \hat{\mathcal{B}}$ is *generating* if its iterates $\hat{f}^n(\mathcal{B}_0)$, $n \in \mathbb{Z}$ generate the whole $\hat{\mathcal{B}}$ mod 0. A *deformation of a smooth cocycle* \hat{F} is a measurable transformation $\tilde{F} : \hat{\mathcal{E}} \rightarrow \hat{\mathcal{E}}$ which is conjugated to \hat{F} ,

$$\tilde{F} = H \circ \hat{F} \circ H^{-1},$$

by an invertible measurable map $H : \hat{\mathcal{E}} \rightarrow \hat{\mathcal{E}}$ of the form $H(\hat{x}, \hat{\xi}) = (\hat{x}, H_{\hat{x}}(\hat{\xi}))$ such that all the $H_{\hat{x}}^{-1}$, $\hat{x} \in \hat{M}$ are Hölder continuous, with uniform Hölder constants: there exist positive constants B and β such that

$$(3) \quad d(\hat{\xi}, \hat{\eta}) \leq B d(H_{\hat{x}}(\hat{\xi}), H_{\hat{x}}(\hat{\eta}))^{\beta} \quad \text{for all } \hat{x} \in \hat{M} \text{ and } \hat{\xi}, \hat{\eta} \in \mathcal{E}_{\hat{x}}.$$

To each \hat{F} -invariant probability measure \hat{m} corresponds an \tilde{F} -invariant probability $\tilde{m} = H_{*}\hat{m}$, and \tilde{m} projects down to $\hat{\mu}$ if and only if \hat{m} does.

Theorem B. *Let \tilde{F} be a deformation of a smooth cocycle \hat{F} . Let $\mathcal{B}_0 \subset \hat{\mathcal{B}}$ be a generating σ -algebra such that both \hat{f} and $x \mapsto \tilde{F}_x$ are \mathcal{B}_0 -measurable mod 0. Let \tilde{m} be an \tilde{F} -invariant probability that projects down to $\hat{\mu}$. If $\lambda_{-}(\tilde{F}, \hat{x}, \hat{\xi}) \geq 0$ for \tilde{m} -almost every $(\hat{x}, \hat{\xi}) \in \hat{\mathcal{E}}$ then any disintegration $x \mapsto \tilde{m}_x$ of the corresponding \tilde{F} -invariant measure \tilde{m} is \mathcal{B}_0 -measurable mod 0.*

We get a dual result assuming that $\lambda_{+}(\hat{F}, \hat{x}, \hat{\xi}) \leq 0$ for \hat{m} -almost every $(\hat{x}, \hat{\xi})$, and considering a σ -algebra \mathcal{B}_0 relative to which both maps \hat{f}^{-1} and $\hat{x} \mapsto \tilde{F}_x^{-1}$ are measurable mod 0. Indeed, it is clear that \hat{F} has the same invariant probabilities

as \hat{F}^{-1} , and \tilde{F} is a deformation of \hat{F} if and only if \tilde{F}^{-1} is a deformation of \hat{F}^{-1} . Since

$$\lambda_+(\hat{F}, \hat{x}, \hat{\xi}) + \lambda_-(\hat{F}^{-1}, \hat{x}, \hat{\xi}) = 0,$$

the new assumption means that $\lambda_-(\hat{F}^{-1}, \hat{x}, \hat{\xi}) \geq 0$ for \hat{m} -almost every $(\hat{x}, \hat{\xi})$. Thus, we may apply Theorem B to the inverse cocycle, to obtain the same conclusion as before under this new assumption. See also Example 3.15 below.

Theorem C. *Let \tilde{F} be a deformation of a smooth cocycle \hat{F} . Let $\mathcal{B}_0 \subset \hat{\mathcal{B}}$ be a generating σ -algebra such that both \hat{f} and $x \mapsto \tilde{F}_x$ are \mathcal{B}_0 -measurable mod 0. Let $(\hat{m}_k)_k$ be a sequence of \hat{F} -invariant probabilities projecting down to $\hat{\mu}$ and converging to some probability \hat{m} in the weak* topology. Assume $\int \min\{0, \lambda_-(\hat{F}, \cdot)\} d\hat{m}_k \rightarrow 0$ when $k \rightarrow \infty$. Then any disintegration $x \mapsto \tilde{m}_x$ of the corresponding \tilde{F} -invariant measure \tilde{m} is \mathcal{B}_0 -measurable mod 0.*

Theorem B may be viewed as the special case when $m_k = m$ for all k : it is clear that $\int \min\{0, \lambda_-(\hat{F}, \cdot)\} d\hat{m} = 0$ if and only if $\lambda_-(\hat{F}, \cdot) \geq 0$ \hat{m} -almost everywhere.

2.3. Hyperbolic homeomorphisms. Next, we are going to derive more concrete versions of these results for continuous cocycles over hyperbolic homeomorphisms.

Let \hat{M} be a metric space. Let $\hat{\mathcal{E}}$ be a continuous fiber bundle (the local coordinates (1) are defined on open sets and the coordinate changes (2) are homeomorphisms) where the diffeomorphisms $g_{\hat{x}}$ vary continuously with $\hat{x} \in \hat{M}$. Assume that a Riemannian metric has been chosen on each fiber, varying continuously with the base point. Moreover, let \hat{F} be a smooth cocycle such that the diffeomorphisms $\hat{F}_{\hat{x}}$ vary continuously with $\hat{x} \in \hat{M}$.

We call a homeomorphism $\hat{f} : \hat{M} \rightarrow \hat{M}$ *hyperbolic* if there exist $\varepsilon > 0$, $\delta > 0$, $K > 0$, $\tau > 0$, and positive functions $\nu(\cdot)$ and $\nu_-(\cdot)$ such that

- (h1) $d(\hat{f}(\hat{y}_1), \hat{f}(\hat{y}_2)) \leq \nu(\hat{x})d(\hat{y}_1, \hat{y}_2)$ for all $\hat{y}_1, \hat{y}_2 \in W_\varepsilon^s(\hat{x})$, $\hat{x} \in \hat{M}$;
- (h2) $d(\hat{f}^{-1}(\hat{z}_1), \hat{f}^{-1}(\hat{z}_2)) \leq \nu_-(\hat{x})d(\hat{z}_1, \hat{z}_2)$ for all $\hat{z}_1, \hat{z}_2 \in W_\varepsilon^u(\hat{x})$, $\hat{x} \in \hat{M}$;
- (h3) $\nu_n(\hat{x}) := \nu(\hat{f}^{n-1}(\hat{x})) \cdots \nu(\hat{x}) \leq Ke^{-\tau n}$ for all $\hat{x} \in \hat{M}$ and $n \geq 1$;
- (h4) $\nu_{-n}(\hat{x}) := \nu_-(\hat{f}^{-n+1}(\hat{x})) \cdots \nu_-(\hat{x}) \leq Ke^{-\tau n}$ for all $\hat{x} \in \hat{M}$ and $n \geq 1$;
- (h5) if $d(\hat{x}_1, \hat{x}_2) \leq \delta$ then $W_\varepsilon^u(\hat{x}_1)$ and $W_\varepsilon^s(\hat{x}_2)$ intersect at exactly one point, denoted $[\hat{x}_1, \hat{x}_2]$, and this point depends continuously on (\hat{x}_1, \hat{x}_2) ;

where $W_\varepsilon^s(\hat{x})$ is the set of all $\hat{y} \in \hat{M}$ such that $d(\hat{f}^n(\hat{x}), \hat{f}^n(\hat{y})) \leq \varepsilon$ for all $n \geq 0$, and $W_\varepsilon^u(\hat{x})$ is defined analogously, with $n \leq 0$ instead. Then the stable and unstable sets of \hat{x} are given by

$$W^s(\hat{x}) = \bigcup_{n \geq 0} \hat{f}^{-n}(W_\varepsilon^s(\hat{f}^n(\hat{x}))) \quad \text{and} \quad W^u(\hat{x}) = \bigcup_{n \geq 0} \hat{f}^n(W_\varepsilon^u(\hat{f}^{-n}(\hat{x}))).$$

Example 2.1. Let $\hat{f} : \hat{M} \rightarrow \hat{M}$ be the shift map on $\hat{M} = X^{\mathbb{Z}}$ where (X, d_X) is a complete metric space, and the metric $d(\cdot, \cdot)$ on \hat{M} is defined by

$$d(\hat{x}, \hat{y}) = \sum_{n \in \mathbb{Z}} e^{-\tau|n|} \min\{1, d_X(x_n, y_n)\} \quad \text{for } \hat{x} = (x_n)_n \text{ and } \hat{y} = (y_n)_n.$$

Take $\varepsilon \in (0, 1)$, $\delta \in (0, 1)$, $K = 1$, and $\nu(\hat{x}) = \nu_-(\hat{x}) = e^{-\tau}$ for all $\hat{x} \in \hat{M}$.

Then there exist relative neighborhoods $B^s(\hat{x}) \subset W_\varepsilon^s(\hat{x})$ and $B^u(\hat{x}) \subset W_\varepsilon^u(\hat{x})$ of every $\hat{x} \in \hat{M}$ such that $\iota : (\hat{x}_1, \hat{x}_2) \mapsto [\hat{x}_1, \hat{x}_2]$ defines a homeomorphism from $B^s(\hat{x}) \times B^u(\hat{x})$ to some neighborhood $B(\hat{x})$ of every $\hat{x} \in \hat{M}$. We always consider

\hat{f} -invariant probabilities $\hat{\mu}$ with *local product structure*: for every \hat{x} in the support there exist measures μ^s and μ^u on $B^s(x)$ and $B^u(x)$, respectively, such that

$$(4) \quad \hat{\mu} \mid B(x) \sim \iota_*(\mu^u \times \mu^s),$$

meaning that the two measures have the same zero sets. This implies that the support is *su-saturated*, meaning it consists of entire stable leaves (*s-saturated set*) and of entire unstable leaves (*u-saturated set*). Moreover, $\hat{\mu}$ is locally ergodic, that is, its ergodic components are essentially open sets.

A measure $\hat{\mu}$ is called σ -compact if it gives full weight to some countable union of compact sets. If M is a Polish space, that is, a separable completely metrizable topological space then every Borel measure in it is σ -compact.

2.4. Invariance Principle - topological. An *s-holonomy* for \hat{F} is a family h^s of β -Hölder homeomorphisms $h_{\hat{x}, \hat{y}}^s : \hat{\mathcal{E}}_{\hat{x}} \rightarrow \hat{\mathcal{E}}_{\hat{y}}$, with uniform Hölder constant $\beta > 0$, defined for all $\hat{y} \in W^s(\hat{x})$ and satisfying

- (sh1) $h_{\hat{y}, \hat{z}}^s \circ h_{\hat{x}, \hat{y}}^s = h_{\hat{x}, \hat{z}}^s$ and $h_{\hat{x}, \hat{x}}^s = \text{id}$
- (sh2) $\hat{F}_{\hat{y}} \circ h_{\hat{x}, \hat{y}}^s = h_{\hat{f}(\hat{x}), \hat{f}(\hat{y})}^s \circ \hat{F}_{\hat{x}}$
- (sh3) $(\hat{x}, \hat{y}, \xi) \mapsto h_{\hat{x}, \hat{y}}^s(\xi)$ is continuous.

In the last condition (\hat{x}, \hat{y}) varies in the space of pairs of points in the same local stable set. A disintegration $\{\hat{m}_{\hat{x}} : \hat{x} \in \hat{M}\}$ of an \hat{F} -invariant probability \hat{m} is *s-invariant* if

$$(5) \quad (h_{\hat{x}, \hat{y}}^s)_* \hat{m}_{\hat{x}} = \hat{m}_{\hat{y}} \quad \text{for every } \hat{y} \in W^s(\hat{x})$$

with \hat{x} and \hat{y} in the support of the projection of \hat{m} . Replacing \hat{f} and \hat{F} by their inverses, one obtains dual notions of *u-holonomy* h^u and *u-invariant disintegration*.

Theorem D. *Assume $\hat{F} : \hat{\mathcal{E}} \rightarrow \hat{\mathcal{E}}$ admits s-holonomy and u-holonomy. Let $(\hat{m}_k)_k$ be a sequence of \hat{F} -invariant probability measures whose projection $\hat{\mu}$ is σ -compact and has local product structure. Assume the sequence converges to some probability measure \hat{m} in the weak* topology and $\int |\lambda_{\pm}(\hat{F}, \cdot)| d\hat{m}_k \rightarrow 0$ when $k \rightarrow \infty$. Then \hat{m} admits a disintegration $\{\hat{m}_{\hat{x}} : \hat{x} \in \hat{M}\}$ which is s-invariant and u-invariant and whose conditional probabilities $\hat{m}_{\hat{x}}$ vary continuously with \hat{x} on the support of $\hat{\mu}$.*

An extension for cocycles over certain partially hyperbolic maps will be given in Theorem 5.10. A first application of Theorem D is given in the proposition that follows. It will be clear from the arguments that the hypotheses can be relaxed considerably.

Corollary E. *Let $\hat{f} : \hat{M} \rightarrow \hat{M}$ be the shift map on $\hat{M} = X^{\mathbb{Z}}$, where X is a complete metric space. Let $\hat{\mathcal{E}} = \hat{M} \times \mathbb{S}^1$ and $\hat{F} : \hat{\mathcal{E}} \rightarrow \hat{\mathcal{E}}$ be a continuous smooth cocycle over \hat{f} admitting invariant holonomies. Suppose \hat{f} admits an invariant probability measure $\hat{\mu}$ and fixed points p and q in the support of $\hat{\mu}$ such that*

- $\hat{F}_p : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ has exactly two fixed points, an attractor a_p and a repeller r_p
- $\hat{F}_q : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ has no periodic points of period less than 3.

Then $\lambda_{\pm}(\hat{F}, \hat{m})$ are bounded away from zero, over all ergodic \hat{F} -invariant measures \hat{m} that project down to $\hat{\mu}$.

2.5. Volume preserving cocycles. We apply the previous ideas to area preserving cocycles over hyperbolic homeomorphisms satisfying certain partial hyperbolicity conditions.

From now on we take the fiber manifold N to be compact. Assume the continuous fiber bundle is Lipschitz, in the sense that the diffeomorphisms $g_{\hat{x}}$ in (2) depend in a Lipschitz fashion on the base point. Assume that the continuous cocycle is Lipschitz, in the sense that $\hat{F}_{\hat{x}}$ depends in a Lipschitz fashion on the point \hat{x} . We shall consider the following topology: two Lipschitz cocycles are close if they admit the same Lipschitz constant, they are uniformly close, and their actions on the fibers are close in the uniform C^1 norm.

Remark 2.2. For all our purposes it suffices to assume Hölder continuity, for some Hölder constant $\nu > 0$: up to replacing the metric on \hat{M} , one may always reduce the situation to the Lipschitz case $\nu = 1$.

We take the cocycle $\hat{F} : \hat{\mathcal{E}} \rightarrow \hat{\mathcal{E}}$ to satisfy a normal hyperbolicity property similar to the center bunching condition of Burns, Wilkinson [18] and which was first introduced in [14] in the context of linear cocycles. We say that a Lipschitz smooth cocycle \hat{F} is *dominated* if there exist $\ell \geq 1$ and $\theta < 1$ such that

$$(6) \quad \|(D\hat{F}_{\hat{x}}^{\ell}(\xi))^{-1}\| \nu_{\ell}(\hat{x}) \leq \theta \quad \text{and} \quad \|D\hat{F}_{\hat{x}}^{\ell}(\xi)\| \nu_{-\ell}(\hat{x}) \leq \theta$$

for every $(\hat{x}, \xi) \in \mathcal{E}$, and we say \hat{f} is *fiber bunched* if, in addition to (6),

$$(7) \quad \|D\hat{F}_{\hat{x}}^{\ell}(\xi)\| \|(D\hat{F}_{\hat{x}}^{\ell}(\xi))^{-1}\| \nu_{\pm\ell}(\hat{x}) \leq \theta$$

for every $(\hat{x}, \xi) \in \mathcal{E}$. Interpretations of these conditions will be provided in Section 5. Let $\mathcal{B}(\hat{f})$ be the set of fiber bunched cocycles over \hat{f} . Observe that this is an open subset of Lipschitz cocycles, relative to the topology introduced above.

Let $\hat{m}_{\hat{g}}$ denote the normalized Riemannian volume on each fiber $\hat{\mathcal{E}}_{\hat{g}}$. We also take the cocycle $\hat{F} : \hat{\mathcal{E}} \rightarrow \hat{\mathcal{E}}$ to be *volume preserving*, meaning that each

$$\hat{F}_{\hat{x}} : \hat{\mathcal{E}}_{\hat{x}} \rightarrow \hat{\mathcal{E}}_{\hat{f}(\hat{x})} \quad \text{maps } \hat{m}_{\hat{x}} \text{ to } \hat{m}_{\hat{f}(\hat{x})}.$$

Then the following probability measure \hat{m} on $\hat{\mathcal{E}}$ is \hat{F} -invariant:

$$(8) \quad \hat{m}(B) = \int \hat{m}_{\hat{x}}(B \cap \hat{\mathcal{E}}_{\hat{x}}) d\hat{\mu}(\hat{x}).$$

Let $\mathcal{B}_{\text{vol}}(\hat{f})$ denote the subset of volume preserving fiber bunched cocycles.

2.6. Continuity of Lyapunov exponents. From now on we take $\hat{\mathcal{E}}$ to be compact and the fiber N to be a surface. Area preserving yields $\lambda_{-}(\hat{F}, \hat{x}, \xi) + \lambda_{+}(\hat{F}, \hat{x}, \xi) = 0$ at \hat{m} -almost every point. We call $\hat{F} \in \mathcal{B}_{\text{vol}}(\hat{f})$ a *continuity point* for Lyapunov exponents if the functions

$$\mathcal{B}_{\text{vol}}(\hat{f}) \ni \hat{G} \mapsto \lambda_{\pm}(\hat{G}, \hat{m})$$

are continuous at \hat{F} . Otherwise, \hat{F} is a discontinuity point for Lyapunov exponents.

By analogy with Pugh, Shub [40], we say that a cocycle is *accessible* if any two points in the fiber bundle are joined by some *su-path*, consisting of a finite number of legs each of which is either an *s-holonomy* path or a *u-holonomy* path (assuming the cocycle admits *s-holonomy* and *u-holonomy*).

Theorem F. *Let $\hat{F} : \hat{\mathcal{E}} \rightarrow \hat{\mathcal{E}}$ be fiber bunched, area preserving, and ergodic. Let \hat{F} be a discontinuity point for Lyapunov exponents. Then $\lambda_-(\hat{F}, \hat{m}) < 0 < \lambda_+(\hat{F}, \hat{m})$ and both Oseledets subspaces $E_{\hat{x}, \hat{\xi}}^-$ and $E_{\hat{x}, \hat{\xi}}^+$ are essentially invariant under the s-holonomy and the u-holonomy of the projective extension. If, in addition, the cocycle \hat{F} is accessible then the Oseledets subspaces vary continuously with $(\hat{x}, \hat{\xi}) \in \hat{\mathcal{E}}$.*

Our methods also reveal a remarkable connection between the behavior of Lyapunov exponents and the topology of the fiber, at least when the cocycle \hat{F} is accessible. This is illustrated by the next corollary, which will follow from the more detailed statement in Theorem 6.6; see also Remark 6.7.

Corollary G. *Let $\hat{F} : \hat{\mathcal{E}} \rightarrow \hat{\mathcal{E}}$ be fiber bunched, area preserving, and accessible. Assume the genus of the fiber N of $\hat{\mathcal{E}}$ is at least 2. Then $\lambda_-(\hat{F}, \hat{m}) < 0 < \lambda_+(\hat{F}, \hat{m})$ and \hat{F} is a continuity point for the Lyapunov exponents $\lambda_{\pm}(\cdot, \hat{m})$.*

Remark 2.3. It is an important problem to characterize those cocycles which are continuity points for the Lyapunov exponents. The results of Bochi [10, 11] and Bochi, Viana [12] show that continuity depends quite subtly on topology in the space of cocycles. In a positive direction, J. Yang (personal communication) uses ideas from the present paper to conclude that the set of cocycles with zero Lyapunov exponents is closed in the Lipschitz topology among fiber bunched $\text{SL}(2, \mathbb{R})$ -cocycles. Furthermore, Bocker, Viana [13] prove that for locally constant $\text{SL}(2, \mathbb{C})$ -cocycles over Bernoulli shifts the Lyapunov exponents always vary continuously with the cocycle relative to the L^∞ norm.

For the next theorem, let $\hat{f} : \hat{M} \rightarrow \hat{M}$ be a C^r Anosov diffeomorphism on a compact manifold, for some $r \geq 1$. Moreover, take the fiber bundle to be trivial, that is, $\hat{\mathcal{E}} = \hat{M} \times N$, and the cocycle $\hat{F} : \hat{\mathcal{E}} \rightarrow \hat{\mathcal{E}}$ to be C^r . Recall we take N to be a compact surface. Let $\mathcal{B}_{\text{vol}}^r(\hat{f})$ be the space of area preserving fiber bunched C^r cocycles, endowed with the uniform C^r topology. For simplicity, we also assume that the fiber N is orientable and \hat{F} preserves the orientation of the fibers (the non-orientable case can be treated by considering a double cover).

Theorem H. *There is an open and dense set $\mathcal{U} \subset \mathcal{B}_{\text{vol}}^r(\hat{f})$ such that every $\hat{G} \in \mathcal{U}$ is ergodic for \hat{m} and the Lyapunov exponents $\hat{G} \mapsto \lambda_{\pm}(\hat{G}, \hat{m})$ vary continuously and never vanish on \mathcal{U} .*

2.7. Rigidity for symplectic diffeomorphisms. Theorem A is contained in the following result for symplectic toral automorphisms in any dimension.

Let $M = \mathbb{T}^d$ for some even integer $d \geq 4$ and $A : M \rightarrow M$ be a linear automorphism with exactly two eigenvalues on the unit circle. Assume A is *pseudo-Anosov*, that is, A is ergodic (equivalently, no eigenvalue is a root of unity) and the characteristic polynomial $p_A(t)$ is irreducible over the integers and can not be written as a polynomial of t^n for any $n \geq 2$; when $d = 4$ ergodicity implies the other two conditions, cf. [26, Corollary A.7]. It was shown by Rodriguez-Hertz [26] that every pseudo-Anosov linear automorphism with 2-dimensional center is stably ergodic: all nearby volume preserving diffeomorphisms $f : M \rightarrow M$ are ergodic.

Here we also assume that A is *symplectic*, meaning that it preserves some symplectic form ω on the torus M . This implies that the stable (E^s) and unstable (E^u) subspaces of A have the same dimension, and the center subspace E^c is symplectic orthogonal to $E^s \oplus E^u$.

Theorem I. *There exists a neighborhood \mathcal{U} of A in the space of C^∞ symplectic diffeomorphisms on M such that every $f \in \mathcal{U}$ either is non-uniformly hyperbolic or is conjugate to A by a volume preserving homeomorphism. In particular, every $f \in \mathcal{U}$ is Bernoulli.*

The statement remains true for C^k diffeomorphisms, as long as k is large enough for the conclusions of Rodriguez-Hertz [26] to hold ($k \geq 22$ suffices).

2.8. Structure of this paper. In Section 3 we prove Theorems B and C. In Section 4 we apply them to continuous cocycles with holonomies, to deduce Theorem D and Corollary E. In Section 5 we show that holonomies do exist if the cocycle is fiber bunched. Theorem F and Corollary G are proved in Section 6. In Section 7 we prove Theorem H and in Section 8 we explain how to obtain Theorem I. The latter contains Theorem A as a special case.

3. INVARIANCE PRINCIPLE

In this section we prove Theorems B and C. It is no restriction to suppose that the fiber bundle \mathcal{E} is trivial, since the measurable trivialization domains U_n in (2) may always be chosen to be disjoint.

The first step is to reduce the proof to a natural extension situation similar to Example 3.13. As observed by Rokhlin [41, §1-§2], one may find a Lebesgue space (M, \mathcal{B}, μ) and a projection $\pi : \hat{M} \rightarrow M$ such that $\mathcal{B} = \pi_* \mathcal{B}_0$ and $\mu = \pi_* \hat{\mu}$. Here M is the quotient space obtained by identifying any two points of \hat{M} which are not distinguished by any $B_0 \in \mathcal{B}_0$ and \mathcal{B} and μ are characterized by the properties we just stated: $B \in \mathcal{B}$ if and only if $\pi^{-1}(B) \in \mathcal{B}_0$ and then $\mu(B) = \hat{\mu}(\pi^{-1}(B))$. Since \hat{f} is \mathcal{B}_0 -measurable mod 0, there exists a \mathcal{B} -measurable mod 0 transformation $f : M \rightarrow M$ such that $\pi \circ \hat{f} = f \circ \pi$. This transformation, which is usually non-invertible, preserves μ . Let $\mathcal{E} = M \times N$ and $P : \mathcal{E} \rightarrow M$ be the canonical projection. Since the deformation \tilde{F} is \mathcal{B}_0 -measurable mod 0, it may be written as $\tilde{F}_{\hat{x}} = F_{\pi(\hat{x})}$ for some \mathcal{B} -measurable mod 0 fiber bundle morphism $F : \mathcal{E} \rightarrow \mathcal{E}$ over f . Since $\tilde{m} = H_* \hat{m}$ is a \tilde{F} -invariant probability projecting down to $\hat{\mu}$, the probability $m = (\pi \times \text{id})_* \tilde{m}$ is F -invariant and projects down to μ .

Let κ be the dimension of N . Let

$$(9) \quad (F_x^{-1})_* m_{f(x)} = J(x, \cdot) m_x + \eta_x$$

be the Lebesgue decomposition of $(F_x^{-1})_* m_{f(x)}$ relative to m_x : the function

$$(10) \quad J(x, \xi) = \frac{d(F_x^{-1})_* m_{f(x)}(\xi)}{dm_x}$$

is integrable for m_x and the measure η_x is singular with respect to m_x . We call $J : \mathcal{E} \rightarrow [0, \infty)$ the *fibered Jacobian*, and define the *fibered entropy* to be

$$(11) \quad h = h(\tilde{F}, \tilde{m}) = \int -\log J \, dm.$$

The definition (9) implies $\int_{\{J>0\}} J \, dm = \int J \, dm \leq 1$. Then, by Jensen's inequality,

$$(12) \quad \int_{\{J>0\}} -\log J \, dm \geq 0.$$

The definition (11) means that h is the sum of this integral with the term $(+\infty) \cdot m(\{J = 0\})$ with the usual convention that the latter vanishes if $m(\{J = 0\}) = 0$.

Thus, h is always well-defined and non-negative. In our context h is finite, as we shall see later, and so $\{J = 0\}$ always has zero measure.

Proposition 3.1. *Let \hat{m} be an \hat{F} -invariant probability measure projecting down to $\hat{\mu}$ and let $\tilde{m} = H_*\hat{m}$. Then*

$$0 \leq \beta h(\tilde{F}, \tilde{m}) \leq -\kappa \int \min\{0, \lambda_-(\hat{F}, \cdot)\} d\hat{m}.$$

This result may be seen as a coarse fibered version of the Ruelle inequality [42]. Indeed, Ruelle showed that the entropy of a diffeomorphism is bounded above by the integrated sum of the positive Lyapunov exponents. Considering the inverse map, we get that the entropy is also bounded by minus the integrated sum of the negative Lyapunov exponents. In view of this, Proposition 3.1 can probably be refined replacing $\kappa \min\{0, \lambda_-(\hat{F}, \cdot)\}$ by the sum of all negative exponents.

Proposition 3.2. *If $h(\tilde{F}, \tilde{m}) = 0$ then $\hat{x} \mapsto \tilde{m}_{\hat{x}}$ is \mathcal{B}_0 -measurable mod 0.*

Theorem B is an immediate consequence of Propositions 3.1 and 3.2. Indeed, the assumption $\lambda_-(\hat{F}, \cdot) \geq 0$ means that $\min\{0, \lambda_-(\hat{F}, \cdot)\}$ vanishes identically. Then Proposition 3.1 yields $h(\tilde{F}, \tilde{m}) = 0$ and, by Proposition 3.2, it follows that the disintegration $\hat{x} \mapsto \tilde{m}_{\hat{x}}$ is \mathcal{B}_0 -measurable mod 0, as claimed. This reduces the proof of Theorem B to proving Propositions 3.1 and 3.2.

For Theorem C we need the following version of Proposition 3.2 for sequences of measures. In what follows it is understood that $\tilde{m}_k = H_*\hat{m}_k$ and $m_k = (\pi \times \text{id})_*\tilde{m}_k$.

Proposition 3.3. *Let $(\hat{m}_k)_k$ be a sequence of \hat{F} -invariant probability measures on \hat{E} that project down to $\hat{\mu}$ and converge to some probability \hat{m} in the weak* topology. If $h(\tilde{F}_k, \tilde{m}_k)$ converges to 0 when $k \rightarrow \infty$ then the disintegration $\hat{x} \mapsto \tilde{m}_{\hat{x}}$ of $\tilde{m} = H_*\hat{m}$ is \mathcal{B}_0 -measurable mod 0.*

In view of Proposition 3.1, the hypothesis of Theorem C implies that $h(\tilde{F}, \tilde{m}_k)$ converges to 0 when k goes to ∞ . Then we may apply Proposition 3.3 to conclude that the disintegration $\hat{x} \mapsto \tilde{m}_{\hat{x}}$ is \mathcal{B}_0 -measurable mod 0, as claimed. This reduces the proof of Theorem C to proving Propositions 3.1 and 3.3.

3.1. Entropy zero means deterministic. Let us prove Propositions 3.2 and 3.3.

Lemma 3.4. *The disintegrations $\{\tilde{m}_{\hat{x}} : \hat{x} \in \hat{M}\}$ and $\{m_x : x \in M\}$ of \tilde{m} and $m = (\pi \times \text{id})_*\tilde{m}$, respectively, are related by*

$$\tilde{m}_{\hat{x}} = \lim_{n \rightarrow \infty} (F_{x(n)}^n)_* m_{x(n)} \text{ where } x(n) = \pi(\hat{f}^{-n}(\hat{x})), \text{ at } \hat{\mu}\text{-almost every } \hat{x} \in \hat{M}.$$

Proof. Let m_0 be the probability defined on \mathcal{B}_0 by $\pi_* m_0 = m$. The disintegration of m_0 is just $\hat{x} \mapsto m_{\pi(\hat{x})}$. The relation $\pi_* \tilde{m} = m$ implies that $\tilde{m} | \mathcal{B}_0 = m_0$ or, in other words, $E(\hat{x} \mapsto \tilde{m}_{\hat{x}} | \mathcal{B}_0) = [\hat{x} \mapsto m_{\pi(\hat{x})}]$. Next, the relation $\tilde{F}_* \tilde{m} = \tilde{m}$ implies that

$$E(\hat{x} \mapsto \tilde{m}_{\hat{x}} | \hat{f}^n(\mathcal{B}_0)) = E(\hat{x} \mapsto (\hat{F}_{\hat{x}(n)}^n)_* \tilde{m}_{\hat{x}(n)} | \mathcal{B}_0),$$

with $\hat{x}(n) = \hat{f}^{-n}(\hat{x})$, and so

$$E(\hat{x} \mapsto \tilde{m}_{\hat{x}} | \hat{f}^n(\mathcal{B}_0)) = [\hat{x} \mapsto (F_{x(n)}^n)_* m_{x(n)}].$$

Any of these expressions defines a martingale of probability measures, relative to the sequence of σ -algebras $\hat{f}^n(\mathcal{B}_0)$. Since \mathcal{B}_0 is generating and the sequence $\hat{f}^n(\mathcal{B}_0)$ is increasing, the limit of the left hand side is

$$[\hat{x} \mapsto \tilde{m}_{\hat{x}}] = E(\hat{x} \mapsto \tilde{m}_{\hat{x}} \mid \hat{\mathcal{B}}).$$

It follows that $(F_{x(n)}^n)_* m_{x(n)}$ converges and the limit coincides with $\tilde{m}_{\hat{x}}$ at $\hat{\mu}$ -almost every point. \square

Lemma 3.5. *If $h(\tilde{F}, \tilde{m}) = 0$ then $(F_x)_* m_x = m_{f(x)}$ for μ -almost every $x \in M$.*

Proof. The definition (9) implies that $\int J(x, \xi) dm_x(\xi) \leq 1$ for μ -every x . So, by Jensen's inequality, $\int -\log J(x, \xi) dm_x(\xi) \geq 0$ for μ -every x . Moreover, the equalities hold if and only if $J(x, \xi) = 1$ for m_x -almost every ξ . This implies that $h \geq 0$, and $h = 0$ if and only if $J(x, \xi) = 1$ for m_x -almost every ξ and μ -almost every x . In particular, $h = 0$ implies $m_{f(x)} = (F_x)_* m_x$ for μ -almost every x , as claimed. \square

Lemma 3.5 implies $(F_{x(n)}^n)_* m_{x(n)} = m_{x(0)}$ for every $n \geq 0$ and $\hat{\mu}$ -almost every \hat{x} . Then Lemma 3.4 yields $\tilde{m}_{\hat{x}} = m_{x(0)}$ for $\hat{\mu}$ -almost every \hat{x} . Since $x(0) = \pi(\hat{x})$, this implies that $\hat{x} \mapsto \tilde{m}_{\hat{x}}$ is \mathcal{B}_0 -measurable. The proof of Proposition 3.2 is complete.

Next, we prove Proposition 3.3. Let $(F_x^{-1})_* m_{k, f(x)} = J_k(x, \cdot) m_{k, x} + \eta_{k, x}$ be the Lebesgue decomposition for each m_k : in particular, $J_k : \mathcal{E} \rightarrow \mathbb{R}$ is the fibered Jacobian. We denote by $\|\xi\|$ the total variation of a signed measure ξ .

Lemma 3.6. *$\int |J_k(x, \xi) - 1| dm_k(x, \xi) \rightarrow 0$ and $\int \|\eta_{k, x}\| d\mu(x) \rightarrow 0$ when $k \rightarrow \infty$.*

Proof. Since the $m_{k, y}$ are probabilities,

$$\eta_{k, x}(\mathcal{E}_x) = m_{k, f(x)}(\mathcal{E}_{f(x)}) - \int J_k(x, \cdot) dm_{k, x} = \int (1 - J_k(x, \cdot)) dm_{k, x}.$$

Integrating with respect to μ , we obtain $\int \|\eta_{k, x}\| d\mu = \int (1 - J_k) dm_k$ and so the second claim is a consequence of the first one. Next, define $\phi(x) = x - \log(1 + x)$ for $x > -1$. Then $\phi(x) \geq 0$ for all x and, given any $\delta > 0$, there exists $c(\delta) > 0$ such that $\phi(x) \geq c(\delta)|x|$ whenever $|x| \geq \delta$. Let $\delta > 0$ be fixed. Denote $a_k = \int J_k dm_k$ for each $k \geq 0$. Using Jensen's inequality,

$$h(\tilde{F}, \tilde{m}_k) \geq -\log a_k \geq 0,$$

and so a_k converges to 1 when $n \rightarrow \infty$. Assume k is large enough that $h(\tilde{F}, \tilde{m}_k)$ and $a_k - 1$ are both less than $\delta c(\delta)$. Then, by the definition of ϕ ,

$$\int -\log J_k dm_k = \int (1 - J_k) dm_k + \int \phi(J_k - 1) dm_k.$$

The first integral is less than $\delta c(\delta)$ and the second one is $1 - a_k > -\delta c(\delta)$. The third integral is bounded below by

$$\int_{\{|J_k - 1| > \delta\}} \phi(J_k - 1) dm_k \geq c(\delta) \int_{\{|J_k - 1| > \delta\}} |J_k - 1| dm_k.$$

This implies

$$\int |J_k - 1| dm_k \leq \delta + \int_{\{|1 - J_k| > \delta\}} |J_k - 1| dm_k \leq 3\delta$$

for all large k . This completes the proof of the lemma. \square

For each $k \geq 1$, let \check{m}_k be the probability measure on \mathcal{E} that projects down to $\hat{\mu}$ and whose conditional measures along the fibers are given by

$$\check{m}_{k,\hat{x}} = m_{k,x} \quad \text{for all } x = \pi(\hat{x}).$$

Up to taking a subsequence, we may assume \check{m}_k to converge to some measure \check{m} , whose disintegration $\hat{x} \mapsto \check{m}_x$ along the fibers is \mathcal{B}_0 -measurable mod 0. Clearly, $(\pi \times \text{id})_* \check{m}_k = m_k$ for every k . Taking the limit as $k \rightarrow \infty$, we conclude that $\pi_* \check{m} = m$.

Lemma 3.7. *The total variation $\|\tilde{F}_*^{-1} \check{m}_k - \check{m}_k\|$ converges to 0 as $k \rightarrow \infty$.*

Proof. Given any measurable set $B \subset \hat{\mathcal{E}}$, we denote $B_{\hat{x}} = B \cap \hat{\mathcal{E}}_{\hat{x}}$ for each $\hat{x} \in \hat{M}$. Then

$$\begin{aligned} (\tilde{F}_*^{-1} \check{m}_k - \check{m}_k)(B) &= \int \check{m}_{k,\hat{y}}(\tilde{F}(B)_{\hat{y}}) d\hat{\mu}(\hat{y}) - \int \check{m}_{k,\hat{x}}(B_{\hat{x}}) d\hat{\mu}(\hat{x}) \\ &= \int \check{m}_{k,\hat{f}(\hat{x})}(\tilde{F}_{\hat{x}}(B_{\hat{x}})) d\hat{\mu}(\hat{x}) - \int \check{m}_{k,\hat{x}}(B_{\hat{x}}) d\hat{\mu}(\hat{x}) \end{aligned}$$

because $\hat{\mu}$ is invariant under \hat{f} . Since $\tilde{F}_{\hat{x}}$ and $\check{m}_{k,\hat{x}}$ are both \mathcal{B}_0 -measurable, the last term may be rewritten as

$$\begin{aligned} \int m_{k,f(x)}(F_x(B_{\hat{x}})) d\hat{\mu}(\hat{x}) - \int m_{k,x}(B_{\hat{x}}) d\hat{\mu}(\hat{x}) \\ = \int \left(\int_{B_{\hat{x}}} (J_k(x, \cdot) - 1) dm_{k,x} + \eta_{k,x}(B_{\hat{x}}) \right) d\hat{\mu}(\hat{x}) \end{aligned}$$

These relations imply that

$$|(\tilde{F}_*^{-1} \check{m}_k - \check{m}_k)(B)| \leq \int \left(\int |J_k(x, \cdot) - 1| dm_{k,x} + \|\eta_{k,x}\| \right) d\hat{\mu}(\hat{x})$$

for every $B \subset \hat{\mathcal{E}}$, and so $\|\tilde{F}_*^{-1} \check{m}_k - \check{m}_k\| \leq \int |J_k - 1| dm_k + \int \|\eta_{k,x}\| d\mu(x)$. Now the claim follows from Lemma 3.6. \square

Taking the limit in Lemma 3.7 we conclude that the measure \check{m} is invariant under \tilde{F} . It follows that $\check{m} = \tilde{m}$: any two \tilde{F} -invariant measures that project down to m under π must coincide, because the σ -algebra \mathcal{B}_0 is generating. This proves that the disintegration of \check{m} is \mathcal{B}_0 -measurable mod 0, as claimed. The proof of Proposition 3.3 is complete.

3.2. Entropy is smaller than exponents. We are left to prove Proposition 3.1. We begin by reducing the proof to the ergodic case. Let $\{\hat{m}_\alpha\}$ be the ergodic decomposition of \hat{m} and $d\alpha$ denote the corresponding quotient measure:

$$\int \varphi d\hat{m} = \int \left(\int \varphi d\hat{m}_\alpha \right) d\alpha$$

for any integrable function φ . Then $\tilde{m}_\alpha = H_* \hat{m}_\alpha$ and $m_\alpha = (\pi \times \text{id})_* \tilde{m}_\alpha$ define the ergodic decompositions of $\tilde{m} = H_* \hat{m}$ and $m = (\pi \times \text{id})_* \tilde{m}$, respectively, with the same quotient measure. If $\lambda_-(\hat{F}, \hat{x}, \xi) \geq 0$ at \hat{m} -almost every point then the same is true at \hat{m}_α -almost every point, for $d\alpha$ -almost every ergodic component. Assuming the proposition holds for ergodic measures, it follows that

$$0 \leq \beta \int -\log J dm_\alpha \leq -\kappa \int \min\{0, \lambda_-(\hat{F}, \cdot)\} d\hat{m}_\alpha$$

for $d\alpha$ -almost every α . Integrating with respect to $d\alpha$, we obtain that

$$0 \leq \beta h(\tilde{F}, \tilde{m}) \leq -\kappa \int \min\{0, \lambda_-(\hat{F}, \cdot)\} d\tilde{m},$$

as claimed. Hence, it is no restriction to assume that \hat{m} is ergodic for \hat{F} , and we do so in what follows. Then \tilde{m} and m are ergodic for \tilde{F} and F , respectively. Moreover, $\min\{0, \lambda_-(\hat{F}, \cdot)\}$ is constant \hat{m} -almost everywhere. Let $-\lambda$ denote this constant.

Now we begin the proof of the proposition in the ergodic case. Given $\varepsilon > 0$, define $J_\varepsilon = J + \varepsilon$ and $h_\varepsilon = -\int \log J_\varepsilon dm$. Notice that $h_\varepsilon \rightarrow h$ as $\varepsilon \rightarrow 0$, by the monotone convergence theorem. Our goal is to prove that $h \leq \kappa\beta^{-1}\lambda$. The proof is by contradiction. Assume this inequality is false. Then we may choose some small $\varepsilon > 0$ such that

$$(13) \quad h_\varepsilon - 10\varepsilon \geq \kappa\beta^{-1}(\lambda + 10\varepsilon).$$

Lemma 3.8. *There exists a sequence of countable partitions $P_n(x, \cdot)$ of each fiber \mathcal{E}_x , depending measurably on $x \in M$, a sequence of measurable subsets W_n of \mathcal{E} with $m(W_n) \rightarrow 1$ such that, for every large n ,*

- (a) $\text{diam } P_n(x, \xi) \leq e^{-\beta^{-1}(\lambda+5\varepsilon)n}$ for every $(x, \xi) \in \mathcal{E}$
- (b) each $W_n \cap \mathcal{E}_x$ is covered by not more than $e^{\kappa\beta^{-1}(\lambda+8\varepsilon)n}$ atoms of $P_n(x, \cdot)$
- (c) $m_x(\partial P_n(x, \xi)) = 0$ for every $(x, \xi) \in \mathcal{E}$.

Proof. Since N is a manifold, we may choose a sequence of countable partitions Q_n with relatively compact atoms with diameter bounded by $e^{-\beta^{-1}(\lambda+6\varepsilon)n}$, and an increasing sequence of subsets V_n exhausting N and such that V_n is covered by not more than $e^{\kappa\beta^{-1}(\lambda+8\varepsilon)n}$ atoms of Q_n . Of course, we may take these to be the first atoms of Q_n with respect to some ordering of the partition. This defines ordered countable partitions $Q_n(x, \cdot)$ of each fiber \mathcal{E}_x , and sets $W_n \subset \mathcal{E}$ exhausting every fiber, such that $\text{diam } Q_n(x, \xi) \leq \text{const } e^{-\beta^{-1}(\lambda+6\varepsilon)n}$ for every (x, ξ) , and every $W_n \cap \mathcal{E}_x$ is covered by the first $e^{\kappa\beta^{-1}(\lambda+8\varepsilon)n}$ atoms of $Q_n(x, \cdot)$. For each atom Q of $Q_n(x, \cdot)$, let B_1, \dots, B_k be a finite covering of the boundary of Q by open sets with diameter less than $e^{-\beta^{-1}(\lambda+6\varepsilon)n}$ and such that $m_x(\partial B_j) = 0$ for all j . Let $\tilde{Q}_n(x, \cdot)$ be the family of all $\tilde{Q} = Q \cup B_1 \cup \dots \cup B_k$ obtained in this way. Notice that $m_x(\partial \tilde{Q}) = 0$. Removing from each \tilde{Q} the union of the elements of $\tilde{Q}_n(x, \cdot)$ that precede it, relative to the ordering inherited from $Q_n(x, \cdot)$, one obtains a new ordered partition $P_n(x, \cdot)$ such that the diameter of its atoms is bounded by $\text{const } e^{-\beta^{-1}(\lambda+6\varepsilon)n}$, the first $e^{\kappa\beta^{-1}(\lambda+8\varepsilon)n}$ atoms cover $W_n \cap \mathcal{E}_x$, and the boundary of every atom has zero m_x -measure. Replacing 6ε by 5ε in the exponent and assuming n is large, one gets rid of the constant. This finishes the construction. \square

For $0 \leq k < n$, define $P_{n,k}(\cdot, \cdot)$ as the pullback of $P_n(\cdot, \cdot)$ by F^{n-k} , that is,

$$P_{n,k}(x, \xi) = (F_x^{n-k})^{-1}(P_n(F^{n-k}(x, \xi))).$$

Let also $P_{n,n}(\cdot, \cdot) = P_n(\cdot, \cdot)$. For each $0 \leq k < n$, define

$$J_n(x, \xi) = \frac{m_{f^n(x)}(P_n(F^n(x, \xi)))}{m_x(P_{n,0}(x, \xi))} \quad \text{and} \quad J_{n,k}(x, \xi) = \frac{m_{f(x)}(P_{n,k+1}(F(x, \xi)))}{m_x(P_{n,k}(x, \xi))}.$$

Then $J_n(x, \xi) = \prod_{k=0}^{n-1} J_{n,k}(F^k(x, \xi))$. Moreover, let

$$J_{n,k,\varepsilon} = J_{n,k} + \varepsilon \quad \text{and} \quad J_{n,\varepsilon}(x, \xi) = \prod_{k=0}^{n-1} J_{n,k,\varepsilon}(F^k(x, \xi)).$$

Notice that $J_{n,\varepsilon} \geq J_n$ because $J_{n,k,\varepsilon} \geq J_{n,k}$ for every k . The key ingredient in the proof of Proposition 3.1 is the following lemma, whose proof we postpone for a while:

Lemma 3.9. *We have $\lim_{n \rightarrow \infty} \sup_{0 \leq k < n} \|\log J_{n,k,\varepsilon} - \log J_\varepsilon\|_{L^1(m)} = 0$.*

As a consequence of this lemma and the ergodic theorem,

$$\lim \frac{1}{n} \log J_{n,\varepsilon} = \lim \frac{1}{n} \sum_{k=0}^{n-1} \log J_\varepsilon \circ F^k = \int \log J_\varepsilon dm = -h_\varepsilon$$

in $L^1(m)$ and, hence, in measure. In particular, for every large n there exists $E_n \subset \mathcal{E}$ with $m(E_n) \geq 1 - \varepsilon$ such that

$$\frac{1}{n} \log J_n(x, \xi) \leq \frac{1}{n} \log J_{n,\varepsilon}(x, \xi) \leq -h_\varepsilon + 5\varepsilon \quad \text{for all } (x, \xi) \in E_n.$$

Using Lemma 3.8 and the definition of J_n , we conclude that the fiber of $F^n(E_n) \cap W_n$ over $f^n(x)$ is covered by at most $e^{\kappa\beta^{-1}(\lambda+8\varepsilon)n}$ atoms of $P_n(f^n(x), \cdot)$ all with $m_{f^n(x)}$ -measure at most $e^{(-h_\varepsilon+5\varepsilon)n}$. By (13), this implies $m(F^n(E_n) \cap W_n)$ goes to zero as $n \rightarrow \infty$, contradicting the fact that both $m(W_n)$ and $m(E_n)$ are close to 1. This contradiction reduces the proof of Proposition 3.1 to proving Lemma 3.9.

For every $l \geq 1$, define $\omega_l(\hat{x}, \xi) = \log \|(D\hat{F}_x^l(\xi))^{-1}\|^{-1}$ and

$$\Omega_l(\hat{x}, \xi) = \liminf_{n \rightarrow \infty} \inf_{0 \leq k < n} \frac{1}{n} \sum_{j=k}^{n-1} \frac{1}{l} \omega_l(\hat{F}^{jl}(\hat{x}, \xi)).$$

Lemma 3.10. *We have $\sup_{l \geq 1} \Omega_l(\hat{x}, \xi) \geq -\lambda$ for every (\hat{x}, ξ) in some full \hat{m} -measure set $\hat{Z} \subset \hat{\mathcal{E}}$.*

Proof. We begin by claiming that $\sup_{l \geq 1} \Omega_l$ is constant along orbits. Indeed, since the norms of $DF^{\pm 1}$ are uniformly bounded, there exists some constant $A > 0$ such that $|\omega_l(\hat{F}(\hat{y}, \eta)) - \omega_l(\hat{y}, \eta)| \leq A$ for every (\hat{y}, η) . This implies

$$(14) \quad |\Omega_l(\hat{F}(\hat{x}, \xi)) - \Omega_l(\hat{x}, \xi)| \leq \frac{A}{l} \quad \text{for every } (\hat{x}, \xi).$$

Similarly, since $\omega_{2l}(\hat{y}, \eta) \geq \omega_l(\hat{y}, \eta) + \omega_l(\hat{F}(\hat{y}, \eta))$ for every (\hat{y}, η) , we have $\Omega_{2l}(\hat{x}, \xi) \geq \Omega_l(\hat{x}, \xi)$ for every (\hat{x}, ξ) and every $l \geq 1$. This implies

$$(15) \quad \sup_l \Omega_l(\hat{x}, \xi) = \limsup_{l \rightarrow \infty} \Omega_l(\hat{x}, \xi) \quad \text{for every } (\hat{x}, \xi).$$

The relations (14) and (15) imply our claim.

Next, by ergodicity and the definition of smallest Lyapunov exponent,

$$\lambda_- = \lim_l \frac{1}{l} \omega_l(\hat{x}, \xi) = \sup_l \frac{1}{l} \omega_l(\hat{x}, \xi) \quad \text{for } \hat{m}\text{-almost every } (\hat{x}, \xi).$$

Given $\varepsilon > 0$, fix $s \geq 1$ large enough so that $\hat{m}(E_{s,\varepsilon}) > 1 - \varepsilon$, where

$$E_{s,\varepsilon} = \{(\hat{y}, \eta) : \frac{1}{s} \omega_s(\hat{y}, \eta) \geq \lambda_- - \varepsilon\}.$$

By ergodicity, for \hat{m} -almost every (\hat{x}, ξ) the number of iterates $0 \leq i < ns$ for which $F^i(\hat{x}, \xi) \notin E_{s,\varepsilon}$ is less than $2\varepsilon ns$, assuming n is large enough. Then there exists $0 \leq r < s$ such that the number of iterates $0 \leq j < n$ for which $F^{js+r}(\hat{x}, \xi) \notin E_{s,\varepsilon}$ is less than $2\varepsilon n$. Let $B > 0$ be an upper bound for the absolute value of $\log \|D\hat{F}^{-1}\|$. Then $|\omega_l(\hat{y}, \eta)| \leq Bl$ for every (\hat{y}, η) and every $l \geq 1$. It follows that, given any $0 \leq k < n$,

$$\begin{aligned} \frac{1}{n} \sum_{j=k}^{n-1} \frac{1}{s} \omega_s(\hat{F}^{js+r}(\hat{x}, \xi)) &\geq \frac{1}{n} \left[(\lambda_- - \varepsilon) \#\{k \leq j < n : \hat{F}^{js+r}(\hat{x}, \xi) \in E_{s,\varepsilon}\} - 2n\varepsilon B \right] \\ &\geq -\lambda - \varepsilon(1 + 2B). \end{aligned}$$

Since this holds for every $0 \leq k < n$ and every n sufficiently large, we conclude that

$$\sup_l \Omega_l(\hat{F}^r(\hat{x}, \xi)) \geq \Omega_s(\hat{F}^r(\hat{x}, \xi)) \geq -\lambda - \varepsilon(1 + 2B).$$

So, in view of the claim in the first paragraph, $\sup_l \Omega_l(\hat{x}, \xi) \geq -\lambda - \varepsilon(1 + 2B)$. Since $\varepsilon > 0$, this completes the proof of the lemma. \square

The next results provides the main estimate for the proof of Lemma 3.9. Let

$$d_x(\xi, \eta) = \sup\{d(H_{\hat{x}}^{-1}(\xi), H_{\hat{x}}^{-1}(\eta)) : \hat{x} \in \pi^{-1}(x)\}$$

for each $x \in M$ and $\xi, \eta \in \mathcal{E}_x$. This defines a metric d_x on each fiber \mathcal{E}_x which, by (3), relates to the Riemannian distance d through $d_x(\xi, \eta) \leq Bd(\xi, \eta)^\beta$. Then let $\Delta_{n,k}(x, \xi)$ denote the d_x -diameter of each atom $P_{n,k}(x, \xi)$.

Lemma 3.11. *We have $\lim_{n \rightarrow \infty} \sup_{0 \leq k < n} \Delta_{n,k} = 0$ at m -almost every point.*

Proof. It suffices to show that $\lim_{n \rightarrow \infty} \sup_{0 \leq k < n} \Delta_{n,k}(x, \xi) = 0$ holds for every (x, ξ) in the full m -measure set $Z = (\pi \times \text{id})(\hat{Z})$. To this end, consider any $\hat{x} \in \pi^{-1}(x) \cap \hat{Z}$. We claim that, given any $\delta > 0$, there exists $m_0 \geq 1$ such that (balls are with respect to the Riemannian metric along the fiber)

$$(16) \quad \hat{F}^m(B_\delta(\hat{x}, \xi)) \supset B_{e^{-(\lambda+4\varepsilon)m}}(\hat{F}^m(\hat{x}, \xi))$$

for every $m > m_0$. Assume this fact for a while. It implies that

$$(17) \quad \hat{F}^{n-k}(B_\delta(\hat{x}, \xi)) \supset B_{e^{-(\lambda+4\varepsilon)n}}(\hat{F}^{n-k}(\hat{x}, \xi))$$

for all $0 \leq k < n$, as long as n is large enough: when $n - k \geq m_0$ this is a direct consequence of (16); otherwise, use the fact that $\hat{F}^{\pm j}$, $1 \leq j \leq m_0$ are uniformly continuous along fibers, and take n to be large enough. By the Hölder property (3) and Lemma 3.8, we also have $H_{\hat{x}}^{-1}(P_n(F^{n-k}(x, \xi))) \subset B_{e^{-(\lambda+4\varepsilon)n}}(\hat{F}^{n-k}(\hat{x}, \xi))$, as long as n is large enough (to make the radius of the last neighborhood sufficiently small). Combined with (17), this gives

$$P_n(F^{n-k}(x, \xi)) \subset F_x^{n-k}(H_{\hat{x}}(B_\delta(\hat{x}, \xi))), \text{ that is, } P_{n,k}(x, \xi) \subset H_{\hat{x}}(B_\delta(\hat{x}, \xi))$$

for all $0 \leq k < n$, as long as n is large enough. Since $H_{\hat{x}}$ is continuous, this implies the conclusion of the lemma.

To prove the claim (16), begin by fixing $l \geq 1$ such that $\Omega_l(\hat{x}, \xi) \geq -(\lambda + \varepsilon)$. Then define $\delta_{k,n}$, $0 \leq k \leq n$ by

$$\log \delta_{n,n} = -(\lambda + 3\varepsilon)ln \quad \text{and} \quad \log \delta_{k,n} = \log \delta_{k+1,n} + \varepsilon - \omega_l(\hat{F}^{kl}(\hat{x}, \xi)).$$

Then $\log \delta_{k,n} \leq -\varepsilon n$ for all $0 \leq k \leq n$, because

$$\log \delta_{k,n} = \log \delta_{n,n} + \sum_{j=k}^{n-1} \varepsilon - \omega_l(\hat{F}^{jl}(\hat{x}, \xi)) \leq -(\lambda + 3\varepsilon)ln + (\varepsilon - \Omega_l(\hat{x}, \xi))ln.$$

Since the derivatives $D\hat{F}_y^{\pm l}$ are uniformly bounded and uniformly continuous, we conclude from the definition of ω_l that

$$\hat{F}^l(B_{\delta_{k,n}}(\hat{F}^{kl}(\hat{x}, \xi))) \supset B_{\delta_{k+1,n}}(\hat{F}^{(k+1)l}(\hat{x}, \xi))$$

for every $0 \leq k < n$, as long as n is large enough (to make $e^{-\varepsilon n}$ sufficiently small). In particular,

$$\hat{F}^{ln}(B_{\delta_{n,0}}(\hat{x}, \xi)) \supset B_{\delta_{n,n}}(\hat{F}^n(\hat{x}, \xi)).$$

This gives a version of (16) for the iterates that are multiples of l : given any $\delta > 0$ there exists $n_0 \geq 1$ such that

$$\hat{F}^{ln}(B_\delta(\hat{x}, \xi)) \supset B_{e^{-(\lambda+3\varepsilon)ln}}(\hat{F}^{ln}(x, \xi))$$

for every $n \geq n_0$. To complete the proof it suffices to note that, since the derivatives $D\hat{F}_y^{\pm j}$, $0 \leq j < l$ are bounded,

$$\hat{F}^j(B_{e^{-(\lambda+3\varepsilon)ln}}(\hat{F}^{ln}(x, \xi))) \supset B_{e^{-(\lambda+4\varepsilon)ln}}(\hat{F}^{ln+j}(x, \xi))$$

for all $0 \leq j < l$, as long as n is large enough. This finishes the proof of (16) and of the lemma. \square

We also need the following abstract result:

Lemma 3.12. *Let K be a complete metric space and μ_0 and μ_1 be probability measures on K with $\mu_1 \geq \alpha\mu_0$ for some $\alpha > 0$. Let $\phi = d\mu_1/d\mu_0$ and, given any countable partition P of K , define*

$$\phi_P(x) = \frac{\mu_1(P(x))}{\mu_0(P(x))}.$$

Then $\int \log \phi(x) d\mu_0(x) \leq \int \log \phi_P(x) d\mu_0(x) \leq 0$. Moreover, given $\varepsilon > 0$ there exists $\delta > 0$ such that $\|\log \phi_P - \log \phi\|_{L^1(\mu_0)} \leq \varepsilon$ for any countable partition P of K such that the total measure of the atoms with diameter larger than δ is smaller than δ .

Proof. By convexity, $\int \log \phi_P d\mu_0 \leq \log \int \phi_P d\mu_0 = 0$. Similarly,

$$\int_{P(x)} \log \phi d\mu_0 \leq \mu_0(P(x)) \log \phi_P(x)$$

for every atom $P(x)$, and so $\int \log \phi d\mu_0 \leq \int \log \phi_P d\mu_0$. This proves the first claim.

Next, notice that the functions ϕ_P satisfy a uniform integrability condition: for all $X \subset K$ with $\mu_0(X) < 1/e$,

$$(18) \quad \int_X |\log \phi_P| d\mu_0 \leq -\mu_0(X)(\log \mu_0(X) + \log \alpha).$$

Indeed, the assumption implies $-\log \phi_P \leq -\log \alpha$ and so the claim is trivial if $\log \phi_P$ happens to be negative on X . When $\log \phi_P \geq 0$ on the set X , the claim follows from convexity:

$$\int_X \log \phi_P d\frac{\mu_0}{\mu_0(X)} \leq \log \int_X \phi_P d\frac{\mu_0}{\mu_0(X)} \leq \log \frac{\mu_1(\tilde{X})}{\mu_0(X)} \leq \log \frac{1}{\mu_0(X)}$$

(\tilde{X} denotes the union of all atoms of P that intersect X). The general case is handled by splitting X into two subsets where $\log \phi_P$ has constant sign.

We also use the following fact: if R refines Q then

$$(19) \quad \|\log \phi_R - \log \phi_Q\|_{L^1(\mu_0)} \leq \|\log \phi - \log \phi_Q\|_{L^1(\mu_0)}.$$

To see that this is so, write

$$\int |\log \phi_R - \log \phi_Q| d\mu_0 = \sum_{r \subset q} \int_r |\log \phi_R - \log \phi_Q| d\mu_0,$$

where the sum is over the pairs of atoms $r \in R$ and $q \in Q$ with $r \subset q$. Since ϕ_R and ϕ_Q are constant on r , this may be rewritten as

$$\begin{aligned} \sum_{r \subset q} \mu_0(r) |\log \phi_R - \log \phi_Q| &= \sum_{r \subset q} \left| \int_r \log \phi d\mu_0 - \mu_0(r) \log \phi_Q \right| \\ &\leq \sum_{r \subset q} \int_r |\log \phi - \log \phi_Q| d\mu_0. \end{aligned}$$

The combination of these two relations proves (19).

Let $(Q_n)_n$ be any refining sequence of partitions with diameter decreasing to zero. Then $\phi_{Q_n} \rightarrow \phi$ at μ_0 -almost every point (martingale convergence theorem). By uniform integrability (18), it follows that $\log \phi_{Q_n} \rightarrow \log \phi$ in $L^1(\mu_0)$. Assume, in what follows, that the sequence was chosen so that $\mu_0(\partial Q_n(x)) = 0$ for every x and every n (this can be obtained using the argument in Lemma 3.8(c)). Given $\varepsilon > 0$, fix n sufficiently large so that

$$\|\log \phi_{Q_n} - \log \phi\|_{L^1(\mu_0)} < \varepsilon/4.$$

Let $R = P \vee Q_n$ (the coarsest partition that refines both P and Q_n) and let Δ be the set of all x such that $P(x) \not\subset Q_n(x)$. By (19),

$$\|\log \phi_R - \log \phi\|_{L^1(\mu_0)} < \varepsilon/2.$$

Choosing $\delta > 0$ small, we also ensure that the measure of Δ is small, so that $-\mu_0(\Delta)(\log \mu_0(\Delta) + \log \alpha) < \varepsilon/4$. Clearly, $P(x) = R(x)$ for every x in the complement of Δ . So, using (18),

$$\|\log \phi_P - \log \phi_R\|_{L^1(\mu_0)} = \int_{\Delta} |\log \phi_P - \log \phi_R| d\mu_0 < \varepsilon/2.$$

From these two relations it follows that $\|\log \phi_P - \log \phi\|_{L^1(\mu_0)} < \varepsilon$, as claimed in the second part of the lemma. \square

We are ready to prove Lemma 3.9. Lemmas 3.8 and 3.11 ensure that the hypotheses of Lemma 3.12 are satisfied for $K = N$ and $\mu_0 = m_x$ and $\mu_1 = (F_x^{-1})_* m_{f(x)} + \varepsilon m_x$ ($\alpha = \varepsilon$) and $P = P_{n,k}(x, \cdot)$. Notice that

$$\phi = J(x, \cdot) + \varepsilon = J_\varepsilon \quad \text{and} \quad \phi_P = J_{n,k,\varepsilon}(x, \cdot).$$

From Lemma 3.12 we conclude that

$$\sup_k \|\log J_{n,k,\varepsilon} - \log J_\varepsilon\|_{L^1(m_x)} \rightarrow 0$$

for μ -almost every x . Since $\int |\log J_{n,k,\varepsilon} - \log J_\varepsilon| dm_x \leq -2 \log \varepsilon$ for μ -almost every x , this implies that $\|\log J_{n,k,\varepsilon} - \log J_\varepsilon\|_{L^1(m)} \rightarrow 0$, as claimed in Lemma 3.9.

The proof of Proposition 3.1 is complete, finishing the proofs of Theorems B/C.

3.3. Examples. A few simple examples illustrate the contents of Theorems B/C.

Example 3.13. Given any (non-invertible) measure-preserving map $f : M \rightarrow M$ in a probability space (M, \mathcal{B}, μ) , define \hat{M} to be the space of all sequences $(x_n)_{n \leq 0}$ in M such that $f(x_n) = x_{n+1}$ for all $n < 0$, and consider the *natural extension* of f ,

$$\hat{f} : \hat{M} \rightarrow \hat{M}, \quad \hat{f}(\dots, x_n, \dots, x_0) = (\dots, x_n, \dots, x_0, f(x_0)).$$

Then \hat{f} is invertible and $\pi \circ \hat{f} = f \circ \pi$, where $\pi : \hat{M} \rightarrow M$ is the projection to the zeroth term. Denote $\mathcal{B}_0 = \pi^{-1}(\mathcal{B})$ and let $\hat{\mathcal{B}}$ be the σ -algebra on \hat{M} generated by the iterates $\hat{f}^n(\mathcal{B}_0)$, $n \geq 0$. Then \hat{f} is measurable with respect to \mathcal{B}_0 and to $\hat{\mathcal{B}}$. Let μ_0 be the probability measure defined on \mathcal{B}_0 by $\pi_*\mu_0 = \mu$. There is a unique \hat{f} -invariant probability $\hat{\mu}$ on $(\hat{M}, \hat{\mathcal{B}})$ such that $\pi_*\hat{\mu} = \mu$: it is characterized by

$$(20) \quad E(\hat{\mu} \mid \hat{f}^n(\mathcal{B}_0)) = \hat{f}_*^n \mu_0 \quad \text{for every } n \geq 0.$$

To any smooth cocycle $F : \mathcal{E} \rightarrow \mathcal{E}$ over f , defined on a fiber bundle $P : \mathcal{E} \rightarrow M$, we may associate the smooth cocycle $\hat{F} : \hat{\mathcal{E}} \rightarrow \hat{\mathcal{E}}$ over \hat{f} defined by $\hat{\mathcal{E}}_{\hat{x}} = \mathcal{E}_{\pi(\hat{x})}$ and $\hat{F}_{\hat{x}} = F_{\pi(\hat{x})}$. Their extremal Lyapunov exponents are related by

$$\lambda_{\pm}(\hat{F}, \hat{x}, \hat{\xi}) = \lambda_{\pm}(F, \pi(\hat{x}), \hat{\xi}).$$

Clearly, $\hat{x} \mapsto \hat{F}_{\hat{x}}$ is \mathcal{B}_0 -measurable. We denote by $\pi \times \text{id}$ the natural projection from $\hat{\mathcal{E}}$ to \mathcal{E} (this terminology is motivated by the case when $\hat{\mathcal{E}} = \hat{M} \times N$ and $\mathcal{E} = M \times N$). Given any F -invariant probability m , there is exactly one \hat{F} -invariant probability \hat{m} with $(\pi \times \text{id})_*\hat{m} = m$: it is characterized by

$$(21) \quad E(\hat{x} \mapsto \hat{m}_{\hat{x}} \mid \hat{f}^n(\mathcal{B}_0)) = [\hat{x} \mapsto (\hat{F}_{\hat{x}}^n)_* m_{\pi(\hat{x})}] \quad \text{for every } n \geq 0$$

(see Lemma 3.4 below), where $\{\hat{m}_{\hat{x}} : \hat{x} \in \hat{M}\}$ and $\{m_x : x \in M\}$ are the disintegrations of \hat{m} and m , respectively. If $P_*m = \mu$ then $\hat{P}_*\hat{m} = \hat{\mu}$.

Example 3.14. Ledrappier [33] deals with the particular case when the cocycle is actually linear or, more precisely, projective: $\mathcal{E} = \hat{M} \times \mathbb{P}(\mathbb{R}^d)$ and each $\hat{F}_{\hat{x}}$ is the diffeomorphism induced on the projective space $N = \mathbb{P}(\mathbb{R}^d)$ by some linear map $A(\hat{x}) \in \text{GL}(d, \mathbb{R})$. Denote

$$\lambda_+(\hat{x}) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \|A^n(\hat{x})\| \quad \text{and} \quad \lambda_-(\hat{x}) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \|A^n(\hat{x})^{-1}\|^{-1}.$$

The subadditive ergodic theorem [31] ensures that these two limits exist almost everywhere, and it is clear that $\lambda_+(\hat{x}) \geq \lambda_-(\hat{x})$ at μ -almost every \hat{x} . Theorem 1 in [33] assumes that

$$(22) \quad \int \lambda_+ d\hat{\mu} = \int \lambda_- d\hat{\mu}$$

or, equivalently, $\lambda_+(\hat{x}) = \lambda_-(\hat{x})$ for $\hat{\mu}$ -almost every \hat{x} . This implies the hypothesis of Theorem B. To see this, notice that, locally, the points of $\mathbb{P}(\mathbb{R}^d)$ may be represented by unit vectors $\hat{\xi}$. Then

$$\hat{F}_{\hat{x}}^n(\hat{\xi}) = \frac{A^n(\hat{x})\hat{\xi}}{\|A^n(\hat{x})\hat{\xi}\|}$$

for every \hat{x} , $\hat{\xi}$, and n . It follows that,

$$D\hat{F}_{\hat{x}}^n(\hat{\xi})\hat{\xi} = \frac{\text{proj}_{A^n(\hat{x})\hat{\xi}}(A^n(\hat{x})\hat{\xi})}{\|A^n(\hat{x})\hat{\xi}\|},$$

where $\text{proj}_u v = v - u(u \cdot v)/(u \cdot u)$ is the projection of v to the orthogonal complement of u . This implies that

$$(23) \quad \|D\hat{F}_x^n(\hat{\xi})\| \leq \|A^n(\hat{x})\|/\|A^n(\hat{x})\hat{\xi}\| \leq \|A^n(\hat{x})\|\|A^n(\hat{x})^{-1}\|$$

for every $x, \hat{\xi}$, and n . Replacing n by $-n$ and $(\hat{x}, \hat{\xi})$ by an appropriate iterate, it also follows that

$$(24) \quad \|D\hat{F}_x^n(\hat{\xi})^{-1}\| \leq \|A^n(\hat{x})^{-1}\|\|A^n(\hat{x})\|$$

for every $x, \hat{\xi}$, and n . The last two inequalities imply that

$$\lambda_-(\hat{x}) - \lambda_+(\hat{x}) \leq \lambda_-(\hat{F}, \hat{x}, \hat{\xi}) \leq \lambda_+(\hat{F}, \hat{x}, \hat{\xi}) \leq \lambda_+(\hat{x}) - \lambda_-(\hat{x}).$$

Hence, (22) implies $\lambda_+(\hat{F}, \hat{x}, \hat{\xi}) = \lambda_-(\hat{F}, \hat{x}, \hat{\xi}) = 0$ for m -almost every $(\hat{x}, \hat{\xi})$.

Similar observations apply in the more general case of projective cocycles on *Grassmannian bundles*, such as considered in [5, 6].

Example 3.15. Let $F : M \times \mathbb{P}(\mathbb{R}^2) \rightarrow M \times \mathbb{P}(\mathbb{R}^2)$ be the projective cocycle defined by some $A : M \rightarrow \text{SL}(2, \mathbb{R})$ over a non-invertible system (f, μ) . Let \hat{F} and $(\hat{f}, \hat{\mu})$ be the natural extensions and the σ -algebra \mathcal{B}_0 be as in Example 3.13. Assume the Lyapunov exponents are distinct

$$\lambda_-(\hat{x}) < \lambda_+(\hat{x}) \quad \text{at almost every point}$$

and let $E_{\hat{x}}^-$ and $E_{\hat{x}}^+$ be the Oseledets subspaces, viewed as elements of the projective space. Notice that

$$\lambda_-(\hat{F}, \hat{x}, \hat{\xi}) = \lambda_+(\hat{F}, \hat{x}, \hat{\xi}) = \begin{cases} \lambda_+(\hat{x}) - \lambda_-(\hat{x}) & \text{for } \hat{\xi} = E_{\hat{x}}^- \\ \lambda_-(\hat{x}) - \lambda_+(\hat{x}) & \text{for } \hat{\xi} = E_{\hat{x}}^+. \end{cases}$$

Consider the \hat{F} invariant measures m^- and m^+ whose conditional probabilities along the fibers are the Dirac masses at $\hat{\xi} = E_{\hat{x}}^-$ and $\hat{\xi} = E_{\hat{x}}^+$, respectively, and whose projections down to M coincide with $\hat{\mu}$. Then $\lambda_-(\hat{F}, \hat{x}, \hat{\xi}) > 0$ at m^- -almost every point and so we may use Theorem D to conclude that the disintegration $\hat{x} \mapsto \delta_{E_{\hat{x}}^-}$ is \mathcal{B}_0 -measurable. This conclusion is also an immediate consequence of the observation that the contracting subspace $E_{\hat{x}}^-$ depends only on the future iterates. On the other hand, $\lambda_+(\hat{F}, \hat{x}, \hat{\xi}) < 0$ at m^- -almost every point and yet $\hat{x} \mapsto \delta_{E_{\hat{x}}^+}$ is usually not \mathcal{B}_0 -measurable: the expanding subspace is determined by the past, not the future iterates of the cocycle.

4. COCYCLES WITH INVARIANT HOLONOMIES

For simplicity, **from now on we write** $M, \mathcal{B}, \mu, f, \mathcal{E}, P, F, m$ **in the place of** $\hat{M}, \hat{\mathcal{B}}, \hat{\mu}, \hat{f}, \hat{\mathcal{E}}, \hat{P}, \hat{F}, \hat{m}$. In this section we prove Theorem D and Corollary E.

Let M be a metric space, \mathcal{E} be a continuous fiber bundle endowed with a continuous Riemannian metric, and F be a continuous smooth cocycle. We assume that μ is a Borel measure on M which gives full measure to some σ -compact set, that is, some countable union of compact subsets of M . This is automatic in several cases of interest, for instance when M is a Polish space.

4.1. Holonomy invariance. An s -lamination for the transformation $f : M \rightarrow M$ is a partition $W^s = \{W^s(x) : x \in M\}$ of M such that there exist $\varepsilon, K, \tau > 0$, a family $W_\varepsilon^s = \{W_\varepsilon^s(x) : x \in M\}$ of closed subsets of M , and a function $\nu(\cdot)$ such that

- (sl1) $W_\varepsilon^s(x) = \{y \in W^s(x) : d(f^n(x), f^n(y)) \leq \varepsilon \text{ for all } n \geq 0\}$
- (sl2) $W^s(x) = \cup_{n \geq 0} f^{-n}(W_\varepsilon^s(f^n(x)))$;
- (sl3) $\text{graph}(W_\varepsilon^s) := \{(x, y) : y \in W_\varepsilon^s(x)\}$ is a closed subset of $M \times M$;
- (sl4) $d(f(y_1), f(y_2)) \leq \nu(x)d(y_1, y_2)$ for all $y_1, y_2 \in W_\varepsilon^s(x)$;
- (sl5) $\nu_n(x) := \nu(f^{n-1}(x)) \cdots \nu(x) \leq Ke^{-\tau n}$ for all $x \in M$ and $n \geq 1$.

Suppose f admits some s -lamination W^s . A subset of M is s -saturated (relative to W^s) if it consists of entire leaves of W^s . An s -holonomy for F (relative to W^s) is a family h^s of homeomorphisms $h_{x,y}^s : \mathcal{E}_x \rightarrow \mathcal{E}_y$ defined for all $y \in W^s(x)$ and satisfying conditions (sh1), (sh2), (sh3) in Section 2.4.

Let $\mathcal{M}(\mu)$ be the set of probabilities on \mathcal{E} that project down to μ . Suppose F admits an s -holonomy h^s (relative to W^s). An F -invariant probability measure $m \in \mathcal{M}(\mu)$ is called an s -state (relative to h^s) if it admits some disintegration $\{m_x : x \in \text{supp } \mu\}$ which is *essentially s -invariant*, meaning that

$$(25) \quad (h_{x,y}^s)_* m_x = m_y \quad \text{for every } y \in W^s(x)$$

with x and y in some full μ -measure subset E . Then the same is true for any other disintegration of m . The full measure set may always be taken to be s -saturated: just consider the union E' of all W^s leaves through E and modify the disintegration on the (zero measure) set $E' \setminus E$ so as to enforce (25) for all points of E' .

Replacing f and F by their inverses, one obtains dual notions of u -lamination W^u , u -saturated set, u -holonomy h^u , u -invariant disintegration, and u -state. We say that the cocycle F admits *invariant holonomies* if it admits both s -holonomy and u -holonomy. Then we call an invariant probability an su -state if it is both an s -state and a u -state.

Remark 4.1. In some cases existence of s -states or u -states can be ensured a priori. For instance, if the fiber N is compact then one may start with a product measure $\mu \times \nu$ on \mathcal{E} and consider Cesaro limits of its forward/backward iterates: any such limit is a u -state/ s -state. See [16, Proposition 4.2] for a similar construction.

Proposition 4.2. *Assume F admits s -holonomy. Let $m \in \mathcal{M}(\mu)$ be such that there exists a sequence $(m_k)_k$ of F -invariant probability measures converging to m in the weak* topology, such that $\int \min\{0, \lambda_-(F, \cdot)\} dm_k \rightarrow 0$ as $k \rightarrow \infty$. Then m is an s -state.*

The proof of Proposition 4.2 occupies Sections 4.2 and 4.4. Let us also register the following particular case, corresponding to $m_k \equiv m$:

Corollary 4.3. *Assume F admits s -holonomy. Let $m \in \mathcal{M}(\mu)$ be an F -invariant probability measure such that $\lambda_-(F, x, \xi) \geq 0$ at m -almost every point. Then m is an s -state.*

Replacing F by its inverse one obtains dual statements for cocycles admitting u -holonomy. In particular: if $m \in \mathcal{M}(\mu)$ is an F -invariant probability measure such that $\lambda_+(F, x, \xi) \leq 0$ at m -almost every point then m is a u -state. Compare Example 3.15.

4.2. Local Markov property. We need the following local Markov property of the s -lamination. Fix $\ell \geq 1$ such that $Ke^{-\tau\ell} < 1/4$ and let $g = f^\ell$. We restrict considerations to an f -invariant σ -compact subset of full measure, which for simplicity we still denote by M .

Proposition 4.4. *There exists $\delta > 0$ and for every $x \in M$ there exists a partition $Q = Q_x$ of M such that*

- (1) *if $g^j(Q(z))$ intersects $Q(w)$ then $g^j(Q(z)) \subset Q(w)$*
- (2) *every $Q(y)$ is contained in some $W_\varepsilon^s(z)$*
- (3) *$W_\varepsilon^s(y) \cap \overline{B}(x, \delta) \subset Q(y) \subset W_\varepsilon^s(y)$ for every $y \in \overline{B}(x, \delta)$.*

Proof. Pick $\delta < \varepsilon/4$ and write $V = \overline{B}(x, \delta)$. For the sake of clearness, we split the proof into three steps:

Step 1: We claim that, for any $z, w \in V$ and $k \geq 0$, if $g^k(W_\varepsilon^s(w))$ intersects $W_\varepsilon^s(z) \cap V$ then either $k = 0$ and $W_\varepsilon^s(w) \cap V = W_\varepsilon^s(z) \cap V$ or $g^k(W_\varepsilon^s(w)) \subset W_\varepsilon^s(z)$. We call this the pre-Markov property. For the proof, consider any $z, w, p \in V$ such that $p \in g^k(W_\varepsilon^s(w)) \cap W_\varepsilon^s(z) \cap V$. Suppose first that $k = 0$. Let q be any point in $W_\varepsilon^s(w) \cap V$. Since $W^s(z)$ and $W^s(w)$ intersect each other (at p), they must coincide. It follows that $q \in W^s(z)$. Given our choices of ℓ and δ , and using that $p \in W_\varepsilon^s(z) \cap V$ and $p, q \in W_\varepsilon^s(w) \cap V$,

$$\begin{aligned} d(g^j(z), g^j(q)) &\leq d(g^j(z), g^j(p)) + d(g^j(p), g^j(q)) \\ &\leq Ke^{-j\ell}d(z, p) + Ke^{-j\ell}d(p, q) \leq 4^{-j}4\delta \leq \varepsilon \end{aligned}$$

for all $j \geq 0$. By (s11), this proves that $q \in W_\varepsilon^s(z)$. So, we have shown that $W_\varepsilon^s(w) \cap V \subset W_\varepsilon^s(z) \cap V$. The converse inclusion follows as well, by symmetry. Now suppose $k \geq 1$. Let q be any point in $g^k(W_\varepsilon^s(w))$. Since $W^s(g^k(w))$ and $W^s(z)$ intersect each other, they must coincide. It follows that $q \in W^s(z)$. Since $p \in W_\varepsilon^s(z)$ and $p, q \in g^k(W_\varepsilon^s(w))$

$$\begin{aligned} d(g^j(z), g^j(q)) &\leq d(g^j(z), g^j(p)) + d(g^j(p), g^j(q)) \\ &\leq 4^{-j}2\delta + 4^{-(j+k)}2\varepsilon \leq \varepsilon \end{aligned}$$

for all $j \geq 0$. This shows that $q \in W_\varepsilon^s(z)$, which completes the proof of the claim.

Step 2: Call P a stable pre-piece of rank $k \geq 0$ if $P = g^k(W_\varepsilon^s(y) \cap V)$ for some $y \in V$. By the pre-Markov property, two stable pre-pieces of rank 0 either coincide or are disjoint. Since g is invertible, the same holds for any two stable pre-pieces of the same rank $k \geq 0$. Call P_0, \dots, P_n a chain if the P_i are stable pre-pieces and $P_i \cap P_{i-1} \neq \emptyset$ for $1 \leq i \leq n$. Denote by k_j the rank of each P_j and call n the length of the chain. We claim that if P_0, \dots, P_n is a chain then

$$(26) \quad \bigcup_{j=0}^n P_j \subset g^{k_s}(W_\varepsilon^s(g^{-k_s}(z)))$$

for any $z \in P_s$ with rank $k_s = \min_{0 \leq j \leq n} k_j$. To see this, we argue by induction on n . The case $n = 0$ is obvious. Assume the claim is true for every $m < n$, and let P_0, \dots, P_n be any chain as above. If $0 < s < n$ then both P_0, \dots, P_s and P_s, \dots, P_n are chains with smaller lengths, and so the conclusion follows immediately from the induction hypothesis. Hence, we may suppose either $s = 0$ or $s = n$. In what follows we deal with the former case, the latter being entirely analogous. We may also assume that $k_s = 0$, up to replacing z and all the P_j by their pre-images under

g^{k_s} . The definition of chain implies that $P_0 = W_\varepsilon^s(z) \cap V$ intersects the union of the other P_j , $1 \leq j \leq n$. By the induction hypothesis,

$$\bigcup_{j=1}^n P_j \subset g^{k_r}(W_\varepsilon^s(g^{-k_r}(\zeta)))$$

for some $\zeta \in P_r$, $1 \leq r \leq n$ with rank $k_r = \min_{1 \leq j \leq n} k_j$. If $k_r = 0$ then the pre-Markov property implies that $W_\varepsilon^s(\zeta) \cap V = W_\varepsilon^s(z) \cap V$, and so the union of all P_j , $0 \leq j \leq n$ is indeed contained in $W_\varepsilon^s(z)$. Similarly, if $k_r > 0$ then the pre-Markov property implies

$$g^{k_r}(W_\varepsilon^s(g^{-k_r}(\zeta))) \subset W_\varepsilon^s(z)$$

and so the union of all P_j , $0 \leq j \leq n$ is again contained in $W_\varepsilon^s(z)$. This completes the induction step.

Step 3: Define the stable piece of a point $y \in M$ to be the set $Q(y)$ all $\xi \in M$ such that there exists a chain P_0, \dots, P_n with $y \in P_0$ and $\xi \in P_n$. If no such point ξ exists, just let $Q(y) = \{y\}$ instead. It is clear that the image under g of a stable piece is contained in a stable piece, and any two stable pieces that intersect must coincide. This implies property (1) in the lemma. Next, the property in (26) gives that $Q(y) \subset g^k(W_\varepsilon^s(g^{-k}(z))) \subset W_\varepsilon^s(z)$ for any z in a pre-piece with minimum rank k . This gives (2). If $y \in V$ then $Q(y) \supset W_\varepsilon^s(y) \cap V$, because one may always take $P_0 = W_\varepsilon^s(y) \cap V$. Moreover, in this case $k = 0$ and $Q(y) \subset W_\varepsilon^s(y)$. This proves (3) and the proposition. \square

4.3. Measurability. In this section we prove that the partition Q constructed in Proposition 4.4 admits a *measurable section*:

Proposition 4.5. *There exists a measurable map $\pi : M \rightarrow M$ constant on every $Q(y)$ and such that $\pi(y) \in Q(y)$ for every $y \in M$. In particular, $d(y, \pi(y)) \leq 2\varepsilon$ for every $y \in M$.*

For the proof, we need the following abstract result. Recall that a partition \mathcal{P} of a probability space is *measurable* (Rokhlin [41]) if it is the limit modulo 0 of an increasing sequence of finite partitions into measurable sets.

Lemma 4.6. *Let (X, \mathcal{A}, ν) be a Lebesgue space and \mathcal{P} be a partition of X into measurable sets. If \mathcal{P} is a measurable partition then there exists a measurable map $\pi : X \rightarrow X$ that is constant on every partition element and satisfies $\pi(x) \in \mathcal{P}(x)$ for every x .*

Proof. We begin by recalling that (Rokhlin [41, §1–§3]) every Lebesgue space is isomorphic (modulo 0) to the interval $I = [0, 1]$, endowed with the σ -algebra of Lebesgue measurable sets and a probability measure whose nonatomic part is a multiple of Lebesgue measure. After appropriate restrictions to full measure sets and consideration of this isomorphism, we may assume that $X = I$ and that \mathcal{P} is the limit (not just in the modulo 0 sense) of an increasing subsequence $(\mathcal{P}_n)_n$ of finite partitions of I into Lebesgue measurable sets. For each atom $P_n \in \mathcal{P}_n$, let $K(P_n) \subset P_n$ be a compact subset such that $\nu(P_n \setminus K(P_n)) \leq 2^{-n}\nu(P_n)$. Let $K'_n = \bigcap_{m \geq n} \bigcup_{P_m \in \mathcal{P}_m} K(P_m) \times K(P_m) \subset X \times X$. Then the projection of K'_n on the first coordinate is $K_n = \bigcap_{m \geq n} \bigcup_{P_m \in \mathcal{P}_m} K(P_m)$. Notice that K_n and K'_n are compact sets, and $\nu(K_n) \geq 1 - 2^{1-n}$. Let $\phi_n : K_n \rightarrow X$ be defined so that $\phi_n(y)$ is the infimum of all z with $(y, z) \in K'_n$. For $n \geq 2$, let Y_n be the projection

on the first coordinate of $K'_n \cap (K_n \times K_{n-1})$, and let $\psi_n : Y_n \rightarrow K_{n-1}$ be such that $\psi_n(y)$ is the infimum of all z with $(y, z) \in K'_n \cap (K_n \times K_{n-1})$. Define by induction $\pi_n : K_n \rightarrow K_n$ so that $\pi_1 = \phi_1$ and for $n \geq 2$ let $\pi_n(y) = \phi_n(y)$ if $y \notin Y_n$ and $\pi_n(y) = \pi_{n-1}(\psi_n(y))$ if $y \in Y_n$. Then π_n is a Borel function. Notice that $\pi_n(y) = \pi_{n-1}(z)$ whenever $y \in \mathcal{P}(z)$, and for $n \geq 2$ we have $\pi_n|_{K_{n-1}} = \pi_{n-1}$. Define $\pi = \lim \pi_n$ on $\bigcup_n K_n$, and extend it to the whole X in an arbitrary way to satisfy $\pi(y) = \pi(z)$ whenever $y \in \mathcal{P}(z)$. Then π is the desired measurable section. \square

As a criterion for measurability of partitions, we use:

Lemma 4.7. *Let \mathcal{P} be a partition in a σ -compact metric space X and suppose that $\text{graph}(\mathcal{P}) = \{(x, y) : y \in \mathcal{P}(x)\}$ is a closed subset of $X \times X$. Then \mathcal{P} is the limit of an increasing sequence of finite partitions of X into Borel sets.*

Proof. Let $Z \subset X$ be a countable dense subset. For each $x \in Z$ and $i \geq 1$, let $U_{x,i}$ be the set of all $z \in X$ such that $\mathcal{P}(z)$ intersects the closed ball of radius $1/i$ centered at x . Then $U_{x,i}$ is σ -compact, since it is the projection of a closed subset of $X \times X$. Since the atoms of \mathcal{P} are closed, for any two different ones there exists (x, i) such that $U_{x,i}$ contains one of the atoms and is disjoint from the other. Thus, the countably many partitions $\{U_{x,i}, X \setminus U_{x,i}\}$ generate \mathcal{P} . This proves the lemma. \square

Proof of Proposition 4.5. Let $G_k \subset M \times M$ be the union of the diagonal $\{(y, y) : y \in M\}$ with the set of all $(y, z) \in V \times V$ such that $z \in W_\epsilon^s(y)$, and for $k \geq 1$ let G_k be the set of all $(f^j(y), f^j(z)) \in M \times M$ with $(y, z) \in G_0$ and $0 \leq j \leq k$. Then G_k is closed for every $k \geq 0$. Let $G_{(k)} \subset M^{k+1} = M \times \cdots \times M$ be the set of all (y_1, \dots, y_{k+1}) such that $(y_i, y_{i+1}) \in G_{(k)}$ for $1 \leq i \leq k$. If we denote by $Q_k(y)$ the set of all $z \in M$ such that there exists $y_2, \dots, y_k \in M$ with $(y, y_2, \dots, y_k, z) \in G_{(k)}$, then it is clear that $Q_k(y) \subset Q_{k+1}(y)$ for every $k \geq 0$ and $\bigcup_k Q_k(y) = Q(y)$. Given a Borel subset $Z \subset M$, let $Q(Z) = \bigcup_{y \in Z} Q(y)$. Notice that $Q(Z)$ is measurable since $Q(Z) = \bigcup_k Q_k(Z)$ where $Q_k(Z) = \bigcup_{y \in Z} Q_k(y)$ and each $Q_k(Z)$, being the projection in the last coordinate of the Borel subset $G_{(k)} \cap (Z \times M^k) \subset M^{k+1}$, is measurable. Moreover, this argument shows that if Z is σ -compact then $Q(Z)$ is σ -compact as well. Notice that if $y \notin Q(V)$ then $f(Q(f^{-1}(y))) = Q(y)$. Thus, if we can construct a measurable section $\pi' : Q(V) \rightarrow Q(V)$ for the restriction \mathcal{P} of Q to $Q(V)$, then we can define the section π by $\pi(y) = f^n(\pi'(f^{-n}(y)))$, when there exists a minimal $n \geq 0$ such that $f^{-n}(y) \in Q(V)$, and $\pi(y) = y$ when no such n exists (as in this case we clearly have $Q(y) = \{y\}$). If $Q(V)$ has zero μ -measure, any section will be measurable. So assume that $Q(V)$ has positive μ -measure. Since $Q(V)$ is σ -compact, it is a Lebesgue space with respect to the completion of any Borel probability measure. So, by Lemma 4.6, it is enough to show that \mathcal{P} is measurable. Let \mathcal{P}' be the restriction of Q to V . Its atoms are of the form $Q(y) \cap V = W_\epsilon^s(y) \cap V$, $y \in V$. So, by Lemma 4.7, it is the limit of an increasing sequence \mathcal{P}'_n of finite partitions of V into Borel subsets. Then \mathcal{P} is the limit of partitions $\mathcal{P}_n = \{Q(P) : P \in \mathcal{P}'_n\}$. In particular \mathcal{P} is measurable, as desired. \square

4.4. Lyapunov exponents and holonomy invariance. In this section we conclude the proof of Proposition 4.2. Let $\ell \geq 1$ and $\delta > 0$ be as in Proposition 4.4. The main remaining step is to show that every disintegration of m is essentially s -invariant restricted to the δ -neighborhood of any point $x \in M$. This will be done by applying Theorem C to a deformation of the cocycle $G = F^\ell$ over $g = f^\ell$, more precisely, to a cocycle which is conjugate to G via s -holonomies. Covering M with

these neighborhoods we obtain a disintegration of m which is essentially s -invariant on the whole space.

Clearly, m is invariant under G and $\lambda_-(G, x, \xi) = \ell\lambda_-(F, x, \xi) \geq 0$ for m -almost every point. Given any $x \in M$, let V be its δ -neighborhood, and let $Q = Q_x$ be the partition constructed in Proposition 4.4. Let \mathcal{B}_0 be the σ -algebra of measurable subsets of M that are unions of entire atoms of Q . In other words, a measurable subset E belongs to \mathcal{B}_0 if and only if every stable piece is either contained in or disjoint from E . Notice that g is \mathcal{B}_0 -measurable, because the image of any stable piece is contained in a stable piece. Let $\pi : M \rightarrow M$ be as in Proposition 4.5.

Let $\tilde{G} = \tilde{G}(x) : \mathcal{E} \rightarrow \mathcal{E}$ be the transformation defined by

$$\tilde{G}_y = h_{g(\pi(y)), \pi(g(y))}^s \circ G_{\pi(y)}.$$

Then \tilde{G} is \mathcal{B}_0 -measurable, because π is constant on stable pieces and the image of every stable piece is contained in a stable piece. Property (sh2) applied to G (the properties in the definition of s -holonomy remain valid when one replaces the cocycle by any forward iterate) yields

$$(27) \quad \tilde{G}_y = h_{g(y), \pi(g(y))}^s \circ G_y \circ h_{\pi(y), y}^s$$

for every $y \in M$. This relation can be rewritten as $\tilde{G} = \Phi \circ G \circ \Phi^{-1}$, where $\Phi : \mathcal{E} \rightarrow \mathcal{E}$ is given by $\Phi(y, \xi) = (y, h_{y, \pi(y)}^s(\xi))$. Then \tilde{G} is a deformation of G , since Φ and its inverse are β -Hölder continuous on every fiber. Let \tilde{m} be the probability measure on \mathcal{E} defined by $\tilde{m} = \Phi_*(m)$. Clearly, it is invariant under \tilde{G} , it projects down to μ , and its conditional probabilities along the fibers are given by

$$(28) \quad \tilde{m}_y = (h_{y, \pi(y)}^s)_* m_y.$$

So, we are in a position to apply Theorem C to conclude that the disintegration $\{\tilde{m}_y\}$ is \mathcal{B}_0 -measurable: there exists a full μ -measure subset restricted to which \tilde{m}_y is constant on stable pieces $Q(y)$. Through (28), this gives rise to a disintegration $\{m_y\}$ of m which is s -invariant on each stable piece:

$$m_z = (h_{\pi(z), z}^s)_* \tilde{m}_z = (h_{\pi(w), z}^s)_* \tilde{m}_w = (h_{w, z}^s)_* (h_{\pi(w), w}^s)_* \tilde{m}_w = (h_{w, z}^s)_* m_w$$

for any z and w in any $Q(y)$. In particular, the disintegration of m is essentially s -invariant restricted to the ball $V = V(x)$ of radius δ around every $x \in M$. Now, consider any countable set $\{x_n\} \subset \text{supp } \mu$ such that the balls of radius $\delta/2$ around these x_n cover the support of μ . For each n , let B_n be a zero μ -measure subset of the ball $V(x_n)$ of radius δ around x_n such that

$$(29) \quad m_w = (h_{z, w}^s)_* m_z \quad \text{for all } z, w \in V(x_n) \setminus B_n \text{ in the same stable piece.}$$

Let E be the set of all point $x \in \text{supp } \mu$ whose orbits never meet $\cup_n B_n$ and such that

$$(30) \quad m_{f^l(x)} = (F_x^l)_* m_x \quad \text{for all } l \in \mathbb{Z}.$$

Then E has full μ -measure in M . By properties (sl1)-(sl5) in the definition of s -lamination, given any $x, y \in E$ with $y \in W^s(x)$ there exists $k \geq 1$ such that $d(f^l(x), f^l(y)) \leq \delta/2$ ($\leq \varepsilon$) for all $l \geq k$. Fix n such that $f^k(x) \in B(x_n, \delta/2)$. Then $f^k(y) \in V(x_n) \cap W_\varepsilon^s(f^k(x))$. So, by Proposition 4.4, the two points $z = f^k(x)$ and $w = f^k(y)$ belong to the same stable piece (associated to x_n). Since they are outside B_n , we may combine (29) and (30) with property (sh2) of s -holonomies

to conclude that $m_y = (h_{x,y}^s)_* m_x$. This proves that the disintegration of m is essentially s -invariant, as claimed. This finishes the proof of Proposition 4.2.

4.5. Product structure and continuity. In particular, if the cocycle F admits invariant holonomies, then any invariant measure $m \in \mathcal{M}(\mu)$ for which Lyapunov exponents vanish almost everywhere is an su -state. This means that m has some disintegration which is s -invariant on a full measure s -saturated set and some disintegration which is u -invariant on a full measure u -saturated set. We are now going to discuss additional conditions under which these two disintegrations may be taken to coincide. Based on this we prove Theorem D and Corollary E.

Assume the s -lamination and the u -lamination of $f : M \rightarrow M$ satisfy a local product condition like (h5) in Section 2.3: *there is $\delta > 0$ such that if $d(x_1, x_2) \leq \delta$ then $W_\varepsilon^u(x_1)$ and $W_\varepsilon^s(x_2)$ intersect at exactly one point, and this point $[x_1, x_2]$ depends continuously on (x_1, x_2) .* Given any $x \in M$, let $B^s(x)$ and $B^u(x)$ be the balls of radius $\delta/2$ around x inside $W_\varepsilon^s(x)$ and $W_\varepsilon^u(x)$, respectively. Then

$$\phi : B^s(x) \times B^u(x) \rightarrow M, \quad (x_1, x_2) \mapsto [x_1, x_2]$$

defines a homeomorphism from $B^s(x) \times B^u(x)$ to a neighborhood $B(x)$ of x . Indeed, assumption (h5) ensures that ϕ is injective, its image covers a neighborhood of x , and its inverse is continuous. Assume also that μ has local product structure. As observed before, this implies that the support of μ is su -saturated.

Proposition 4.8. *Assume F admits s -holonomy and u -holonomy. Assume f and μ have local product structure, as described above. If m is an su -state then it admits a disintegration which is s -invariant and u -invariant and whose conditional probabilities m_x vary continuously with x in the support of μ .*

Proof. It suffices to prove that given any $z \in \text{supp } \mu$ there exists a disintegration of m which satisfies the conclusion restricted to $B(z)$. By assumption, there exists some s -invariant disintegration $\{m_x^s\}$ and some u -invariant disintegration $\{m_x^u\}$. By essential uniqueness, $m_x^u = m_x^s$ for every x in some full μ -measure set $E \subset M$. Using product structure, there exists $x^u \in B^u(z)$ such that E intersects $\{x^u\} \times B^s(z)$ on a full μ^s -measure set. Define

$$m_x = (h_{y,x}^u)_* m_y^s \quad \text{for every } x \in B(z), \quad y = [x^u, x].$$

Note that $y \mapsto m_y^s$ is continuous on $\{x^u\} \times B^s(z)$, because m_x^s is s -invariant and s -holonomies vary continuously with the point. Using that u -holonomies also vary continuously with the point, we get that $x \mapsto m_x$ is continuous on $B(z)$. By construction, m_x is u -invariant. Moreover, it coincides with m_x^u almost everywhere, due to product structure and the choice of x^u . Hence, m_x also coincides with m_x^s almost everywhere. By continuity, it follows that m_x is also s -invariant. \square

Proof of Theorem D. Clearly, the assumption $\int |\lambda_\pm(F, \cdot)| dm_k \rightarrow 0$ is stronger than $\int \min\{0, \lambda_-(F, \cdot)\} dm_k \rightarrow 0$. So, by Proposition 4.2, the measure m is an s -state. Since the assumption is symmetric under time reversion, we may apply the proposition to F^{-1} as well, to conclude that m is also a u -state. Now the conclusion of the theorem follows from Proposition 4.8. \square

Remark 4.9. The same arguments yield a more local version Theorem D. Let $U \subset M$ be an invariant, s -saturated, u -saturated set, with positive Lebesgue measure. An su -state over U is an invariant measure that projects down to the normalized

restriction of the Lebesgue measure to U and which admits some essentially s -invariant disintegration and some essentially u -invariant disintegration. Then it admits a disintegration which is su -invariant and continuous.

Proof of Corollary E. Since the fiber is 1-dimensional, $\lambda_-(F, \cdot) = \lambda_+(F, \cdot)$ wherever they are defined. Suppose there is a sequence $(m_k)_k$ of ergodic probability measures projecting down to μ and such that $\lambda_{\pm}(F, m_k) \rightarrow 0$ as $k \rightarrow \infty$. By ergodicity, this is the same as the condition in the assumption of Theorem D. Since the fiber is compact, the sequence must have some accumulation point. Every accumulation point m is also an invariant measure that projects down to μ . By Theorem D, the measure m admits a disintegration $\{m_x : x \in M\}$ which is s -invariant, u -invariant, and continuous on the support of μ . By invariance and continuity, this disintegration satisfies

$$(F_x)_* m_x = m_{f(x)}$$

for every point in $\text{supp } \mu$. In particular,

$$(31) \quad (F_p)_* m_p = m_p \quad \text{and} \quad (F_q)_* m_q = m_q.$$

The first equality implies that the support of m_p is contained in the subset $\{a_p, r_p\}$ of the circle. Let $z \in W^u(p) \cap W^s(q)$. By invariance of the conditional probabilities,

$$m_q = (h_{z,q}^s \circ h_{p,z}^u)_* m_p.$$

Consequently, $\text{supp } m_q$ contains at most two points. The second equality in (31) implies that the support is invariant under F_q . It follows that F_q has periodic points of period 1 or 2, which contradicts the assumption of the corollary. This contradiction proves that the exponent is indeed bounded away from zero. \square

5. DOMINATION AND FIBER BUNCHING

Here we introduce a number of ideas that will be useful for analyzing the dependence of Lyapunov exponents on the cocycle. We take the fiber manifold N to be compact; towards the end of the section, we assume the fiber bundle \mathcal{E} itself to be compact. In addition to the conditions in the previous section, we assume the fiber bundle and the cocycle to be Lipschitz, in the sense of Section 2.5. We also consider the Lipschitz topology in the space of Lipschitz cocycles introduced in Section 2.5.

5.1. Existence of holonomies. Assume f admits an s -lamination W^s . We call a cocycle F *s -dominated* (relative to W^s) if there exist $\ell \geq 1$ and $\theta < 1$ such that

$$(32) \quad \|DF_x^\ell(\xi)^{-1}\| \nu_\ell(x) \leq \theta \quad \text{for all } (x, \xi) \in \mathcal{E}.$$

In other words, the strongest contraction of F^ℓ along the fibers is strictly weaker than the weakest contraction of f^ℓ along the leaves of W^s . Replacing f and F by its inverses, we obtain the dual notion of *u -dominated cocycle*. Denote by $\mathcal{D}^s(f)$ the subset of s -dominated cocycles and by $\mathcal{D}^u(f)$ the subset of u -dominated cocycles. It is clear from the definition that these are open sets for the topology we have just introduced. When f admits both an s -lamination and a u -lamination, we let $\mathcal{D}(f) = \mathcal{D}^s(f) \cap \mathcal{D}^u(f)$ be the subset of *dominated* cocycles.

The s -domination condition is designed so that the usual graph transform argument yields a “strong-stable” lamination for the map F (there is a dual statement for u -dominated cocycles):

Proposition 5.1. *If the cocycle F is s -dominated then there exists a unique partition $\mathcal{W}^s = \{\mathcal{W}^s(x, \xi) : (x, \xi) \in \mathcal{E}\}$ of the fiber bundle \mathcal{E} such that*

- (1) *every $\mathcal{W}^s(x, \xi)$ is a Lipschitz graph over $W^s(x)$, with Lipschitz constant uniform on x ;*
- (2) *$F(\mathcal{W}^s(x, \xi)) \subset \mathcal{W}^s(F(x, \xi))$ for all $(x, \xi) \in \mathcal{E}$;*
- (3) *the map $h_{x,y}^s : \mathcal{E}_x \rightarrow \mathcal{E}_y$ defined by $(y, h_{x,y}^s(\xi)) \in \mathcal{W}^s(x, \xi)$, for $y \in W^s(x)$, coincides with the uniform limit of $(F_y^n)^{-1} \circ F_x^n$ as $n \rightarrow \infty$;*
- (4) *the family of maps $h_{x,y}^s : \mathcal{E}_x \rightarrow \mathcal{E}_y$ is an s -holonomy for F .*

Proof. The claims follow from the same partial hyperbolicity methods (see Hirsch, Pugh, Shub [27]) used before to obtain similar results for linear cocycles [14, 16, 43], and so we just sketch the main ingredients. Existence (1) and invariance (2) of the family \mathcal{W}^s follow from a standard application of the graph transform argument [27].

Notice that, for every x and y in the same stable manifold, and every $n \geq 0$,

$$(33) \quad h_{x,y}^s = (F_y^n)^{-1} \circ h_{f^n(x), f^n(y)}^s \circ F_x^n$$

and the uniform distance from $h_{f^n(x), f^n(y)}^s$ to the identity map is bounded by $Cd(f^n(x), f^n(y))$, where C is the uniform Lipschitz constant in (1). Putting these observations together, we find that

$$\begin{aligned} d_{C^0}(h_{x,y}^s, (F_y^n)^{-1} \circ F_x^n) &\leq \text{Lip}((F_y^n)^{-1}) d_{C^0}(h_{f^n(x), f^n(y)}^s, \text{id}) \\ &\leq C \sup_{\xi} \|DF_y^n(\xi)^{-1}\| d(f^n(x), f^n(y)). \end{aligned}$$

Fix ℓ as in the domination condition (32) and write $k = [n/\ell]$. Clearly,

$$d(f^n(x), f^n(y)) \leq Kd(f^{k\ell}(x), f^{k\ell}(y)) \leq K \prod_{i=0}^{k-1} \nu_{\ell}(f^{i\ell}(y))d(x, y)$$

and $\|DF_y^n(\xi)^{-1}\|$ is similarly bounded above by a product of norms of the derivative of $(F^{\ell})^{-1}$ along the orbit of y . Using the domination condition (32) we conclude that

$$d_{C^0}(h_{x,y}^s, (F_y^n)^{-1} \circ F_x^n) \leq \text{const } \theta^k \leq \text{const } \theta^{n/\ell},$$

where the constants are independent of n, x, y . This proves (3).

Conditions (sh1)-(sh3) in the definition of s -holonomy are direct consequences of the definition of $h_{x,y}^s$. Thus, to prove (4) we only have to check that the maps $h_{x,y}^s$ are Hölder continuous, with uniform exponential Hölder constant. The arguments are quite standard, see for instance [1, 27]. In view of (33), and the fact that the F_z and their inverses are Lipschitz, it is no restriction to assume that x and y are in the same local stable set. For each $n \geq 0$, denote $x_n = f^n(x)$ and $y_n = f^n(y)$. Given $\xi, \eta \in \mathcal{E}_x$, denote $\xi_n = F_x^n(\xi)$ and $\eta_n = F_x^n(\eta)$. By domination, there exists $K > 0$ and

$$(34) \quad n \leq -K \log d(\xi, \eta)$$

such that $d(x_n, y_n) \leq d(\xi_n, \eta_n)$. Then

$$d(h_{x_n, y_n}^s(\xi), h_{x_n, y_n}^s(\eta)) \leq d(\xi_n, \eta_n) + 2Cd(x_n, y_n) \leq 3Cd(\xi_n, \eta_n),$$

where C is the uniform Lipschitz constant in (1). Let L be a uniform upper bound for the norms of $\|DF_z^{\pm 1}\|$. The previous inequality yields

$$L^{-n}d(h_{x,y}^s(\xi), h_{x,y}^s(\eta)) \leq 3CL^n d(\xi, \eta).$$

In view of (34), we have $L^{2n} \leq d(\xi, \eta)^{-\theta}$ for some uniform constant $\theta \in (0, 1)$. Then the previous inequality gives

$$d(h_{x,y}^s(\xi), h_{x,y}^s(\eta)) \leq 3Cd(\xi, \eta)^{1-\theta},$$

which proves our claim. \square

5.2. Continuity of holonomies. We are going to see that s -holonomies vary continuously with the cocycle on $\mathcal{D}^s(f)$. Of course, there is a dual statement for u -holonomies on $\mathcal{D}^u(f)$. Let $\mathcal{W}^s(G) = \{\mathcal{W}^s(G, x, \xi) : (x, \xi) \in \mathcal{E}\}$ denote the ‘‘strong-stable’’ lamination of a cocycle $G \in \mathcal{D}^s$, as in Proposition 5.1, and $h_G^s = h_{G,x,y}^s$ be the corresponding s -holonomy:

$$(35) \quad (y, h_{G,x,y}^s(\xi)) \in \mathcal{W}^s(G, x, \xi).$$

Recall $\mathcal{W}^s(G, x, \xi)$ is a graph over $W^s(x)$. We also denote by $\mathcal{W}_\varepsilon^s(G, x, \xi)$ the part of the graph located over $W_\varepsilon^s(x)$, that is, the set of all points $(y, h_{G,x,y}^s(\xi))$ with $y \in W_\varepsilon^s(x)$.

Proposition 5.2. *Let $(F_k)_k$ be a sequence of cocycles converging to F in $\mathcal{D}^s(f)$. Then*

- (1) *every $\mathcal{W}^s(F_k, x, \xi)$ is a Lipschitz graph, with Lipschitz constant uniform on x, ξ , and k*
- (2) *$\mathcal{W}_\varepsilon^s(F_k, x, \xi)$ converges to $\mathcal{W}_\varepsilon^s(F, x, \xi)$, as graphs over the same domain, uniformly on $(x, \xi) \in \mathcal{E}$*
- (3) *$h_{F_k,x,y}^s(\xi)$ converges to $h_{F,x,y}^s(\xi)$ for every $x \in M, y \in W^s(x)$, and $\xi \in \mathcal{E}_x$, and the converse is uniform over all $y \in W_\varepsilon^s(x)$.*

Proof. This is another standard consequence of the classical graph transform argument [27]. Indeed, the assumptions imply that the graph transform of F_k converges to the graph transform of F in an appropriate sense, so that the corresponding fixed points converge as well. This yields (1) and (2). Part (3) is a direct consequence of (2) and the definition (35), in the case $y \in W_\varepsilon^s(y)$. The general statement follows, using the invariance property (sh2):

$$H_{F_k,x,y}^s = (F_{k,y}^n)^{-1} \circ h_{F_k, f^n(x), f^n(y)} \circ F_{k,x}^n.$$

See also [43, Section 4] where stronger results are proved in detail using similar methods, in the context of linear cocycles. \square

Corollary 5.3. *The subset of cocycles admitting some su -state is closed in $\mathcal{D}(f)$.*

Proof. If $F_k \rightarrow F$ in $\mathcal{D}(f)$ and m_k are su -states for F_k projecting down to μ then any weak limit m of the sequence m_k is an su -state for F projecting down to μ . \square

5.3. Projective extension. Let $F : \mathcal{E} \rightarrow \mathcal{E}$ be a dominated cocycle. It will be convenient to think of F as a transformation in its own right, and to consider a certain smooth cocycle $\mathbb{P}(F)$ over F that we call projective extension. Here we define $\mathbb{P}(F)$ and discuss a stronger domination condition, called fiber bunching, that ensures robust existence of holonomies for $\mathbb{P}(F)$.

The partition $\mathcal{W}^s = \mathcal{W}^s(F)$ given by Proposition 5.1 is an s -lamination for F : in particular, since strong-stable leaves are Lipschitz graphs (Proposition 5.1) and local charts are Lipschitz, we have

$$(36) \quad d(F^n(y, \eta), F^n(z, \zeta)) \leq C_0 \nu_n(x) d((y, \xi), (z, \zeta))$$

for every $(y, \eta), (z, \zeta) \in \mathcal{W}^s(x, \xi)$, where $C_0 > 1$ is a uniform constant. Analogously, the “strong-unstable” lamination $\mathcal{W}^u = \mathcal{W}^u(F)$ is a u -lamination for F . In addition, we consider the c -lamination

$$\mathcal{W}^c = \{\mathcal{W}^c(x, \xi) = \mathcal{E}_x : (x, \xi) \in \mathcal{E}\}.$$

Let $\mathbb{P}(\mathcal{E})$ be the projective tangent bundle of \mathcal{E} , that is, the fiber bundle over \mathcal{E} such that the fiber of each (x, ξ) is the projectivization of the tangent space $T_\xi \mathcal{E}_x$. The *projective extension* of F is the smooth cocycle $\mathbb{P}(F) : \mathbb{P}(\mathcal{E}) \rightarrow \mathbb{P}(\mathcal{E})$ over $F : \mathcal{E} \rightarrow \mathcal{E}$ defined by

$$\mathbb{P}(F)(x, \xi, [v]) = (f(x), F_x(\xi), [DF_x(\xi)v]), \quad \text{for each } [v] \in \mathbb{P}(T_\xi \mathcal{E}_x).$$

Notice $\mathbb{P}(\mathcal{E})$ is also a fiber bundle over M , with fiber $\mathbb{P}(T\mathcal{E}_x)$, and one may think of $\mathbb{P}(F)$ as a cocycle over $f : M \rightarrow M$ instead. However, this will usually not be our point of view: instead, most of the time, we think of $\mathbb{P}(F)$ as a cocycle over F itself.

Assume the cocycle F is s -dominated. We say that F is *s-fiber bunched* if there exist $\ell \geq 1$ and $\theta < 1$ such that

$$(37) \quad \|DF_x^\ell(\xi)\| \|(DF_x^\ell(\xi))^{-1}\| \nu_\ell(x) \leq \theta \quad \text{for every } (x, \xi) \in \mathcal{E}.$$

The product of the first two factors bounds the norm of the derivative $\mathbb{P}(F)^\ell$ and its inverse. Recall (23) and (24). Thus, this condition means that the strongest contraction of $\mathbb{P}(F)^\ell$ along the fibers $\mathbb{P}(T_\xi \mathcal{E}_x)$ is strictly weaker than the weakest contraction of f^ℓ along the leaves of W^s .

It is easy to see that s -fiber bunching implies that $\mathbb{P}(F)$ is s -dominated relative to the s -lamination \mathcal{W}^s of F and, consequently, has s -holonomy. Indeed, (37) implies

$$\|DF_x^{k\ell}(\xi)\| \|(DF_x^{k\ell}(\xi))^{-1}\| C_0 \nu_{k\ell}(x) \leq C_0 \theta^k$$

and so, in view of (36), it suffices to fix $k \geq 1$ such that $C_0 \theta^k < 1$.

Remark 5.4. Under condition (37), a computation similar to Proposition 5.1(3) shows that $(F_y^n)^{-1} \circ F_x^n$ converges to $h_{x,y}^s : \mathcal{E}_x \rightarrow \mathcal{E}_y$ in the C^1 topology. In particular, in this case the s -holonomy maps are diffeomorphisms between the fibers of \mathcal{E} . The projectivizations

$$\mathbb{P}(Dh_{x,y}^s(\xi)) : \mathbb{P}(T_\xi \mathcal{E}_x) \rightarrow \mathbb{P}(T_\eta \mathcal{E}_y), \quad \eta = h_{x,y}^s(\xi)$$

of the derivatives are precisely the s -holonomy maps of $\mathbb{P}(F)$.

A u -dominated cocycle F is *u-fiber bunched* if its inverse F^{-1} is s -fiber bunched. Then $\mathbb{P}(F)$ is u -dominated (relative to the u -lamination \mathcal{W}^u of F), and so it admits u -holonomies. Let $\mathcal{B}^s(f) \subset \mathcal{D}^s(f)$ be the subspace of s -fiber bunched cocycles, and $\mathcal{B}^u(f) \subset \mathcal{D}^u(f)$ be the subspace of u -fiber bunched cocycles. We call a dominated cocycle *fiber bunched* if it belongs to $\mathcal{B}(f) = \mathcal{B}^s(f) \cap \mathcal{B}^u(f)$.

Remark 5.5. Let $F : \mathcal{E} \rightarrow \mathcal{E}$ be a fiber bunched cocycle and m be an F -invariant probability such that $\lambda_-(F, x, \xi) = \lambda_+(F, x, \xi) = 0$ at m -almost every $x \in M$. Then every $\mathbb{P}(F)$ -invariant probability η projecting down to m is an su -state. This follows directly from Corollary 4.3 applied to the cocycle $\mathbb{P}(F)$ over the transformation F , and to its inverse.

5.4. Accessibility. In order to handle the construction in the previous section we shall need a few facts about cocycles over partially hyperbolic systems, that we present in here. Propositions 5.6 and 5.7 below are special versions of much more general results of Pugh, Shub [40] and Avila, Santamaria, Viana [3], respectively. We include the proofs since the arguments are much simpler in our setting, namely skew-products with differentiable stable and unstable holonomies.

Let F be a smooth cocycle admitting stable and unstable holonomies. The *accessibility class* of a point $(x, \xi) \in \mathcal{E}$ is the set of all $(y, \eta) \in \mathcal{E}$ such that there exist $(z_0, \zeta_0) = (x, \xi)$, $(z_1, \zeta_1), \dots, (z_{n-1}, \zeta_{n-1}), (z_n, \zeta_n) = (y, \eta)$ in \mathcal{E} satisfying

$$(z_{j+1}, \zeta_{j+1}) \in \mathcal{W}^s(z_j, \zeta_j) \cup \mathcal{W}^u(z_j, \zeta_j) \quad \text{for every } j = 0, \dots, n-1.$$

It is easy to see that any accessibility class with non-empty interior is open. We say that F is *accessible* if the whole \mathcal{E} is an accessibility class.

Proposition 5.6. *If F is a fiber bunched volume preserving cocycle and Z is an accessibility class with positive m -measure then there exists $n \geq 1$ such that $F^n(Z) = Z$ and $F^n \upharpoonright Z$ is ergodic for m . In particular, if F is accessible then it is ergodic.*

Proof. The first claims are immediate: Z must intersect $F^n(Z)$ for some $n \geq 1$, since $m(Z) > 0$, and then the two sets must coincide. We are left to prove that, given any continuous function $\varphi : \mathcal{E} \rightarrow \mathbb{R}$, the time averages

$$\varphi^\pm = \lim_n \frac{1}{n} \sum_{j=0}^{n-1} \varphi \circ F^{\pm j}$$

are constant m -almost everywhere on Z . Given $c \in \mathbb{R}$, let A_c be the set of points $z \in Z$ for which $\varphi^\pm(z)$ are well-defined and satisfy $\varphi^+(z) = \varphi^-(z) \leq c$. All we have to do is prove that every A_c has either zero or full m -measure in Z . Let c be such that A_c has positive m -measure and let m^c be the normalized restriction of m to A_c . Since φ^+ is constant on s -leaves, the set A_c is essentially s -saturated; for similar reasons it is also essentially u -saturated. This implies that m^c is an su -state and projects down to μ (which we assume to have local product structure). Then, by Proposition 4.8, the measure m^c admits a continuous s -invariant and u -invariant disintegration $\{m_x^c : x \in \text{supp } \mu\}$. Using also that the holonomies of F are area preserving diffeomorphisms, we obtain that the density of m_x^c with respect to Lebesgue measure on the fiber is constant along s -leaves and along u -leaves, over the support of μ . It follows that the density is constant on the whole accessibility class Z , over the support of μ . This can only happen if A_c has full m -measure in Z . \square

Let $\mathcal{M} \rightarrow \mathcal{E}$ denote the fiber bundle where the fiber of each $z = (x, \xi) \in \mathcal{E}$ is the space of probability measures in the projective fiber $\mathbb{P}(\mathcal{E})_z = \mathbb{P}(T_\xi \mathcal{E}_x)$. Let $H_{z,w}^s : \mathbb{P}(\mathcal{E})_z \rightarrow \mathbb{P}(\mathcal{E})_w$ be the s -holonomy maps of the projective extension $\mathbb{P}(F)$: if $w = (y, \eta)$ with $\eta = h_{x,y}^s(\xi)$ then $H_{z,w}^s : \mathbb{P}(T_\xi \mathcal{E}_x) \rightarrow \mathbb{P}(T_\eta \mathcal{E}_y)$ is the projectivization of the derivative of $h_{x,y}^s : \mathcal{E}_x \rightarrow \mathcal{E}_y$ at the point ξ .

Through the end of this section, we assume the ambient space \mathcal{E} to be compact.

Proposition 5.7. *Let F be a fiber bunched accessible volume preserving cocycle. Then any invariant su -state of $\mathbb{P}(F)$ projecting down to m admits a disintegration which is s -invariant and u -invariant and whose conditional probabilities vary continuously with the base point on the support of μ .*

Proof. Let ζ be any invariant su -state of $\mathbb{P}(F)$ projecting down to m . We begin by thinking of $\mathbb{P}(F)$ as a cocycle over the hyperbolic transformation $f : M \rightarrow M$. It is clear that ζ is an su -state of this cocycle as well. Then, by Proposition 4.8, there exists a disintegration $\{\zeta_x : x \in M\}$, along the fibers of $\mathbb{P}(\mathcal{E}) \rightarrow M$ which is su -invariant and continuous. To proceed with the proof, let $\{\zeta_{(x,\xi)} : \xi \in \mathcal{E}_x\}$ be any disintegration of ζ_x along the fibers of $\mathbb{P}(T\mathcal{E}_x) \rightarrow \mathcal{E}_x$, for every $x \in M$. Consider the section $\psi : \mathcal{E} \rightarrow \mathcal{M}$ defined by $\psi(x, \xi) = \zeta_{(x,\xi)}$. We call $z = (x, \xi) \in \mathcal{E}$ a point of measurable continuity for ψ if there exists some probability measure ν on $\mathbb{P}(T_x\mathcal{E}_x)$ such that z is a Lebesgue density point of $\psi^{-1}(U)$ for any neighborhood U of ν (use any local trivialization of the fiber bundle \mathcal{E} ; the definition does not depend on the particular choice). Notice that ν is unique when it exists, and the set $\text{MC}(\psi)$ of points of measurable continuity has full m -measure. In that case define $\tilde{\psi}(z) = \nu$.

Lemma 5.8. *MC(ψ) is su -saturated and the section $\tilde{\psi} : \text{MC}(\psi) \rightarrow \mathcal{M}$ is su -invariant on $\text{MC}(\psi)$.*

Proof. The fact that $\{\zeta_x : x \in M\}$ is s -invariant means that $(\hat{H}_{x,y}^s)_*\zeta_x = \zeta_y$ for every x and y on the same stable leaf of f , where \hat{H}^s denotes the s -holonomy of the cocycle $\mathbb{P}(F)$ over f (which fibers over the s -holonomy H^s of the cocycle $\mathbb{P}(F)$ over F). Consequently, $(H_{z,w}^s)_*\zeta_z = \zeta_w$ for Lebesgue almost every $\xi \in \mathcal{E}_x$, where $w = (y, \eta)$ with $\eta = h_{x,y}^s(\xi)$. Since the holonomy maps $h_{x,y}^s$ are diffeomorphisms, and the $H_{z,w}^s$ are the fibers of the continuous map $\hat{H}_{x,y}^s$, it follows that measurable continuity points of the section ψ are preserved by the H^s . This proves s -saturation and s -invariance; the arguments for u -saturation and u -invariance are analogous. \square

Since we assume accessibility, this gives that $\text{MC}(\psi)$ is the whole \mathcal{E} and $\tilde{\psi}$ is su -invariant. Since $\tilde{\psi}$ coincides with ψ almost everywhere, it defines a disintegration of ζ . To conclude the proof we only have to check that $\tilde{\psi}$ is continuous. Given $z \in \mathcal{E}$, let us denote by $\mathcal{B}(z, N) \subset \mathcal{E}$ the set of points which are accessible from z through an su -path with not more than N legs, all of them contained in local stable or unstable manifolds.

Lemma 5.9. *There exists $N \geq 1$ such that $\mathcal{B}(z, N) = \mathcal{E}$ for every $z \in \mathcal{E}$.*

Proof. First, notice that, given any $\varepsilon > 0$ there exists $N(\varepsilon) \geq 1$ such that $\mathcal{B}(z, N)$ is ε -dense in \mathcal{E} . Indeed, otherwise there would exist $\varepsilon > 0$ and sequences z_N and w_N such that $\mathcal{B}(z_N, N)$ avoids the ball $B(w_N, \varepsilon)$ for every N . By compactness, it would follow that there exist z and w such that $w \notin \mathcal{B}(z, N)$ for every N . This would contradict the assumption of accessibility. Now fix $z_0 \in \mathcal{E}$. Clearly from the definition, $\mathcal{B}(z_0, N)$ is compact for every $N \geq 1$. Since $\bigcup_{N \geq 1} \mathcal{B}(z_0, N) = \mathcal{E}$, and \mathcal{E} is compact, there exists N_0 such that $\mathcal{B}(z_0, N)$ has non-empty interior. Hence it contains some ε -ball for some $\varepsilon > 0$. Thus, $\mathcal{B}(z, N(\varepsilon)) \cap \mathcal{B}(z_0, N_0) \neq \emptyset$ for every $z \in \mathcal{E}$. It follows that $\mathcal{B}(z, N) = \mathcal{E}$ for every $z \in \mathcal{E}$, with $N = 2(N_0 + N(\varepsilon))$. \square

Using this lemma, we can now upgrade measurable continuity to *uniform* measurable continuity, as follows. Fix any metric on the fibers of \mathcal{M} compatible with the weak* topology. We claim that for every $\varepsilon > 0$ and every sufficiently small ball B on any fiber \mathcal{E}_x (with respect to a fixed, but arbitrary Riemannian metric depending continuously on the fiber) there exists a subset W of B , with $\text{Leb}(W) > (1 - \varepsilon) \text{Leb}(B)$, such that $\tilde{\psi}(W)$ is contained in the ε -ball around $\tilde{\psi}(z)$ in \mathcal{M} . Indeed, for any two points $z \in \mathcal{E}_x$ and $w \in \mathcal{E}_y$, there exists a composition

$H : \mathcal{E}_x \rightarrow \mathcal{E}_y$ of at most N local holonomy maps such that $H(z) = w$. It follows that H has uniformly bounded derivative, and the corresponding projective extension $\hat{H} : \mathbb{P}(\mathcal{E}_x) \rightarrow \mathbb{P}(\mathcal{E}_y)$ is uniformly continuous. So, the quantifiers for measurable continuity at any two points are related with bounded distortion, yielding the claimed uniformity. Finally, it is easy to see that any uniformly measurable continuous function is in fact uniformly continuous in the fiber. Thus, the image under $\tilde{\psi}$ of any small ball in any fiber has small diameter in \mathcal{M} . Since ζ_x depends continuously on x , it follows that $\tilde{\psi}$ is continuous. \square

Combining Remark 5.5 with Proposition 5.7 one immediately obtains

Theorem 5.10. *Let F be a fiber bunched accessible volume preserving cocycle. If $\lambda_-(F, x, \xi) = \lambda_+(F, x, \xi) = 0$ at m -almost every $(x, \xi) \in \mathcal{E}$ then every $\mathbb{P}(F)$ -invariant probability that projects down to m admits a disintegration which is s -invariant, u -invariant, and whose conditional probabilities vary continuously with the base point on the support of μ .*

6. CONTINUITY AND POSITIVITY OF EXPONENTS

Here we start our analysis of area preserving cocycles, to prove Theorem F and Corollary G. Let us begin by observing that *every cocycle volume preserving cocycle admits some su -state*, namely, the measure m defined by (8). Indeed, it is clear that m is an F -invariant probability. Moreover, its disintegration m_x is invariant under s -holonomy and u -holonomy because, by part (4) of Proposition 5.1, all holonomy maps are volume preserving if the cocycle is. This means that, unlike the situation in Corollary E for instance, the methods we developed in the previous sections can not be applied directly to cocycles $F \in \mathcal{B}_{\text{vol}}(f)$.

Nevertheless, we are going to show that those criteria remain useful to obtain information on the Lyapunov exponents of F . The strategy is to apply them to the projective extension $\mathbb{P}(F)$ instead. As observed in Section 5.3, the fiber bunching condition ensures that $\mathbb{P}(F)$ is dominated and, hence, admits holonomies in a robust fashion. A fiber bunched cocycle F is called *bundle free* if its projective extension admits no su -states. Corollary 5.3 implies that this is an open condition (recall that at this point we take the fiber N to be compact). More generally, given any invariant su -saturated set U with positive Lebesgue measure, we say that F is *bundle free over U* if the projective extension has no su -state over U .

6.1. Discontinuity points. We are going to prove Theorem F. Let $F \in \mathcal{B}_{\text{vol}}(f)$ be ergodic for m and a discontinuity point for the Lyapunov exponents $\lambda_{\pm}(F, m)$. Recall that $\lambda_+(F, m) + \lambda_-(F, m) = 0$. It is well-known that the upper exponent $\lambda_+(\cdot, m)$ is upper semi-continuous and the lower exponent $\lambda_-(\cdot, m)$ is lower semi-continuous. Thus, if F is a discontinuity point then we must have

$$\lambda_-(F, m) < 0 < \lambda_+(F, m).$$

By ergodicity, this means that $\lambda_-(F, x, \xi) < 0 < \lambda_+(F, x, \xi)$ for m -almost every (x, ξ) . Let $T_{x, \xi} \mathcal{E} = E_{x, \xi}^s \oplus E_{x, \xi}^u$ be the Oseledets decomposition of F . For $* \in \{s, u\}$, denote by η_* the probability measure on $\mathbb{P}(\mathcal{E})$ which projects down to m under the fibration $\mathbb{P}(\mathcal{E}) \rightarrow \mathcal{E}$ and whose conditional probability measure on the fiber of each (x, ξ) is the Dirac mass at the Oseledets space $E_{x, \xi}^*$. Equivalently,

$$\eta_*(B) = m(\{(x, \xi) : (x, \xi, E_{x, \xi}^*) \in B\})$$

for every measurable set $B \subset \mathbb{P}(\mathcal{E})$. Notice that η_u is an invariant u -state and η_s is an invariant s -state of $\mathbb{P}(F)$. Let $\mathcal{M}(m)$ denote the space of probability measures η on $\mathbb{P}(\mathcal{E})$ that are mapped to m under the fibration $\mathbb{P}(\mathcal{E}) \rightarrow \mathcal{E}$ and, hence, project down to μ under $\mathbb{P}(\mathcal{E}) \rightarrow M$.

Lemma 6.1. *A measure $\eta \in \mathcal{M}(m)$ is $\mathbb{P}(F)$ -invariant if and only if it is a convex combination of η_u and η_s , that is, if $\eta = \alpha\eta_u + \beta\eta_s$ for some F -invariant functions $\alpha, \beta : M \rightarrow [0, 1]$ such that $\alpha + \beta = 1$.*

Proof. The ‘if’ part is trivial. For the converse just notice that every compact subset of $\mathbb{P}(\mathcal{E})$ disjoint from $\{E^u, E^s\}$ accumulates on E^u in the future and on E^s in the past. \square

Lemma 6.2. *The exponent $\lambda_+(F, m)$ coincides with the maximum of*

$$\int \log \|DF_x(\xi)v\| d\eta(x, \xi, v)$$

over all $\mathbb{P}(F)$ -invariant probability measures $\eta \in \mathcal{M}(m)$. When $\lambda_+(F, m) > 0$, the probability measure $\eta = \eta^u$ realizes the maximum.

Proof. Clearly, for any probability η that projects down to m ,

$$\frac{1}{n} \int \log \|DF_x^n(\xi)v\| d\eta(x, \xi, v) \leq \frac{1}{n} \int \log \|DF_x^n(\xi)\| dm(x, \xi).$$

The right hand side converges to $\lambda_+(F, m)$ when $n \rightarrow \infty$. The left hand side coincides with

$$\frac{1}{n} \int \sum_{j=0}^{n-1} \log \|DF_{x_j}(\xi_j)v_j\| d\eta(x, \xi, v) = \int \log \|DF_x(\xi)v\| d\eta(x, \xi, v),$$

where $(x_j, \xi_j, v_j) = \mathbb{P}(F)^j(x, \xi, v)$ and we take η to be $\mathbb{P}(F)$ -invariant. Combining these observations, one obtains the upper bound in the statement.

Now we only have to check that η_u realizes the maximum. To this end, notice

$$\frac{1}{n} \int \log \|DF_x^n(\xi)v\| d\eta^u(x, \xi, v) = \frac{1}{n} \int \log \|DF_x^n(\xi)v_u\| dm(x, \xi),$$

where $v_u = v_u(x, \xi)$ is a unit representative of $E_{x, \xi}^u$. By the previous arguments, the left hand side coincides with $\int \log \|DF_x(\xi)v\| d\eta^u(x, \xi, v)$, for every $n \geq 1$. By dominated convergence, the right hand side goes to

$$\int \lim_{n \rightarrow \infty} \frac{1}{n} \log \|DF_x^n(\xi)v_u\| dm(x, \xi) = \int \lambda_+(F, x, \xi) dm(x, \xi) = \lambda_+(F, m)$$

when $n \rightarrow \infty$. This proves our claim. \square

Proposition 6.3. *Let F be fiber bunched and ergodic. If F is a point of discontinuity for the Lyapunov exponent then every $\mathbb{P}(F)$ -invariant probability $\eta \in \mathcal{M}(m)$ is an su -state for $\mathbb{P}(F)$. In particular, F is not bundle free.*

Proof. The assumption implies there exists a sequence $(F_k)_k$ of cocycles converging to F in $\mathcal{B}_{\text{vol}}(f)$ such that $\lim_k \lambda_+(F_k, m) < \lambda_+(F, m)$ (the other inequality always holds, by semi-continuity of the largest exponent). Then, by Lemma 6.2, there exists some invariant u -state η_k for each $\mathbb{P}(F_k)$, such that

$$\lim_k \int \log \|DF_{k,z}(\xi)v\| d\eta_k < \lambda_+(F, m).$$

We may assume that $(\eta_k)_k$ converges to some probability measure η . Clearly, η is an invariant u -state for $\mathbb{P}(F)$. By Lemma 6.1,

$$\eta = \alpha\eta_u + \beta\eta_s$$

where α and β are constants (by ergodicity). Moreover,

$$\int \log \|DF_z(\xi)v\| d\eta = \lim_k \int \log \|DF_{k,z}(\xi)v\| d\eta_k < \lambda_+(F, m).$$

This implies that $\eta \neq \eta_u$ and, thus, β is not zero. It follows that η_s is a u -state for $\mathbb{P}(F)$, since η and η_u are. Analogously, η_u is an s -state for $\mathbb{P}(F)$. Therefore, η is an su -state for $\mathbb{P}(F)$. \square

Corollary 6.4. *If F is fiber bunched, ergodic, and bundle free then it is a point of continuity for the Lyapunov exponents and satisfies $\lambda_-(F, m) < 0 < \lambda_+(F, m)$.*

Remark 6.5. In the non-ergodic case we find that if F is a discontinuity point for the Lyapunov exponents then there exists an s -saturated positive measure set $Z^s \subset \mathcal{E}$ where $E_{x,\xi}^-$ is s -invariant and a u -saturated positive measure set $Z^u \subset \mathcal{E}$ where $E_{x,\xi}^+$ is u -invariant.

We are ready to finish the proof of Theorem F. We have seen in Proposition 6.3 that, under the theorem's assumptions, every $\mathbb{P}(F)$ -invariant probability in $\mathcal{M}(m)$ is an su -state. From Proposition 5.7 we conclude that it admits some disintegration which is su -invariant and continuous. This completes the proof.

6.2. Topological obstructions. In this section we observe that the topology of the fiber imposes certain restrictions on the behavior of the Lyapunov exponents. Corollary G is a consequence of the following result:

Theorem 6.6. *Let $F : \mathcal{E} \rightarrow \mathcal{E}$ be a fiber bunched area preserving cocycle admitting some open accessibility class C . If F is not bundle free over $U = \cup_{n \in \mathbb{Z}} F^n(C)$ then either*

- (1) *F is accessible, $N = \mathbb{S}^2$ or $N = \mathbb{T}^2$, and there exists a continuous Riemannian metric on the fibers, inducing the same area form, and which is invariant under both F and the invariant holonomies,*
- (2) *or F admits either an invariant continuous line field over U or an invariant pair of transverse continuous line fields over U .*

Proof. Let η be an su -state for the projective extension, and $\{\eta_z : z \in \text{supp } m\}$ be a continuous, $\mathbb{P}(F)$ -invariant and su -invariant disintegration of η . Observe that $\text{supp } m = \text{supp } \mu \times N$. For each $x \in M$, let U_x be the intersection of U with each fiber $\mathcal{E}_x = N$. Then U_x is an open subset of N . The definition implies that, given any points (x_0, ξ_0) and (x_1, ξ_1) in U there exist homeomorphisms $U_{x_0} \rightarrow U_{x_1}$ obtained by concatenating cocycle iterates and stable and unstable holonomies and mapping ξ_0 to ξ_1 . These homeomorphisms preserve the family of conditional probabilities.

Suppose first that for some (and, hence, for any) $z \in \text{supp } m$, the probability η_z admits some atom with mass at least $1/2$. Either such an atom is unique or there exist exactly two, that exhaust the total mass of the conditional probability. In the first case, the family of conditional probabilities defines a continuous map assigning to each point in U_x a point in projective space, that is, a continuous line field on U_x . Moreover, the line field is preserved by the cocycle and its invariant

holonomies. The second case is analogous, except that one gets an invariant pair of line fields instead. This gives part (2) of the theorem.

Now suppose that every η_z admits no atom with mass $1/2$ or larger. Then, by Douady, Earle [20, Section 2], the conditional measure has a well-defined conformal barycenter $\xi(z) \in \mathbb{D}$ and, consequently, it defines a conformal structure on the tangent space to the fiber at z . This endows every U_x with a Riemann surface structure. Together with the area form, this conformal structure defines a Riemannian metric on the tangent space to the leaves, which is invariant under the cocycle and its holonomies. In particular, the group of isometries acts transitively on every U_x . Thus (see Farkas, Kra [21, Theorem V.4]), U_x must be one of five exceptional surfaces: the sphere \mathbb{S} , the plane \mathbb{C} , the punched plane \mathbb{C}^* , the hyperbolic disk \mathbb{D} , or the torus \mathbb{T} . Moreover, the plane, the punched plane, and the disk may be excluded, since U_x has finite area. It follows that U_x is either the sphere or the torus and, in either case, coincides with the whole fiber N . In particular, F is accessible. \square

Remark 6.7. Part (2) of Theorem 6.6 can be strengthened considerably, if the cocycle is sufficiently regular: the fiber is $N = \mathbb{T}^2$ and the cocycle is conjugate to a shear $(\xi, \eta) \mapsto (\xi + t(x)\eta, \eta)$ on the fibers. This fact is neither proved nor used in this paper.

Proof of Corollary G. The assumptions of the corollary ensure we are in the setting of Theorem 6.6, with $C = U = \mathcal{E}$. The hypothesis on the genus excludes alternative 1 in the conclusion of the theorem. Alternative 2 is also similarly excluded: since the Euler characteristic of the fiber is non-zero, there can be no continuous vector field, nor pair of vector fields, over the whole \mathcal{E} . This proves that F must be bundle free. Now the conclusion follows from Corollary 6.4. \square

7. GENERIC AREA PRESERVING COCYCLES

Here we prove Theorem H: every $F \in \mathcal{B}_{\text{vol}}(f)$ is approximated by open sets where the Lyapunov exponents vary continuously and do not vanish. We begin with an outline of the arguments.

We have seen in Corollary 6.4 that if a cocycle F is bundle free and ergodic then its Lyapunov exponents are non-zero and they are continuous at F . We also know, from Corollary 5.3, that every bundle free cocycle is stably bundle free. In [4] we prove that every accessible cocycle with 2-dimensional fiber is stably accessible. By Proposition 5.6, every accessible fiber bunched volume preserving cocycle is ergodic. In [4] we also prove that every fiber bunched cocycle with 2-dimensional fiber is approximated by a (stably) accessible one. Thus, it suffices to prove that every accessible $F \in \mathcal{B}_{\text{vol}}(f)$ is approximated by a (stably) bundle free cocycle.

By Theorem 6.6, if the fiber N is a hyperbolic surface then F itself is bundle free, and so there is nothing to prove. Indeed, to finish the proof we only have to explain how to perturb the cocycle in each of the situations left open by Theorem 6.6, in order to make it bundle free. We use a simple mechanism to ensure the bundle free property: creation of non-degenerate elliptic periodic points of F on some periodic fiber. A few explanations are in order, before giving the details.

A periodic point p of an area preserving map $h : N \rightarrow N$ is *elliptic* if the eigenvalues of $Dh^n(p)$ are not real, where n denotes the period. We call the elliptic periodic point ζ *non-degenerate* if there exists $\kappa \neq 0$, $\epsilon > 0$, and a Diophantine

number $\alpha \in \mathbb{R}$, such that h^n is locally conjugate to

$$r \exp(2\pi i\theta) \mapsto r \exp(2\pi i\theta + \alpha + \kappa r^2) + O(r^5)$$

by some C^∞ diffeomorphism mapping ζ to $0 \in \mathbb{R}^2$. Then, by the Kolmogorov-Arnold-Moser theorem, there are arbitrarily small neighborhoods V of p which are C^∞ embedded disks invariant under h^n such that $h^n|_{\partial V}$ is conjugate to an irrational rotation and $\|Dh^n(x)\|$ grows linearly with n for every $x \in \partial V$. Consequently, h can not be an isometry with respect to any continuous Riemannian metric, and h can not preserve any continuous line field on N either.

Choose some periodic point $p \in M$ of the transformation f once and for all. Note that periodic points do exist, indeed they are dense in the support of μ : this follows from the Poincaré recurrence theorem, using the shadowing lemma (see Bowen [17]) to close recurrent trajectories. For simplicity we take the period to be 1.

Let us consider first the case when $N = \mathbb{S}^2$ and the cocycle and its holonomies are isometries with respect to some continuous Riemannian metric on the fibers. Since $F_p : \mathcal{E}_p \rightarrow \mathcal{E}_p$ is an orientable homeomorphism of the sphere, it has some fixed point $\zeta \in \mathcal{E}_p$. Since F_p is an isometry, this fixed point must be elliptic and degenerate. Perturb F near the fiber of p so as to make ζ non-degenerate. By the previous observations and Theorem 6.6, the new cocycle is bundle free.

Now let us consider the case when $N = \mathbb{T}^2$ and the cocycle and its holonomies are isometries. We claim that, perturbing the cocycle if necessary, the map F_p has some periodic point $\zeta \in \mathcal{E}_p$. If F_p is not homotopic to the identity then existence of a periodic point follows for topological reasons. If F_p is homotopic to the identity, then consider the rotation number

$$\rho(F_p) = \int (F_p - \text{id}) d\text{Leb}.$$

Perturbing F near the fiber over p in such a way that F_p is replaced by $F_p + v$ for some convenient $v \in \mathbb{T}^2$, we can ensure that the rotation number is rational. Then, by Franks [22], the map F_p has some periodic point. This proves the claim. From now on the argument is analogous to the sphere case: perturbing the cocycle once more, we can make the periodic point non-degenerate, and then the new cocycle must be bundle free.

Next, assume $N = \mathbb{T}^2$ and the cocycle admits a continuous invariant line field. Let V be the (open) set of all $x \in \mathbb{T}^2$ such that there exists K_x and ϵ_x such that for every w which is ϵ_x close to x , and every $k \geq 0$ such that $F_p^k(w)$ is ϵ_x -close to w , we have $\|DF_p^k(w)\| < K_x$. If there exists a periodic point of F_p in V , it must be elliptic and we can argue as before. So, assume that there is no periodic point in V . Similarly to what we did in the proof of Theorem 6.6, we can define on V a locally bounded measurable Riemannian metric inducing the same area form, which is F_p -invariant. Thus, V gets the structure of a one-dimensional complex manifold, possibly disconnected, on which F_p acts holomorphically. By Poincaré recurrence, all connected components of V are periodic. From the classification of conformal automorphisms (see [21, Chapter V]) we see that any automorphism of a Riemann surface which satisfies Poincaré recurrence admits a periodic point, unless the Riemann surface is an annulus and the automorphism is an irrational rotation, or the Riemann surface is the torus and the automorphism is not periodic. The torus case is covered by previous arguments (F_p is necessarily homotopic to the

identity and then a periodic point can be created by a small perturbation). So, we are left with the annulus case only.

For every $x \notin V$, choose sequences $w_n \rightarrow x$ and $k_n \rightarrow \infty$ such that $F_p^{k_n}(w_n) \rightarrow x$ and $\|DF_p^{k_n}(w_n)\| \rightarrow \infty$, the direction s_n most contracted under $DF_p^{k_n}(w_n)$ converges to some limit $s(x)$, and the image u_n of the direction most expanded under $DF_p^{k_n}(w_n)$ converges to some limit $u(x)$. Though the choice is not canonical, we fix it once and for all. Then $l(x) \in \{s(x), u(x)\}$, because every line bundle is attracted to $u(\cdot)$ under iteration, unless it coincides with $s(\cdot)$. Let h be the holonomy map associated to an arbitrary homoclinic loop of p . Since h preserves area, there exists a connected component V_0 of V and some $k \geq 1$ such that $h^k(V_0) \cap V_0 \neq \emptyset$. Using area preservation again, we conclude that $h^k(\partial V_0) \cap \partial V_0 \neq \emptyset$, which implies that there exists $z \in \mathbb{T}^2 \setminus V$ such that $h^k(z) \in \mathbb{T}^2 \setminus V$. Up to perturbing the dynamics without touching F_p , we may assume that $Dh^k(z) \cdot \{s(z), u(z)\} \cap \{s(h^k(z)), u(h^k(z))\} \neq \emptyset$. For the perturbed system, no line field or pair of transverse invariant line fields is invariant under both the dynamics and the invariant holonomies.

Finally, assume $N = \mathbb{T}^2$ and the cocycle admits a pair of transverse continuous invariant line fields $\{l_1(\xi), l_2(\xi)\}$, but no continuous invariant line field. Then F_p lifts to a map $\tilde{F}_p : \tilde{\mathcal{E}}_p \rightarrow \tilde{\mathcal{E}}_p$, where $\tilde{\mathcal{E}}_p$ is the set of all (x, ξ) with $x \in \mathbb{T}^2$ and $\xi \in \{l_1(x), l_2(x)\}$. The assumption that there is no invariant line field ensures that $\tilde{\mathbb{T}}^2$ is connected, and so it is a torus. Let $\pi : \tilde{\mathbb{T}}^2 \rightarrow \mathbb{T}^2$ be the projection on the first coordinate. Let \tilde{V} be the set of all $x \in \tilde{\mathbb{T}}^2$ such that there exists K_x and ϵ_x such that for every w which is ϵ_x close to x , and every $k \geq 0$ such that $\tilde{F}_p^k(w)$ is ϵ_x close to w we have $\|D\tilde{F}_p^k(w)\| < K_x$. Then the proof proceeds just as in the previous case, with $V = \pi(\tilde{V})$. In the present situation one gets $\{l_1(x), l_2(x)\} = \{s(x), u(x)\}$. This completes the proof of Theorem H.

8. RIGIDITY AND CENTER LYAPUNOV EXPONENTS

Here we prove Theorem I. The argument has two main parts, corresponding to Theorem 8.1 and Proposition 8.2 below. As mentioned before, all the arguments hold in finite differentiability, as long as it is large enough for [26] to hold.

Theorem 8.1. *There exists a neighborhood \mathcal{U}_0 of A in the space of C^∞ volume preserving diffeomorphisms on M such that if $f \in \mathcal{U}_0$ is accessible then its center Lyapunov exponents are distinct.*

The proof will appear in Sections 8.1 and 8.2. Here we just give an outline. Assume, by contradiction, that f is accessible and the two center Lyapunov exponents coincide. We deduce that there exists a translation surface structure on the center leaves which is invariant under s -holonomy, u -holonomy, and the dynamics. This is a consequence of the Invariance Principle, although we can not use Theorem D directly because the relevant base dynamics, the map f itself, is only *partially* hyperbolic (and not necessarily a skew-product, so that Theorem 5.10 is also not sufficient here). Instead, we use an extension for cocycles over volume preserving partially hyperbolic diffeomorphisms which is proven in [3]. In particular, this translation structure gives rise to an invariant \mathbb{R}^2 -action on the center leaves. Using accessibility once more, we promote this to a transitive action of some commutative group G by homeomorphisms of M . Then $G \simeq M$ and we also check that $f : M \rightarrow M$ corresponds to some automorphism of G . Up to the identification

$G \simeq M$, this automorphism must coincide with A . It follows that f is topologically conjugate to A , and we deduce that f is not accessible. This is a contradiction.

Proposition 8.2. *There exists a neighborhood $\mathcal{U} \subset \mathcal{U}_0$ of A in the space of C^∞ symplectic diffeomorphisms on M such that if $f \in \mathcal{U}$ and its center Lyapunov exponents coincide then f is conjugate to A by a volume preserving homeomorphism.*

The proof will appear in Section 8.3. Here is an outline. By Theorem 8.1, the hypothesis implies that f is not accessible. Then by Rodriguez-Hertz [26], the strong stable and strong unstable subbundles are jointly integrable and, in fact, the su -foliation of f is smoothly conjugate to the su -foliation of A . Moreover, the two maps are conjugate by some homeomorphism h . Using the fact that E^c is symplectic orthogonal to $E^s \oplus E^u$, we conclude that the center foliation is also smooth. We show that the su -holonomy preserves the family of area measures defined on the center leaves by the symplectic form ω , and we deduce that the conjugacy h is absolutely continuous along center leaves. Similarly, the center holonomy preserves the volume measures induced by ω on the su -leaves, and this implies that h is absolutely continuous along su -leaves as well. We deduce that h is absolutely continuous. Since both A and f ergodic, it follows that h preserves volume.

Remark 8.3. In dimension $d = 4$ the conjugacy h is a C^∞ diffeomorphism. This can be shown using ideas from Avila, Viana, Wilkinson [7] as we now outline; detailed arguments will appear in [7]. Let $h(x) = h^s(x) + h^c(x) + h^u(x)$ be the expression of the conjugacy (lifted to the universal cover) with respect to the partially hyperbolic splitting of A . It is observed in [26, Section 6] that h^c is a C^1 diffeomorphism from each center leaf of f to the center subspace of A . Moreover, h^s and h^u are constant on every center leaf of f . In fact, the same (KAM-type) arguments give much better regularity for h^c as long as f is sufficiently regular. In particular, in our setting h^c is C^∞ , and so h is C^∞ on every center leaf of f . Next, since h preserves area along the $E^s \oplus E^u$ direction and it also preserves the strong stable and the strong unstable foliations, it must preserve the disintegrations of Lebesgue measure along strong stable and strong unstable leaves. It is well known that the conditional measures are given by smooth densities (C^{k-1} if f is C^k , $k \geq 2$) which are positive and finite at every point. Since the leaves are 1-dimensional, the fact that h preserves these densities implies that h is C^∞ along strong stable leaves and along strong unstable leaves. Since strong stable, strong unstable, and center leaves span complementary directions, it follows (Journé [28]) that h is C^∞ in the ambient space, as stated.

To deduce Theorem I, consider any $f \in \mathcal{U}$. If the center Lyapunov exponents coincide then, by Proposition 8.2, the diffeomorphism f is volume preserving conjugate to A . In particular, f is Bernoulli since A is (Katznelson [30]). If the center Lyapunov exponents are distinct then, as f is symplectic, they must be non-zero. Then, f is non-uniformly hyperbolic. Since all the positive iterates of f are ergodic, we may use Ornstein, Weiss [36] to conclude that f is Bernoulli. So, we have indeed reduced the proof of Theorem I to proving Theorem 8.1 and Proposition 8.2.

8.1. Translation structures. Let us start the proof of Theorem 8.1. We think of A as a partially hyperbolic diffeomorphism on $M = \mathbb{T}^d$, with invariant splitting

$$TM = E^u \oplus E^c \oplus E^s, \quad \dim E^c = 2.$$

Then every diffeomorphism $f : M \rightarrow M$ in a neighborhood is also partially hyperbolic, with 2-dimensional center direction. Using that the center eigenvalues of A have norm 1, one easily gets that f is center bunched in the sense of [18, 3]: for some choice of the Riemannian structure

$$\begin{aligned} & \sup_{x \in M} \|Df|_{E_x^c}\| \|(Df|_{E_x^c})^{-1}\| \|Df|_{E_x^s}\| < 1 \\ \text{and} \quad & \sup_{x \in M} \|Df|_{E_x^c}\| \|(Df|_{E_x^c})^{-1}\| \|(Df|_{E_x^u})^{-1}\| < 1. \end{aligned}$$

Moreover, f is dynamically coherent, meaning that there exist invariant foliations W^{cs} and W^{cu} with C^1 leaves tangent to the subbundles $E^s \oplus E^c$ and $E^u \oplus E^c$, respectively, at every point. That is because A is dynamically coherent with smooth invariant foliations; see Theorems 7.1 and 7.2 in [27]. Throughout, we take f to be volume preserving and accessible and its center Lyapunov exponents to be equal.

Let $x, y \in M$ be any two points in the same strong stable leaf. Then $W^c(x)$ and $W^c(y)$ are contained in the same center stable leaf and the strong stable leaf through every $z \in W^c(x)$ intersects $W^c(y)$ at exactly one point. This defines a *strong stable holonomy* $h_{x,y}^s : W^c(x) \rightarrow W^c(y)$ with $h_{x,y}^s(x) = y$. Moreover, $h_{x,y}^s$ is a C^1 diffeomorphism. See [26, Appendix B]. Analogously, one constructs a *strong unstable holonomy* $h_{x,y}^u : W^c(x) \rightarrow W^c(y)$ for every x and y in the same strong unstable leaf.

Proposition 8.4. *There exists a conformal structure on the center leaves of f that is continuous and invariant under the diffeomorphism, strong stable holonomies, and strong unstable holonomies.*

Proof. Let $F : E^c \rightarrow E^c$ be the restriction of the derivative Df to the center bundle and $\mathbb{P}(F) : \mathbb{P}(E^c) \rightarrow \mathbb{P}(E^c)$ be the projectivization of F . We think of F and $\mathbb{P}(F)$ as cocycles over the partially hyperbolic diffeomorphism f . The previous observations ensure these are cocycles with holonomies, in the sense of [3]: the s -holonomies of F and $\mathbb{P}(F)$ are given by

$$DH_{x,y}^s(x) : E_x^c \rightarrow E_y^c \quad \text{and} \quad \mathbb{P}(DH_{x,y}^s(x)) : \mathbb{P}(E_x^c) \rightarrow \mathbb{P}(E_y^c),$$

for every x, y in the same strong stable leaf of f ; the u -holonomies are defined analogously. Then, since f is accessible and the center Lyapunov exponents are equal, we may apply [3, Theorem B] to conclude that any $\mathbb{P}(F)$ -invariant probability measure η on $\mathbb{P}(E^c)$ projecting down to Lebesgue measure on M admits an su -invariant continuous disintegration $\{\eta_x : x \in M\}$ into conditional probabilities along the fibers $\mathbb{P}(E_x^c)$. Let η be fixed (arbitrarily).

The hypotheses on A imply that no eigenvalue is a root of unity, so $D_0A|_{E^c} = A|_{E^c}$ is an irrational rotation. Every nearby diffeomorphism has a unique fixed point near $0 \in M$ and, up to conjugating f with a small translation, we may suppose it to sit at 0. Then the derivative of f along the center direction at 0 has no periodic points with small period. In particular, η_0 has no atoms of mass $1/2$ or larger and then, by holonomy invariance and accessibility, the same is true for every conditional probability η_x , $x \in M$. Then, by Douady, Earle [20, Section 2], the conditional measure η_x determines a unique conformal structure on E_x^c .¹ By

¹Recall that one can see $\mathbb{P}\mathbb{R}^2$ as the boundary of the Poincaré half-plane $\mathbb{H} \subset \mathbb{P}\mathbb{C}^2$. The construction of Douady-Earle associates to a probability measure μ on $\mathbb{P}\mathbb{R}^2$ (with no atom of mass $1/2$ or larger), a ‘‘conformal barycenter’’ $z \in \mathbb{H}$ so that $\mu \mapsto z$ is continuous (with respect to the weak-* topology on its domain) and equivariant with respect to conformal automorphisms of

construction, this conformal structure depends continuously on the point x and is invariant under strong stable holonomies, strong unstable holonomies, and the diffeomorphism f itself. \square

The next step is to upgrade this conformal structure to a translation structure, that is, a system of local coordinates on each center leaf such that all coordinate changes are translations.

Proposition 8.5. *There exists a translation structure on the center leaves of f that is continuous and invariant under strong stable and strong unstable holonomies.*

Proof. The idea is quite simple. As explained before, it is no restriction to suppose that $f(0) = 0$. We will show in a moment that the center leaf $W^c(0)$ through the fixed point is conformally equivalent to \mathbb{C} . Then an arbitrary choice of a conformal isomorphism determines a translation structure on $W^c(0)$, which we push around by strong stable and strong unstable holonomies. More precisely, given any $x \in M$ we choose an su -path $\gamma = (0, x_1, \dots, x_{n-1}, x)$ connecting 0 to x and we endow $W^c(x)$ with the translation structure transported from $W^c(0)$ through

$$h_\gamma : W^c(0) \rightarrow W^c(x), \quad h_\gamma = h_{x_{n-1}, x}^{*n} \circ \dots \circ h_{x_{i-1}, x_i}^{*i} \circ \dots \circ h_{0, x_1}^{*1}$$

where $*_i \in \{s, u\}$. The following lemma ensures that the definition does not depend on the choice of the path γ :

Lemma 8.6. *For any su -path $\gamma = (0, x_1, \dots, x_{n-1}, x)$ with $x \in W^c(0)$, the holonomy $h_\gamma : W^c(0) \rightarrow W^c(0)$ is a translation for the chosen translation structure.*

Before starting the proof of the lemma, let us recall some useful facts from [26, Appendix B]. Let $p : \mathbb{R}^d \rightarrow \mathbb{T}^d$ be the universal cover and $F : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be the lift of f that fixes the origin. Each f -invariant foliation W^* , $*$ $\in \{s, cs, c, cu, u\}$ lifts to an F -invariant foliation \hat{W}^* on \mathbb{R}^d with embedded leaves, and p restricts to an injective immersion on each leaf.

The leaves may be written in the form

$$\hat{W}^*(\hat{x}) = \hat{x} + \text{graph}(\gamma_{\hat{x}}^*), \quad \text{with} \quad \begin{array}{ll} \gamma_{\hat{x}}^s : E^s \rightarrow E^{cu} & \gamma_{\hat{x}}^u : E^u \rightarrow E^{cs} \\ \gamma_{\hat{x}}^c : E^c \rightarrow E^{su} & \\ \gamma_{\hat{x}}^{cs} : E^{cs} \rightarrow E^u & \gamma_{\hat{x}}^{cu} : E^{cu} \rightarrow E^s. \end{array}$$

This induces a natural parametrization

$$(38) \quad E^* \ni v \mapsto \hat{x} + v + \gamma_{\hat{x}}^*(v)$$

of each leaf $\hat{W}^*(\hat{x})$ in \mathbb{R}^d and, composing with the projection $p : \mathbb{R}^d \rightarrow \mathbb{T}^d$, of each leaf $W^*(x)$ in \mathbb{T}^d . The center leaves are uniformly close to the corresponding subbundles: there is $\kappa = \kappa(f)$ such that $\kappa \rightarrow 0$ when $f \rightarrow A$ and

$$(39) \quad \|\gamma_{\hat{x}}^*(v)\| \leq \kappa \quad \text{for all } \hat{x} \in \mathbb{R}^d, v \in E^*, \text{ and } * \in \{cs, c, cu\}.$$

In addition,

$$(40) \quad \|\gamma_{\hat{x}}^*(v)\| \leq \kappa \|v\| \quad \text{for all } \hat{x} \in \mathbb{R}^d, v \in E^*, \text{ and } * \in \{s, u\}.$$

\mathbb{H} (the real projective transformations). The stabilizer of z in $\text{PSL}(2, \mathbb{R})$ is a maximal compact subgroup, i.e., a conformal structure in \mathbb{R}^2 . By projective equivariance, the conformal barycenter construction extends canonically to the level of two-dimensional real vector spaces, so it can be applied to E^c .

Moreover, center leaves through nearby points are uniformly close to each other: given $\varepsilon > 0$ and $* \in \{cs, c, cu\}$ there exists δ such that

$$(41) \quad d(\hat{x}, \hat{y}) \leq \delta \quad \text{implies} \quad \sup\{\|\gamma_{\hat{x}}^*(v) - \gamma_{\hat{y}}^*(v)\| : v \in E^*\} \leq \varepsilon.$$

This is easily read out from the expression of γ^* in the proof of [26, Lemma B3].

Using that the leaves of the \hat{W}^c are graphs over E^c we get:

Lemma 8.7. *$W^c(0)$ is conformally equivalent to \mathbb{C} .*

Proof. Fix an (\mathbb{R} -linear) isomorphism $\mathbb{C} \rightarrow E^c$. It defines in particular a conformal structure on E^c , which can be used to define an alternative conformal structure along the leaves of the central foliation \hat{W}^c : restricted to each $\hat{W}^c(x)$, we declare the projection on E^c along E^{su} (which is a C^1 diffeomorphism) to be conformal. It descends to an alternative conformal structure along the leaves of the central foliation W^c . Since p is injective along leaves, $W^c(0)$ is conformally equivalent to \mathbb{C} with respect to this alternative conformal structure. But both the original and the alternative conformal structures are continuous, so the Beltrami coefficient which relates both defines a continuous function on \mathbb{T}^d , which, by compactness, must be bounded away from 1. So the original and the alternative conformal structures on $W^c(0)$ are quasiconformally equivalent. This shows that $W^c(0)$ (with the original conformal structure) is quasiconformally equivalent to \mathbb{C} , and by the uniformization theorem it must be actually conformally equivalent to \mathbb{C} . \square

For every $\hat{x}, \hat{z} \in \mathbb{R}^d$ the strong stable leaf $\hat{W}^s(\hat{x})$ intersects $\hat{W}^{cu}(\hat{z})$ at exactly one point \hat{y} . Varying \hat{x} inside its center leaf this defines a homeomorphism

$$\hat{h}_{\hat{x}, \hat{y}}^s : \hat{W}^c(\hat{x}) \rightarrow \hat{W}^c(\hat{y})$$

that lifts the stable holonomy $h_{x,y}^s : W^c(x) \rightarrow W^c(y)$, $x = p(\hat{x})$, $y = p(\hat{y})$. Lifts of the unstable holonomies are constructed in the same way.

A special family of holonomies is defined as follows. For each $\mathbf{n} \in \mathbb{Z}^d$, let $x_{\mathbf{n}}$ be the unique point in $\hat{W}^u(\mathbf{n}) \cap \hat{W}^{cs}(0)$ and $y_{\mathbf{n}}$ be the unique point in $\hat{W}^s(x_{\mathbf{n}}) \cap \hat{W}^c(0)$. Let

$$\hat{t}_{\mathbf{n}} : \hat{W}^c(\mathbf{n}) \rightarrow \hat{W}^c(0), \quad \hat{t}_{\mathbf{n}} = \hat{h}_{x_{\mathbf{n}}, y_{\mathbf{n}}} \circ \hat{h}_{0, x_{\mathbf{n}}}^u$$

and $t_{\mathbf{n}} : W^c(0) \rightarrow W^c(0)$ be its projection down to \mathbb{T}^d . By [26, Corollary 2.4], the expression of these holonomy maps in the parametrization (38) has the form

$$(42) \quad t_{\mathbf{n}}(v) = v + \mathbf{n}^c + \phi(\mathbf{n}, v),$$

where \mathbf{n}^c is the component of \mathbf{n} in the direction of E^c and ϕ is close to zero, uniformly on \mathbf{n} and $v \in E^c$, if f is close to A . If \mathbf{n} is non-zero then so is \mathbf{n}^c and then, assuming f is close to A , this expression implies that $t_{\mathbf{n}}$ has no fixed points.

Proof of Lemma 8.6. We use two different global coordinates $z \in \mathbb{C}$ and $v \in E^c$ on the center leaf $W^c(0)$: the former arises from the chosen uniformization $\mathbb{C} \rightarrow W^c(0)$, whereas the latter corresponds to the parametrization (38). Accordingly, each $W^c(0) \rightarrow W^c(0)$ may be viewed as a transformation in either \mathbb{C} or E^c .

Since the conformal structure is holonomy invariant, h_{γ} is a conformal automorphism, that is, an affine transformation $h_{\gamma}(z) = az + b$. We want to show that $a = 1$. We begin by showing that h_{γ} displaces points by a bounded amount along the center leaf. Given any $\hat{x}, \hat{y} \in \mathbb{R}^d$ in the same strong stable leaf there exists $C(\hat{x}, \hat{y}) > 0$ such that $\|\hat{h}_{\hat{x}, \hat{y}}^s(\xi) - \xi\|_{\mathbb{R}^d} \leq C(\hat{x}, \hat{y})$ for every $\xi \in \hat{W}^c(\hat{x})$. That is because center leaves are at uniformly bounded distance from the direction of E^c and strong stable

leaves are uniformly close to the direction of E^s on compact parts; recall (39) and (40). For the same reasons, we have an analogous estimate for strong unstable holonomies. Consequently, there is $C(\gamma) > 0$ such that $\|\hat{h}_\gamma(\xi) - \xi\|_{\mathbb{R}^d} \leq C(\gamma)$ for every $\xi \in \hat{W}^c(0)$. Since (38) is a bi-Lipschitz embedding, this translates to the coordinate v :

$$(43) \quad \sup_{v \in E^c} \|h_\gamma(v) - v\|_{E^c} < \infty.$$

Now consider $t_{\mathbf{n}} : W^c(0) \rightarrow W^c(0)$ for any non-zero $\mathbf{n} \in \mathbb{Z}^d$. As observed before, $t_{\mathbf{n}}$ has no fixed points if f is close enough to A . Hence, $t_{\mathbf{n}}(z) = z + c$ for some $c \in \mathbb{C}$ different from zero. Consider the map

$$\phi_k = t_{\mathbf{n}}^k \circ h_\gamma : W^c(0) \rightarrow W^c(0) \quad \text{for } k \geq 1.$$

On the one hand, $\phi_k(z) = az + b + kc$ for every k . On the other hand, (42) and (43) imply that ϕ_k has no fixed points if k is large enough. This can only happen if $a = 1$. So the proof of the lemma is complete. \square

This ensures that the definition of the translation structure on every center leaf is indeed well-defined. By construction, this structure is holonomy invariant and, hence, continuous. This finishes the proof of Proposition 8.5. \square

8.2. Continuous group actions. In the previous section we defined a translation structure on the center leaves of f . This gives rise to an \mathbb{R}^2 -action

$$T : \mathbb{R}^2 \times M \rightarrow M, \quad (v, x) \mapsto T_v(x) = x + v$$

where $x + v$ denotes the v -translate of x along its center leaf. Since the translation structure is continuous and su -invariant, each T_v is a homeomorphism of M and commutes with both strong stable and strong unstable holonomy.

Proposition 8.8. *The closure G of $\{T_v : v \in \mathbb{R}^2\}$ is a compact subgroup of the group of homeomorphisms of M .*

Lemma 8.9. *For every $\varepsilon > 0$ there is $\delta > 0$ such that given any $x, z \in M$ with $d(x, z) \leq \delta$ there is some su -path γ such that $h_\gamma(x)$ belongs to the local central leaf of z with $d(h_\gamma(x), z) \leq \varepsilon$ and $d(h_\gamma(\xi), \xi) \leq \varepsilon$ for every $\xi \in W^c(x)$.*

Proof. By transversality of the stable, unstable, and center directions, one can find $\delta > 0$ such that for any x and z with $d(x, z) \leq \delta$ there exists two-legs su -paths $\gamma = (x, y, \zeta)$ with ζ in the local central leaf of z . Clearly, we may always choose γ such that y is in the strong unstable leaf through x and ζ is in the strong stable leaf through y . Moreover, the lengths of the legs and the distance from z to ζ can be made uniformly small by reducing δ . Then both $h_{x,y}^u : W^c(x) \rightarrow W^c(y)$ and $h_{y,\zeta}^s : W^c(y) \rightarrow W^c(z)$ are uniformly close to the identity, meaning that

$$\sup_{\xi \in W^c(x)} d(h_{x,y}^u(\xi), \xi) \quad \text{and} \quad \sup_{\eta \in W^c(x)} d(h_{y,\zeta}^s(\eta), \eta)$$

are small. That is because the center leaves are uniformly close to each other and strong (stable or unstable) leaves are uniformly transverse to the direction of E^c on compact parts; recall (41) and (40). In particular, $h_\gamma = h_{y,\zeta}^s \circ h_{x,y}^u$ is uniformly ε -close to the identity and $\zeta = h_\gamma(x)$ is ε -close to z , as stated. \square

Lemma 8.10. *For every $\varepsilon > 0$ there exists $\delta > 0$ such that, given any $p \in M$ and any $u \in \mathbb{R}^2$, if $d(T_u(p), p) \leq \delta$ then $d(T_u(x), x) \leq \varepsilon$ for every $x \in M$.*

Proof. Given $\varepsilon > 0$ take $\delta > 0$ as in Lemma 8.9. Then there exists some su -path γ such that $h_\gamma : W^c(p) \rightarrow W^c(p)$ satisfies

$$(44) \quad \begin{aligned} & h_\gamma(T_u(p)) \text{ belongs to the local center leaf of } p \text{ and } d(h_\gamma(T_u(p)), p) \leq \varepsilon \\ & \text{and } d(h_\gamma(\xi), \xi) \leq \varepsilon \text{ for all } x \in W^c(p). \end{aligned}$$

Before proceeding, let us observe that translations are uniformly Lipschitz on center leaves: there exists $\kappa > 0$ such that

$$d(T_v(x), T_v(y)) \leq \kappa d(x, y) \quad \text{for any } v \in \mathbb{R}^2 \text{ and } x, y \text{ in the same local center leaf.}$$

Indeed, translations are isometries relative to the flat metric defined on the center leaves by the translation structure. Since this flat metric is continuous on the whole compact M , it must be uniformly equivalent to the one obtained by restricting to center leaves the Riemannian structure of M . This gives the claim. Now, any point $q \in W^c(p)$ may be written as $q = T_v(p)$ for some $v \in \mathbb{R}^2$. Then,

$$d(T_u(q), q) \leq d(T_u(q), h_\gamma(T_u(q))) + d(h_\gamma(T_{u+v}(p)), T_v(p)).$$

The first term on the right is $\leq \varepsilon$, by the last part of (44). As for the second term, since the translation structure is holonomy invariant,

$$d(h_\gamma(T_{u+v}(p)), T_v(p)) = d(T_v(h_\gamma(T_u(p))), T_v(p)) \leq \kappa d(h_\gamma(T_u(p)), p) \leq \kappa \varepsilon.$$

This proves that $d(T_u(q), q) \leq (1 + \kappa)\varepsilon$ for all $q \in W^c(p)$. Since $W^c(p)$ is dense in M , this gives the desired estimate (up to an irrelevant factor of $1 + \kappa$). \square

Corollary 8.11. *For every $\varepsilon > 0$ there is $\rho > 0$ such that for every $v \in \mathbb{R}^2$ there exists $w \in \mathbb{R}^2$ with $\|w\| \leq \rho$ and $d(T_v(x), T_w(x)) \leq \varepsilon$ for all $x \in M$.*

Proof. Given $\varepsilon > 0$, let $\delta > 0$ be as given by Lemma 8.10. Since $W^c(0)$ is dense in M , there exists $\rho > 0$ such that $\{T_w(0) : \|w\| \leq \rho\}$ is δ -dense in M . Then, given any $v \in \mathbb{R}^2$ we can find $w \in \mathbb{R}^2$ such that $\|w\| \leq \rho$ and $d(T_v(0), T_w(0)) \leq \delta$. By Lemma 8.10, with $u = v - w$ and $p = T_w(0)$, it follows that $d(T_v(x), T_w(x)) \leq \varepsilon$ for all $x \in M$. This proves the corollary. \square

Proof of Proposition 8.8. Let $(v_n)_n$ be any sequence in \mathbb{R}^2 . We want to show that the sequence $(T_{v_n})_n$ has some convergent subsequence. Taking $\varepsilon = 2^{-k}$, $k \geq 1$ in Corollary 8.11, we find $\rho_k > 0$ and for each $n \geq 1$ we find $w_{k,n} \in \mathbb{R}^2$ with $\|w_{k,n}\| \leq \rho_k$ and

$$(45) \quad d(T_{v_n}(x), T_{w_{k,n}}(x)) \leq 2^{-k} \quad \text{for every } x \in M.$$

Since $(w_{k,n})_n$ is bounded, for every $k \geq 1$, we may use a diagonal argument to find a sequence $(n_j)_j \rightarrow \infty$ such that $w_k = \lim_j w_{k,n_j}$ exists for every $k \geq 1$. The triangle inequality gives

$$d(T_{w_{k,n_j}}(x), T_{w_{l,n_j}}(x)) \leq 2^{-k} + 2^{-l} \quad \text{for all } j, k, l \geq 1, \text{ and } x \in M.$$

Taking the limit as $j \rightarrow \infty$, we get $d(T_{w_k}(x), T_{w_l}(x)) \leq 2^{-k} + 2^{-l}$ for all $k, l \geq 1$, and $x \in M$. That implies $(T_{w_k})_k$ is a Cauchy sequence, relative to the uniform norm. Let T be the limit. It is clear that T is a homeomorphism of M : the inverse is the limit of $(T_{-w_k})_k$. Moreover, (45) implies that $T_{v_{n_j}}$ also converges to T . This proves the proposition. \square

Lemma 8.12. *The action of G on M is commutative, transitive, and free. Consequently, the map $G \ni \phi \mapsto \phi(0) \in M$ is a homeomorphism.*

Proof. The first claim is clear: since the \mathbb{R}^2 -action is commutative, so is its closure. The second claim is also easy: since the orbits of the \mathbb{R}^2 -action coincide with the center leaves, and these are dense in M , the action of the closure G is transitive in M . Finally, Lemma 8.10 implies that if an element of G has some fixed point then it must be the identity. This means that the action is free. As for the last claim in the lemma, transitivity implies $\phi \mapsto \phi(0)$ is onto, and freeness implies it is injective. Thus, it is a homeomorphism, as claimed. \square

Let M_f denote the torus \mathbb{T}^d endowed with the group structure transported from G via the homeomorphism in Lemma 8.12: the group operation corresponds to the composition in G . In the sequel we think of M as the torus \mathbb{T}^d endowed with the standard group structure.

Lemma 8.13. *The diffeomorphism f is a group automorphism of M_f . Moreover, $f : M_f \rightarrow M_f$ is conjugate to $A : M \rightarrow M$ by some group isomorphism $M \rightarrow M_f$.*

Proof. We begin by noting that the holonomy map $h_\gamma : W^c(x) \rightarrow W^c(y)$ associated to any su -path $\gamma = (x, \dots, y)$ extends to a homeomorphism of \mathbb{T}^d , namely, the unique element $\phi_{x,y}$ of G that maps x to y . Indeed, $h_\gamma(\xi) = \phi_{x,y}(\xi)$ for every $\xi \in W^c(x)$, because the equality holds for $\xi = x$ and both maps commute with all translations. This correspondence $\gamma \mapsto \phi_{x,y}$ is a representation of the groupoid of su -paths in the group G : concatenation $*$ of su -paths corresponds to composition of the associated homeomorphisms of \mathbb{T}^d . It is clear that this representation is surjective. We have $f \circ \phi_{x,y} = \phi f(x), f(y) \circ f$ for every x and y , because $f \circ h_\gamma = h_{f(\gamma)} \circ f$ for any su -path γ . Thus, since $f(0) = 0$,

$$f(x + y) = f \circ \phi_{0,y}(x) = \phi_{0,f(y)}(f(x)) = f(x) + f(y)$$

for any $x, y \in M_f$. This proves the first claim.

To prove the second one it suffices to show that the lift $\hat{f} : \hat{M}_f \rightarrow \hat{M}_f$ of f to the universal cover is conjugate to A . Since f is an automorphism of M_f the lift is determined by its action on the fiber O of the fixed point 0 . Given any homology class $[\gamma] \in H_1(M, \mathbb{Z})$ consider a representative γ through the fixed point $0 \in M_f$ and let $\hat{\gamma}$ be the its lift starting at $0 \in \hat{M}_f$. The map

$$\psi : H_1(M, \mathbb{Z}) \rightarrow O, \quad \psi([\gamma]) = \text{endpoint of } \hat{\gamma}$$

is a group isomorphism (accessibility ensures surjectivity). Moreover, $\psi([f(\gamma)])$ coincides with $\hat{f}(\psi([\gamma]))$ for every $[\gamma]$. In other words, the isomorphism ψ conjugates \hat{f} to the action of f on the homology, that is, to A . This completes the proof. \square

The strong stable, strong unstable, and center foliations of $A : M \rightarrow M$ are dynamically defined, and so they are preserved by the conjugacy in Lemma 8.13. It follows that f has an su -foliation W^{su} whose leaves are subfoliated by strong stable leaves and by strong unstable leaves. That implies that f is not accessible, contradicting the hypotheses. This contradiction completes the proof of Theorem 8.1.

8.3. Integrable case. By Rodriguez-Hertz [26], any non-accessible volume preserving diffeomorphism f in a neighborhood of A is conjugate to A by some homeomorphism H . Our goal is to show that H preserves the volume on M . This will prove Proposition 8.2.

Clearly, H conjugates the strong stable, strong unstable, and center foliations of f to the corresponding invariant foliations of A . Moreover, the accessibility

classes of f form an invariant foliation W^{su} by smooth submanifolds tangent to the subbundle $E^c \oplus E^u$, and [26] also proves that this foliation is conjugate to the su -foliation of A by some C^1 diffeomorphism. Consequently, the su -foliation W^{su} is C^1 . Since E^c is the symplectic orthogonal of $E^{su} = E^s \oplus E^u$, it follows that the center foliation W^c of f is also C^1 .

Lemma 8.14. *Let ω be a non-degenerate continuous 2-form on a manifold M such that $\int_{\partial A} \omega = 0$ for every C^1 embedded image A of $[-1, 1]^3$. Let \mathcal{F}^1 and \mathcal{F}^2 be C^1 transverse foliations of M whose tangent spaces are ω -orthogonal to each other at every point. Then the restriction of ω to the leaves of \mathcal{F}^2 is invariant under the holonomy maps of \mathcal{F}^1 .*

Proof. We want to prove that if x and y are nearby points in the same \mathcal{F}^1 -leaf and h denotes the projection along \mathcal{F}^1 -leaves from a neighborhood of x inside \mathcal{F}_x^2 to a neighborhood of y inside \mathcal{F}_y^2 then

$$(46) \quad Dh(x)_*(\omega(x) | T_x \mathcal{F}^2 \times T_x \mathcal{F}^2) = (\omega(y) | T_y \mathcal{F}^2 \times T_y \mathcal{F}^2).$$

Let $I = [-1, 1]$ and denote $\hat{x} = (-1, 0, 0)$ and $\hat{y} = (1, 0, 0)$. Given any linearly independent vectors $v, w \in T_x \mathcal{F}^2$ consider a C^1 embedding $\psi : I^3 \rightarrow M$ such that

- $\psi(\hat{x}) = x$ and $\psi(\hat{y}) = y$
- $D\psi(\hat{x}) \cdot (\{0\} \times \mathbb{R}^2)$ coincides with the plane generated by v and w
- every $\psi(\{a\} \times I^2)$, with $a \in I$ is contained in some \mathcal{F}^2 -leaf
- every $\psi(I \times \{b\} \times \{c\})$, with $b, c \in I$ is contained in some \mathcal{F}^1 -leaf.

Let ω_ψ denote the pull-back of ω under ψ . Observe that the pull-back of the holonomy h under ψ is just the trivial projection $\{-1\} \times I^2 \rightarrow \{1\} \times I^2$. So, to prove (46) it suffices to show that

$$(47) \quad \omega_\psi(\hat{x}) | \{0\} \times \mathbb{R}^2 = \omega_\psi(\hat{y}) | \{0\} \times \mathbb{R}^2$$

for every x, y, u, v , and any choice of ψ . To this end, observe that the hypothesis of the lemma implies

$$(48) \quad \int_{\partial(I \times (\varepsilon I)^2)} \omega_\psi = 0 \quad \text{for every } \varepsilon > 0, \text{ where } \varepsilon I = [-\varepsilon, \varepsilon].$$

By construction, every $D\psi(\xi)$ maps $\mathbb{R}^2 \times \{0\}$ either to a line, or to a plane V_ξ that intersects both the tangent space to $\mathcal{F}_{\psi(\xi)}^1$ and the tangent space to $\mathcal{F}_{\psi(\xi)}^2$. Since the two tangent spaces are ω -orthogonal, it follows that $\omega(\psi)$ vanishes on V_ξ . This means that $\omega_\psi(\xi) | \mathbb{R}^2 \times \{0\} = 0$ at every $\xi \in I^3$. Analogously, $\omega_\psi(\xi) | \mathbb{R} \times \{0\} \times \mathbb{R} = 0$ at every $\xi \in I^3$. Thus, (48) may be rewritten as

$$(49) \quad \int_{\{-1\} \times (\varepsilon I)^2} \omega_\psi | \{0\} \times \mathbb{R}^2 = \int_{\{1\} \times (\varepsilon I)^2} \omega_\psi | \{0\} \times \mathbb{R}^2 \quad \text{for every } \varepsilon > 0.$$

Taking the limit as ε goes to 0 we get (47). This proves the lemma. \square

Let ω be the symplectic form on M . From Lemma 8.14 we get that the restriction of ω to center leaves is invariant under the holonomy maps of the su -foliation, and then so is the family η of measures defined by ω on center leaves. Then the push-forward $H_*\eta$ is a family of measures on the center leaves of A invariant under the su -holonomy maps of A . By unique ergodicity of irrational foliations on the torus, $H_*\eta$ must coincide, up to scaling, with the family of Lebesgue measures on the center leaves. This implies that the homeomorphism H has a continuous Jacobian along

every center leaf. Exchanging the roles of the center and su -foliations throughout this argument we get that H has a continuous Jacobian along su -leaves as well. Since these foliations are C^1 , it follows that $H : M \rightarrow M$ is absolutely continuous. Thus, the push-forward H_*m of the normalized volume measure in M is an A -invariant measure absolutely continuous with respect to m . Since A is ergodic, H_*m must coincide with m . This finishes the proof of Proposition 8.2 and Theorem I.

REFERENCES

- [1] F. Abdenur and M. Viana. Flavors of partial hyperbolicity. Preprint www.preprint.impa.br 2009.
- [2] J. F. Alves, C. Bonatti, and M. Viana. SRB measures for partially hyperbolic systems whose central direction is mostly expanding. *Invent. Math.*, 140:351–398, 2000.
- [3] A. Avila, J. Santamaria, and M. Viana. Cocycles over partially hyperbolic maps. Preprint www.preprint.impa.br 2008.
- [4] A. Avila and M. Viana. On stable accessibility with 2-dimensional center. Preprint www.preprint.impa.br 2009.
- [5] A. Avila and M. Viana. Simplicity of Lyapunov spectra: a sufficient criterion. *Port. Math.*, 64:311–376, 2007.
- [6] A. Avila and M. Viana. Simplicity of Lyapunov spectra: proof of the Zorich-Kontsevich conjecture. *Acta Math.*, 198:1–56, 2007.
- [7] A. Avila, M. Viana, and A. Wilkinson. Absolute continuity, Lyapunov exponents, and rigidity. Preprint www.preprint.impa.br 2009.
- [8] L. Barreira, Ya. Pesin, and J. Schmeling. Dimension and product structure of hyperbolic measures. *Ann. of Math.*, 149:755–783, 1999.
- [9] P. Baxendale. Lyapunov exponents and relative entropy for a stochastic flow of diffeomorphisms. *Probab. Theory Related Fields*, 81:521–554, 1989.
- [10] J. Bochi. Genericity of zero Lyapunov exponents. *Ergod. Th. & Dynam. Sys.*, 22:1667–1696, 2002.
- [11] J. Bochi. C^1 -generic symplectic diffeomorphisms: partial hyperbolicity and zero centre Lyapunov exponents. *J. Inst. Math. Jussieu*, 8, 2009.
- [12] J. Bochi and M. Viana. The Lyapunov exponents of generic volume-preserving and symplectic maps. *Ann. of Math.*, 161:1423–1485, 2005.
- [13] C. Bocker and M. Viana. *Continuity of Lyapunov exponents for 2D random matrices*. PhD thesis, IMPA, 2009.
- [14] C. Bonatti, X. Gómez-Mont, and M. Viana. Généricité d’exposants de Lyapunov non-nuls pour des produits déterministes de matrices. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 20:579–624, 2003.
- [15] C. Bonatti and M. Viana. SRB measures for partially hyperbolic systems whose central direction is mostly contracting. *Israel J. Math.*, 115:157–193, 2000.
- [16] C. Bonatti and M. Viana. Lyapunov exponents with multiplicity 1 for deterministic products of matrices. *Ergod. Th. & Dynam. Sys.*, 24:1295–1330, 2004.
- [17] R. Bowen. A horseshoe with positive measure. *Invent. Math.*, 29:203–204, 1975.
- [18] K. Burns and A. Wilkinson. On the ergodicity of partially hyperbolic systems. *Annals of Math.*
- [19] A. Carverhill. Furstenberg’s theorem for nonlinear stochastic systems. *Probab. Theory. Related Fields*, 74:529–534, 1987.
- [20] A. Douady and J. C. Earle. Conformally natural extension of homeomorphisms of the circle. *Acta Math.*, 157:23–48, 1986.
- [21] H. Farkas and I. Kra. *Riemann surfaces*, volume 71 of *Graduate Texts in Mathematics*. Springer-Verlag, second edition, 1992.
- [22] J. Franks. Realizing rotation vectors for torus homeomorphisms. *Trans. Amer. Math. Soc.*, 311:107–115, 1989.
- [23] H. Furstenberg. Non-commuting random products. *Trans. Amer. Math. Soc.*, 108:377–428, 1963.
- [24] I. Ya. Gol’dsheid and G. A. Margulis. Lyapunov indices of a product of random matrices. *Uspekhi Mat. Nauk.*, 44:13–60, 1989.

- [25] Y. Guivarc'h and A. Raugi. Products of random matrices : convergence theorems. *Contemp. Math.*, 50:31–54, 1986.
- [26] F. Rodriguez Hertz. Stable ergodicity of certain linear automorphisms of the torus. *Ann. of Math.*, 162:65–107, 2005.
- [27] M. Hirsch, C. Pugh, and M. Shub. *Invariant manifolds*, volume 583 of *Lect. Notes in Math.* Springer Verlag, 1977.
- [28] J.-L. Journé. A regularity lemma for functions of several variables. *Rev. Mat. Iberoamericana*, 4:187–193, 1988.
- [29] A. Katok. Lyapunov exponents, entropy and periodic points of diffeomorphisms. *Publ. Math. IHES*, 51:137–173, 1980.
- [30] Yi. Katznelson. Ergodic automorphisms of T^n are Bernoulli shifts. *Israel J. Math.*, 10:186–195, 1971.
- [31] J. Kingman. The ergodic theorem of subadditive stochastic processes. *J. Royal Statist. Soc.*, 30:499–510, 1968.
- [32] F. Ledrappier. Propriétés ergodiques des mesures de Sinaï. *Publ. Math. I.H.E.S.*, 59:163–188, 1984.
- [33] F. Ledrappier. Positivity of the exponent for stationary sequences of matrices. In *Lyapunov exponents (Bremen, 1984)*, volume 1186 of *Lect. Notes Math.*, pages 56–73. Springer, 1986.
- [34] F. Ledrappier and L.-S. Young. The metric entropy of diffeomorphisms. I. Characterization of measures satisfying Pesin's entropy formula. *Ann. of Math.*, 122:509–539, 1985.
- [35] F. Ledrappier and L.-S. Young. The metric entropy of diffeomorphisms. II. Relations between entropy, exponents and dimension. *Ann. of Math.*, 122:540–574, 1985.
- [36] D. Ornstein and B. Weiss. On the Bernoulli nature of systems with some hyperbolic structure. *Ergodic Theory Dynam. Systems*, 18:441–456, 1998.
- [37] V. I. Oseledets. A multiplicative ergodic theorem: Lyapunov characteristic numbers for dynamical systems. *Trans. Moscow Math. Soc.*, 19:197–231, 1968.
- [38] Ya. B. Pesin. Families of invariant manifolds corresponding to non-zero characteristic exponents. *Math. USSR. Izv.*, 10:1261–1302, 1976.
- [39] Ya. B. Pesin. Characteristic Lyapunov exponents and smooth ergodic theory. *Russian Math. Surveys*, 32:4:55–114, 1977.
- [40] C. Pugh and M. Shub. Stably ergodic dynamical systems and partial hyperbolicity. *J. Complexity*, 13:125–179, 1997.
- [41] V. A. Rokhlin. On the fundamental ideas of measure theory. *A. M. S. Transl.*, 10:1–52, 1952. Transl. from *Mat. Sbornik* 25 (1949), 107–150.
- [42] D. Ruelle. An inequality for the entropy of differentiable maps. *Bull. Braz. Math. Soc.*, 9:83–87, 1978.
- [43] M. Viana. Almost all cocycles over any hyperbolic system have nonvanishing Lyapunov exponents. *Ann. of Math.*, 167:643–680, 2008.
- [44] M. Viana and J. Yang. Physical measures and absolute continuity for one-dimensional center directions. Preprint IMPA 2010.
- [45] L.-S. Young. Statistical properties of dynamical systems with some hyperbolicity. *Annals of Math.*, 147:585–650, 1998.
- [46] L.-S. Young. Recurrence times and rates of mixing. *Israel J. of Math.*, 110:153–188, 1999.

E-mail address: artur.avila@gmail.com

URL: www.impa.br/~avila/

E-mail address: viana@impa.br

URL: www.impa.br/~viana/