

isid/ms/2007/04

March 28, 2007

<http://www.isid.ac.in/~statmath/eprints>

Extreme Points of the Convex Set of Joint Probability Distributions with Fixed Marginals

K. R. PARTHASARATHY

Indian Statistical Institute, Delhi Centre
7, SJSS Marg, New Delhi-110 016, India

Extreme Points of the Convex Set of Joint Probability Distributions with Fixed Marginals

by

K. R. Parthasarathy

Indian Statistical Institute, Delhi Centre,

7, S. J. S. Sansanwal Marg,

New Delhi - 110 016, India.

e-mail : krp@isid.ac.in

Summary : By using a quantum probabilistic approach we obtain a description of the extreme points of the convex set of all joint probability distributions on the product of two standard Borel spaces with fixed marginal distributions.

Key words : C^* algebra, covariant bistochastic maps, completely positive map, Stinespring's theorem, extreme points of a convex set

AMS Subject Classification Index 46L53, 15A51

1 Introduction

It is a well-known theorem of Garret Birkhoff [3] and von Neumann [6], [1], [2] that the extreme points in the convex set of all $n \times n$ bistochastic (or doubly stochastic) matrices are precisely the n -th order permutation matrices. Here we address the following problem: If G is a standard Borel group acting measurably on two standard probability spaces $(X_i, \mathcal{F}_i, \mu_i)$, $i = 1, 2$ where μ_i is invariant under the G -action for each i then what are the extreme points of the convex set of all joint probability distributions on the product Borel space $(X_1 \times X_2, \mathcal{F}_1 \otimes \mathcal{F}_2)$ which are invariant under the diagonal action $(x_1, x_2) \mapsto (gx_1, gx_2)$ where $x_i \in X_i$, $i = 1, 2$ and $g \in G$?

Our approach to the problem mentioned above is based on a quantum probabilistic method arising from Stinespring's [5] description of completely positive maps on C^* algebras. We obtain a necessary and sufficient condition for the extremality of a joint distribution in the form of a regression condition. This leads to examples of extremal nongraphic joint distributions in the unit square with uniform marginal distributions on the unit interval. The Birkhoff-von Neumann theorem is deduced as a corollary of the main theorem.

2 The convex set of covariant bistochastic maps on C^* algebras

For any complex separable Hilbert space \mathcal{H} , express its scalar product in the Dirac notation $\langle \cdot | \cdot \rangle$ and denote by $\mathcal{B}(\mathcal{H})$ the C^* algebra of all bounded operators on \mathcal{H} . Let G be a group with fixed unitary representations $g \mapsto U_g$, $g \mapsto V_g$, $g \in G$ in Hilbert spaces \mathcal{H}_1 , \mathcal{H}_2 respectively and let $\mathcal{A}_i \subset \mathcal{B}(\mathcal{H}_i)$, $i = 1, 2$ be unital C^* algebras invariant under respective conjugations by U_g , V_g for every g in G . Let ω_i be a fixed state in \mathcal{A}_i for each i , satisfying the invariance conditions:

$$\omega_1(U_g X U_g^{-1}) = \omega_1(X), \omega_2(V_g Y V_g^{-1}) = \omega_2(Y) \quad \forall X \in \mathcal{A}_1, Y \in \mathcal{A}_2, g \in G. \quad (2.1)$$

Consider a linear, unital and completely positive map $T : \mathcal{A}_1 \rightarrow \mathcal{A}_2$ satisfying the following:

$$\omega_2(T(X)) = \omega_1(X) \quad \forall X \in \mathcal{A}_1, \quad (2.2)$$

$$T(U_g X U_g^{-1}) = V_g T(X) V_g^{-1} \quad \forall X \in \mathcal{A}_1, g \in G. \quad (2.3)$$

Then we say that T is a G -covariant bistochastic map with respect to the pair of states ω_1, ω_2 and representations U, V . Denote by \mathbb{K} the convex set of all such covariant bistochastic maps from \mathcal{A}_1 into \mathcal{A}_2 . We shall now present a necessary and sufficient condition for an element T in \mathbb{K} to be an extreme point of \mathbb{K} .

To any $T \in \mathbb{K}$ we can associate a Stinespring triple (\mathcal{K}, j, Γ) where \mathcal{K} is a Hilbert space, j is a C^* homomorphism from \mathcal{A}_1 into $\mathcal{B}(\mathcal{K})$ and Γ is an isometry from \mathcal{H}_2 into \mathcal{K} satisfying the following properties:

- (i) $\Gamma^\dagger j(X) \Gamma = T(X) \quad \forall X \in \mathcal{A}_1$;
- (ii) The linear manifold generated by $\{j(X) \Gamma u \mid u \in \mathcal{H}_2, X \in \mathcal{A}_1\}$ is dense in \mathcal{K} .

Such a Stinespring triple is unique upto a unitary isomorphism, i.e., if $(\mathcal{K}', j', \Gamma')$ is another triple satisfying the properties (i) and (ii) above then there exists a unitary isomorphism $\theta : \mathcal{K} \rightarrow \mathcal{K}'$ such that $\theta j(X) = j'(X)\theta \forall X \in \mathcal{A}_1$ and $\theta \Gamma v = \Gamma' v \forall v \in \mathcal{H}_2$. (See [5].)

We now claim that the covariance property of T ensures the existence of a unitary representation $g \mapsto W_g$ of G in \mathcal{K} satisfying the relations:

$$W_g j(X) \Gamma u = j(U_g X U_g^{-1}) \Gamma V_g u \quad \forall X \in \mathcal{A}_1, g \in G, u \in \mathcal{H}_2, \quad (2.4)$$

$$W_g j(X) W_g^{-1} = j(U_g X U_g^{-1}) \quad \forall X \in \mathcal{A}_1, g \in G. \quad (2.5)$$

Indeed, for any X, Y in \mathcal{A}_1 $u, v \in \mathcal{H}_2$ and $g \in G$ we have from the properties (i) and (ii) above and (2.3)

$$\begin{aligned} & \langle j(U_g X U_g^{-1}) \Gamma V_g u | j(U_g Y U_g^{-1}) \Gamma V_g v \rangle \\ &= \langle u | V_g^{-1} \Gamma^\dagger j(U_g X^\dagger Y U_g^{-1}) \Gamma V_g v \rangle \\ &= \langle u | V_g^{-1} T(U_g X^\dagger Y U_g^{-1}) V_g v \rangle \\ &= \langle u | T(X^\dagger Y) | v \rangle \\ &= \langle j(X) \Gamma u | j(Y) \Gamma v \rangle. \end{aligned}$$

In other words, the correspondence $j(X) \Gamma u \mapsto j(U_g X U_g^{-1}) \Gamma V_g u$ is a scalar product preserving map on a total subset of \mathcal{K} , proving the claim.

Theorem 2.1 Let $T \in \mathbb{K}$ and let (\mathcal{K}, j, Γ) be a Stinespring triple associated to T . Let $g \mapsto W_g$ be the unique unitary representation of G satisfying the relations (2.4) and (2.5). Then T is an extreme point of \mathbb{K} if and only if there exists no nonzero hermitian operator Z in the commutant of the set $\{j(X), X \in \mathcal{A}_1\} \cup \{W_g, g \in G\}$ satisfying the following two conditions:

- (i) $\Gamma^\dagger Z \Gamma = 0$;
- (ii) $\Gamma^\dagger Z j(X) \Gamma \in \mathcal{A}_2$ and $\omega_2(\Gamma^\dagger Z j(X) \Gamma) = 0 \quad \forall X \in \mathcal{A}_1$.

Proof Suppose T is not an extreme point of \mathbb{K} . Then there exist $T_1, T_2 \in \mathbb{K}$, $T_1 \neq T_2$ such that $T = \frac{1}{2}(T_1 + T_2)$. Let $(\mathcal{K}_1, j_1, \Gamma_1)$ be a Stinespring triple associated to T_1 . Then by the argument outlined in the proof of Proposition 2.1 in [4] there exists a bounded operator $J : \mathcal{K} \rightarrow \mathcal{K}_1$ satisfying the following properties:

- (i) $J j(X) \Gamma u = j_1(X) \Gamma_1 u \quad \forall X \in \mathcal{A}_1, u \in \mathcal{H}_2$;
- (ii) The positive operator $\rho := J^\dagger J$ is in the commutant of $\{j(X), X \in \mathcal{A}\}$ in $\mathcal{B}(\mathcal{K})$;
- (iii) $T_1(X) = \Gamma^\dagger \rho j(X) \Gamma$.

Since $T_1 \neq T_2$ it follows that $T_1 \neq T$ and hence ρ is different from the identity operator. We now claim that ρ commutes with W_g for every g in G . Indeed, for any X, Y in \mathcal{A}_1 , u, v in \mathcal{H}_2 we have from the definition of ρ and J , equation (2.4) and the covariance of T_1

$$\begin{aligned}
& \langle j(X)\Gamma u | \rho W_g | j(Y)\Gamma v \rangle \\
&= \langle j(X)\Gamma u | J^\dagger J | j(U_g Y U_g^{-1})\Gamma V_g v \rangle \\
&= \langle j_1(X)\Gamma_1 u | j_1(U_g Y U_g^{-1})\Gamma_1 V_g v \rangle \\
&= \langle u | \Gamma_1^\dagger j_1(X^\dagger U_g Y U_g^{-1})\Gamma_1 | V_g v \rangle \\
&= \langle u | T_1(X^\dagger U_g Y U_g^{-1}) | V_g v \rangle \\
&= \langle u | V_g T_1(U_g^{-1} X^\dagger U_g Y) | v \rangle.
\end{aligned}$$

On the other hand, by the same arguments, we have

$$\begin{aligned}
& \langle j(X)\Gamma u | W_g \rho | j(Y)\Gamma v \rangle \\
&= \langle j(U_g^{-1} X U_g)\Gamma V_g^{-1} u | J^\dagger J | j(Y)\Gamma v \rangle \\
&= \langle j_1(U_g^{-1} X U_g)\Gamma_1 V_g^{-1} u | j_1(Y)\Gamma_1 v \rangle \\
&= \langle u | V_g T_1(U_g^{-1} X^\dagger U_g Y) | v \rangle
\end{aligned}$$

Comparing the last two identities and using property (ii) of the Stinespring triple we conclude that ρ commutes with W_g . Putting $Z = \rho - I$ we have

$$\Gamma^\dagger Z j(X)\Gamma = T_1(X) - T(X) \quad \forall X \in \mathcal{A}_1. \quad (2.6)$$

Clearly, the right hand side of this equation is an element of \mathcal{A}_2 and

$$\omega_2(\Gamma^\dagger Z j(X)\Gamma) = \omega_1(X) - \omega_1(X) = 0 \quad \forall X \in \mathcal{A}_1.$$

Putting $X = I$ in (2.6) we have $\Gamma^\dagger Z \Gamma = 0$. Then Z satisfies properties (i) and (ii) in the statement of the theorem, proving the sufficiency part.

Conversely, suppose there exists a nonzero hermitian operator Z in the commutant of $\{j(X), X \in \mathcal{A}_1\} \cup \{W_g, g \in G\}$ satisfying properties (i) and (ii) in the theorem. Choose and fix a positive constant ε such that the operators $I \pm \varepsilon Z$ are positive. Define the maps $T_\pm : \mathcal{A}_1 \rightarrow \mathcal{A}_2$ by

$$T_\pm(X) = \Gamma^\dagger (I \pm \varepsilon Z) j(X)\Gamma, \quad X \in \mathcal{A}_1. \quad (2.7)$$

Since

$$(I \pm \varepsilon Z)j(X) = \sqrt{I \pm \varepsilon Z} j(X) \sqrt{I \pm \varepsilon Z}$$

it follows that T_\pm are completely positive. By putting $X = I$ in (2.7) and using property (i) of Z in the theorem we see that T_\pm are unital. Furthermore, we have from equations (2.4) and (2.5), for any $g \in G$, $X \in \mathcal{A}_1$,

$$\begin{aligned}
T_\pm(U_g X U_g^{-1}) &= \Gamma^\dagger (I \pm \varepsilon Z) W_g j(X) W_g^{-1} \Gamma \\
&= V_g \Gamma^\dagger (I \pm \varepsilon Z) j(X) \Gamma V_g^{-1} \\
&= V_g T_\pm(X) V_g^{-1}.
\end{aligned}$$

Also, by property (ii) in the theorem we have

$$\omega_2(T_{\pm}(X)) = \omega_2(T(X)) = \omega_1(X) \quad \forall X \in \mathcal{A}_1.$$

Thus $T_{\pm} \in \mathbb{K}$. Note that

$$\langle u | \Gamma^\dagger Z j(X^\dagger Y) \Gamma | v \rangle = \langle j(X) \Gamma u | Z | j(Y) \Gamma v \rangle$$

cannot be identically zero when X and Y vary in \mathcal{A}_1 and u and v vary in \mathcal{H}_2 . Thus $\Gamma^\dagger Z j(X) \Gamma \neq 0$ and hence $T_+ \neq T_-$. But $T = \frac{1}{2}(T_+ + T_-)$. In other words T is not an extreme point of \mathbb{K} . This proves necessity. \square

3 The convex set of invariant joint distributions with fixed marginal distributions

Let $(X_i, \mathcal{F}_i, \mu_i)$, $i = 1, 2$ be standard probability spaces and let G be a standard Borel group acting measurably on both X_1 and X_2 preserving μ_1 and μ_2 . Denote by $\mathbb{K}(\mu_1, \mu_2)$ the convex set of all joint probability distributions on the product Borel space $(X_1, \times X_2, \mathcal{F}_1 \otimes \mathcal{F}_2)$ invariant under the diagonal G action $(g, (x_1, x_2)) \mapsto (gx_1, gx_2)$, $x_i \in X_i$, $g \in G$ and having the marginal distribution μ_i in X_i for each i . Choose and fix $\omega \in \mathbb{K}(\mu_1, \mu_2)$. Our present aim is to derive from the quantum probabilistic result in Theorem 2.1, a necessary and sufficient condition for ω to be an extreme point of $\mathbb{K}(\mu_1, \mu_2)$. To this end we introduce the Hilbert spaces $\mathcal{H}_i = L^2(\mu_i)$, $\mathcal{K} = L^2(\omega)$ and the abelian von Neumann algebras $\mathcal{A}_i \subset \mathcal{B}(\mathcal{H}_i)$ where $\mathcal{A}_i = L^\infty(\mu_i)$ is also viewed as the algebra of operators of multiplication by functions from $L^\infty(\mu_i)$. For any $\varphi \in L^\infty(\mu_i)$ we shall denote by the same symbol φ the multiplication operator $f \mapsto \varphi f$, $f \in L^2(\mu_i)$. For any $\varphi \in \mathcal{A}_1$ define the operator $j(\varphi)$ in \mathcal{K} by

$$(j(\varphi)f)(x_1, x_2) = \varphi(x_1)f(x_1, x_2), \quad f \in \mathcal{K}, x_i \in X_i. \quad (3.1)$$

Then the correspondence $\varphi \mapsto j(\varphi)$ is a von Neumann algebra homomorphism from \mathcal{A}_1 into $\mathcal{B}(\mathcal{K})$. Define the isometry $\Gamma : \mathcal{H}_2 \rightarrow \mathcal{K}$ by

$$(\Gamma v)(x_1, x_2) = v(x_2), \quad v \in \mathcal{H}_2. \quad (3.2)$$

Then, for $f \in \mathcal{K}$, $v \in \mathcal{H}_2$ we have

$$\begin{aligned} \langle f | \Gamma v \rangle &= \int_{X_1 \times X_2} \bar{f}(x_1, x_2) v(x_2) \omega(dx_1 dx_2) \\ &= \int_{X_2} \mu_2(dx_2) [\bar{f}(x_1, x_2) \nu(dx_1, x_2)] v(x_2) \end{aligned}$$

where $\nu(E, x_2)$, $E \in \mathcal{F}_1$, $x_2 \in X_2$ is a measurable version of the conditional probability distribution on \mathcal{F}_1 given the sub σ -algebra $\{X_1 \times F, F \in \mathcal{F}_2\} \subset \mathcal{F}_1 \otimes \mathcal{F}_2$. Thus the adjoint $\Gamma^\dagger : \mathcal{K} \rightarrow \mathcal{H}_2$ of Γ is given by

$$(\Gamma^\dagger f)(x_2) = \int_{X_1} f(x_1, x_2) \nu(dx_1, x_2). \quad (3.3)$$

Hence

$$(j(\varphi)\Gamma v)(x_1, x_2) = \varphi(x_1)v(x_2), \quad \varphi \in \mathcal{A}_1, \quad v \in \mathcal{H}_2, \quad (3.4)$$

$$(\Gamma^\dagger j(\varphi)\Gamma v)(x_2) = \left[\int \varphi(x_1)\nu(dx_1, x_2) \right] v(x_2). \quad (3.5)$$

In other words

$$\Gamma^\dagger j(\varphi)\Gamma = T(\varphi) \quad (3.6)$$

where $T(\varphi) \in \mathcal{A}_2$ is given by

$$T(\varphi)(x_2) = \int_{X_1} \varphi(x_1)\nu(dx_1, x_2). \quad (3.7)$$

Equations (3.1)-(3.7) imply that T is a linear, unital and positive (and hence completely positive) map from the abelian von Neumann algebra \mathcal{A}_1 into \mathcal{A}_2 and (\mathcal{K}, j, Γ) is, indeed, a Stinespring triple for T . Furthermore, the unitary operators U_g, V_g and W_g in $\mathcal{H}_1, \mathcal{H}_2$ and \mathcal{K} respectively defined by

$$\begin{aligned} (U_g u)(x_1) &= u(g^{-1}x_1), \quad u \in \mathcal{H}_1, \\ (V_g v)(x_2) &= v(g^{-1}x_2), \quad v \in \mathcal{H}_2, \\ (W_g f)(x_1, x_2) &= f(g^{-1}x_1, g^{-1}x_2), \quad f \in \mathcal{K} \end{aligned}$$

satisfy the relations (2.4) and (2.5).

Our next lemma describes operators of the form Z occurring in Theorem 2.1.

Lemma 3.1 Let Z be a bounded hermitian operator in \mathcal{K} satisfying the following conditions:

- (i) $Zj(\varphi) = j(\varphi)Z \quad \forall \varphi \in \mathcal{A}_1,$
- (ii) $ZW_g = W_gZ \quad \forall g \in G,$
- (ii) $\Gamma^\dagger Zj(\varphi)\Gamma \in \mathcal{A}_2 \quad \forall \varphi \in \mathcal{A}_1.$

Then there exists a function $\zeta \in L^\infty(\omega)$ satisfying the following properties:

- (a) $\zeta(gx_1, gx_2) = \zeta(x_1, x_2)$ a.e. $(\omega) \quad \forall g \in G,$
- (a) $(Zf)(x_1, x_2) = \zeta(x_1, x_2)f(x_1, x_2) \quad \forall f \in \mathcal{K}$

Proof Let

$$\zeta(x_1, x_2) = (Z1)(x_1, x_2)$$

where the symbol 1 also denotes the function identically equal to unity. For functions u, v on X_1, X_2 respectively denote by $u \otimes v$ the function on $X_1 \times X_2$ defined by $u \otimes v(x_1, x_2) = u(x_1)v(x_2)$. By property (i) of Z in the lemma we have

$$\begin{aligned} (Z\varphi \otimes 1)(x_1, x_2) &= (Zj(\phi)1)(x_1, x_2) \\ &= (j(\phi)Z1)(x_1, x_2) \\ &= \varphi(x_1)\zeta(x_1, x_2) \quad \forall \varphi \in \mathcal{A}_1. \end{aligned} \quad (3.8)$$

If $\varphi \in \mathcal{A}_1$, $v \in \mathcal{H}_2$, we have

$$\begin{aligned}
(Z\varphi \otimes v)(x_1, x_2) &= (Zj(\varphi)\Gamma v)(x_1, x_2) \\
&= (j(\varphi)Z\Gamma v)(x_1, x_2) \\
&= \varphi(x_1)(Z1 \otimes v)(x_1, x_2)
\end{aligned} \tag{3.9}$$

From properties (i) and (iii) of Z in the lemma and equations (3.3), (3.8) and (3.9) we have

$$\begin{aligned}
(\Gamma^\dagger Zj(\varphi)\Gamma v)(x_2) &= \int (Z\varphi \otimes v)\nu(dx_1, x_2) \\
&= \int \varphi(x_1)(Z1 \otimes v)(x_1, x_2)\nu(dx_1, x_2)
\end{aligned}$$

whereas the left hand side is of the form $R(\varphi)(x_1)v(x_2)$ for some $R(\varphi) \in L^\infty(\mu_2)$. Thus

$$R(\varphi)(x_2)v(x_2) = \int \varphi(x_1)(Z1 \otimes v)(x_1, x_2)\nu(dx_1, x_2).$$

Choosing $v = 1$ we have from the definition of ζ

$$R(\varphi)(x_2) = \int \varphi(x_1)\zeta(x_1, x_2)\nu(dx_1, x_2).$$

Thus, for every $\varphi \in \mathcal{A}_1$

$$\int \varphi(x_1)\zeta(x_1, x_2)v(x_2)\nu(dx_1, x_2) = \int \varphi(x_1)(Z1 \otimes v)(x_1, x_2)\nu(dx_1, x_2)$$

and hence

$$(Z1 \otimes v)(x_1, x_2) = \zeta(x_1, x_2)v(x_2) \text{ a.e. } x_1(\nu(\cdot, x_2)) \text{ a.e. } x_2(\mu_2).$$

Applying $j(\varphi)$ on both sides we get

$$(Z\varphi \otimes v)(x_1, x_2) = \zeta(x_1, x_2)\varphi(x_1)v(x_2) \text{ a.e. } (\omega).$$

In other words Z is the operator of multiplication by ζ and it follows that $\zeta \in L^\infty(\omega)$. Now property (ii) of Z implies property (a) in the lemma. \square

Theorem 3.2 Let $\omega \in \mathbb{K}(\mu_1, \mu_2)$. Then ω is an extreme point of $\mathbb{K}(\mu_1, \mu_2)$ if and only if there exists no nonzero real-valued function $\zeta \in L^\infty(\omega)$ satisfying the following conditions:

- (i) $\zeta(gx_1, gx_2) = \zeta(x_1, x_2)$ a.e. $\omega \forall g \in G$;
- (ii) $\mathbb{E}(\zeta(\xi_1, \xi_2)|\xi_1) = 0$, $\mathbb{E}(\zeta(\xi_1, \xi_2)|\xi_2) = 0$ where (ξ_1, ξ_2) is an $X_1 \times X_2$ -valued random variable with distribution ω .

Proof Let Z be a bounded selfadjoint operator in the commutant of $\{j(\varphi), \varphi \in \mathcal{A}_1\} \cup \{W_g, g \in G\}$ such that $\Gamma^\dagger Zj(\varphi)\Gamma \in \mathcal{A}_2 \forall \varphi \in \mathcal{A}_1$. Then by Lemma 3.1 it follows that Z is of the form

$$(Zf)(x_1, x_2) = \zeta(x_1, x_2)f(x_1, x_2)$$

where $\zeta \in L^\infty(\omega)$ and $\zeta(gx_1, gx_2) = \zeta(x_1, x_2)$ a.e. (ω) . Note that

$$(\Gamma^\dagger Z \Gamma v)(x_2) = \left[\int_{X_1} \zeta(x_1, x_2) \nu(dx_1, x_2) \right] v(x_2) \text{ a.e. } (\mu_2), v \in \mathcal{H}_2.$$

Thus $\Gamma^\dagger Z \Gamma = 0$ if and only if $\mathbb{E}(\zeta(\xi_1, \xi_2) | \xi_2) = 0$. Now we evaluate

$$(\Gamma^\dagger Z j(\varphi) \Gamma v)(x_2) = \int \varphi(x_1) v(x_2) \zeta(x_1, x_2) \nu(dx_1, x_2) \text{ a.e. } (\mu_2).$$

Looking upon $\Gamma^\dagger Z j(\varphi) \Gamma$ as an element of \mathcal{A}_2 and evaluating the state μ_2 on this element we get

$$\begin{aligned} \mu_2(\Gamma^\dagger Z j(\varphi) \Gamma) &= \int \varphi(x_1) \zeta(x_1, x_2) \nu(dx_1, x_2) \mu(dx_2) \\ &= \int \varphi(x_1) \zeta(x_1, x_2) \omega(dx_1, dx_2) \\ &= \mathbb{E}_\omega \varphi(\xi_1) \zeta(\xi_1, \xi_2) \\ &= \mathbb{E}_{\mu_1} \varphi(\xi_1) \mathbb{E}(\zeta(\xi_1, \xi_2) | \xi_1). \end{aligned}$$

Thus $\mu_2(\Gamma^\dagger Z j(\varphi) \Gamma) = 0 \forall \varphi \in \mathcal{A}_1$ if and only if $\mathbb{E}(\zeta(\xi_1, \xi_2) | \xi_1) = 0$. Now an application of Theorem 2.1 completes the proof of the theorem. \square

We shall now look at the special case when G is the trivial group consisting of only the identity element. Let $(X_i, \mathcal{F}_i, \mu_i)$, $i = 1, 2$ be standard probability spaces and let $T : X_1 \rightarrow X_2$ be a Borel map such that $\mu_2 = \mu_1 T^{-1}$. Consider an X_1 -valued random variable ξ with distribution μ_1 . Then the joint distribution ω of the pair $(\xi, T \circ \xi)$ is an element of $\mathbb{K}(\mu_1, \mu_2)$ and by Theorem 2.1 is an extreme point. Similarly, if $T : X_2 \rightarrow X_1$ is a Borel map such that $\mu_2 T^{-1} = \mu_1$ and η is an X_2 -valued random variable with distribution μ_2 then $(T \circ \eta, \eta)$ has a joint distribution which is an extreme point of $\mathbb{K}(\mu_1, \mu_2)$. Such extreme points are called *graphic* extreme points. Thus there arises the natural question whether there exist nongraphic extreme points. Our next lemma facilitates the construction of nongraphic extreme points.

Lemma 3.3 Let $(X, \mathcal{F}, \lambda)$, (Y, \mathcal{G}, μ) , (Z, \mathcal{K}, ν) be standard probability spaces and let ξ, η, ζ be random variables on a probability space with values in X, Y, Z and distribution λ, μ, ν respectively. Suppose ζ is independent of (ξ, η) and the joint distribution ω of (ξ, η) is an extreme point of $\mathbb{K}(\lambda, \mu)$. Let $\tilde{\lambda}, \tilde{\mu}, \tilde{\omega}$ be the distributions of (ξ, ζ) , (η, ζ) and $((\xi, \zeta), (\eta, \zeta))$ respectively in the spaces $X \times Z$, $Y \times Z$ and $(X \times Z) \times (Y \times Z)$. Then $\tilde{\omega}$ is an extreme point of $\mathbb{K}(\tilde{\lambda}, \tilde{\mu})$.

Proof Let f be a bounded real-valued measurable function on $(X \times Z) \times (Y \times Z)$ satisfying the relations

$$\begin{aligned} \mathbb{E} \{ f((\xi, \zeta), (\eta, \zeta)) | (\eta, \zeta) \} &= 0, \\ \mathbb{E} \{ f((\xi, \zeta), (\eta, \zeta)) | (\xi, \zeta) \} &= 0. \end{aligned}$$

If we write

$$F_z(x, y) = f((x, z), (y, z)) \quad \text{where} \quad (x, y, z) \in X \times Y \times Z$$

then we have

$$\mathbb{E}(F_z(\xi, \eta)|\eta) = 0, \quad \mathbb{E}(F_z(\xi, \eta)|\xi) = 0 \quad \text{a.e. } z(\nu).$$

Since ω is extremal it follows that $F_z(\xi, \eta) = 0$ a.e. $z(\nu)$ and therefore $f((\xi, \zeta), (\eta, \zeta)) = 0$. By Theorem 3.1 it follows that $\tilde{\omega}$ is, indeed, an extreme point of $\mathbb{K}(\tilde{\lambda}, \tilde{\mu})$. \square

Example 3.4 Let λ be the uniform distribution in the unit interval $[0, 1]$. We shall use Lemma 3.3 and construct nongraphic extreme points of $\mathbb{K}(\lambda, \lambda)$ which are distributions in the unit square. To this end we start with the two points space $\mathbb{Z}_2 = \{0, 1\}$ with the probability distribution P where

$$P(\{0\}) = p, \quad P(\{1\}) = q, \quad 0 < p < q < 1, \quad p + q = 1.$$

Now consider \mathbb{Z}_2 -valued random variables ξ, η with the joint distribution given by

$$P(\xi = 0, \eta = 0) = 0, \quad P(\xi = 0, \eta = 1) = P(\xi = 1, \eta = 0) = p, \quad P(\xi = 1, \eta = 1) = q - p.$$

Note that the joint distribution of (ξ, η) is a nongraphic extreme point of $\mathbb{K}(P, P)$. Now consider an i.i.d sequence ζ_1, ζ_2, \dots of \mathbb{Z}_2 -valued random variables independent of (ξ, η) and having the same distribution P . Put

$$\zeta = (\zeta_1, \zeta_2, \dots).$$

Then by Lemma 3.3 the joint distribution ω of $((\xi, \zeta), (\eta, \zeta))$ is an extreme point of $\mathbb{K}(\nu, \nu)$ where $\nu = P \otimes P \otimes \dots$ in $\mathbb{Z}_2^{\{0,1,2,\dots\}}$. Furthermore, since (ξ, η) is nongraphic so is $((\xi, \zeta), (\eta, \zeta))$. Denote by F_p the common probability distribution function of the random variables

$$\tilde{\xi} = \frac{\xi}{2} + \sum_{j=1}^{\infty} \frac{\zeta_j}{2^{j+1}}, \quad \tilde{\eta} = \frac{\eta}{2} + \sum_{j=1}^{\infty} \frac{\zeta_j}{2^{j+1}}.$$

Then F_p is a strictly increasing and continuous function on the unit interval and therefore the correspondence $t \rightarrow F_p(t)$ is a homeomorphism of $[0, 1]$. Put $\xi' = F_p(\tilde{\xi}), \eta' = F_p(\tilde{\eta})$. Then the joint distribution ω of (ξ', η') is a nongraphic extreme point of $\mathbb{K}(\lambda, \lambda)$.

Now we consider the case when X_1 and X_2 are finite sets, G is a finite group acting on each X_i , the number of G -orbits in X_1, X_2 and $X_1 \times X_2$ are respectively m_1, m_2 and m_{12} and μ_i is a G -invariant probability distribution in X_i with support X_i for each $i = 1, 2$. For any probability distribution λ in any finite set denote by $S(\lambda)$ its support set. We first note that Theorem 3.2 assumes the following form.

Theorem 3.5 A probability distribution $\omega \in \mathbb{K}(\mu_1, \mu_2)$ is an extreme point if and only if there is no nonzero real-valued function ζ on $S(\omega)$ satisfying the following conditions:

$$(i) \quad \zeta(gx_1, gx_2) = \zeta(x_1, x_2) \quad \forall (x_1, x_2) \in S(\omega), g \in G;$$

- (ii) $\sum_{x_2 \in X_2} \zeta(x_1, x_2)\omega(x_1, x_2) = 0 \quad \forall x_1 \in X_1;$
- (iii) $\sum_{x_1 \in X_1} \zeta(x_1, x_2)\omega(x_1, x_2) = 0 \quad \forall x_2 \in X_2.$

Proof Immediate. \square

Corollary 3.6 Let ω_1, ω_2 be extreme points of $\mathbb{K}(\mu_1, \mu_2)$ and $S(\omega_1) \subseteq S(\omega_2)$. Then $\omega_1 = \omega_2$. In particular, any extreme point ω of $\mathbb{K}(\mu_1, \mu_2)$ is uniquely determined by its support set $S(\omega)$.

Proof Suppose $\omega_1 \neq \omega_2$. Then put $\omega = \frac{1}{2}(\omega_1 + \omega_2)$. Then $\omega \in \mathbb{K}(\mu_1, \mu_2)$ and ω is not an extreme point. By Theorem 3.5 there exists a nonzero real-valued function ζ satisfying conditions (i)-(iii) of the theorem. By hypothesis $S(\omega) = S(\omega_2)$. Define

$$\zeta'(x_1, x_2) = \frac{\zeta(x_1, x_2)\omega(x_1, x_2)}{\omega_2(x_1, x_2)} \quad \text{where } (x_1, x_2) \in S(\omega_2).$$

Then conditions (i)-(iii) of Theorem 3.5 are fulfilled when the pair ζ, ω is replaced by ζ', ω_2 contradicting the extremality of ω_2 . \square

Corollary 3.7 For any $\omega \in \mathbb{K}(\mu_1, \mu_2)$ let $N(\omega)$ denote the number of G -orbits in its support set $S(\omega)$. If ω is an extreme point of $\mathbb{K}(\mu_1, \mu_2)$ then

$$\max(m_1, m_2) \leq N(\omega) \leq m_1 + m_2.$$

In particular, the number of extreme points in $\mathbb{K}(\mu_1, \mu_2)$ does not exceed

$$\sum_{\max(m_1, m_2) \leq r \leq m_1 + m_2} \binom{m_{12}}{r}.$$

Proof Let ω be an extreme point of $\mathbb{K}(\mu_1, \mu_2)$. Suppose $N(\omega) > m_1 + m_2$. Observe that all G -invariant real-valued functions on $S(\omega)$ constitute a linear space of cardinality $N(\omega)$. Functions ζ satisfying conditions (i)-(iii) of the theorem constitute a subspace of dimension $\geq N(\omega) - (m_1 + m_2)$, contradicting the extremality of ω . For any distribution ω in $\mathbb{K}(\mu_1, \mu_2)$ we have $N(\omega) \geq m_i, i = 1, 2$. This proves the first part. The second part is now immediate from Corollary 3.6. \square

Corollary 3.8 (Birkhoff-von Neumann Theorem) Let $X_1 = X_2 = X, \#X = m, \mu_1 = \mu_2 = \mu$ where $\mu(x) = \frac{1}{m} \forall x \in X$. Then any extreme point ω in $\mathbb{K}(\mu, \mu)$ is of the form

$$\omega(x, y) = \frac{1}{m} \delta_{\sigma(x)y} \quad \forall x, y \in X$$

where σ is a permutation of the elements of X .

Proof Without loss of generality we assume that $X = \{1, 2, \dots, m\}$ and view ω as a matrix of order m with nonnegative entries with each row or column total being $1/m$. First assume that in each row or column there are at least two nonzero entries. Then ω has at least $2m$ nonzero entries and by Corollary 3.7 it follows that every row or column has exactly two nonzero entries. We claim that for any $i \neq i', j \neq j'$ in the set $\{1, 2, \dots, m\}$ at least one among $\omega_{ij}, \omega_{ij'}, \omega_{i'j}, \omega_{i'j'}$ vanishes. Suppose this is not true for some $i \neq i', j \neq j'$. Put

$$p = \min \{ \omega_{rs} \mid (r, s) : \omega_{rs} > 0 \}.$$

Define

$$\omega_{rs}^{\pm} = \begin{cases} \omega_{rs} \pm p & \text{if } r = i, s = j \text{ or } r = i', s = j', \\ \omega_{rs} \mp p & \text{if } r = i', s = j \text{ or } r = i, s = j', \\ \omega_{rs} & \text{otherwise.} \end{cases}$$

Then $\omega^{\pm} \in \mathbb{K}(\mu, \mu)$, $\omega^+ \neq \omega^-$ and $\omega = \frac{1}{2}(\omega^+ + \omega^-)$, a contradiction to the extremality of ω . Now observe that permutation of columns as well as rows of ω lead to extreme points of $\mathbb{K}(\mu, \mu)$. By appropriate permutations of columns and rows ω reduces to a tridiagonal matrix of the form

$$\tilde{\omega} = \begin{bmatrix} p_{11} & p_{12} & 0 & 0 & \dots & \dots & \dots & \dots & 0 \\ p_{21} & 0 & p_{23} & 0 & \dots & \dots & \dots & \dots & 0 \\ 0 & p_{32} & 0 & p_{34} & \dots & \dots & \dots & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \dots & \dots & 0 & p_{n-1n-2} & 0 & p_{n-1n} \\ 0 & 0 & \dots & \dots & \dots & 0 & 0 & p_{nn-1} & p_{nn} \end{bmatrix}$$

where the p 's with suffixes are all greater than or equal to p . Now consider the matrices

$$\lambda^{\pm} = \begin{bmatrix} p_{11} \pm p & p_{12} \mp p & 0 & 0 & 0 & \dots \\ p_{21} \mp p & 0 & p_{23} \pm p & 0 & 0 & \dots \\ 0 & p_{32} \pm p & 0 & p_{34} \mp p & 0 & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \end{bmatrix}$$

Then $\lambda^{\pm} \in \mathbb{K}(\mu, \mu)$ and $\tilde{\omega} = \frac{1}{2}(\lambda^+ + \lambda^-)$, contradicting the extremality of $\tilde{\omega}$ and therefore of ω . In other words any extreme point ω of $\mathbb{K}(\mu, \mu)$ must have at least one row with exactly one nonzero entry. Then by permutations of rows and columns ω can be brought to the form

$$\omega_1 = \left[\begin{array}{c|ccc} 1/m & 0 & 0 & \dots & 0 \\ \hline 0 & & & & \\ \vdots & & \hat{\omega} & & \\ 0 & & & & \end{array} \right]$$

where $\frac{m}{m-1}\hat{\omega}$ is an extreme point of $\mathbb{K}(\hat{\mu}, \hat{\mu})$ where $\hat{\mu}$ is the uniform distribution on a set of $m - 1$ points. Now an inductive argument completes the proof. \square

We conclude with the remark that it is an interesting open problem to characterize the support sets of all extreme points of $\mathbb{K}(\mu_1, \mu_2)$ in terms of μ_1 and μ_2 .

References

- [1] R. B. Bapat and T. E. S. Raghavan, *Nonnegative Matrices and Applications*, Cambridge University Press, Cambridge 1997.
- [2] R. Bhatia, *Matrix Analysis*, Springer Verlag, New York 1996.
- [3] G. Birkhoff, *Tres observaciones sobre el algebra lineal*, University. Nac. Tucuman Rev, Ser. A5 (1946) 147-150.
- [4] K. R. Parthasarathy, *Extreme points of the convex set of stochastic maps on a C^* algebra*, Inf. Dim. Analysis, Quantum Probab. and Rel. Topics, 1 (1998) 599-609.
- [5] W. F. Stinespring, *Positive functions on C^* algebras*, Proc. Amer. Math. Soc., 6 (1955) 211-216.
- [6] J. von Neumann, *A certain zero-sum two-person game equivalent to an optimal assignment problem*, Ann. Math. Studies, 28 (1953) 5-12.