Extreme statistics of superdiffusive Lévy flights and every other Lévy subordinate Brownian motion

Sean D. Lawley^{*}

March 16, 2021

Abstract

The search for hidden targets is a fundamental problem in many areas of science, engineering, and other fields. Studies of search processes often adopt a probabilistic framework, in which a searcher randomly explores a spatial domain for a randomly located target. There has been significant interest and controversy regarding optimal search strategies, especially for superdiffusive processes. The optimal search strategy is typically defined as the strategy that minimizes the time it takes a given single searcher to find a target, which is called a first hitting time (FHT). However, many systems involve multiple searchers and the important timescale is the time it takes the fastest searcher to find a target, which is called an extreme FHT. In this paper, we study extreme FHTs for any stochastic process that is a random time change of Brownian motion by a Lévy subordinator. This class of stochastic processes includes superdiffusive Lévy flights in any space dimension, which are processes described by a Fokker-Planck equation with a fractional Laplacian. We find the short-time distribution of a single FHT for any Lévy subordinate Brownian motion and use this to find the full distribution and moments of extreme FHTs as the number of searchers grows. We illustrate these rigorous results in several examples and numerical simulations.

1 Introduction

What is the best way to search for a target whose location is *a priori* unknown? This basic search problem arises at various spatial and temporal scales in many areas of science, engineering, and other fields [1]. Examples include rescuers searching for castaways [2], military forces searching for enemy targets [3], animals searching for food, shelter, or a mate [1,4,5], proteins searching for DNA binding sites [6], and computers searching a database [7].

^{*}Department of Mathematics, University of Utah, Salt Lake City, UT 84112 USA (lawley@math.utah.edu).

Empirical and theoretical studies of search processes often adopt a probabilistic framework, in which the searcher randomly explores a spatial domain for a randomly located target [1, 5, 8]. The random movement of the searcher is often classified as diffusive, subdiffusive, or superdiffusive, depending respectively on whether the square of its displacement scales linearly, sublinearly, or superlinearly in time. While subdiffusive and superdiffusive motion are termed "anomalous" diffusion, they have been observed in many physical and biological systems [9].

Mathematical models of random search often assume that searchers explore space via a continuous-time random walk. In this framework, a searcher waits at its current location for a random time chosen from some waiting time probability density w(t), and then moves to a new location by jumping a random distance chosen from some jump length probability density l(y). The searcher repeats these two steps indefinitely or until it reaches the target. This process can be diffusive, subdiffusive, or superdiffusive, depending on the tails of the waiting time density w(t) and the jump length density l(y). In particular, if the mean waiting time is finite and the jump length density has the following slow power law decay,

$$l(y) \propto y^{-1-\alpha}$$
 as $y \to \infty$ for some $\alpha \in (0,2)$, (1)

then the process is superdiffusive and is often called a Lévy flight [9, 10]. In a certain scaling limit, the probability density p(x,t) for the position of a Lévy flight in \mathbb{R}^d satisfies the space fractional Fokker-Planck equation [11].

$$\partial_t p = -K(-\Delta)^{\alpha/2} p, \quad x \in \mathbb{R}^d, \, t > 0, \tag{2}$$

where $(-\Delta)^{\alpha/2}$ denotes the fractional Laplacian [12] and K > 0 is the generalized diffusion coefficient. Similar models of superdiffusive search involving long relocation events with distances chosen from a power law density akin to (1) are Lévy walks [13] and truncated Lévy flights [5].

Many have argued that superdiffusion is a more efficient search method compared to normal diffusion, since superdiffusive processes spend less time in previously explored regions of space [14–16]. Signatures of superdiffusion have been found in movement data for many different animal species [4], including albatrosses [15], spider monkeys [17,18], jackals [19], sharks [20], microorganisms [21], and also within biological cells [22]. In addition, superdiffusive search methods involving Lévy flights are employed in computational algorithms such as simulated annealing [23]. However, there has been controversy regarding the empirical evidence of superdiffusion in animal foraging, as some have questioned the accuracy of the statistical methods used in some of these studies [24].

Furthermore, the theoretical optimality of superdiffusive search models has also been controversial. Indeed, the seminal work of Viswanathan et al. [16] in 1999 claimed that the search time of a single searcher is minimized if the searcher employs an inverse square Lévy walk (corresponding to $\alpha = 1$ in (1)). This result forms the core of the very influential Lévy flight foraging hypothesis, which states that biological organisms must have evolved to perform such Lévy walks because of their optimality [5]. However, a recent analysis proved that this founding result of the Lévy flight foraging hypothesis is incorrect [25–27].

Search time is often quantified in terms of a so-called first hitting time (FHT), which is the first time the searcher reaches the target. If $X = \{X(t)\}_{t\geq 0}$ denotes the position of the searcher as a function of time $t \geq 0$, then the FHT is

$$\tau := \inf\{t > 0 : X(t) \in U\},\tag{3}$$

where U denotes the position of the target(s) (or the region of space in which the searcher can detect the target). There have been many studies of FHTs of Lévy flights, using both computational and analytical approaches [28–33]. An interesting aspect of these studies is the discrepancy between first hitting events (the searcher reaches or hits the target) and first passage events (the searcher moves beyond the target). While these two notions are equivalent for standard diffusion processes, paths of Lévy flights are discontinuous and thus may jump across a target without actually hitting it, which is called a "leapover" [29, 30, 32, 33].

Studies of optimal search strategies generally ask what search method minimizes the FHT of a single searcher [1, 5, 8]. However, many systems involve multiple searchers and the relevant timescale is the time it takes the fastest searcher to find the target. Indeed, this can be the case for many of the traditional search scenarios referenced above, such as the search for missing persons and castaways, military searches for enemy targets, and computer search processes. In the context of biology, cellular events are often triggered when the first of many searchers finds a target [34], and cooperative foraging involves multiple animals working together to find a target [35–42]. If τ_1, \ldots, τ_N are the respective FHTs of N parallel searchers, then the fastest searcher finds the target at time

$$T_N := \min\{\tau_1, \dots, \tau_N\},\tag{4}$$

which is often called an extreme FHT, fastest FHT [43], or parallel FHT [44, 45]. Hence, in these scenarios the relevant question is not what search strategy minimizes the single searcher FHT τ in (3), but rather what search strategy minimizes the extreme FHT T_N in (4).

In this paper, we investigate the extreme FHTs of a general class of stochastic processes which includes superdiffusive Lévy flights in \mathbb{R}^d for any $d \geq 1$. The class of stochastic processes is called Lévy subordinate Brownian motions, as the processes are obtained by random time changes of Brownian motion. We find the short-time distribution of a single FHT τ and then use this to find the full distribution and moments of the extreme FHT T_N as the number of searchers N grows.

To summarize our results, let $B = \{B(s)\}_{s\geq 0}$ be a *d*-dimensional Brownian motion with unit diffusivity, and let $S = \{S(t)\}_{t\geq 0}$ be an independent subordinator, which means that S is a one-dimensional, nondecreasing Lévy process




Figure 1: Lévy flights are subordinate Brownian motions. Left: The thin gray trajectory is the path of a Brownian motion B(s), and the thick black trajectory is the path of a Lévy flight X(t) = B(S(t)) obtained as a random time change of B(s) according to the $(\alpha/2)$ -stable subordinator S(t). Right: The path of the $(\alpha/2)$ -stable subordinator S(t). We take $\alpha = 1.5$ in this plot.

with S(0) = 0. Define the path of a single searcher $X = \{X(t)\}_{t>0}$ by

$$X(t) := B(S(t)) + X(0) \in \mathbb{R}^{d}, \quad t \ge 0,$$
(5)

where X(0) is some initial position independent of B and S. That is, X is a random time change of Brownian motion (see Figure 1 for the special case that X is a Lévy flight). Assuming that X(0) cannot lie in the target U, we prove that the FHT in (3) has the universal short-time distribution,

$$\mathbb{P}(\tau \le t) \sim \mathbb{P}(X(t) \in U) \sim \rho t \quad \text{as } t \to 0+, \tag{6}$$

where $\rho \in (0, \infty)$ is the rate,

$$\rho := \int_0^\infty \mathbb{P}(B(s) + X(0) \in U) \,\nu(\mathrm{d}s),\tag{7}$$

and $\nu(ds)$ is the Lévy measure of S. Throughout this paper, " $f \sim g$ " means $f/g \rightarrow 1$. If we set X(0) = 0, then the Gaussianity of B(s) means the integrand in (7) is

$$\mathbb{P}(B(s) + X(0) \in U) = \frac{1}{(4\pi s)^{d/2}} \int_U \exp\left(\frac{-\|x\|^2}{4s}\right) \mathrm{d}x.$$

We prove (6) for any nondeterministic subordinator S and any target set $U \subset \mathbb{R}^d$ that is nonempty and is the closure of its interior.

Furthermore, if τ_1, \ldots, τ_N are independent and identically distributed (iid) realizations of the FHT τ , then we prove that $(\rho N)T_N$ converges in distribution to a unit rate exponential random variable as N grows, which means

$$\mathbb{P}\Big(T_N > \frac{z}{\rho N}\Big) \to e^{-z} \quad \text{as } N \to \infty \text{ for each } z \ge 0.$$
(8)




Figure 2: Fastest FHTs of Brownian motions versus Lévy flights. Starting from x_0 (green ball), the thick blue path illustrates a Brownian motion that is the first to hit the target (red regions) out of many iid Brownian motions. Such fastest Brownian motions tend to follow the shortest path to the target. The thick black path illustrates the fastest Lévy flight out of many iid Lévy flights, which does not take the shortest path to the target. This fastest Lévy flight is obtained as a random time change of a typical Brownian motion (thin gray path) that wanders around and moves in and out of targets. This illustration of the fastest Lévy flight is characteristic of any subordinate Brownian motion.

Hence, T_N is approximately exponentially distributed with rate ρN . Furthermore, if $\mathbb{E}[T_N] < \infty$ for some $N \ge 1$, then we obtain all the moments of T_N for large N. In particular, we prove that

$$\mathbb{E}[T_N] \sim \sqrt{\text{Variance}[T_N]} \sim \frac{1}{\rho N} \quad \text{as } N \to \infty.$$
(9)

We also extend (8) and (9) to the kth fastest FHT for any $1 \le k \ll N$. In the case that X is a superdiffusive Lévy flight whose probability density satisfies the fractional Fokker-Planck equation in (2), the Lévy measure ν used in the rate ρ in (7) is

$$\nu(ds) = K \frac{\alpha/2}{\Gamma(1 - \alpha/2)} \frac{1}{s^{1 + \alpha/2}} ds, \quad s > 0.$$
(10)

We emphasize that our results hold for any nondeterministic subordinator S (meaning we exclude only the trivial case that S(t) is a deterministic function bt for some $b \ge 0$). In particular, the Lévy measure ν of the subordinator S need not have the slow power law decay in (10) which gives rise to long jumps of X in (5). Examples of other subordinators commonly used in modeling are given in section 4.3.

Before outlining the rest of the paper, we comment on how our results on subordinate Brownian motions relate to extreme statistics and large deviation theory for standard diffusion processes (i.e. processes satisfying a standard driftdiffusion Itô stochastic differential equation). First, the 1/N decay in (9) is much faster than the well-known $1/\ln N$ decay for diffusion processes [46]. Indeed, the extreme FHT for diffusion processes has the following rather slow decay in mean as the number of searchers N grows [47],

$$\mathbb{E}[T_N^{\text{diff}}] \sim \frac{L^2}{4D\ln N} \quad \text{as } N \to \infty, \tag{11}$$

where L > 0 is a certain geodesic distance from the possible searcher starting locations to the target and D > 0 is the characteristic diffusivity. Further contrasting (9) and (11), a salient feature of extreme FHTs of diffusion processes is that they only depend on the shortest path to the target since the fastest searchers follow this geodesic path [47]. In particular, extreme FHTs of diffusion are unaffected by changes to the problem outside of this path, such as altering the size of the target, the size of the domain, or even the space dimension $d \ge 1$. In contrast, it is evident from the formula for the rate ρ in (7) that the extreme FHTs of subordinate Brownian motion depend on all these global properties of the problem, which reflects the fact that the fastest subordinate Brownian searchers do not take a direct path to the closest part of the target. These results are illustrated in Figure 2 for the case of a Lévy flight (though the illustration is characteristic of any subordinate Brownian motion).

These differences stems from the difference between our result in (6) for subordinate Brownian motion and Varadhan's formula for diffusion processes [48]. Varadhan's formula is a celebrated result in large deviation theory which implies that if $X^{\text{diff}} = \{X^{\text{diff}}(t)\}_{t\geq 0}$ is a diffusion process, then

$$\lim_{t \to 0+} t \ln \mathbb{P}(X^{\text{diff}}(t) \in U) = -L^2/(4D) < 0, \tag{12}$$

where L and D are as in (11). The result in (6) can thus be interpreted as a type of Varadhan's formula for subordinate Brownian motion.

The rest of the paper is organized as follows. In section 2, we review some definitions and results from probability theory. In section 3, we present our general mathematical results. In section 4, we illustrate our results in several examples and compare the theory to numerical simulations. We conclude by discussing relations to prior work. Proofs are presented in the appendix.

2 Preliminaries

We begin by reviewing properties of subordinators, subordinate Brownian motions, Lévy flights, fractional Laplacians, and related concepts.

2.1 Subordinators

A Lévy process is a continuous-time stochastic process that has iid increments and satisfies certain technical conditions [49]. A subordinator is a one-dimensional, nondecreasing Lévy process $S = \{S(t)\}_{t\geq 0}$ with S(0) = 0. The distribution of S is determined by its Laplace exponent $\Phi(\beta)$, which satisfies

$$\mathbb{E}\left[e^{-\beta S(t)}\right] = e^{-t\Phi(\beta)}, \quad \text{for all } t \ge 0 \text{ and } \beta \ge 0,$$
$$\Phi(\beta) = b\beta + \int_0^\infty (1 - e^{-\beta s}) \nu(\mathrm{d}s), \quad \text{for all } \beta \ge 0, \tag{13}$$

where $b \ge 0$ is the drift and ν is the Lévy measure. In particular, ν satisfies

$$\nu((-\infty,0])=0 \quad \text{and} \quad \int_0^\infty \min\{1,s\}\,\nu(\mathrm{d} s)<\infty.$$

A Lévy measure $\nu(ds)$ can be interpreted as the rate that S increases by s > 0.

A subordinator S is called an $(\alpha/2)$ -stable subordinator for $\alpha \in (0,2)$ if it satisfies the following self-similarity or scaling property,

$$t^{-2/\alpha}S(t) =_{\rm d} S(1) \quad \text{for all } t > 0,$$
 (14)

where $=_{d}$ denotes equality in distribution. In this case, S is a pure jump process (i.e. zero drift b = 0) with Laplace exponent $\Phi(\beta) = K\beta^{\alpha/2}$ and Lévy measure in (10) for some K > 0. Examples of other subordinators are given in section 4.3.

2.2 Subordinate Brownian motion

For any dimension $d \ge 1$, let $B = \{B(s)\}_{s\ge 0}$ be a d-dimensional Brownian motion with mean-squared displacement

$$\mathbb{E}\left[\|B(s)\|^2\right] = 2ds \quad \text{for all } s \ge 0.$$
(15)

It is well-known that B satisfies the diffusive scaling property,

$$s^{-1/2}B(s) =_{d} B(1) \text{ for all } s > 0.$$
 (16)

If S is an independent subordinator with Laplace exponent Φ , then the Lévy process $X = \{X(t)\}_{t \ge 0}$ defined by

$$X(t) := B(S(t)) + X(0), \quad t \ge 0,$$
(17)

is called a subordinate Brownian motion [50]. That is, X is a random time change of Brownian motion. We assume that the possibly random initial condition $X(0) \in \mathbb{R}^d$ is independent of B and S. The Lévy exponent of X is $\Phi(|\xi|^2)$, meaning

$$\mathbb{E}\left[e^{i\xi\cdot(X(t)-X(0))}\right] = e^{-t\Phi(|\xi|^2)}, \quad \xi \in \mathbb{R}^d, \, t \ge 0.$$

Subordinate Brownian motions are said to be isotropic since their Lévy exponent depends only on $|\xi|^2$. The infinitesimal generator of X can be written as $-\Phi(-\Delta)$, where Δ is the Laplacian in \mathbb{R}^d [50]. It follows immediately from (15)-(17) that the mean-squared displacement of X is

$$\mathbb{E}\left[\|X(t) - X(0)\|^2\right] = 2d\mathbb{E}[S(t)] \quad \text{for all } t \ge 0.$$

2.3 Lévy flights

If S is an $(\alpha/2)$ -stable subordinator with $\alpha \in (0, 2)$ as in (10), then we call the corresponding subordinate Brownian motion $X = \{X(t)\}_{t\geq 0}$ in (17) a Lévy flight [10]. It follows immediately from (14)-(16) that a Lévy flight X satisfies the superdiffusive scaling property,

$$t^{-1/\alpha}X(t) =_{d} X(1) \text{ for all } t > 0.$$
 (18)

Lévy flights arise as a scaling limit of a random walk with heavy-tailed, power law jumps [9]. The probability density function for the position of the Lévy flight satisfies the space fractional Fokker-Planck equation in (2) [11].

2.4 First hitting times (FHTs)

Let $\tau > 0$ denote the FHT of the subordinate Brownian motion X in (17) to some target set $U \subset \mathbb{R}^d$,

$$\tau := \inf\{t > 0 : X(t) \in U\},\tag{19}$$

and let $\sigma > 0$ denote the FHT of the Brownian motion B to U,

$$\sigma := \inf\{s > 0 : B(s) \in U\}.$$
(20)

We are not interested in the behavior of X after time τ , and thus it is enough to consider the so-called stopped subordinate Brownian motion,

$$X(\min\{\tau, t\}) = B(S(\min\{\tau, t\})).$$
(21)

In (21), we first subordinate Brownian motion and then stop the process when it hits the target. Reversing the order of these two operations gives the so-called subordinate stopped Brownian motion,

$$X(t) := B(\min\{\sigma, S(t)\}) \quad t \ge 0.$$

$$(22)$$

The FHT of (22) to U is,

$$\widetilde{\tau} := \inf\{t > 0 : \widetilde{X}(t) \in U\} = \inf\{t > 0 : S(t) > \sigma\}.$$
(23)

While we are primarily interested in τ in (19) rather than $\tilde{\tau}$ in (23), the fact that $\tilde{\tau} \leq \tau$ almost surely plays an important role in studying τ .

3 General analysis

In this section, we present our general analysis and results on subordinate Brownian motions. We begin with two propositions.

3.1 Two useful propositions

The first proposition computes the generator of a subordinator in a case that is useful for our analysis.

Proposition 1. Assume $F : [0, \infty) \to [0, 1]$ is Lipschitz continuous and satisfies

$$F(0) = 0$$
 and $F'(0) := \lim_{s \to 0+} \frac{F(s)}{s} \in [0, \infty).$ (24)

If $S = \{S(t)\}_{t \ge 0}$ is a subordinator with drift $b \ge 0$ and Lévy measure ν , then

$$\lim_{t \to 0+} \frac{\mathbb{E}[F(S(t))]}{t} = \rho := bF'(0) + \int_0^\infty F(s)\,\nu(\mathrm{d}s) < \infty.$$
(25)

Proposition 1 is useful for finding the short-time distribution of functionals of subordinated processes. The next proposition shows how the short-time distribution of a single FHT yields the asymptotic behavior of extreme FHTs.

Before stating the proposition, we recall a few definitions. A random variable T has an exponential distribution with rate $\lambda > 0$ if $\mathbb{P}(T \leq t) = 1 - e^{-\lambda t}$ for $t \geq 0$. If $\{T_i\}_{i=1}^k$ are $k \geq 1$ iid exponential random variables with rate $\lambda > 0$, then their sum has an Erlang distribution with rate $\lambda > 0$ and shape $k \in \{1, 2, 3, ...\}$, which means

$$\mathbb{P}\Big(\sum_{i=1}^{k} T_i \le t\Big) = 1 - \frac{\Gamma(k, \lambda t)}{\Gamma(k)}, \quad t \ge 0.$$

where $\Gamma(a, z) := \int_{z}^{\infty} u^{a-1} e^{-u} du$ is the upper incomplete gamma function. A sequence of random variables $\{Z_N\}_{N\geq 1}$ converges in distribution to Z as $N \to \infty$ if

$$\mathbb{P}(Z_N \le z) \to \mathbb{P}(Z \le z) \quad \text{as } N \to \infty,$$

for all points $z \in \mathbb{R}$ such that $F(z) := \mathbb{P}(Z \leq z)$ is continuous. If $\{Z_N\}_{N\geq 1}$ converges in distribution to an Erlang random variable with rate λ and shape k, then we write $Z_N \to_d \operatorname{Erlang}(\lambda, k)$, and if k = 1, then we write $Z_N \to_d \operatorname{Erlang}(\lambda)$.

Proposition 2. Let $\{\tau_n\}_{n\geq 1}$ be an iid sequence of random variables with

$$\mathbb{P}(\tau_n \le t) \sim \rho t \quad as \ t \to 0+, \tag{26}$$

for some rate $\rho > 0$. Let $T_{k,N}$ be the kth order statistic,

$$T_{k,N} := \min\left\{\{\tau_1, \dots, \tau_N\} \setminus \bigcup_{j=1}^{k-1} \{T_{j,N}\}\right\}, \quad k \in \{1, \dots, N\},$$
(27)

where $T_{1,N} := \min\{\tau_1, \ldots, \tau_N\}$. The following rescaling of $T_{k,N}$ converges in distribution to an Erlang random variable with unit rate and shape k,

$$(\rho N)T_{k,N} \to_{\mathrm{d}} \mathrm{Erlang}(1,k) \quad as \ N \to \infty.$$

If we assume further that $\mathbb{E}[T_{1,N}] < \infty$ for some $N \ge 1$, then

$$\mathbb{E}[(T_{k,N})^m] \sim \frac{\Gamma(k+m)}{\Gamma(k)} \frac{1}{(\rho N)^m} \quad \text{for each moment } m \in (0,\infty) \text{ as } N \to \infty.$$

Proposition 2 is a special case of Theorems 5 and 6 in [51] which were proven for the case $\mathbb{P}(\tau \leq t) \sim \rho t^q$ as $t \to 0$ for some $\rho > 0$ and q > 0.

3.2 Subordinated processes

Before considering subordinate Brownian motion, we first analyze subordinate processes when the "parent" process is not necessarily Brownian. Let $S = \{S(t)\}_{t\geq 0}$ be a subordinator with drift $b \geq 0$ and Lévy measure ν as in section 2.1. Let $Y = \{Y(s)\}_{s\geq 0}$ be a stochastic process independent of S. Define the FHT to a set U in the state space of Y,

$$\sigma := \inf\{s > 0 : Y(s) \in U\}.$$

Define the two subordinations of the "parent" process Y,

$$X(t) := Y(S(t)), \quad X(t) := Y(\min\{\sigma, S(t)\}), \quad t \ge 0.$$

Define the FHTs of X and \widetilde{X} to U,

$$\tau := \inf\{t > 0 : X(t) \in U\}, \quad \widetilde{\tau} := \inf\{t > 0 : \widetilde{X}(t) \in U\} = \inf\{t > 0 : S(t) > \sigma\}.$$

Since Y and S are independent, conditioning on the value of S(t) gives

$$\mathbb{P}(\widetilde{\tau} \leq t) = \mathbb{P}(\sigma \leq S(t)) = \mathbb{E}[\widetilde{F}(S(t))], \quad t \geq 0,$$

where $\widetilde{F}(s) := \mathbb{P}(\sigma \leq s)$. Therefore, if $\widetilde{F}(s)$ is merely Lipschitz and satisfies (24), then Proposition 1 yields the short-time behavior of the distribution of $\widetilde{\tau}$,

$$\lim_{t \to 0+} \frac{\mathbb{P}(\tilde{\tau} \le t)}{t} = \tilde{\rho} := b\tilde{F}'(0) + \int_0^\infty \tilde{F}(s)\,\nu(\mathrm{d}s) < \infty.$$
(28)

Furthermore, if $\tilde{\rho} > 0$ and $\tilde{T}_{k,N}$ is the *k*th fastest FHT of *N* iid realizations of $\tilde{\tau}$ (see (27)), then Proposition 2 yields the large *N* distribution of $\tilde{T}_{k,N}$ in terms of an Erlang random variable. Furthermore, if $\mathbb{E}[\tilde{T}_{1,N}] < \infty$ for some $N \geq 1$, then Proposition 2 also yields the large *N* behavior of the *m*th moment of $\tilde{T}_{k,N}$.

Next, notice that we have the following bounds on the distribution of the FHT τ ,

$$\mathbb{P}(X(t) \in U) \le \mathbb{P}(\tau \le t) \le \mathbb{P}(\tilde{\tau} \le t) \quad \text{for all } t \ge 0,$$
(29)

since $\tilde{\tau} \leq \tau$ almost surely and $X(t) \in U$ implies $\tau \leq t$. Since Y and S are independent, we again condition on the value of S(t) to obtain

$$\mathbb{P}(X(t) \in U) = \mathbb{E}[F(S(t))], \quad t \ge 0,$$

where $F(s) := \mathbb{P}(Y(s) \in U)$. Therefore, if F(s) is Lipschitz and satisfies (24), then Proposition 1 yields

$$\lim_{t \to 0+} \frac{\mathbb{P}(X(t) \in U)}{t} = \rho := bF'(0) + \int_0^\infty F(s)\,\nu(\mathrm{d}s) < \infty.$$
(30)

Therefore, the bounds in (29) and the limits in (28) and (30) yield the following bounds on the short-time behavior of the distribution of τ ,

$$\rho t + o(t) \le \mathbb{P}(\tau \le t) \le \tilde{\rho}t + o(t) \quad \text{as } t \to 0+,$$

where f(t) = o(t) means $f(t)/t \to 0$. If $T_{k,N}$ is the kth fastest FHT of N iid realizations of τ (see (27)), $\rho \tilde{\rho} > 0$, and $\mathbb{E}[T_{1,N}] < \infty$ for some $N \ge 1$, then it follows from Proposition 2 that we can bound the decay of the *m*th moment of $T_{k,N}$ as $N \to \infty$,

$$\frac{\Gamma(k+m)}{\Gamma(k)}\frac{1}{(\tilde{\rho}N)^m} + o(N^{-m}) \le \mathbb{E}[(T_{k,N})^m] \le \frac{\Gamma(k+m)}{\Gamma(k)}\frac{1}{(\rho N)^m} + o(N^{-m}).$$

Summarizing, if X is defined by subordinating some process Y, then Proposition 1 yields information about the short-time distribution of X and FHTs of X. Then, Proposition 2 translates this short-time distribution of a single FHT into the behavior of extreme FHTs. Importantly, these conclusions require only mild assumptions on the parent process Y. In the next subsection, we consider the case that the parent process is a Brownian motion.

3.3 Subordinate Brownian motion

Let $S = \{S(t)\}_{t \ge 0}$ be a subordinator as in section 2.1 and assume that S has nontrivial Lévy measure,

$$\nu((0,\infty)) > 0,\tag{31}$$

to exclude the trivial case in which S is the deterministic function S(t) = btfor all $t \ge 0$. Let $B = \{B(s)\}_{s\ge 0}$ be an independent, d-dimensional Brownian motion for any $d \ge 1$ as in (15). Define $X = \{X(t)\}_{t\ge 0}$ as the random time change of B,

$$X(t) := B(S(t)) + X(0), \quad t \ge 0,$$
(32)

where $X(0) \in \mathbb{R}^d$ is a possibly random initial position independent of S and B.

Let τ be the FHT of X to some target set $U \subset \mathbb{R}^d$ (see (19)). Assume U is nonempty and is the closure of its interior, which precludes trivial cases such as the target having zero Lebesgue measure. Assume that the distribution of X(0)is a probability measure with compact support $U_0 \subset \mathbb{R}^d$ that does not intersect the target,

$$U_0 \cap U = \emptyset. \tag{33}$$

Note that U_0 and U are both closed sets, and thus (33) ensures that U_0 and U are separated by a strictly positive distance. As two examples, the initial distribution could be a Dirac mass at a point $X(0) = x_0 = U_0 \in \mathbb{R}^d$ if $x_0 \notin U$ or it could be uniform on a set U_0 satisfying (33).

Theorem 3. Under the assumptions of section 3.3, we have that

$$\mathbb{P}(\tau \le t) \sim \mathbb{P}(X(t) \in U) \sim \rho t \quad as \ t \to 0+,$$
(34)

where
$$\rho := \int_0^\infty \mathbb{P}(B(s) + X(0) \in U) \,\nu(\mathrm{d}s) \in (0, \infty).$$
 (35)

Furthermore, if $T_N := \min\{\tau_1, \ldots, \tau_N\}$, where $\{\tau_n\}_{n\geq 1}$ is an iid sequence of realizations of τ , then

$$(\rho N)T_N \to_{\mathrm{d}} \mathrm{Exponential}(1) \quad as \ N \to \infty.$$
 (36)

More generally, if $T_{k,N}$ is the kth fastest FHT in (27), then

$$(\rho N)T_{k,N} \to_{\mathrm{d}} \mathrm{Erlang}(1,k) \quad as \ N \to \infty.$$
 (37)

If $\mathbb{E}[T_N] < \infty$ for some $N \ge 1$, then

$$\mathbb{E}[(T_{k,N})^m] \sim \frac{\Gamma(k+m)}{\Gamma(k)} \frac{1}{(\rho N)^m} \quad \text{for each moment } m \in (0,\infty) \text{ as } N \to \infty.$$
(38)

Before applying Theorem 3 to some examples in section 4, we make several comments. First, the asymptotic equality $\mathbb{P}(\tau \leq t) \sim \mathbb{P}(X(t) \in U)$ in (34) means that paths which hit the target before a short time t are much more likely to stay in the target than to leave before t. While this is intuitive, it does not hold for Brownian motion, except on a logarithmic scale (the assumption in (31) means that X cannot be a Brownian motion). Second, (36) means that T_N is approximately exponentially distributed with rate ρN is N is large, and similarly $T_{k,N}$ is approximately Erlang distributed with rate ρN and shape k. Third, the asymptotics in (34) and (38) differ markedly from the case of diffusion. Further, the exponential distribution in (36) differs from the typically Gumbel distributed extreme FHTs of diffusion [52]. See the Introduction section for more on how Theorem 3 differs from the diffusion case. Finally, while (34) gives the short-time distributions, these are equivalent to the "small noise" distributions in the case of a Lévy flight. Indeed, if X is a Lévy flight with generalized diffusion coefficient K, then (34) implies

$$\mathbb{P}(X(t)\in U)\sim Kt\int_0^\infty \mathbb{P}(B(s)+X(0)\in U)\frac{\alpha/2}{\Gamma(1-\frac{\alpha}{2})}\frac{1}{s^{1+\frac{\alpha}{2}}}\,\mathrm{d}s\quad \text{as }K\to 0+.$$

4 Examples and numerical simulation

We now apply Theorem 3 for various choices of the space dimension $d \ge 1$, the target U, and the subordinator S.

4.1 Half-line

Consider a one-dimensional Lévy flight X in \mathbb{R} that starts at X(0) = 0 with $\alpha \in (0, 2)$. That is, X is defined in (32) and S is an $(\alpha/2)$ -stable subordinator defined in section 2.1. Suppose the target is $U = (-\infty, -L]$ for some L > 0. Theorem 3 implies that τ has the short-time distribution in (34) with rate

$$\rho = K \int_0^\infty \mathbb{P}(B(s) \in U) \frac{\alpha/2}{\Gamma(1 - \alpha/2)} \frac{1}{s^{1 + \alpha/2}} \, \mathrm{d}s = \frac{\Gamma(\alpha) \sin(\alpha \pi/2)}{\pi} \frac{K}{L^\alpha} \in (0, \infty),$$

since $\mathbb{P}(B(s) \in U) = \mathbb{P}(B(s) \leq -L) = \frac{1}{2}[1 + \operatorname{erf}(-L/\sqrt{4s})]$ for s > 0. This result for this example was derived formally in [32]. Theorem 3 further implies the convergence in distribution in (36)-(37). In addition, the Sparre-Anderson theorem [29] implies that $\mathbb{P}(\tau > t) = \mathcal{O}(t^{-1/2})$ as $t \to \infty$ which implies

$$\mathbb{E}[T_N] = \int_0^\infty \mathbb{P}(T_N > t) \, \mathrm{d}t = \int_0^\infty (\mathbb{P}(\tau > t))^N \, \mathrm{d}t < \infty \quad \text{if } N \ge 3.$$

Hence, Theorem 3 implies $\mathbb{E}[(T_N)^m] \sim \Gamma(m+1)(\rho N)^{-m}$ as $N \to \infty$ for any m > 0.

These conclusions of Theorem 3 about the asymptotic behavior of T_N as $N \to \infty$ are illustrated in Figure 3 using stochastic simulations (simulation details are given in section 4.6 below). In the top left panel, we plot the empirical probability density of $(\rho N)T_N$ obtained from stochastic simulations with $\alpha = 1.5$. As implied by Theorem 3, $(\rho N)T_N$ converges in distribution to a unit rate exponential random variable. In the top right panel, we plot the maximum difference between the empirical distribution of $(\rho N)T_N$ and a unit rate exponential random variable,

$$\sup_{z\geq 0} \left| \mathbb{P}((\rho N)T_N > z) - \exp(-z) \right|,\tag{39}$$

as a function of N for different choices of α . The difference (39) is the Kolmogorov-Smirnov distance. This plot shows that the convergence of $(\rho N)T_N$ to an exponential random variable is faster for small α . In the bottom two plots, we plot the absolute errors between the simulations and the theory for the mean and standard deviation,

$$\left|\mathbb{E}[T_N] - (\rho N)^{-1}\right|, \quad \left|\sqrt{\operatorname{Variance}[T_N]} - (\rho N)^{-1}\right|, \tag{40}$$

as functions of N for $\alpha = 1.5$ (bottom left panel) and $\alpha = 1$ (bottom right panel). As implied by Theorem 3, these errors decay faster than N^{-1} as N grows.

4.2 Escape from a *d*-dimensional sphere

Consider a Lévy flight X in \mathbb{R}^d with $d \ge 1$ starting at $X(0) = 0 \in \mathbb{R}^d$ with $\alpha \in (0, 2)$. Suppose the target is

$$U = \{ x \in \mathbb{R}^d : ||x|| \ge L \},$$
(41)




Figure 3: Comparison of Theorem 3 and empirical results obtained from stochastic simulations for Lévy flights in the one-dimensional geometry in section 4.1. Top left: Empirical probability density of $(\rho N)T_N$ for $\alpha = 1.5$. Top right: Kolmogorov-Smirnov distance in (39) between the empirical probability density of $(\rho N)T_N$ and a unit rate exponential for different choices of α . Bottom: Absolute errors for the mean and standard deviation in (40) for $\alpha = 1.5$ (bottom left) and $\alpha = 1$ (bottom right). In all four plots, we take K = L = 1.

so that τ is the escape time from a *d*-dimensional sphere of radius L > 0 centered at the origin. Theorem 3 implies that (34) holds with

$$\rho = \rho(L) = K \int_0^\infty \mathbb{P}(\|B(s)\| \ge L) \frac{\alpha/2}{\Gamma(1 - \alpha/2)} \frac{1}{s^{1 + \alpha/2}} \,\mathrm{d}s = \frac{2^\alpha \Gamma(\frac{d + \alpha}{2})}{\Gamma(\frac{d}{2})\Gamma(1 - \frac{\alpha}{2})} \frac{K}{L^\alpha},\tag{42}$$

since $\mathbb{P}(||B(s)|| \ge L) = \Gamma(\frac{d}{2}, \frac{L^2}{4s})/\Gamma(\frac{d}{2})$ for s > 0. Theorem 3 further implies the convergence in distribution in (36)-(37). Furthermore,

$$\mathbb{E}[T_N] \le \mathbb{E}[\tau] = \left[\rho \Gamma\left(1 - \frac{\alpha}{2}\right) \Gamma\left(1 + \frac{\alpha}{2}\right)\right]^{-1} < \infty \quad \text{for any } N \ge 1,$$

where the formula for $\mathbb{E}[\tau]$ is due to Getoor [53]. Therefore, Theorem 3 implies that $\mathbb{E}[(T_N)^m] \sim \Gamma(m+1)(\rho N)^{-m}$ as $N \to \infty$ for any moment $m \in (0, \infty)$.

These results are illustrated in Figure 4 for dimension d = 3. In the top left panel, we plot the empirical probability density of $(\rho N)T_N$ obtained from stochastic simulations with $\alpha = 1.5$, which shows that $(\rho N)T_N$ converges in distribution to a unit rate exponential random variable. The top right panel plots the Kolmogorov-Smirnov distance in (39) as a function of N for difference choices of α . The bottom two plots show the absolute errors for the mean and standard deviation in (40) for $\alpha = 1.5$ (bottom left panel) and $\alpha = 1$ (bottom right panel). As implied by Theorem 3, these errors decay faster than N^{-1} as N grows.

We emphasize that the large N decay of the moments of T_N for Lévy flights is much faster than for normal diffusion. To illustrate, let τ^{diff} be the FHT of a pure diffusion process $\{X^{\text{diff}}(t)\}_{t\geq 0}$ to the target, $\tau^{\text{diff}} := \inf\{t>0: \|X^{\text{diff}}(t)\| \geq L\}$. The mean FHT is $\mathbb{E}[\tau^{\text{diff}}] = \frac{L^2}{2dD}$ [53], where D is the diffusivity of X^{diff} . If $T_N^{\text{diff}} := \min\{\tau_1^{\text{diff}}, \ldots, \tau_N^{\text{diff}}\}$ is fastest FHT out of N iid realizations of τ^{diff} , then [46, 47]

$$\mathbb{E}[T_N^{\text{diff}}] \sim \frac{L^2}{4D \ln N} \quad \text{as } N \to \infty.$$

Now, it is straightforward to choose the diffusion coefficient of X^{diff} so that $\mathbb{E}[\tau] = \mathbb{E}[\tau^{\text{diff}}]$. Hence, for these parameters, the mean FHT for a single Lévy flight and a single diffusion process are identical, but the mean fastest FHT for many Lévy flights is much faster than for many diffusion processes.

4.3 Tempered stable subordinator and gamma subordinator

The slow power law decay of the Lévy measure ν of the stable subordinator S means that a Lévy flight X often takes large jumps. This may be undesirable in some modeling situations, and thus it common to "temper" the stable subordinator by multiplying its Lévy measure by a decaying exponential in order




Figure 4: Comparison of Theorem 3 and empirical results obtained from stochastic simulations for the Lévy flight escape problem in section 4.2 with d = 3. Top left: Empirical probability density of $(\rho N)T_N$ for $\alpha = 1.5$. Top right: Kolmogorov-Smirnov distance in (39) between the empirical probability density of $(\rho N)T_N$ and a unit rate exponential for different choices of α . Bottom: Absolute errors for the mean and standard deviation in (40) for $\alpha = 1.5$ (bottom left) and $\alpha = 1$ (bottom right). In all four plots, we take K = L = 1.

to suppress these large jumps. Specifically, the so-called tempered stable subordinator is defined by zero drift and the following Laplace exponent and Lévy measure,

$$\Phi(\beta) = K((\beta + \mu)^{\alpha/2} - \mu^{\alpha/2}), \qquad \frac{\nu(\mathrm{d}s)}{\mathrm{d}s} = K \frac{\alpha/2}{\Gamma(1 - \alpha/2)} \frac{e^{-\mu s}}{s^{1 + \alpha/2}}, \quad s > 0,$$
(43)

for $\alpha \in (0,2)$, K > 0, and $\mu > 0$. Taking $\alpha \to 0$ in the exponent in the Lévy measure of the tempered stable subordinator yields another subordinator commonly used in modeling called the gamma subordinator, which has zero drift and the following Laplace exponent and Lévy measure for some rate C > 0,

$$\Phi(\beta) = C \log\left(\frac{\beta + \mu}{\mu}\right), \qquad \frac{\nu(\mathrm{d}s)}{\mathrm{d}s} = C \frac{e^{-\mu s}}{s}, \quad s > 0.$$
(44)

Suppose $S = \{S(t)\}_{t\geq 0}$ is the gamma subordinator defined by (44) and let X(t) := B(S(t)) where $B = \{B(s)\}_{s\geq 0}$ is a 3-dimensional Brownian motion. Letting the target be as in (41), Theorem 3 implies that (34) holds with

$$\rho = C \int_0^\infty \mathbb{P}(\|B(s)\| \ge L) \frac{e^{-\mu s}}{s} \, \mathrm{d}s = 2C \Big(e^{-L\sqrt{\mu}} + \int_{L\sqrt{\mu}}^\infty \frac{e^{-z}}{z} \, \mathrm{d}z \Big)$$

Theorem 3 further implies the convergence in distribution in (36)-(37) and the moment behavior in (38) (it is straightforward to check that $\mathbb{E}[T_N] \leq \mathbb{E}[\tau] < \infty$). These results are illustrated in Figure 5 using stochastic simulations (see section 4.6). In the left panel, we illustrate the convergence in distribution in (36) by plotting the Kolmogorov-Smirnov distance in (39) as a function of N. The moment convergence in (38) is illustrated in the right panel of Figure 5, where we plot the absolute errors for the mean and standard deviation (see (40)) as functions of N.

4.4 Annular target in \mathbb{R}^d

As in section 4.2, consider a Lévy flight X in \mathbb{R}^d with $X(0) = 0 \in \mathbb{R}^d$. However, now suppose that the target is the annular region,

$$U = \{ x \in \mathbb{R}^d : 0 < L_- \le ||x|| \le L_+ \}, \text{ where } 0 < L_- < L_+.$$

Hence, (34) holds with $\rho = \rho(L_+) - \rho(L_-) > 0$, where $\rho(L_{\pm})$ is defined in (42) since

$$\mathbb{P}(B(s) \in U) = \mathbb{P}(\|B(s)\| \ge L_{-}) - \mathbb{P}(\|B(s)\| \ge L_{+}).$$

This example illustrates some features not seen in the examples above. First, the FHT τ to U is not the same as the first passage time, $\tau_{\text{fpt}} := \inf\{t > 0 : \|X(t)\| \ge L_{-}\}$. This is because, in contrast to normal diffusion, X is a jump process, and therefore it may "leapover" the annulus U so that $\tau_{\text{fpt}} < \tau$. Second,




Figure 5: Comparison of Theorem 3 and empirical results obtained from stochastic simulations for the problem in section 4.3 of a Brownian motion subordinated by a gamma subordinator. Left: Kolmogorov-Smirnov distance in (39) between the empirical probability density of $(\rho N)T_N$ and a unit rate exponential for different choices of α . Right: Absolute errors for the mean and standard deviation in (40). In both plots, we take $\mu = C = L = 1$.

the FHT is infinite with positive probability in dimensions $d \ge 3$. That is, there exists q(d) > 0 so that

$$\mathbb{P}(\tau = \infty) = q(d) > 0 \quad \text{in dimension } d \ge 3.$$
(45)

To see why (45) holds, note that X may leap over U with positive probability. After leaping over U, the process starts at some radius larger than L_+ and may never return to a radius less than L_+ , as a result of the strong Markov property and the fact that Brownian motion is transient if $d \ge 3$. Third, (45) implies that $\mathbb{P}(T_N = \infty) = (q(d))^N > 0$ if $d \ge 3$. Therefore, the mean fastest FPT is infinite if $d \ge 3$,

$$\mathbb{E}[T_N] = \infty$$
 for every $N \ge 1$ if $d \ge 3$.

Hence, Theorem 3 ensures that the convergence in distribution in (36)-(37) holds, but the moment asymptotics in (38) do not hold.

4.5 Poisson distributed targets in \mathbb{R}^d

Consider again a Lévy flight X in \mathbb{R}^d . Studies of the efficiency of superdiffusive search often consider Poisson distributed targets [25]. To illustrate, suppose $\{x_i\}_{i\in\mathbb{N}}$ is a *d*-dimensional Poisson spatial point process with constant density $\lambda > 0$. Fix a realization of $\{x_i\}_{i\in\mathbb{N}}$ and suppose that the target is obtained by making each point $x_i \in \mathbb{R}^d$ into a ball of radius l > 0,

$$U := \{ x \in \mathbb{R}^d : \|x - x_i\| \le l \quad \text{for some } i \in \mathbb{N} \}.$$

Prior work often considers the case of sparse targets, which means that $\lambda l^d V_d \ll 1$, where $V_d := \pi^{d/2} / \Gamma(1+d/2) > 0$ is the *d*-dimensional volume of a unit sphere.

If the support of the initial distribution of X(0) does not intersect the target (see (33)), then Theorem 3 applies. To approximate the rate ρ in (34), we use that $\mathbb{P}(B(s) \in U)$ vanishes exponentially as $s \to 0+$ and $\mathbb{P}(B(s) \in U) \to \lambda l^d V_d \in$ (0,1) as $s \to \infty$, since $\lambda l^d V_d$ is the fraction of space occupied by targets. The characteristic distance between neighboring x_i and x_j is $L := (\lambda V_d)^{-1/d} \gg l$ and so the characteristic timescale when B reaches the target is L^2 (B has unit diffusivity). Hence, if we approximate $\mathbb{P}(B(s) \in U)$ by 0 for $s < L^2$ and by $\lambda l^d V_d$ for $s > L^2$, then we obtain

$$\rho \approx K \int_{L^2}^{\infty} \lambda l^d V_d \frac{\alpha/2}{\Gamma(1-\frac{\alpha}{2})} \frac{1}{s^{1+\alpha/2}} \,\mathrm{d}s = \frac{V_d^{1+\alpha/d}}{\Gamma(1-\frac{\alpha}{2})} K l^d \lambda^{1+\alpha/d}.$$
 (46)

If we define X via X(t) := B(S(t)) where S is the tempered stable subordinator in (43) with $\mu > 0$, then the analysis above holds and the approximation in (46) is

$$\rho \approx K \int_{L^2}^{\infty} \lambda l^d V_d \frac{\alpha/2}{\Gamma(1-\frac{\alpha}{2})} \frac{e^{-\mu s}}{s^{1+\alpha/2}} \, \mathrm{d}s = \frac{K l^d V_d \alpha \lambda \mu^{\alpha/2} \Gamma(-\frac{\alpha}{2}, (V_d \lambda)^{-2/d} \mu)}{2\Gamma(1-\frac{\alpha}{2})}.$$

4.6 Stochastic simulation algorithm

We now give the stochastic simulation algorithm used to generate FHTs of Lévy flights. Given a discrete time step $\Delta t > 0$, we generate a statistically exact path of the $(\alpha/2)$ -stable subordinator $S = \{S(t)\}_{t\geq 0}$ on the discrete time grid $\{t_k\}_{k\in\mathbb{N}}$ with $t_k = k\Delta t$ via

$$S(t_{k+1}) = S(t_k) + (\Delta t)^{2/\alpha} \Theta_k, \quad k \ge 0,$$

where $S(t_0) = S(0) = 0$ and $\{\Theta_k\}_{k \in \mathbb{N}}$ is an iid sequence of realizations of [54]

$$\Theta = \frac{\sin(\gamma(V + \pi/2))}{(\cos(V))^{1/\gamma}} \left(\frac{\cos(V - \gamma(V + \pi/2))}{E}\right)^{\frac{1-\gamma}{\gamma}}, \quad \text{with } \gamma := \alpha/2 \in (0, 1),$$

where V is uniformly distributed on $(-\pi/2, \pi/2)$ and E is an independent exponential random variable with $\mathbb{E}[E] = 1$. This allows us to generate a statistically exact path of the Brownian motion $\{B(s)\}_{s\geq 0}$ on the (random) discrete time grid $\{S(t_k)\}_{k\in\mathbb{N}}$ via

$$B(S(t_{k+1})) = B(S(t_k)) + \sqrt{2(K\Delta t)^{2/\alpha}\Theta_k}\xi_k, \quad k \ge 0,$$

where $\{\xi_k\}_{k\in\mathbb{Z}}$ is an iid sequence of standard *d*-dimensional Gaussian vectors. Finally, we obtain a statistically exact path of the Lévy process $X = \{X(t)\}_{t\geq 0}$ on the discrete time grid $\{t_k\}_{k\in\mathbb{N}}$ via $X(t_k) = B(S(t_k))$ for $k \geq 0$. The FHT τ to $U \subset \mathbb{R}^d$ is then approximated by $\overline{k} := \min\{k\Delta t \geq 0 : X(t_k) \in U\}$.

Paths of the gamma subordinated Brownian motion in section 4.3 are simulated using the same method, except that $\{\Theta_k\}_{k\in\mathbb{N}}$ is an iid sequence of realizations of gamma random variables with shape $C\Delta t > 0$ and rate $\mu > 0$. The data in Figures 3-5 is computed from 10^5 independent trials with $\Delta t = 10^{-5}$.

5 Discussion

Most studies of search processes measure the speed of search in terms of the FHT of a single searcher. In this paper, we considered the scenario in which there are $N \gg 1$ iid searchers and studied the FHT of the fastest searcher to find the target. Our analysis involved finding the short-time distribution of the FHT of a single searcher and using this to find the distribution and moments of the FHT for the fastest searcher. Our results apply to searchers whose paths follow a subordinate Brownian motion, which is any process obtained by composing a Brownian motion with a Lévy subordinator. We were primarily interested in the case that the searchers move by Lévy flights, which is a prototypical model for superdiffusive search [10].

Previous analysis of extreme FHTs has focused on diffusion, which began with the work of Weiss, Shuler, and Lindenberg in 1983 [46]. The 1/N decay of mean extreme FHTs for subordinate Brownian motion contrasts sharply with the well-known $1/\ln N$ decay of extreme FHTs for diffusion (compare (9) and (11)). See the Introduction section for more on how extreme statistics and large deviation theory for subordinate Brownian motion compare to diffusion. Our results also contrast with results on extreme FHTs of subdiffusive processes modeled by a time fractional Fokker-Planck equation [55]. For searchers exploring a discrete space, an interesting recent study analyzed extreme FHTs for Lévy walks on the two-dimensional integer lattice [45], which was motivated by the Lévy flight foraging hypothesis described in the Introduction section above. Other works investigating extreme FHTs on discrete state networks include [42, 56] in discrete time and [57] in continuous-time.

Biological search processes are often modeled by superdiffusive Lévy walks [4], which are similar to Lévy flights but move with a finite velocity [14]. In particular, Lévy walks follow ballistic flights of uniformly distributed random directions and constant speed, and the lengths of the flights are chosen from a probability density with the slow power law decay in (1). Lévy walks are thus similar to run-and-tumble processes, except run-and-tumble models typically assume the distance of each ballistic flight (i.e. a "run") is chosen from an exponential distribution. The choice of an exponential distribution makes a run-and-tumble a piecewise deterministic Markov process. While Lévy walks are not Markovian, they are nonetheless piecewise deterministic in the sense that the motion is deterministic (constant velocity in a fixed direction) between turns. Extreme FHTs of piecewise deterministic processes were analyzed in [58], and it would be interesting to apply that theory to Lévy walks.

6 Appendix

In this appendix, we prove the results in the main text.

Lemma 4. Assume $S = \{S(t)\}_{t\geq 0}$ is a compound Poisson process plus a drift, meaning its Laplace exponent is in (13) with $b \geq 0$ and $\int_0^\infty \nu(dz) \in (0, \infty)$. If $F: [0, \infty) \to [0, 1]$ is continuous and satisfies (24), then (25) holds. Proof of Lemma 4. By assumption, we have that $S(t) = bt + \sum_{m=1}^{M(t)} Z_m$, where $M = \{M(t)\}_{t\geq 0}$ is a Poisson process with rate $\lambda = \int_0^\infty \nu(dz) \in (0,\infty)$ and $\{Z_m\}_{m\geq 1}$ are iid nonnegative random variables independent of M. In this case, the probability measure of Z_m is $\nu(dz)/\lambda$. Decomposing the mean based on the value of M(t) yields

$$\mathbb{E}[F(S(t))] = \mathbb{E}[F(S(t))1_{M(t)=0}] + \mathbb{E}[F(S(t))1_{M(t)=1}] + \mathbb{E}[F(S(t))1_{M(t)\geq 2}],$$

where 1_A denotes the indicator function on an event A. Since M(t) is a Poisson random variable with mean λt and F is bounded, we have that $\mathbb{E}[F(S(t))1_{M(t)\geq 2}] = o(t)$ as $t \to 0+$. Furthermore, since M and Z_1 are independent, we have that

$$\mathbb{E}[F(S(t))1_{M(t)=1}] = \mathbb{P}(M(t)=1)\mathbb{E}[F(bt+Z_1)] = \lambda t e^{-\lambda t} \mathbb{E}[F(bt+Z_1)],$$
$$\mathbb{E}[F(S(t))1_{M(t)=0}] = \mathbb{P}(M(t)=0)\mathbb{E}[F(bt)] = e^{-\lambda t}F(bt).$$

Since F is bounded, F is continuous, and $\int_0^\infty \nu(ds) < \infty$, we complete the proof by applying the Lebesgue dominated convergence to conclude

$$\mathbb{E}[F(bt+Z_1)] = \frac{1}{\lambda} \int_0^\infty F(bt+s)\,\nu(\mathrm{d}s) \to \frac{1}{\lambda} \int_0^\infty F(s)\,\nu(\mathrm{d}s) \quad \text{as } t \to 0+.$$

Proof of Proposition 1. The boundedness of F and (24) ensure that the integral in (25) is finite. Let $\varepsilon = 2^{-j} > 0$ for some $j \in \{0, 1, 2, ...\}$ and define

$$S_{[\varepsilon,\infty)}(t) := bt + \iint_{z \in [\varepsilon,\infty), t' \in [0,t]} z \mathbf{N}(\mathrm{d}t', \mathrm{d}z),$$

$$S_{(0,\varepsilon)}(t) := \iint_{z \in (0,\varepsilon), t' \in [0,t]} z \mathbf{N}(\mathrm{d}t', \mathrm{d}z),$$
(47)

where **N** is a Poisson point process on the first quadrant with intensity measure $dt' \nu(dz)$. The process S can then be written as $S(t) = S_{[\varepsilon,\infty)}(t) + S_{(0,\varepsilon)}(t)$. Since F is Lipschitz, there exists a constant $\kappa > 0$ so that

$$\mathbb{E}[F(S_{[\varepsilon,\infty)}(t))] - \kappa \mathbb{E}[S_{(0,\varepsilon)}(t))] \le \mathbb{E}[F(S(t))] \\ \le \mathbb{E}[F(S_{[\varepsilon,\infty)}(t))] + \kappa \mathbb{E}[S_{(0,\varepsilon)}(t))] \quad \text{for all } t > 0.$$
(48)

Since $S_{[\varepsilon,\infty)}$ is a compound Poisson process plus a drift, Lemma 4 implies that

$$\lim_{t \to 0+} t^{-1} \mathbb{E}[F(S_{[\varepsilon,\infty)}(t))] = \rho_{\varepsilon} := bF'(0) + \int_{\varepsilon}^{\infty} F(s)\,\nu(\mathrm{d}s) < \infty.$$
(49)

To handle the terms in (48) involving $S_{(0,\varepsilon)}(t)$, recall that $\varepsilon = 2^{-j}$ and observe that a dyadic partitioning of the interval $(0,\varepsilon)$ yields

$$S_{(0,\varepsilon)}(t) := \iint_{z \in (0,\varepsilon), t' \in [0,t]} z \, \mathbf{N}(\mathrm{d}t', \mathrm{d}z) \le \sum_{k=j}^{\infty} 2^{-k} \mathbf{N}([0,t] \times [2^{-k-1}, 2^{-k}]).$$

Since ${\bf N}$ is a Poisson point process, we have that

$$2^{-k}\mathbb{E}\big[\mathbf{N}([0,t]\times[2^{-k-1},2^{-k}])\big] = 2t\int_{2^{-k-1}}^{2^{-k}} 2^{-k-1}\nu(\mathrm{d} z) \le 2t\int_{2^{-k-1}}^{2^{-k}} z\,\nu(\mathrm{d} z).$$

Therefore,

$$\mathbb{E}[S_{(0,\varepsilon)}(t)] \le 2t \int_0^\varepsilon z \,\nu(\mathrm{d}z).$$
(50)

Combining (48) with (49) and (50) yields

$$\rho_{\varepsilon} - 2\kappa \int_0^{\varepsilon} z \,\nu(\mathrm{d} z) \leq \liminf_{t \to 0+} \frac{\mathbb{E}[F(S(t))]}{t} \leq \limsup_{t \to 0+} \frac{\mathbb{E}[F(S(t))]}{t} \leq \rho_{\varepsilon} + 2\kappa \int_0^{\varepsilon} z \,\nu(\mathrm{d} z)$$

Since these bounds converge to ρ as $\varepsilon \to 0+$, the proof is complete.

Lemma 5. Let $H : [0, \infty) \to [0, 1]$ be nondecreasing and satisfy (24). Then $\limsup_{t\to 0+} \mathbb{E}[H(S(t))]/t < \infty$.

Proof of Lemma 5. Using the definitions in (47), we have that

$$H(S(t)) = H(S_{[\varepsilon,\infty)}(t) + S_{(0,\varepsilon)}(t)) \le H(2S_{[\varepsilon,\infty)}(t)) + H(2S_{(0,\varepsilon)}(t)).$$

Since $2S_{[\varepsilon,\infty)}(t)$ is a compound Poisson process plus a drift, Lemma 4 ensures that $\lim_{t\to 0+} \mathbb{E}[H(2S_{[\varepsilon,\infty)}(t))]/t < \infty$. Since H satisfies (24), there exists an $s_0 \in (0,1]$ and a $\theta \ge 1$ so that $H(s) \le \theta s$ for all $s \in (0,s_0]$. Therefore, $H(s) \le \theta s/s_0$ for all $s \ge 0$. The proof is complete since (50) implies

$$\mathbb{E}[H(2S_{(0,\varepsilon)}(t))] \le \frac{2\theta}{s_0} \mathbb{E}[S_{(0,\varepsilon)}(t)] \le \frac{4\theta t}{s_0} \int_0^\varepsilon z \,\nu(\mathrm{d}z).$$

Proof of Theorem 3. Define $F(s) := \mathbb{P}(B(s) + X(0) \in U) \in [0, 1]$ for $s \ge 0$. Using the independence of B and X(0), we have

$$F(s) = \frac{1}{(4\pi s)^{d/2}} \iint_{U \times U_0} \exp\left(\frac{-\|x - x_0\|^2}{4s}\right) \mu_0(\mathrm{d}x_0) \,\mathrm{d}x, \quad \text{if } s > 0, \tag{51}$$

where μ_0 is the probability measure of X(0) with support $U_0 \subset \mathbb{R}^d$. Using standard results for interchanging differentiation with integration (for example, see Theorem A.5.3 in [59]), F(s) is infinitely differentiable and each derivative is bounded. Furthermore, (33) ensures that F(0) = F'(0) = 0 and thus Proposition 1 implies

$$\lim_{t \to 0+} \frac{\mathbb{P}(X(t) \in U)}{t} = \lim_{t \to 0+} \frac{\mathbb{E}[F(s)]}{t} = \rho := \int_0^\infty \mathbb{P}(B(s) + X(0) \in U) \,\nu(\mathrm{d}s).$$
(52)

In the first equality in (52), we have used the independence of B, S, and X(0). Note that $\rho \in (0, \infty)$. Indeed, Proposition 1 implies $\rho < \infty$. Further, $\rho > 0$ by (i) the assumption in (31), (ii) the fact that $B(s) \in \mathbb{R}^d$ is a Gaussian random variable with variance proportional to s > 0, and (iii) U has strictly positive Lebesgue measure (since U is nonempty and the closure of its interior).

To complete the proof, we therefore need to show that

$$\lim_{t \to 0+} t^{-1} \mathbb{P}(\tau \le t) = \lim_{t \to 0+} t^{-1} \mathbb{P}(X(t) \in U).$$
(53)

For t > 0, define the enlarged target $U^{\delta(t)} := \{x \in \mathbb{R}^d : \inf_{y \in U} ||x - y|| \le \delta(t)\}$, where we set $\delta(t) := t^{1/4} > 0$ in order to satisfy

$$\lim_{t \to 0+} \delta(t) = 0 \quad \text{and} \quad \lim_{t \to 0+} \delta(t) t^{-1/2} = \infty.$$
(54)

Decomposing the event $\tau \leq t$ based on the position of X(t) yields

$$\mathbb{P}(\tau \leq t) = \mathbb{P}(X(t) \in U) + \mathbb{P}(\tau \leq t, X(t) \in U^{\delta(t)} \setminus U) + \mathbb{P}(\tau \leq t, X(t) \notin U^{\delta(t)}).$$

Therefore, showing (53) amounts to showing that

$$\lim_{t \to 0+} t^{-1} \mathbb{P}(\tau \le t, X(t) \in U^{\delta(t)} \setminus U) = 0 = \lim_{t \to 0+} t^{-1} \mathbb{P}(\tau \le t, X(t) \notin U^{\delta(t)}).$$
(55)

We first prove the first equality in (55). Since X(t) = B(S(t)) + X(0) and B, X(0), and S are independent, integrating over the possible values of S(t) yields

$$\mathbb{P}(\tau \le t, X(t) \in U^{\delta(t)} \setminus U) \le \mathbb{P}(X(t) \in U^{\delta(t)} \setminus U) = \mathbb{E}[F_0(S(t); t)],$$

where $F_0(s;t) := \mathbb{P}(B(s) + X(0) \in U^{\delta(t)} \setminus U)$. By the assumption in (33), we may take t_0 sufficiently small so that $U^{\delta(t_0)} \cap U_0 = \emptyset$. Therefore, if $t \in (0, t_0]$, then $F_0(s;t)$ satisfies the assumptions of Proposition 1 (by the same argument used for F(s) in (51)). Therefore, Proposition 1 implies that we may take tsufficiently small so that,

$$t^{-1}\mathbb{P}(\tau \le t, X(t) \in U^{\delta(t)} \setminus U) \le 2\int_0^\infty \mathbb{P}(B(s) + X(0) \in U^{\delta(t_0)} \setminus U) \,\nu(\mathrm{d}s) < \infty.$$

Now, it is immediate that $\mathbb{P}(B(s) + X(0) \in U^{\delta(t_0)} \setminus U) \to 0$ as $t_0 \to 0$ for each $s \ge 0$. Hence, the Lebesgue dominated convergence theorem implies

$$\lim_{t_0 \to 0+} \int_0^\infty \mathbb{P}(B(s) + X(0) \in U^{\delta(t_0)} \setminus U) \,\nu(\mathrm{d}s) = 0,$$

and thus the first equality in (55) holds. Turning to the second equality in (55), conditioning that $\tau \leq t$ implies

$$\mathbb{P}(\tau \le t, X(t) \notin U^{\delta(t)}) = \mathbb{P}(X(t) \notin U^{\delta(t)} \mid \tau \le t) \mathbb{P}(\tau \le t),$$

and the fact that $\tilde{\tau} \leq \tau$ almost surely and Lemma 5 imply

$$\limsup_{t \to 0+} t^{-1} \mathbb{P}(\tau \le t) \le \limsup_{t \to 0+} t^{-1} \mathbb{P}(\widetilde{\tau} \le t) < \infty,$$

since $\mathbb{P}(\tilde{\tau} \leq t) = \mathbb{E}[H(S(t)] \text{ where } H(s) := \mathbb{P}(\sigma \leq s) \text{ is nondecreasing. Next, it follows from the strong Markov property [49] that}$

$$\mathbb{P}(X(t) \notin U^{\delta(t)} \mid \tau \le t) \le \sup_{r \in (0,t]} \mathbb{P}_0(\|X(r)\| \ge \delta(t)),$$

where \mathbb{P}_0 denotes the probability measure conditioned that X(0) = 0. Again using that B and S are independent, we have that

$$\mathbb{P}_{0}(\|X(r)\| \ge \delta(t)) = \mathbb{E}[F_{1}(S(r);t)] \le \mathbb{E}[F_{1}(S(t);t)], \quad \text{if } r \in [0,t], \tag{56}$$

since $F_1(s;t) := \mathbb{P}(||B(s)|| \ge \delta(t))$ is an increasing function of s and S is almost surely nondecreasing. Define

$$\delta_1(t) := (1+b)t > 0, \tag{57}$$

and observe that (56) implies that for $r \in (0, t]$,

$$\mathbb{P}_0(\|X(r)\| \ge \delta(t)) \le \mathbb{E}[F_1(S(t);t)\mathbf{1}_{S(t) < \delta_1(t)}] + \mathbb{E}[F_1(S(t);t)\mathbf{1}_{S(t) \ge \delta_1(t)}].$$

Since S(t)/t converges in probability to $b \ge 0$ as $t \to 0+$ [49], we have that

 $\mathbb{E}[F_1(S(t);t)\mathbf{1}_{S(t)\geq \delta_1(t)}] \leq \mathbb{P}(S(t)\geq \delta_1(t)) = \mathbb{P}(S(t)\geq (1+b)t) \to 0 \quad \text{as } t \to 0+.$

Next, since $F_1(s;t)$ is an increasing function of s, we have that

$$\mathbb{E}[F_1(S(t);t)\mathbf{1}_{S(t)<\delta_1(t)}] \le F_1(\delta_1(t);t) = \mathbb{P}(\|B(\delta_1(t))\| \ge \delta(t)).$$

The Brownian scaling in (16) and the choices of $\delta(t)$ in (54) and $\delta_1(t)$ in (57) imply

$$\mathbb{P}(\|B(\delta_1(t))\| \ge \delta(t)) = \mathbb{P}(\|B(1)\| \ge \delta(t)(\delta_1(t))^{-1/2}) \to 0 \quad \text{as } t \to 0 + .$$

Hence, the second equality in (55) holds and the proof is complete.

References

- O Bénichou, C Loverdo, M Moreau, and R Voituriez. Intermittent search strategies. *Rev Mod Phys*, 83(1):81, 2011.
- [2] JR Frost and Lawrence D Stone. Review of search theory: advances and applications to search and rescue decision support. US Department of Transportation, 2001.
- [3] Phillip M Morse and G Kendall. How to hunt a submarine: The world of mathematics, 1956.

- [4] Andy M Reynolds. Current status and future directions of Lévy walk research. Biology open, 7(1), 2018.
- [5] GM Viswanathan, EP Raposo, and MGE Da Luz. Lévy flights and superdiffusion in the context of biological encounters and random searches. *Physics of Life Reviews*, 5(3):133–150, 2008.
- [6] Michael A Lomholt, Tobias Ambjörnsson, and Ralf Metzler. Optimal target search on a fast-folding polymer chain with volume exchange. *Physical review letters*, 95(26):260603, 2005.
- [7] Ming-Yang Kao, John H Reif, and Stephen R Tate. Searching in an unknown environment: An optimal randomized algorithm for the cow-path problem. *Information and Computation*, 131(1):63–79, 1996.
- [8] Michael F Shlesinger. Search research. Nature, 443(7109):281–282, 2006.
- [9] Ralf Metzler and Joseph Klafter. The restaurant at the end of the random walk: recent developments in the description of anomalous transport by fractional dynamics. *Journal of Physics A: Mathematical and General*, 37(31):R161, 2004.
- [10] Alexander A Dubkov, Bernardo Spagnolo, and Vladimir V Uchaikin. Lévy flight superdiffusion: an introduction. *International Journal of Bifurcation* and Chaos, 18(09):2649–2672, 2008.
- [11] Mark M Meerschaert and Alla Sikorskii. Stochastic models for fractional calculus, volume 43. Walter de Gruyter GmbH & Co KG, 2019.
- [12] Anna Lischke, Guofei Pang, Mamikon Gulian, Fangying Song, Christian Glusa, Xiaoning Zheng, Zhiping Mao, Wei Cai, Mark M Meerschaert, Mark Ainsworth, et al. What is the fractional Laplacian? A comparative review with new results. *Journal of Computational Physics*, 404:109009, 2020.
- [13] V Zaburdaev, S Denisov, and J Klafter. Lévy walks. Reviews of Modern Physics, 87(2):483, 2015.
- [14] Michael F Shlesinger and Joseph Klafter. Lévy walks versus Lévy flights. In On growth and form, pages 279–283. Springer, 1986.
- [15] Gandhimohan M Viswanathan, V Afanasyev, SV Buldyrev, EJ Murphy, PA Prince, and H Eugene Stanley. Lévy flight search patterns of wandering albatrosses. *Nature*, 381(6581):413–415, 1996.
- [16] Gandimohan M Viswanathan, Sergey V Buldyrev, Shlomo Havlin, MGE Da Luz, EP Raposo, and H Eugene Stanley. Optimizing the success of random searches. *nature*, 401(6756):911–914, 1999.

- [17] Gabriel Ramos-Fernández, José L Mateos, Octavio Miramontes, Germinal Cocho, Hernán Larralde, and Barbara Ayala-Orozco. Lévy walk patterns in the foraging movements of spider monkeys (ateles geoffroyi). *Behavioral* ecology and Sociobiology, 55(3):223–230, 2004.
- [18] Denis Boyer, Gabriel Ramos-Fernández, Octavio Miramontes, José L Mateos, Germinal Cocho, Hernán Larralde, Humberto Ramos, and Fernando Rojas. Scale-free foraging by primates emerges from their interaction with a complex environment. *Proceedings of the Royal Society B: Biological Sciences*, 273(1595):1743–1750, 2006.
- [19] RPD Atkinson, CJ Rhodes, DW Macdonald, and RM Anderson. Scale-free dynamics in the movement patterns of jackals. *Oikos*, 98(1):134–140, 2002.
- [20] David W Sims, Matthew J Witt, Anthony J Richardson, Emily J Southall, and Julian D Metcalfe. Encounter success of free-ranging marine predator movements across a dynamic prey landscape. *Proceedings of the Royal Society B: Biological Sciences*, 273(1591):1195–1201, 2006.
- [21] Frederic Bartumeus, Francesc Peters, Salvador Pueyo, Celia Marrasé, and Jordi Catalan. Helical Lévy walks: adjusting searching statistics to resource availability in microzooplankton. *Proceedings of the National Academy of Sciences*, 100(22):12771–12775, 2003.
- [22] Julia F Reverey, Jae-Hyung Jeon, Han Bao, Matthias Leippe, Ralf Metzler, and Christine Selhuber-Unkel. Superdiffusion dominates intracellular particle motion in the supercrowded cytoplasm of pathogenic acanthamoeba castellanii. Scientific reports, 5(1):1–14, 2015.
- [23] Ilya Pavlyukevich. Lévy flights, non-local search and simulated annealing. Journal of Computational Physics, 226(2):1830–1844, 2007.
- [24] Andrew M Edwards, Richard A Phillips, Nicholas W Watkins, Mervyn P Freeman, Eugene J Murphy, Vsevolod Afanasyev, Sergey V Buldyrev, Marcos GE da Luz, Ernesto P Raposo, H Eugene Stanley, et al. Revisiting Lévy flight search patterns of wandering albatrosses, bumblebees and deer. Nature, 449(7165):1044–1048, 2007.
- [25] Nicolas Levernier, Johannes Textor, Olivier Bénichou, and Raphaël Voituriez. Inverse square Lévy walks are not optimal search strategies for $d \geq 2$. Physical review letters, 124(8):080601, 2020.
- [26] SV Buldyrev, EP Raposo, Frederic Bartumeus, S Havlin, FR Rusch, MGE da Luz, and GM Viswanathan. Comment on "Inverse square Lévy walks are not optimal search strategies for $d \ge 2$ ". *Physical Review Letters*, 126(4):048901, 2021.
- [27] Nicolas Levernier, Johannes Textor, Olivier Bénichou, and Raphaël Voituriez. Reply to "Comment on 'Inverse square Lévy walks are not optimal search strategies for $d \ge 2$ ". *Physical Review Letters*, 126(4):048902, 2021.

- [28] Iddo Eliazar and Joseph Klafter. On the first passage of one-sided Lévy motions. *Physica A: Statistical Mechanics and its Applications*, 336(3-4):219– 244, 2004.
- [29] Tal Koren, Michael A Lomholt, Aleksei V Chechkin, Joseph Klafter, and Ralf Metzler. Leapover lengths and first passage time statistics for Lévy flights. *Physical review letters*, 99(16):160602, 2007.
- [30] T Koren, AV Chechkin, and J Klafter. On the first passage time and leapover properties of Lévy motions. *Physica A: Statistical Mechanics and its Applications*, 379(1):10–22, 2007.
- [31] Ting Gao, Jinqiao Duan, Xiaofan Li, and Renming Song. Mean exit time and escape probability for dynamical systems driven by Lévy noises. SIAM Journal on Scientific Computing, 36(3):A887–A906, 2014.
- [32] Vladimir V Palyulin, George Blackburn, Michael A Lomholt, Nicholas W Watkins, Ralf Metzler, Rainer Klages, and Aleksei V Chechkin. First passage and first hitting times of Lévy flights and Lévy walks. *New Journal* of *Physics*, 21(10):103028, 2019.
- [33] Asem Wardak. First passage leapovers of Lévy flights and the proper formulation of absorbing boundary conditions. *Journal of Physics A: Mathematical and Theoretical*, 53(37):375001, 2020.
- [34] Z. Schuss, K. Basnayake, and D. Holcman. Redundancy principle and the role of extreme statistics in molecular and cellular biology. *Physics of Life Reviews*, January 2019.
- [35] Thomas W Schoener. Theory of feeding strategies. Annual review of ecology and systematics, 2(1):369–404, 1971.
- [36] James FA Traniello. Recruitment behavior, orientation, and the organization of foraging in the carpenter and camponotus pennsylvanicus degeer (hymenoptera: Formicidae). *Behavioral Ecology and Sociobiology*, 2(1):61– 79, 1977.
- [37] Bert Hölldobler, Edward O Wilson, et al. The ants. Harvard University Press, 1990.
- [38] John W Wenzel and John Pickering. Cooperative foraging, productivity, and the central limit theorem. Proceedings of the National Academy of Sciences, 88(1):36-38, 1991.
- [39] Jennifer UM Jarvis, Nigel C Bennett, and Andrew C Spinks. Food availability and foraging by wild colonies of damaraland mole-rats (cryptomys damarensis): implications for sociality. *Oecologia*, 113(2):290–298, 1998.
- [40] Colin Torney, Zoltan Neufeld, and Iain D Couzin. Context-dependent interaction leads to emergent search behavior in social aggregates. *Proceedings* of the National Academy of Sciences, 106(52):22055–22060, 2009.

- [41] Colin J Torney, Andrew Berdahl, and Iain D Couzin. Signalling and the evolution of cooperative foraging in dynamic environments. *PLoS Comput Biol*, 7(9):e1002194, 2011.
- [42] Ofer Feinerman, Amos Korman, Zvi Lotker, and Jean-Sébastien Sereni. Collaborative search on the plane without communication. In *Proceedings* of the 2012 ACM symposium on Principles of distributed computing, pages 77–86, 2012.
- [43] S D Lawley and J B Madrid. A probabilistic approach to extreme statistics of brownian escape times in dimensions 1, 2, and 3. J Nonlinear Sci, pages 1–21, 2020.
- [44] S Ro and Y W Kim. Parallel random target searches in a confined space. *Phys Rev E*, 96(1):012143, 2017.
- [45] Andrea Clementi, Francesco d'Amore, George Giakkoupis, and Emanuele Natale. On the search efficiency of parallel Lévy walks on Z². arXiv preprint arXiv:2004.01562, 2020.
- [46] G H Weiss, K E Shuler, and K Lindenberg. Order statistics for first passage times in diffusion processes. J Stat Phys, 31(2):255–278, 1983.
- [47] S D Lawley. Universal formula for extreme first passage statistics of diffusion. Phys Rev E, 101(1):012413, 2020.
- [48] Sathamangalam R Srinivasa Varadhan. Diffusion processes in a small time interval. Commun Pure Appl Math, 20(4):659–685, 1967.
- [49] Jean Bertoin. Lévy processes, volume 121. Cambridge University Press, 1996.
- [50] Panki Kim, Renming Song, and Zoran Vondraček. Two-sided green function estimates for killed subordinate brownian motions. *Proceedings of the London Mathematical Society*, 104(5):927–958, 2012.
- [51] Jacob B Madrid and Sean D Lawley. Competition between slow and fast regimes for extreme first passage times of diffusion. Journal of Physics A: Mathematical and Theoretical, 53(33):335002, 2020.
- [52] S D Lawley. Distribution of extreme first passage times of diffusion. *Journal* of Mathematical Biology, 2020.
- [53] RK Getoor. First passage times for symmetric stable processes in space. Transactions of the American Mathematical Society, 101(1):75–90, 1961.
- [54] Sean Carnaffan and Reiichiro Kawai. Solving multidimensional fractional Fokker–Planck equations via unbiased density formulas for anomalous diffusion processes. *SIAM Journal on Scientific Computing*, 39(5):B886–B915, 2017.

- [55] Sean D Lawley. Extreme statistics of anomalous subdiffusion following a fractional fokker-planck equation: subdiffusion is faster than normal diffusion. Journal of Physics A: Mathematical and Theoretical, 53(38):385005, 2020.
- [56] Tongfeng Weng, Jie Zhang, Michael Small, and Pan Hui. Multiple random walks on complex networks: A harmonic law predicts search time. *Physical Review E*, 95(5):052103, 2017.
- [57] Sean D Lawley. Extreme first-passage times for random walks on networks. *Physical Review E*, 102(6):062118, 2020.
- [58] Sean D Lawley. Extreme first passage times of piecewise deterministic markov processes. arXiv preprint arXiv:1912.03438, 2019.
- [59] R Durrett. *Probability: theory and examples.* Cambridge university press, 2019.