

AD-A120 180

NORTH CAROLINA UNIV AT CHAPEL HILL DEPT OF STATISTICS
EXTREMES AND LOCAL DEPENDENCE IN STATIONARY SEQUENCES. (U)
JUN 82 M R LEADBETTER

F/G 12/1

F49620-82-C-0009

UNCLASSIFIED

TR-12

AFOSR-TR-82-0862

NL


1 OF 1
AD A
120 180

END
DATE
FILMED
11-82
DTIC

CENTER FOR STOCHASTIC PROCESSES

Department of Statistics
University of North Carolina
Chapel Hill, North Carolina

AD A120180



EXTREMES AND LOCAL DEPENDENCE IN STATIONARY SEQUENCES

by
M.R. Leadbetter

DTIC
SELECTED
OCT 13 1982
S D E

TECHNICAL REPORT #12

June 1982

Approved for public release;
distribution unlimited.

DTIC FILE COPY

82 1 12 189

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR- 82-0862	2. GOVT ACCESSION NO. AD-A120180	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) EXTREMES AND LOCAL DEPENDENCE IN STATIONARY SEQUENCES		5. TYPE OF REPORT & PERIOD COVERED TECHNICAL
		6. PERFORMING ORG. REPORT NUMBER TR#12
7. AUTHOR(s) M.R. Leadbetter		8. CONTRACT OR GRANT NUMBER(s) F49620-82-C-0009
		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS PE61102F; 2304/A5
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Statistics University of North Carolina Chapel Hill NC 27514		12. REPORT DATE June 1982
		13. NUMBER OF PAGES 24
11. CONTROLLING OFFICE NAME AND ADDRESS Directorate of Mathematical & Information Sciences Air Force Office of Scientific Research Bolling AFB DC 20332		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Extremes; maxima; stationary processes.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Extensions of classical extreme value theory to apply to stationary sequences generally make use of two types of dependence restriction: (a) a weak 'mixing condition' restricting long range dependence; (b) a local condition restricting the 'clustering' of high level exceedances. The purpose of this paper is to investigate extremal properties when the local condition (b) is omitted. It is found that, under general conditions, the type of the limiting distribution for maxima is unaltered. The precise modifications and degree of clustering of high level exceedances are found to be largely described by a (CONTINUED)		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

ITEM #20, CONTINUED: parameter here called the 'extremal index' of the sequence.



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

EXTREMES AND LOCAL DEPENDENCE IN STATIONARY SEQUENCES

by

M.R. Leadbetter*

Summary: Extensions of classical extreme value theory to apply to stationary sequences generally make use of two types of dependence restriction:

- (a) a weak "mixing condition" restricting long range dependence
- (b) a local condition restricting the "clustering" of high level exceedances.

The purpose of this paper is to investigate extremal properties when the local condition (b) is omitted. It is found that, under general conditions, the type of the limiting distribution for maxima is unaltered. The precise modifications and the degree of clustering of high level exceedances are found to be largely described by a parameter here called the "extremal index" of the sequence.

Key words: Extremes, maxima, stationary processes

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	



* Research supported by the Air Force Office of Scientific Research Grant No. ~~AF~~ F49620 82 C 0009.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DTIC
This technical report has been reviewed and is approved for public release IAW AFR 190-12.
Distribution is unlimited.
MATTHEW J. KEEPER
Chief, Technical Information Division

1. INTRODUCTION

Classical Extreme Value Theory discusses the possible limiting laws for the maximum

$$(1.1) \quad M_n = \max(\xi_1, \xi_2 \dots \xi_n)$$

of n independent identically distributed (i.i.d.) random variables (r.v.) as $n \rightarrow \infty$. Specifically it is shown that if M_n has a non-degenerate distribution G i.e. if

$$(1.2) \quad P\{a_n(M_n - b_n) \leq x\} \xrightarrow{d} G(x)$$

for some constants $a_n > 0$, b_n then G must be one of the following classical types (in the sense that $G(x) = H(ax+b)$ for some $a > 0, b$ where H is one of the listed distributions):

Type I	$H(x) = \exp(-e^{-x})$	$-\infty < x < \infty$	
Type II	$H(x) = \exp(-x^{-\alpha})$	$x > 0$	$(\alpha > 0)$
	$= 0$	$x \leq 0$	
Type III	$H(x) = \exp(-(-x)^{\alpha})$	$x \leq 0$	$(\alpha > 0)$
	$= 1$	$x > 0$	

It may be shown that this result remains true (cf [7], [9]) if the condition that the ξ_i be i.i.d. is replaced by the requirement that they form a stationary sequence satisfying a very weak dependence restriction. This restriction, here referred to as the distributional mixing condition $D(u_n)$ is defined as follows.

Write $F_{i_1 \dots i_n}(x_1 \dots x_n) = P\{\xi_{i_1} \leq x_1 \dots \xi_{i_n} \leq x_n\}$ for the joint distribution function of $\xi_{i_1} \dots \xi_{i_n}$, and, for brevity, $F_{i_1 \dots i_n}(u) = F_{i_1 \dots i_n}(u, u, \dots, u)$ for each $n, i_1 \dots i_n, u$. Let $\{u_n\}$ be a sequence of constants. Then the sequence $\{\xi_n\}$ is said to satisfy $D(u_n)$ if for each n, ℓ and each choice of integers $i_1 \dots i_p, j_1 \dots j_p$, such that $1 \leq i_1 < i_2 \dots < i_p < j_1 \dots < j_p \leq n, j_1 - i_p \geq \ell$ we have

$$|F_{i_1 \dots i_p j_1 \dots j_{p'}}(u_n) - F_{i_1 \dots i_p}(u_n) F_{j_1 \dots j_{p'}}(u_n)| < \alpha_{n, \ell}$$

where $\alpha_{n, \ell} \rightarrow 0$ as $n \rightarrow \infty$ for some sequence $\{\ell_n\}$ with $\ell_n = o(n)$.

In spite of the slightly complicated definition this condition is clearly much weaker than the standard forms of mixing condition (such as strong mixing) in that it requires only approximate independence of events A "from the past" and B "from the future" having the special, simple forms

$$A = \bigcap_{r=1}^p \{\xi_{i_r} \leq u_n\}, \quad B = \bigcap_{s=1}^{p'} \{\xi_{j_s} \leq u_n\}.$$

The specific form of the theorem referred to above (proved in [7], [9]), is as follows.

Theorem 1.1. Let $\{\xi_n\}$ be a stationary sequence such that $M_n = \max\{\xi_1, \dots, \xi_n\}$ has a non-degenerate limiting distribution G as in (1.2) for some constants $a_n > 0, b_n$. Suppose that $D(u_n)$ holds for all sequences u_n given by $u_n = x/a_n + b_n$; $-\infty < x < \infty$. Then G is one of the three classical types given above.

Thus the condition $D(u_n)$ alone is sufficient to guarantee that the central classical result concerning the possible extremal types, holds also for stationary sequences.

It is also shown in [7] that if a further condition holds - there called $D'(u_n)$, viz.

$$(1.3) \quad D'(u_n): \quad \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{j=2}^{[n/k]} P\{\xi_1 > u_n, \xi_j > u_n\} \rightarrow 0 \text{ as } k \rightarrow \infty$$

(for each $u_n = x/a_n + b_n$), then the particular type which applies is the same as if the sequence $\{\xi_n\}$ were independent and identically distributed (i.i.d.) with marginal d.f. F , and the same normalizing constants may be used. In particular this means that the classical criteria for domains of attraction (cf. [9]) may be used to determine (on the basis of the behavior of the tail $1 - F(x)$ for

large x) which limiting law applies. These assertions result from making appropriate identifications (e.g. $u_n = x/a_n + b_n$) in the following theorem (cf. [7]) which generalizes a simple classical result.

Theorem 1.2. Let $\{\xi_n\}$ be a stationary sequence and $\{u_n\}$ a sequence of constants such that $D(u_n)$, $D'(u_n)$ hold. Let $0 \leq \tau < \infty$. Then

$$(1.4) \quad P\{M_n \leq u_n\} \rightarrow e^{-\tau}$$

if and only if

$$(1.5) \quad n[1 - F(u_n)] \rightarrow \tau.$$

Conditions similar to $D'(u_n)$ have been used in virtually all studies of extremes of dependent sequences beginning with the early works of Watson [15] and Loynes [10] who showed in particular that (1.5) implies (1.4), using stronger dependence restrictions than $D(u_n)$. However since Theorem 1.1 does not require $D'(u_n)$ in limiting the extremal distributions to the classical types, it seems worthwhile to investigate the precise role of conditions of this kind.

It has in fact been shown by Chernick [3] (extending a result of Loynes [10]) that if for each $\tau > 0$, $u_n = u_n(\tau)$ is defined to satisfy (1.5), then under $D(u_n)$ conditions alone, any limit (function) for $P\{M_n \leq u_n(\tau)\}$ must be of the form

$$(1.6) \quad P\{M_n \leq u_n(\tau)\} \rightarrow e^{-\theta\tau}$$

for some θ with $0 \leq \theta \leq 1$.

In the present paper we extend this result in various ways. It will then follow, as a consequence, that in virtually all cases of practical interest the condition $D(u_n)$ alone is sufficient to guarantee that any asymptotic distribution for the maximum M_n is of precisely the same type as if the sequence $\{\xi_n\}$ were i.i.d. with the same marginal df. F . In fact the only essential difference which appears in dropping the assumption $D'(u_n)$ is that the normalizing constants in

(1.2) may have to be modified from those applying to the i.i.d. case. In obtaining these results we use some ideas from O'Brien ([12], [13]).

The parameter θ in (1.6) is here (as in [9]) called the extremal index of the sequence $\{\xi_n\}$. The main results concerning its existence are given in Section 2, with particular criteria and examples cited in Section 3. In Section 4 we look briefly at the role of $D'(u_n)$ in obtaining a Poisson limit for the (time-normalized) point process of exceedances of the level u_n by the ξ_j 's. When $D'(u_n)$ does not hold, the exceedances of u_n can occur in clusters, leading to multiple points in a limiting point process. As will be seen from Section 4 the degree of clustering is directly related to the extremal index θ .

2. Extremal results under $D(u_n)$

The basic technique of [7] for extending extremal theory to stationary cases is to show that

$$(2.1) \quad P\{M_n \leq u_n\} - P^k\{M_{r_n} \leq u_n\} \rightarrow 0$$

for each $k = 1, 2, \dots$ when $D(u_n)$ holds, where $r_n = [n/k]$ (the integer part of n/k). This clearly simply reflects a form of approximate independence of the submaxima in the k subsets of $[n/k] = r_n$ integers $(1, 2, \dots, r_n)$, $(r_n + 1, \dots, 2r_n)$... which together comprise essentially all of $(1, 2, \dots, n)$. Here we obtain a somewhat more general version of this result. The notation $M(E)$ will be used (here and subsequently) to denote the maximum of ξ_j for j in the set E of integers.

Lemma 2.1. Let $\{u_n\}$ be a sequence of constants and let $D(u_n)$ be satisfied by the stationary sequence $\{\xi_n\}$. Let $\{k_n\}$ be a sequence of constants such that $k_n = o(n)$ and, in the notation used in stating $D(u_n)$, $k_n \ell_n = o(n)$, $k_n \alpha_n, \ell_n \rightarrow 0$. Then

$$(2.2) \quad P\{M_n \leq u_n\} - P^{k_n}\{M_{r_n} \leq u_n\} \rightarrow 0 \text{ as } n \rightarrow \infty$$

where $r_n = [n/k_n]$.

Proof: This will be sketched only since it is analogous to the proof of (2.1) given e.g. in [7], [9]. We shall also assume that $n[1 - F(u_n)]$ is bounded, which is not necessary (cf. [7]) but simplifying (and holds via (1.5) in the applications to be made).

Let $\{\ell_n\}$ be as in the definition of $D(u_n)$. Divide the integers $1 \dots n$ into intervals (i.e. sets of consecutive integers) $I_1, I_1^*, I_2, I_2^* \dots I_{k_n}, I_{k_n}^*$ where $I_1 = (1, 2 \dots r_n - \ell_n)$, $I_1^* = (r_n - \ell_n + 1 \dots r_n)$, $I_2 = (r_n + 1 \dots 2r_n - \ell_n)$, $I_2^* = (2r_n - \ell_n + 1 \dots 2r_n) \dots I_{k_n} = ((k_n - 1)r_n + 1 \dots k_n r_n - \ell_n)$, $I_{k_n}^* = (k_n r_n - \ell_n + 1 \dots n)$. Thus each interval I_j contains $r_n - \ell_n$ integers, with each I_j^* except $I_{k_n}^*$ having ℓ_n integers, and $I_{k_n}^*$ having $n - k_n r_n + \ell_n \leq k_n + \ell_n$ (since $r_n = [n/k_n]$). It is readily seen that

$$\begin{aligned}
 (2.3) \quad 0 &\leq P\left\{\prod_{j=1}^{k_n} \{M(I_j) \leq u_n\}\right\} - P\{M_n \leq u_n\} \\
 &\leq (k_n - 1)P\{M(I_1^*) > u_n\} + P\{M(I_{k_n}^*) > u_n\} \\
 &\leq [(k_n - 1)\ell_n + (k_n + \ell_n)]P\{\xi_1 > u_n\} \\
 &\leq K \frac{k_n(\ell_n + 1)}{n} \rightarrow 0 \text{ as } n \rightarrow \infty
 \end{aligned}$$

by virtue of the stated assumptions (K being a constant).

It follows from $D(u_n)$ by a straightforward induction (cf. [7, Lemma 2.3]) that

$$(2.4) \quad \left| P\left\{\prod_{j=1}^{k_n} \{M(I_j) \leq u_n\}\right\} - P^{k_n}\{M(I_1) \leq u_n\} \right| \leq k_n \alpha_n, \ell_n$$

which tends to zero by assumption. Finally it is readily checked that

$$\begin{aligned}
 (2.5) \quad &\left| P^{k_n}\{M(I_1) \leq u_n\} - P^{k_n}\{M_{r_n} \leq u_n\} \right| \\
 &\leq k_n [P\{M(I_1) \leq u_n\} - P\{M_{r_n} \leq u_n\}] = k_n P\{M(I_1) < u_n \leq M(I_1^*)\} \\
 &\leq k_n \ell_n P\{\xi_1 > u_n\} \leq K k_n \ell_n / n \rightarrow 0.
 \end{aligned}$$

The result now follows at once by combining (2.3), (2.4) and (2.5). \square

We suppose from now on that for each $\tau > 0$ a sequence $\{u_n(\tau)\}$ is defined to satisfy (1.5), viz.

$$(2.6) \quad n[1-F(u_n(\tau))] \rightarrow \tau$$

This imposes a slight restriction on the marginal d.f. F of the ξ_n , but one which will always be satisfied in the applications made. Of course if F is continuous, $u_n(\tau)$ can be defined to give equality in (2.6). In any case it is necessary and sufficient for (2.6) to hold that

$$(2.7) \quad [1-F(x-)]/[1-F(x)] \rightarrow 1 \text{ as } x \rightarrow \infty$$

(cf. [9]), a condition which always holds for any F in any of the three classical domains of attraction. It is also evident that if there exists $u_n(\tau)$ satisfying (2.6) for one fixed $\tau > 0$, then there exists such a $u_n(\tau)$ for all $\tau > 0$ (e.g. if $u_n(1)$ satisfies (2.6) with $\tau = 1$, define $u_n(\tau) = u_{[n/\tau]}(1)$).

The following result reformulates and extends Theorem 1.2.

Theorem 2.2. Let $\{\xi_n\}$ be a stationary sequence and $\{u_n(\tau)\}$ constants satisfying (2.6) and such that $D(u_n(\tau_0))$ holds for some $\tau_0 > 0$. Then there exist constants $\theta, \theta', 0 \leq \theta \leq \theta' \leq 1$ such that

$$(2.8) \quad \begin{aligned} \limsup_{n \rightarrow \infty} P\{M_n \leq u_n(\tau)\} &= e^{-\theta\tau} \\ \liminf_{n \rightarrow \infty} P\{M_n \leq u_n(\tau)\} &= e^{-\theta'\tau} \end{aligned}$$

for $0 < \tau \leq \tau_0$. Hence if $P\{M_n \leq u_n(\tau)\}$ converges for some $\tau, 0 < \tau \leq \tau_0$, then $\theta = \theta'$ and $P\{M_n \leq u_n(\tau)\} \rightarrow e^{-\theta\tau}$ for all such τ .

Proof: Note first that it is readily shown (cf. [9]) that $D(u_n(\tau))$ holds for $0 < \tau \leq \tau_0$ since it holds for $\tau = \tau_0$. Write $\psi(\tau) = \limsup_{n \rightarrow \infty} P\{M_n \leq u_n(\tau)\}$ and let k be a fixed integer. Then it follows from Lemma 2.1 with $k_n = k$ that

$$(2.9) \quad \limsup_{n \rightarrow \infty} P\{M_n \leq u_n(\tau)\} = \psi^{1/k}(\tau)$$

where $n' = [n/k]$. Now if $u_n(\tau) \geq u_{n'}(\tau/k)$ it follows that

$$\begin{aligned} 0 \leq P\{M_n \leq u_n(\tau)\} - P\{M_{n'} \leq u_{n'}(\tau/k)\} &\leq P\left\{\bigcup_{j=1}^{n'} (u_{n'}(\tau/k) < \xi_j \leq u_n(\tau))\right\} \\ &\leq n' [F_{u_n}(\tau) - F(u_{n'}(\tau/k))] \end{aligned}$$

This together with the corresponding inequality when $u_n(\tau) < u_{n'}(\tau/k)$ show that

$$\begin{aligned} |P\{M_n \leq u_n(\tau)\} - P\{M_{n'} \leq u_{n'}(\tau/k)\}| &\leq n' |F(u_n(\tau)) - F(u_{n'}(\tau/k))| \\ &= n' \left| \frac{\tau/k}{n'} (1+o(1)) - \frac{\tau}{n} (1+o(1)) \right| \end{aligned}$$

by (2.6), and this tends to zero as $n \rightarrow \infty$ since $n' \sim n/k$. But clearly

$$\limsup_{n \rightarrow \infty} P\{M_n \leq u_n(\tau/k)\} = \psi(\tau/k), \text{ and it thus follows that } \limsup_{n \rightarrow \infty} P\{M_n \leq u_n(\tau)\} = \psi(\tau/k).$$

Combining this with (2.9) we see that

$$(2.10) \quad \psi(\tau/k) = \psi^{1/k}(\tau) \quad 0 < \tau \leq \tau_0, \quad k = 1, 2, \dots$$

Now $P\{M_n \leq u_n(\tau)\} \geq 1 - n' P\{\xi_1 > u_n(\tau)\} \rightarrow 1 - \tau/k$ as $n \rightarrow \infty$ so that by taking k th powers and using Lemma 2.1, it follows that $\liminf_{n \rightarrow \infty} P\{M_n \leq u_n(\tau)\} \geq (1 - \tau/k)^k$, and letting $k \rightarrow \infty$ that

$$(2.11) \quad \liminf_{n \rightarrow \infty} P\{M_n \leq u_n(\tau)\} \geq e^{-\tau}$$

In particular this implies that $\psi(\tau)$ is strictly positive. It is also non-increasing since if $\tau' < \tau$ it is clear that $u_n(\tau') > u_n(\tau)$ when n is sufficiently large. But the only strictly positive non-increasing solution to the functional equation (2.10) is $\psi(\tau) = e^{-\theta\tau}$ for some $\theta \geq 0$. That is $\limsup_{n \rightarrow \infty} P\{M_n \leq u_n(\tau)\} = e^{-\theta\tau}$ with $\theta \geq 0$.

Similarly it may be shown that $\liminf_{n \rightarrow \infty} P\{M_n \leq u_n(\tau)\} = e^{-\theta'\tau}$ where clearly $\theta' \geq \theta$.

By (2.11) $\theta' \leq 1$ and hence $0 \leq \theta \leq \theta' \leq 1$ as asserted. Thus the relations (2.8) follow and the final statements of the theorem are immediate from these. \square

If $P\{M_n \leq u_n(\tau)\} \rightarrow e^{-\theta\tau}$ for each $\tau > 0$ with $u_n(\tau)$ satisfying (2.6), we shall say that the sequence $\{\xi_n\}$ has extremal index θ (cf. Section 1). Use of this terminology will simplify statements of later results, and in particular gives the following obvious restatement of part of the above theorem.

Corollary 2.3. Let $\{\xi_n\}$ be stationary and satisfy $D(u_n(\tau))$ for each $\tau > 0$ where $n[1-F(u_n(\tau))] \rightarrow \tau$. If for some $\tau_0 > 0$, $P\{M_n \leq u_n(\tau_0)\}$ converges to a limit α then $\{\xi_n\}$ has extremal index $\theta = -\tau_0^{-1} \log \alpha$ so that $P\{M_n \leq u_n(\tau)\} \rightarrow e^{-\theta\tau}$ for all $\tau > 0$. \square

In the next section we shall show that the addition of the condition $D'(u_n)$ (cf. § 1) implies that $\theta = 1$, and give other criteria determining θ when $0 \leq \theta < 1$. However here we proceed with the more general theory, showing that if $\{\xi_n\}$ has a non-zero extremal index θ , then any limiting distribution for the maximum must be of the same type as if the terms were i.i.d. with the same normalizing constants if $\theta = 1$, and simply modified constants for $0 < \theta < 1$. The basic result generalizes a theorem proved by O'Brien [13] under strong mixing assumptions. Here in addition to the previous notation we write

$$\hat{M}_n = \max(\hat{\xi}_1, \hat{\xi}_2, \dots, \hat{\xi}_n)$$

where $\hat{\xi}_1, \hat{\xi}_2, \dots$ are i.i.d. random variables with the same d.f. F as each of the stationary sequence $\hat{\xi}_1, \hat{\xi}_2, \dots$ (following Loynes [10] we call $\hat{\xi}_1, \hat{\xi}_2, \dots$ the "associated independent sequence"). We note the well known (and easily proved) result that for any $\tau > 0$ and sequence $\{u_n\}$,

$$(2.12) \quad P\{\hat{M}_n \leq u_n\} (=F^n(u_n)) \rightarrow e^{-\tau}$$

if and only if

$$(2.13) \quad n[1-F(u_n)] \rightarrow \tau$$

(i.e. if and only if (2.6) holds with $u_n = u_n(\tau)$).

Theorem 2.4. Suppose that the stationary sequence $\{\xi_n\}$ has extremal index θ , $0 \leq \theta \leq 1$. Let $\{v_n\}$ be any sequence of constants and ρ any constant with $0 \leq \rho \leq 1$.

Then

(i) If $\theta > 0$

$$P\{\hat{M}_n \leq v_n\} \rightarrow \rho \text{ if and only if } P\{M_n \leq v_n\} \rightarrow \rho^\theta$$

(ii) If $\theta = 0$

(a) if $\liminf_{n \rightarrow \infty} P\{\hat{M}_n \leq v_n\} > 0$, then $P\{M_n \leq v_n\} \rightarrow 1$

(b) if $\limsup_{n \rightarrow \infty} P\{M_n \leq v_n\} < 1$, then $P\{\hat{M}_n \leq v_n\} \rightarrow 0$

Proof: (i) Suppose $\theta > 0$ and $P\{\hat{M}_n \leq v_n\} \rightarrow \rho$ where $0 < \rho \leq 1$. Choose $\tau > 0$ such that $e^{-\tau} < \rho$. Then

$$P\{\hat{M}_n \leq u_n(\tau)\} \rightarrow e^{-\tau}, \quad P\{\hat{M}_n \leq v_n\} \rightarrow \rho > e^{-\tau}$$

so that $v_n > u_n(\tau)$ for sufficiently large n , and hence

$$\liminf_{n \rightarrow \infty} P\{M_n \leq v_n\} \geq \lim_{n \rightarrow \infty} P\{M_n \leq u_n(\tau)\} = e^{-\theta\tau}$$

Since this holds for any τ such that $e^{-\tau} < \rho$ it follows that $\liminf_{n \rightarrow \infty} P\{M_n \leq v_n\} \geq \rho^\theta$. It also follows in particular that if $\rho = 1$ then $P\{M_n \leq v_n\} \rightarrow 1 = \rho^\theta$.

Similarly by taking $e^{-\tau} > \rho$ it may be shown that $\limsup_{n \rightarrow \infty} P\{M_n \leq v_n\} \leq \rho^\theta$ when $0 \leq \rho < 1$. Hence $P\{M_n \leq v_n\} \rightarrow 0$ when $\rho = 0$, and for $0 < \rho < 1$, $P\{M_n \leq v_n\} \rightarrow \rho^\theta$ by combining the inequalities for the upper and lower limits. The proof of the converse is similar so that (i) follows.

To prove (ii) we assume $\theta = 0$, so that $P\{M_n \leq u_n(\tau)\} \rightarrow 1$ as $n \rightarrow \infty$ for each $\tau > 0$. If $\liminf_{n \rightarrow \infty} P\{\hat{M}_n \leq v_n\} = \rho > 0$, choose τ with $e^{-\tau} < \rho$ and hence $P\{\hat{M}_n \leq u_n(\tau)\} \rightarrow e^{-\tau} < \rho$ so that $v_n \geq u_n(\tau)$ for sufficiently large n . Thus

$$\liminf_{n \rightarrow \infty} P\{M_n \leq v_n\} \geq \lim_{n \rightarrow \infty} P\{M_n \leq u_n(\tau)\} = 1,$$

giving (a).

To show (b) note that if $\limsup P\{M_n \leq v_n\} < 1$ we must have $v_n < u_n(\tau)$ for sufficiently large n and hence

$$\limsup_{n \rightarrow \infty} P\{\hat{M}_n \leq v_n\} \leq \lim_{n \rightarrow \infty} P\{\hat{M}_n \leq u_n(\tau)\} = e^{-\tau}$$

for each τ . The conclusion (b) follows by letting $\tau \rightarrow \infty$. \square

This result now enables us to give conditions in terms of the extremal index under which M_n has a limiting distribution if and only if \hat{M}_n does. This of course implies that in such cases, the classical domain of attraction criteria may be used in the dependent situation.

Theorem 2.5. Let the stationary sequence $\{\xi_n\}$ have extremal index $\theta > 0$. Then M_n has a non-degenerate limiting distribution if and only if \hat{M}_n does, and these are then of the same type based on the same normalizing constants. In the case $\theta = 1$ the limiting distributions for M_n and \hat{M}_n are identical.

Proof: If $P\{a_n(\hat{M}_n - b_n) \leq x\} \rightarrow G(x)$, non-degenerate, then Theorem 2.4 (i) shows (with $v_n = x/a_n + b_n$) that $P\{a_n(M_n - b_n) \leq x\} \rightarrow G^\theta(x)$. But G is an extreme value distribution and it is well known (and easily checked from the possible functional forms) that G^θ is of the same type as G in the sense of Section 1 that $G^\theta(x) = G(ax+b)$ for some $a > 0, b$.

The converse follows similarly, noting that if $P\{a_n(M_n - b_n) \leq x\} \rightarrow H(x)$, non-degenerate, then $P\{a_n(\hat{M}_n - b_n) \leq x\} \rightarrow H^{1/\theta}(x)$. As a limiting distribution for maxima from an i.i.d. sequence, $H^{1/\theta}$ must be of extreme value type and $H = (H^{1/\theta})^\theta$ must be of the same type as $H^{1/\theta}$.

The final remark for $\theta = 1$ is obvious. \square

For the case $0 < \theta < 1$ the same normalizing constants give limits e.g. $G(x)$,

$G^\theta(x) = G(ax+b)$ for \hat{M}_n and M_n . Of course a simple change of the set of normalizing constants for M_n will lead to the same limit $G(x)$.

It is, of course, also of interest to explore the situation when the extremal index is zero. An argument of R. Davis [4] shows (using also Theorem 2.4 (ii)) that M_n and \hat{M}_n cannot both have non-degenerate limiting distributions based on the same normalizing constants. This is stated precisely as follows, without proof.

Theorem 2.6. Let the stationary sequence $\{\xi_n\}$ satisfy $D(u_n(\tau))$ where for each $\tau > 0$ $u_n(\tau)$ satisfies (2.6). If $\{\xi_n\}$ has extremal index $\theta = 0$, then M_n and \hat{M}_n cannot both have non-degenerate limiting distributions based on the same normalizing constants. That is, it is not possible to have $P\{a_n(\hat{M}_n - b_n) \leq x\} \rightarrow G(x)$, $P\{a_n(M_n - b_n) \leq x\} \rightarrow H(x)$ for non-degenerate G, H . \square

Further comments on the (perhaps somewhat pathological) case when $\theta = 0$ will be given in Section 3.

3. Some criteria for the extremal index, and examples.

The first result has perhaps more theoretical than practical interest but serves as a means of extending the condition $D'(u_n)$ to apply to more dependent cases with $\theta < 1$. By way of convenient notation we again write $n' = [n/k]$ for fixed k , $n = 1, 2, \dots$. Also as previously $F_{i_1 \dots i_r}(u)$ will denote the joint d.f. of $\xi_{i_1} \dots \xi_{i_r}$ evaluated at (u, u, \dots, u) .

Theorem 3.1. Let the stationary sequence $\{\xi_n\}$ satisfy $D(u_n(\tau))$ for each $\tau > 0$ where $u_n(\tau)$ satisfies (2.6). Then $\{\xi_n\}$ has extremal index θ ($0 \leq \theta \leq 1$) if and only if

$$(3.1) \quad k \limsup_{n \rightarrow \infty} |1 - F_{1, \dots, n'}(u_n) - \theta \tau_0/k| \rightarrow 0 \text{ as } k \rightarrow \infty$$

for some $\tau_0 > 0$. Equivalently this holds if and only if

$$(3.2) \quad 1 - F_{1, \dots, n'}(u_n) \rightarrow \theta \tau_0/k + \lambda_k \text{ as } n \rightarrow \infty$$

where $k\lambda_k \rightarrow 0$ as $k \rightarrow \infty$.

Proof: For simplicity of notation we take $\tau_0 = 1$ and write $u_n = u_n(1)$. If $\{\xi_n\}$ has extremal index θ , and since (2.1) holds by Lemma 2.1,

$$F_{1 \dots n'}(u_n) = P\{M_n \leq u_n\} \rightarrow e^{-\theta/k} = 1 - \theta/k + o(1/k)$$

from which (3.2) (and hence obviously (3.1)) follow.

Conversely if (3.1) holds then

$$\begin{aligned} \limsup_{n \rightarrow \infty} P\{M_n \leq u_n\} &= \limsup_{n \rightarrow \infty} [F_{1 \dots n'}(u_n) - 1 + \theta/k] + 1 - \theta/k \\ &\leq 1 - \theta/k + \limsup_{n \rightarrow \infty} [1 - F_{1 \dots n'}(u_n) - \theta/k] \\ &= 1 - \theta/k + o(1/k) \end{aligned}$$

Hence again by (2.1)

$$\limsup_{n \rightarrow \infty} P\{M_n \leq u_n\} \leq \{1 - \theta/k + o(1/k)\}^k$$

for all k giving, on letting $k \rightarrow \infty$,

$$\limsup_{n \rightarrow \infty} P\{M_n \leq u_n\} \leq e^{-\theta}$$

The opposite inequality for $\liminf P\{M_n \leq u_n\}$ follows similarly so that $P\{M_n \leq u_n\} \rightarrow e^{-\theta}$. Thus we have convergence of $P\{M_n \leq u_n(\tau)\}$ to $e^{-\theta\tau}$ at $\tau = \tau_0 = 1$ and the result follows from Corollary 2.3. \square

The condition $D'(u_n)$ given by (1.3) limits the probability of one exceedance of u_n being followed "closely" by another. One obvious generalization is to permit (with high probability) no more than some specified number of exceedances to occur together. One specific such restriction is to limit the quantity

$$(3.2) \quad E_{n,k}^{(r)} = \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq n'} P\{\xi_{i_1} > u_n, \xi_{i_2} > u_n, \dots, \xi_{i_r} > u_n\}$$

for some r . For example the assumption $D'(u_n)$ limits $E_{n,k}^{(2)}$ so that in fact

$\limsup_{n \rightarrow \infty} E_{n,k}^{(2)} \rightarrow 0$ as $k \rightarrow \infty$, from which it follows (cf. [7]) that (1.4) holds so that

$\{\xi_n\}$ has extremal index 1. Generalizations of this are clearly possible to allow a non-zero limit for $\limsup_{n \rightarrow \infty} E_{n,k}^{(r)}$ as $k \rightarrow \infty$ for some values of $r \geq 2$ as the following simplest case beyond $D'(u_n)$ shows.

Corollary 3.2. Let the stationary sequence $\{\xi_n\}$ satisfy $D(u_n(\tau))$ for each $\tau > 0$ where $u_n(\tau)$ satisfies (2.6). Suppose that for some $\tau_0 > 0$, $u_n = u_n(\tau_0)$ and some θ , $0 \leq \theta \leq 1$,

$$(3.3) \quad \limsup_{n \rightarrow \infty} |E_{n,k}^{(2)} - \tau_0(1-\theta)| \rightarrow 0 \text{ as } k \rightarrow \infty$$

and

$$(3.4) \quad \limsup_{n \rightarrow \infty} E_{n,k}^{(3)} \rightarrow 0 \text{ as } k \rightarrow \infty$$

Then $\{\xi_n\}$ has extremal index θ .

Proof: Since $1 - F_{1 \dots n'}(u_n) = P\{\bigcup_{j=1}^{n'} (\xi_j > u_n)\}$ it follows by standard Bonferroni inequalities that

$$\begin{aligned} n'[1-F(u_n)] - k^{-1}E_{n,k}^{(2)} &\leq 1 - F_{1 \dots n'}(u_n) \\ &\leq n'[1-F(u_n)] - k^{-1}E_{n,k}^{(2)} + k^{-1}E_{n,k}^{(3)} \end{aligned}$$

and hence

$$\begin{aligned} kn'[1-F(u_n)] - \tau_0(1-\theta) - |E_{n,k}^{(2)} - \tau_0(1-\theta)| &\leq k[1-F_{1 \dots n'}(u_n)] \\ &\leq kn'[1-F(u_n)] - \tau_0(1-\theta) + |E_{n,k}^{(2)} - \tau_0(1-\theta)| + E_{n,k}^{(3)} \end{aligned}$$

Since $u_n = u_n(\tau_0)$, letting $n \rightarrow \infty$ with k fixed yields

$$\begin{aligned} \theta\tau_0 - \limsup_{n \rightarrow \infty} |E_{n,k}^{(2)} - \tau_0(1-\theta)| &\leq \liminf_{n \rightarrow \infty} k[1-F_{1 \dots n'}(u_n)] \\ &\leq \limsup_{n \rightarrow \infty} k[1-F_{1 \dots n'}(u_n)] \leq \theta\tau_0 + \limsup_{n \rightarrow \infty} |E_{n,k}^{(2)} - \tau_0(1-\theta)| + \limsup_{n \rightarrow \infty} E_{n,k}^{(3)} \end{aligned}$$

from which it follows simply that

$$\limsup_{n \rightarrow \infty} |k[1-F_{1 \dots n'}(u_n)] - \theta\tau_0| \leq \limsup_{n \rightarrow \infty} |E_{n,k}^{(2)} - \tau_0(1-\theta)| + \limsup_{n \rightarrow \infty} E_{n,k}^{(3)}$$

which tends to zero as $k \rightarrow \infty$, giving (3.1) and hence the desired conclusion by the theorem. \square

The condition (3.3) may be restated in an obvious way to give the following alternative version of the corollary.

Corollary 3.3. The result of Corollary 3.2 holds if (3.3) is replaced by

$$(3.5) \quad \sum_{j=2}^{n'} (1-j/n') P\{\xi_j > u_n | \xi_1 > u_n\} \rightarrow 1 - \theta + \lambda_k \text{ as } n \rightarrow \infty,$$

where $\lambda_k \rightarrow 0$ as $k \rightarrow \infty$.

Proof: It is simply checked that

$$\begin{aligned} E_{n,k}^{(2)} - \tau_0(1-\theta) &= n[1-F(u_n)] \sum_{j=2}^{n'} (1-j/n') P\{\xi_j > u_n | \xi_1 > u_n\} - \tau_0(1-\theta) \\ &\rightarrow \lambda_k \text{ as } n \rightarrow \infty, \end{aligned}$$

from which (3.3) follows at once. \square

By way of a very simple illustration of the use of results of this type consider i.i.d. random variables η_1, η_2, \dots with d.f. F and define the sequence $\{\xi_n: n \geq 1\}$ by

$$(\xi_1, \xi_2, \xi_3, \dots) = (\eta_1, \eta_2, \eta_2, \eta_3, \eta_3, \dots)$$

$$\text{or } (\eta_1, \eta_1, \eta_2, \eta_2, \dots)$$

each with probability $\frac{1}{2}$. It is readily checked that $P\{\xi_2 > u_n | \xi_1 > u_n\} \rightarrow \frac{1}{2}$

and $P\{\xi_j > u_n | \xi_1 > u_n\} = 1 - F(u_n)$ for $j > 2$ so that

$$\begin{aligned} \sum_{j=2}^{n'} (1-j/n') P\{\xi_1 > u_n, \xi_j > u_n\} &= \frac{1}{2}(1+o(1)) + (1-F(u_n)) \sum_{j=3}^{n'} (1-j/n') \\ &\rightarrow \frac{1}{2} + \tau_0/(2k) \text{ as } n \rightarrow \infty \end{aligned}$$

so that (3.5) holds with $\theta = \frac{1}{2}$. (3.4) is also clearly satisfied as is $D(u_n(\tau))$

so that $\{\xi_n\}$ has extremal index $\theta = \frac{1}{2}$.

While "repeated limit conditions" such as (3.1) can be useful in practice, it may sometimes be more convenient to use conditions depending on a single limit only, and we shall show briefly how this may be done, giving an alternative form for Theorem 3.1.

The condition $D(u_n)$ requires that the quantity $\alpha_{n, \ell_n} \rightarrow 0$ as $n \rightarrow \infty$ for some $\ell_n = o(n)$. It is clearly possible to obtain $k_n \rightarrow \infty$ such that both

$$(3.6) \quad k_n \alpha_{n, \ell_n} \rightarrow 0$$

and

$$(3.7) \quad k_n \ell_n = o(n)$$

hold (e.g. taking $k_n = \min(\alpha_{n, \ell_n}^{-\frac{1}{2}}, (n/\ell_n)^{\frac{1}{2}})$). Using such a sequence k_n we have the following variant of Theorem 3.1.

Theorem 3.4. Let the stationary sequence $\{\xi_n\}$ satisfy $D(u_n(\tau))$ for each $\tau > 0$ where $u_n(\tau)$ satisfies (2.6). For some $\tau_0 > 0$ let $k_n \rightarrow \infty$ be such that (3.6) and (3.7) hold with $u_n = u_n(\tau_0)$. If, writing $r_n = [n/k_n]$,

$$(3.8) \quad k_n [1 - F_1 \dots_{r_n}(u_n)] \rightarrow \theta \tau_0 \text{ as } n \rightarrow \infty \quad (0 \leq \theta \leq 1)$$

then $\{\xi_n\}$ has extremal index θ . Conversely if $\{\xi_n\}$ has extremal index θ then (3.8) holds for each $\tau_0 > 0$ and each $k_n \rightarrow \infty$ satisfying (3.6) and (3.7) with $u_n = u_n(\tau_0)$.

Proof: If (3.8) holds then

$$P\{M_{r_n} \leq u_n\} = F_1 \dots_{r_n}(u_n) = 1 - \frac{\theta \tau_0}{k_n} (1 + o(1))$$

so that

$$P^{k_n}\{M_{r_n} \leq u_n\} = \left[1 - \frac{\theta \tau_0}{k_n} + o\left(\frac{1}{k_n}\right)\right]^{k_n} \rightarrow e^{-\theta \tau_0}$$

and hence $P\{M_n \leq u_n\} \rightarrow e^{-\theta \tau_0}$ by Lemma 2.1 showing that $\{\xi_n\}$ has extremal index θ by Corollary 2.3.

Conversely if $\{\xi_n\}$ has extremal index θ , and $\tau_0 > 0$, $k_n \rightarrow \infty$ satisfying (3.6) and (3.7) then Lemma 2.1 shows that $P^{k_n}(M_{r_n} \leq u_n) \rightarrow e^{-\theta\tau_0}$ with $u_n = u_n(\tau_0)$ follows simply that $F_1 \dots F_{r_n}(u_n) \rightarrow 1$ and

$$\log[1 - (1 - F_1 \dots F_{r_n}(u_n))] = -\frac{\theta\tau_0}{k_n} (1 + o(1))$$

so that

$$-[1 - F_1 \dots F_{r_n}(u_n)] [1 + o(1)] = -\frac{\theta\tau_0}{k_n} (1 + o(1))$$

giving (3.8) as required. \square

A simply expressed sufficient condition for (3.8) may be given as in the following corollary. In this we write $E_n^{(s)}$ for $E_{n, k_n}^{(s)}$ where this is given by (3.2) i.e.

$$(3.9) \quad E_n^{(s)} = k_n \sum_{1 \leq i_1 < i_2 < \dots < i_s \leq r_n} P\{\xi_{i_1} > u_n \dots \xi_{i_s} > u_n\}$$

(where $r_n = [n/k_n]$).

Corollary 3.5. Let the stationary sequence $\{\xi_n\}$ satisfy $D(u_n(\tau))$ for each $\tau > 0$, where $u_n(\tau)$ satisfies (2.6) For some $\tau_0 > 0$ let $k_n \rightarrow \infty$ be such that (3.6) and (3.7) hold with $u_n = u_n(\tau_0)$. Suppose that for each $s = 1, 2, \dots$ the $E_n^{(s)}$ defined by (3.9) satisfy

$$(3.10) \quad E_n^{(s)} \rightarrow \alpha_s$$

where $\alpha_s \rightarrow 0$ as $s \rightarrow \infty$. Then $\{\xi_n\}$ has extremal index

$$\theta = \tau_0^{-1} \sum_{r=1}^{\infty} (-1)^{r-1} \alpha_r$$

Proof: Write $\lambda_n = k_n [1 - F_1 \dots F_{r_n}(u_n)] = k_n P\{\bigcup_{j=1}^{r_n} (\xi_j > u_n)\}$. Then using Bonferroni Inequalities we have for s odd, $n > s$,

$$E_n^{(1)} - E_n^{(2)} + E_n^{(3)} \dots + E_n^{(s)} \geq \lambda_n \geq E_n^{(1)} - E_n^{(2)} + E_n^{(3)} \dots - E_n^{(s+1)}$$

Writing $\underline{\lambda} = \liminf_{n \rightarrow \infty} \lambda_n$, $\bar{\lambda} = \limsup_{n \rightarrow \infty} \lambda_n$ and letting $n \rightarrow \infty$, we obtain, for each odd s ,

$$(3.11) \quad \alpha_1 - \alpha_2 + \alpha_3 \dots + \alpha_s \geq \bar{\lambda} \geq \underline{\lambda} \geq \alpha_1 - \alpha_2 + \alpha_3 \dots - \alpha_{s+1}$$

Since the extreme terms differ by α_{s+1} which tends to zero as $s \rightarrow \infty$, it follows that

$\bar{\lambda} = \underline{\lambda}$, say. If $T_s = \sum_{r=1}^s (-)^{r-1} \alpha_r$ it further follows from (3.11) that

$$0 \leq T_s - \lambda \leq T_s - T_{s+1} = \alpha_{s+1} \text{ for } s \text{ odd and similarly } 0 \leq \lambda - T_s \leq T_{s-1} - T_s = \alpha_s$$

for s even, so that in both cases

$$|T_s - \lambda| \leq \alpha_s + \alpha_{s+1} \rightarrow 0 \text{ as } s \rightarrow \infty$$

Hence $\sum_{r=1}^{\infty} (-)^{r-1} \alpha_r$ converges to the value λ and (3.8) holds with $\theta \tau_0 = \sum_{r=1}^{\infty} (-)^{r-1} \alpha_r$ giving the desired result. \square

Finally in this section we cite some examples of sequences exhibiting all the possible types of behavior relative to the extremal index. In each of these cases $D(u_n(\tau))$ is satisfied.

The most common case is where $D'(u_n(\tau))$ holds leading to the extremal index $\theta=1$. For example this is so for a stationary normal sequence $\{\xi_n\}$ with covariance sequence $\{r_n\}$ satisfying the condition of S.M. Berman, [2], viz. $r_n \log n \rightarrow 0$ - an obviously weak condition indeed.

We have given a simple example of a case when $\theta = \frac{1}{2}$ in the discussion above. An example where a series of values of θ is possible through parameter choice in an autoregressive scheme, has been given by Chernick [3]. The stable processes considered by Rootzén [14], can have any value of θ in the range $0 < \theta \leq 1$. A simple example due to L. de Haan also exhibiting this behavior is the sequence

$$\xi_n = \max_{k \geq 0} \rho^k \eta_{n-k}$$

where $0 < \rho < 1$ and $\{\eta_n\}$ is an i.i.d. sequence with common d.f. $\exp(-1/x)$. This yields an extremal index $\theta = 1 - \rho$.

An example of Denzel and O'Brien [5] exhibits a "chain dependent" sequence $\{\xi_n\}$ with extremal index $\theta = 0$. In this case \hat{M}_n has a Type II limiting distribution, but

we do not know whether M_n has any sort of limiting distribution.

A further example of L. de Haan, however, provides a case where $\theta = 0$ and M_n, \hat{M}_n both have limiting distributions. Specifically the sequence $\{\xi_n\}$ is defined

$$\text{by } \xi_n = \max_{k \geq 0} (\eta_{n-k}^{-k})$$

where η_n are i.i.d. with common d.f. $\exp(-x^{-\alpha})$ $x > 0$, ($\alpha > 1$). In this case M_n has a Type III limit with parameter α and norming constants $a_n = n^{-1/\alpha}$, $b_n = 0$ whereas \hat{M}_n has a Type II limit with parameter $\alpha - 1$ and norming constants $a_n = n^{-1/(\alpha-1)}$, $b_n = 0$.

Finally an example of O'Brien [12] exhibits a case in which $\{\xi_n\}$ has no extremal index at all. In this each ξ_n is uniform over the interval $[0,1]$, $\xi_1, \xi_3 \dots$ being independent and ξ_{2n} a certain function of ξ_{2n-1} for each n .

4. Point process of clusters.

As noted in Section 1, when $n[1-F(u_n)] \rightarrow \tau$ and $D(u_n)$ and $D'(u_n)$ both hold, the (time normalized) instants at which the sequence exceeds u_n take on a Poisson character as n becomes large. More specifically let N_n denote the point process on the unit interval $(0,1]$ consisting of those points j/n such that $\xi_j > u_n$. Then under the conditions above it may be shown ([8]) that N_n converges weakly to a Poisson Process with intensity τ on $(0,1]$.

When $D'(u_n)$ does not hold, the exceedances of u_n may tend to occur in clusters, leading to the simultaneous occurrence of multiple events i.e. a "compounding" in the limiting point process. A complete description of the limiting point process has been given by Rootzén [14] in the case where the underlying sequence $\{\xi_n\}$ belongs to a class of stable processes (cf. the above discussion in Section 3).

Again under a (multidimensional type of) strengthening of the condition $D(u_n)$, and assuming $D'(u_n)$, it is possible to obtain a "complete Poisson theorem" (cf. [1], [9]). This involves convergence of the point process in the plane with points at

$(j/n, (\xi_j - b_n)/a_n)$, with appropriate a_n, b_n , to a certain Poisson process in the plane. Results of this type allow rather complete descriptions of (joint) asymptotic distributional results for extreme order statistics.

It is also of interest to determine the effect of eliminating the condition $D'(u_n)$ in results of this type. For example Mori [11] has shown that under strong mixing the limiting point processes are confined to a certain class (and it seems likely that this is true also under the weaker $D(u_n)$ -type of condition).

We shall not investigate limiting results of these types in detail here. However it does seem interesting and useful to give the simplest of convergence results - involving the Poisson limit for the point process "positions" of the "clusters" of exceedances of high levels. This is analogous to a result of Rootzén in [14] for stable processes.

One very simple means of defining clusters of exceedances is to take a sequence r_n and consider that events occurring within a distance r_n of each other belong to the same cluster. r_n should of course be chosen so that it is at least as large as (virtually all) cluster "lengths" but small compared with cluster "separation." For many usual situations this still leaves considerable flexibility in the choice of r_n , while leading to unique results as we shall see.

More specifically we shall suppose that $D(u_n)$ holds for $u_n = u_n(\tau)$ satisfying (2.6), a sequence $k_n \rightarrow \infty$ is chosen to satisfy (3.6) and (3.7) and $r_n = [n/k_n]$. A point process N_n is defined on the unit interval $(0,1]$ as follows. If for given $s = 1, 2 \dots k_n$ there is an exceedance of u_n by ξ_j for at least one j such that $(s-1)r_n < j \leq sr_n$, then N_n has a single event at the point $t = sr_n/n$. That is any group of exceedances between $(s-1)r_n$ and sr_n is replaced by a single event - after time-scaling - at sr_n/n , "representing" the original group. We refer to N_n as the "point process of cluster positions." With this construction the

following result holds.

Theorem 4.1. Let the stationary sequence $\{\xi_n\}$ satisfy $D(u_n(\tau))$ for each $\tau > 0$ where $u_n(\tau)$ satisfies (2.6). Let $k_n \rightarrow \infty$ be chosen to satisfy (3.6) and (3.7) and let $\{\xi_n\}$ have extremal index θ ($0 < \theta \leq 1$). Then the point process N_n of cluster positions for exceedances of $u_n(\tau)$ converges in distribution to a Poisson Process N on $(0,1]$ with intensity parameter $\theta\tau$.

Proof: As in previous proofs of similar results (cf. [8]) it is by a theorem of Kallenberg [6] only necessary to show that

$$(4.1) \quad EN_n\{(a,b]\} \rightarrow EN\{(a,b]\} \quad \text{for } 0 < a < b \leq 1$$

and

$$(4.2) \quad P\{N_n(E)=0\} \rightarrow P\{N(E)=0\}$$

for each finite disjoint union E of sets $(a_i, b_i] \subset (0,1]$.

If v_n denotes the number of (integer) intervals $((s-1)r_n, sr_n]$ completely contained in $([na], [nb])$ it is clear that $v_n \sim nr_n^{-1}(b-a) \sim k_n(b-a)$ and further that

$$\begin{aligned} EN_n\{(a,b]\} &\sim v_n P\left\{\bigcup_{i=1}^{v_n} (\xi_i > u_n)\right\} \\ &\sim k_n(b-a) [1 - F_{1 \dots r_n}(u_n)] \\ &\rightarrow (b-a)\theta\tau \end{aligned}$$

by (3.8). But this is just $EN\{(a,b]\}$ so that (4.1) follows.

To show (4.2) we write $E = \bigcup_1^p (a_j, b_j]$ and write B_j for the integers in $([na_j], [nb_j])$. Then it is readily seen that

$$\begin{aligned} P\{N_n(E)=0\} &= P\left\{\bigcap_{j=1}^p (M(B_j) \leq u_n)\right\} + o(1) \\ &= \prod_{j=1}^p \{P(M(B_j) \leq u_n)\} + [P\left\{\bigcap_{j=1}^p (M(B_j) \leq u_n)\right\} - \prod_{j=1}^p P\{M(B_j) \leq u_n\}] + o(1) \end{aligned}$$

By a straightforward induction, the difference in square brackets does not exceed $p\alpha_{n,n\lambda}$ in absolute value where λ is the minimum separation of the intervals $(a_j, b_j]$ (λ can be taken non-zero since abutting intervals can be combined). But $\alpha_{n,\ell}$ may be taken non-increasing in ℓ (cf. [9]) and it follows from $D(u_n)$ that $\alpha_{n,n\lambda} \rightarrow 0$ as $n \rightarrow \infty$. Since $\{\xi_n\}$ has extremal index θ it follows in an obvious way that $P\{M(B_j) \leq u_n\} \rightarrow e^{-\theta\tau(b_j - a_j)}$ and hence

$$\begin{aligned} P\{N_n(E)=0\} &\rightarrow \prod_{j=1}^p P\{N(a_j, b_j]=0\} \\ &= P\{N(E)=0\} \end{aligned}$$

proving 4.2. □

It is of interest to note an intuitively appealing interpretation of the extremal index as the inverse of mean cluster size. This may be seen even in terms of the simple approach above. For the mean cluster size can be interpreted as the (limiting) mean number of exceedances in an interval of length r_n , given at least one exceedance in that interval i.e. if Z denotes the number of exceedances of $u_n(\tau)$ in an interval of length r_n ,

$$\begin{aligned} E\{Z|Z \geq 1\} &= \sum_{s=1}^{\infty} sP\{Z=s|Z \geq 1\} \\ &= EZ/P\{Z \geq 1\} \\ &= r_n[1-F(u_n)]/F_1 \dots r_n(u_n) \\ &\rightarrow \theta^{-1}. \end{aligned}$$

Finally it should be noted that the limiting distributions of extreme order statistics will be affected in a more complicated way by the clustering than the maximum. These distributions would emerge from the more complete limiting result for individual exceedances. However use of the simple Poisson result given above will result in the distributions for the heights of the "kth highest clusters"

rather than the k th extreme order statistics, in an obvious way. This of course is analogous to consideration of k th highest local maxima in continuous parameter situations.

ACKNOWLEDGEMENTS

It is a pleasure to express my gratitude to Holger Rootzén, Georg Lindgren and Laurens de Haan for many stimulating conversations concerning this and related topics.

REFERENCES

- [1] Adler R.J. "Weak convergence results for extremal processes generated by dependent random variables" Ann. Probab. 6 660-667, 1978.
- [2] Berman S.M. "Limiting distribution of the maximum term in a sequence of dependent random variables" Ann. Math. Statist. 33 894-908 1962.
- [3] Chernick M.R. "A limit theorem for the maximum of autoregressive processes with uniform marginal distribution" Ann. Probab. 9 145-149 1981.
- [4] Davis R.A. "Limit laws for the maximum and minimum of stationary sequences" Technical report Mass. Inst. Tech. 1981.
- [5] Denzel G.E. and O'Brien G.L. "Limit theorems for extreme values of chain-dependent processes" Ann. Probab. 3 773-779, 1975.
- [6] Kallenberg O. "Random measures" Akademie-Verlag and Academic Press, 1976.
- [7] Leadbetter M.R. "On extreme values in stationary sequences" Z. Wahrsch. verw. Geb. 28 289-303 1974.
- [8] _____ "Weak convergence of high level exceedances by a stationary sequence" Z. Warsch. verw. Geb. 34 11-15 1976.
- [9] Leadbetter M.R., Lindgren G., Rootzén, H. "Extremes and related properties of random sequences and processes" to appear in Springer Statistics Series 1982.
- [10] Loynes R.M. "Extreme values in uniformly mixing stationary stochastic processes" Ann. Math. Statist. 36, 993-999 1965.
- [11] Mori T. "Limit distributions of two-dimensional point processes generated by strong mixing sequences" Yokohama Math. J. 25 155-168 1977.
- [12] O'Brien, G.L. "The maximum term of uniformly mixing stationary processes" Z. Warsch. verw. Geb. 30 57-63 1974.
- [13] _____ "Limit theorems for the maximum term of a stationary process" Ann. Probab. 2 540-545 1974.
- [14] Rootzén, H. "Extremes of moving averages of stable processes" Ann. Probab. 6 847-869 1978.
- [15] Watson G.S. "Extreme values in samples from m-dependent stationary processes" Ann. Math. Statist. 25 798-800 1954

