

Facile and Efficient Method for Synthesis of Benzimidazole Derivatives Catalyzed by Zinc Triflate

Ramineni Srinivasulu¹, Kannasani Ravi Kumar², Peruri Veera Venkata Satyanarayana³

¹Chalapathi Institute of Engineering and Technology, Department of chemistry, Guntur, India

²RA Chem Pharma Limited, R&D Division, Prasanth Nagar, Hyderabad, India

³Department of Chemistry, Acharya Nagarjuna University, Guntur, India

Email: ramineni29@yahoo.com

Received December 4, 2013; revised January 4, 2014; accepted January 11, 2014

Copyright © 2014 Ramineni Srinivasulu *et al.* This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. In accordance of the Creative Commons Attribution License all Copyrights © 2014 are reserved for SCIRP and the owner of the intellectual property Ramineni Srinivasulu *et al.* All Copyright © 2014 are guarded by law and by SCIRP as a guardian.

ABSTRACT

We report the synthesis of benzimidazole derivatives using zinc triflate as an efficient catalyst. One-pot synthesis of 2-substituted benzimidazole derivatives from *o*-phenylenediamine and substituted aldehydes were developed under zinc triflate in ethanol solvent at reflux temperature.

KEYWORDS

Benzimidazole Derivatives; *O*-Phenylenediamine; Substituted Aldehydes; Zinc Triflate

1. Introduction

Benzimidazole derivatives have received much interest in the field of medicinal chemistry [1,2]. Benzimidazole group of substances has found practical applications in a number of fields. Recently the interest in benzimidazole chemistry has been revived by the discovery that the 5,6-dimethyl benzimidazole moiety is part of the chemical structure of vitamin B12 [3]. Substituted Benzimidazoles display a broad spectrum of potential pharmacological activities and are present in a number of pharmacologically active molecules such as albendazole/mebendazole/thiabendazole (antihelminthic), omeprazole (anti-ulcer), etc. (Figure 1). Considerable interest has been focused on the benzimidazole structure. The discovery of this class of drugs provides an outstanding case history of modern drug development and also points out the unpredictability of pharmacological activity from structural modification of a prototype drug molecule. It is having a variety of medicinal applications. Benzimidazole derivatives carrying different substituent's in the benzimidazole structure were associated with a wide range of biological activities including anticancer, antiviral, antibacterial, antifungal, antihelminthic, anti-inflammatory, antihistaminic, proton pump inhibitor, antioxidant, antihypertensive and anticoagulant activities. Their derivatives were also

found to exhibit cytotoxic activity. Substituted benzimidazole derivatives are evaluated by their ability to inhibit gastric H⁺/K⁺ ATPase and by blocking the gastric acid secretion [4]. Recently, benzimidazoles have also been used as ligands for asymmetric catalysis [5].

Many methods have been reported for the synthesis of these benzimidazole derivatives. The condensation of 1,2-phenylenediamines with carboxylic acids or their derivatives is a common method, but it needs harsh conditions like polyphosphoric acid [6] at 170°C - 180°C. Another alternative approach is the condensation of aldehyde with 1,2-phenylenediamine in presence of different catalysts like Indion 190 resin [7], BF₃.OEt₂ [8], Ceric ammonium nitrate [9], iodine, [10] Silica sulfuric acid [11], In(OTf)₃ [12], SiO₂/ZnCl₂ [13], silica supported sodium hydrogen sulphate [14], PEG [15], H₂O₂/Fe(NO₃)₃ [16]. In recent years, Solvent-free synthesis of benzimidazoles under microwave irradiation using Yb(OTf)₃ [17], KSF clay [18], metal halide supported alumina [19] and solid support [20,21] has been reported. However, many of these methods suffer from one or more drawbacks such as requirement of strong acidic conditions, long reaction times, low yields, tedious work-up procedures, requirement of excess amounts of reagents, and use of toxic reagents, catalysts or solvents.




Figure 1. Established antiulcer agents in clinical practice.

Therefore, there is a strong demand for a highly efficient and environmentally benign method for the synthesis of these heterocycles.


As part of our research program in developing various synthetic methodologies, we report the synthesis of benzimidazoles using zinc triflate as an efficient catalyst (**Scheme 1**). The catalyst is known as an efficient catalyst in the literature for various organic transformations [22–26].

2. Results and Discussions

In order to establish the optimum reaction condition for this reaction, different solvents and various mole ratios of zinc triflate were examined. In our preliminary investigation was carried out on the model reaction of *o*-phenylenediamine and 4-methoxy benzaldehyde. As shown in **Table 1**, different solvents can result in different yields. It was found that ethanol is the best solvent for condensation reaction, with its fast conversion, high yield and low toxicity. Zinc triflate was added in various mole ratios in ethanol at reflux. As shown in **Table 2**. The best yields were obtained with 10 mol% of zinc triflate. The electronic effects of the different substituted aldehydes have been investigated in **Table 3** and it was observed that aldehydes bearing both electron donating and electron withdrawing substituents gave the desired benzimidazoles in good yields. Products were confirmed by comparing with authentic sample (^1H NMR, IR and Mass).

3. Conclusions

In conclusion, Zinc triflate was found to be an efficient catalyst for the formation of benzimidazole from aldehydes and *o*-phenylenediamine. The use of this inexpensive and easily available catalyst makes this protocol practical, environment friendly and economically attractive.



Scheme 1. Synthesis of Benzimidazole derivatives catalyzed by zinc triflate.

Table 1. Effect of Solvent in the synthesis of 2-(4-Methoxyphenyl) benzimidazole.

Entry	Solvent	Temperature(°C)	Time (hr)	Yield (%) ^a
1	CH ₂ Cl ₂	40	12	58
2	CH ₃ OH	65	10	70
3	CH ₃ CH ₂ OH	80	8	95
4	THF	68	12	62
5	CH ₃ CN	85	10	72

^aAll are isolated yields.

Table 2. various mole ratios of zinc triflate for the synthesis of 2-(4-Methoxyphenyl) benzimidazole.

Entry	Zinc triflate (mol%)	Time (hr)	Yield (%) ^a
1	0	12	18
2	5	8	68
3	10	8	95
4	15	8	95
5	20	8	94





















^aAll are isolated yields.

The simple work-up procedure, high yields of products and nontoxic nature of the catalyst are other advantages of the present method.

3.1. Experimental

All ^1H NMR spectra were recorded on 400 MHz Varian FT-NMR spectrometers. All chemical shifts are given as δ

Table 3. synthesis of 2-substituted benzimidazoles from *O*-Phenylenediamine and aldehydes^a.

Entry	aldehyde	Benzimidazole	Yield ^b
1			94
2			84
3			88
4			89
5			89
6			90
7			90
8			95
9			81
10			95

^aReaction conditions: *o*-phenylenediamine (1 mmol), benzaldehyde (1 mmol), Zn(OTf)₂ (10 mol%) were stirred for 8h under reflux in Ethanol, ^b isolated yields.

value with reference to Tetra methyl silane (TMS) as an internal standard. Products were purified by flash chromatography on 100 - 200 mesh silica gel. The chemicals and solvents were purchased from commercial suppliers either from Aldrich, Spectrochem and they were used without purification prior to use.

3.2. Zinc Triflate Catalyzed Synthesis of 2-Substituted Benzimidazole Derivatives from Aldehydes

A mixture of *o*-phenylenediamine (1 mmol), benzaldehyde (1.0 mmol) and Zn(OTf)₂ (10 mol%) in Ethanol (5 ml) was placed in a 50 ml round bottom flask and stirred at reflux for 8 h. The progress of the reaction was monitored by TLC Hexane: EtOAc (8:2) after completion of the reaction, the reaction mixture was cooled and treated by dilution with EtOAc (20 mL). Total organic layer was washed with water, brine solution and dried over Na₂SO₄ and evaporated under vacuum. Obtained crude residue was purified by column chromatography to give 2-substituted benzimidazoles.

2-Phenylbenzimidazole [27]: Off white solid; m.p: 289°C - 291°C; ¹H NMR (DMSO-d₆): δ13.02 (br s, 1H), 8.20 (d, *J* = 7.6 Hz, 2H), 7.67 - 7.65 (m, 1H), 7.56 - 7.49 (m, 4H), 7.22 - 7.18 (m, 2H); (LC-MS) *m/z*: 195.08 [M + H]⁺; IR (KBr, cm⁻¹): 3420, 2920, 2627, 1623, 1410, 1276, 1119, 970, 738.

2-(2-Chlorophenyl) benzimidazole [28]: Light pink red solid; m.p: 231°C - 233°C; ¹H NMR (DMSO-d₆): δ12.80 (br s, 1H), 7.91 - 0.89 (m, 1H), 7.67 - 7.62 (m, 3H), 7.57 - 7.52 (m, 2H), 7.25 - 7.23 (m, 2H); (LC-MS) *m/z*: 229.04 [M + H]⁺

2-(3-Chlorophenyl) benzimidazole [28]: Colourless solid; m.p: 234°C - 236°C; ¹H NMR (DMSO-d₆): δ13.06 (br s, 1H), 8.40 (s, 1H), 8.27 (d, *J* = 6.8 Hz, 1H), 7.81 - 7.72 (m, 4H), 7.49 - 7.47 (m, 2H); (LC-MS) *m/z*: 229.04 [M + H]⁺

2-(4-Chlorophenyl) benzimidazole [29]: Colour less solid; m.p: 289°C - 291°C; ¹H NMR (DMSO-d₆): δ12.9 (br s, 1H), 8.15 (d, *J* = 8 Hz, 2H), 7.64 - 7.49 (m, 4H), 7.20 (d, *J* = 8 Hz, 2H); (LC-MS) *m/z*: 229.04 [M + H]⁺

2-*o*-tolylbenzimidazole [27]: Colour less solid; m.p: 220°C - 222°C; ¹H NMR (DMSO-d₆): δ13.03 (br s, 1H), 7.82 - 7.79 (m, 3H), 7.60 - 7.58 (m, 1H), 7.56 - 7.45 (m, 4H), 2.58 (s, 3H); (LC-MS) *m/z*: 209.10 [M + H]⁺

2-*p*-tolylbenzimidazole [27]: Colourless solid; m.p: 265°C - 267°C; ¹H NMR (DMSO-d₆): δ12.81 (br s, 1H), 8.06 (d, *J* = 8 Hz, 2H), 7.56 (m, 2H), 7.36 (d, *J* = 8 Hz, 2H), 7.19 (m, 2H), 2.38 (s, 3H); (LC-MS) *m/z*: 209.10 [M + H]⁺

2-(2-Methoxyphenyl) benzimidazole [30]: Colourless solid; m.p: 173°C - 175°C; ¹H NMR (DMSO-d₆): δ13.5 (br s, 1H), 8.29 (d, *J* = 7.2 Hz, 1H), 7.76 - 7.74 (m, 2H), 7.63 - 7.59 (m, 1H), 7.39 - 7.32 (m, 3H), 7.22 - 7.18 (m, 1H), 4.06 (s, 3H); (LC-MS) *m/z*: 225.07 [M + H]⁺

2-(4-Methoxyphenyl) benzimidazole [27]: Colourless solid; m.p: 218°C - 221°C; ¹H NMR (DMSO-d₆): δ12.90 (br s, 1H), 8.21 (d, *J* = 8.4 Hz, 2H), 7.70 - 7.68 (m, 2H), 7.38 - 7.36 (m, 2H), 7.21 (d, *J* = 8.8 Hz, 2H), 3.88 (s, 3H); (LC-MS) *m/z*: 225.07 [M + H]⁺

2-(3-nitrophenyl) benzimidazole [29]: Off-white so-

lid; m.p: 203°C - 205°C; ¹H NMR (DMSO-d₆): δ13.2 (br s, 1H), 9.02 (s, 1H), 8.60 (d, *J* = 7.6 Hz, 1H), 8.33 (d, *J* = 7.9 Hz, 1H), 7.85 (t, *J* = 7.9 Hz, 1H), 7.7 - 7.52 (m, 2H), 7.25 (t, *J* = 6.8 Hz, 2H); (LC-MS) *m/z*: 240.06 [M + H]⁺

2-benzylbenzimidazole [27]: Off white solid; m.p: 177 - 179°C; ¹H NMR (DMSO-d₆): δ13.0 (br s, 1H), 7.52 - 7.50 (m, 2H), 7.34 - 7.16 (m, 7H), 4.21 (s, 2H); (LC-MS) *m/z*: 209.10 [M + H]⁺

Acknowledgements

The authors are very much grateful to the management of Chalapathi Institute of Engineering and Technology, Guntur, A.P, India, for providing moral support in carrying out this work.

REFERENCES

- [1] H. Zarrinmayeh, A. M. Nunes, P. L. Ornstein, D. M. Zimmerman, M. B. Arnold, D. A. Schober, S. L. Gackenheim, R. F. Bruns, P. A. Hipskind, T. C. Britton, B. E. Cantrell and D. R. Gehlert, "Synthesis and Evaluation of a Series of Novel 2-[(4-Chlorophenoxy)methyl]-benzimidazoles as Selective Neuropeptide Y Y1 Receptor Antagonists," *Journal of Medical Chemistry*, Vol. 41, No. 15, 1998, pp. 2709-2719. <http://dx.doi.org/10.1021/jm9706630>
- [2] Z. Zhu, B. Lippa, J. C. Drach and L. B. Towersend, "Design, Synthesis, and Biological Evaluation of Tricyclic Nucleosides (Dimensional Probes) as Analogues of Certain Antiviral Polyhalogenated Benzimidazole Ribonucleosides," *Journal of Medical Chemistry*, Vol. 43, No. 12, 2000, pp. 2430-2437. <http://dx.doi.org/10.1021/jm990290y>
- [3] H. A. Barker, R. D. Smyth, H. Weissbach, J. I. Toohey, J. N. Ladd and B. E. Volcani, "Isolation and Properties of Crystalline Cobamide Coenzymes Containing Benzimidazole or 5,6-Dimethylbenzimidazole," *Journal of Biological Chemistry*, Vol. 235, No. 2, 1960, pp. 480-488.
- [4] J. C. Sih, W. B. Im, A. Robert, D. R. Graber and D. P. Blackmann, "Studies on (H⁺-K⁺)-ATPase Inhibitors of Gastric acid Secretion. Prodrugs of 2-[(2-Pyridinylmethyl)sulfinyl]benzimidazole Proton-Pump Inhibitors," *Journal of Medical Chemistry*, Vol. 34, No. 3, 1991, pp. 1049-1062. <http://dx.doi.org/10.1021/jm00107a026>
- [5] A. Figge, H. J. Altenbach, D. J. Brauer and P. Tielmann, "Synthesis and Resolution of 2-(2-Diphenylphosphinyl-naphthalen-1-yl)-1-isopropyl-1*H*-benzimidazole: A New Atropisomeric *P,N*-Chelating Ligand for Asymmetric Catalysis," *Tetrahedron: Asymmetry*, Vol. 13, No. 2, 2002, pp. 137-144. [http://dx.doi.org/10.1016/S0957-4166\(02\)00079-4](http://dx.doi.org/10.1016/S0957-4166(02)00079-4)
- [6] E. Alcalde, I. Dinares, L. P. Garcia and T. Roca, "An Advantageous Synthesis of 2-Substituted Benzimidazoles Using Polyphosphoric Acid. 2-(Pyridyl)-1*H*-benzimidazoles, 1-Alkyl-(1*H*-benzimidazol-2-yl)pyridinium Salts, Their Homologues and Vinyllogues," *Synthesis*, Vol. 1992, No. 4, 1992, pp. 395-398. <http://dx.doi.org/10.1055/s-1992-26121>
- [7] V. S. Padalkar, V. D. Gupta, K. R. Phatangare, V. S. Patil, P. G. Umape and N. Sekar, "Indion-190 Resin: Efficient, Environmentally Friendly and Reusable Catalyst for Synthesis of Benzimidazoles, Benzoxazoles and Benzothiazoles," *Green Chemistry Letters and Reviews*, Vol. 5, No. 2, 2012, pp. 139-145. <http://dx.doi.org/10.1080/17518253.2011.585666>
- [8] R. R. Nagawade and D. B. Shinde, "BF₃.OEt₂ Promoted Solvent Free Synthesis of Benzimidazole Derivatives," *Chinese Chemical Letters*, Vol. 17, No. 4, 2006, pp. 453-456.
- [9] M. Kidwai, A. Jahan and D. Bhatnagar, "Polyethylene Glycol: A Recyclable Solvent System for the Synthesis of Benzimidazole Derivatives Using CAN as Catalyst," *Journal of Chemical Sciences*, Vol. 122, No. 4, 2012, pp. 607-612. <http://dx.doi.org/10.1007/s12039-010-0095-7>
- [10] P. Sun and Z. Hu, "The Convenient Synthesis of Benzimidazole Derivatives Catalyzed by I₂ in Aqueous Media," *Journal of Heterocyclic Chemistry*, Vol. 43, No. 3, 2006, pp. 773-775. <http://dx.doi.org/10.1002/jhet.5570430338>
- [11] B. Sadeghi and M. G. Nejad, "Silica Sulfuric Acid: An Eco Friendly and Reusable Catalyst for Synthesis of Benzimidazole Derivatives," *Journal of Chemistry*, Vol. 2013, 2013, Article ID: 581465.
- [12] R. Trivedi, S. K. De and R. A. Gibbs, "A Convenient One-Pot Synthesis of 2-Substituted Benzimidazoles," *Journal of Molecular Catalysis A: Chemical*, Vol. 245, No. 1-2, 2006, pp. 8-11. <http://dx.doi.org/10.1016/j.molcata.2005.09.025>
- [13] R. G. Jacob, L. G. Dutra, C. S. Radatz, S. R. Mendes, G. Perin and E. Lenardao, "Synthesis of 1,2-Disubstituted Benzimidazoles Using SiO₂/ZnCl₂," *Tetrahedron Letters*, Vol. 50, No. 13, 2009, pp. 1495-1497. <http://dx.doi.org/10.1016/j.tetlet.2009.01.076>
- [14] R. K. Kumar, P. V. V. Satyanarayana and S. B. Reddy, "NaHSO₄-SiO₂ Promoted Synthesis of Benzimidazole Derivatives," *Archives of Applied Science Research*, Vol. 4, No. 3, 2012, pp. 1517-1521.
- [15] C. Mukhopadhyay and P. K. Tapaswi, "PEG-Mediated Catalyst-Free Expedient Synthesis of 2-Substituted Benzimidazoles and Bis-Benzimidazoles under Solvent-Less Conditions," *Tetrahedron Letters*, Vol. 49, No. 43, 2008, pp. 6237-6240. <http://dx.doi.org/10.1016/j.tetlet.2008.08.041>
- [16] K. Bahrami, M. M. Khodaei and F. Naali, "H₂O₂/Fe(NO₃)₃-Promoted Synthesis of 2-Arylbenzimidazoles and 2-Arylbenzothiazoles" *Synlett*, Vol. 2009, No. 4, 2009, pp. 569-572. <http://dx.doi.org/10.1055/s-0028-1087911>
- [17] L. Wang, J. Sheng, H. Tian, *et al.*, "An Efficient Procedure for the Synthesis of Benzimidazole Derivatives Using Yb(OTf)₃ as Catalyst Under Solvent-Free Conditions," *Synthetic Communications*, Vol. 34, No. 23, 2004, pp. 4265-4272. <http://dx.doi.org/10.1081/SCC-200039340>
- [18] A. Loupy, A. Petit, J. Hamelin, *et al.*, "New Solvent-Free Organic Synthesis Using Focused Microwaves," *Synthesis*, Vol. 1998, No. 9, 1998, pp. 1213-1234. <http://dx.doi.org/10.1055/s-1998-6083>
- [19] G. V. Reddy, V. V. V. N. S. Ramarao, B. Narsaiah, *et al.*,

- "A Simple and Efficient Method for the Synthesis of Novel Trifluoromethyl Benzimidazoles under Microwave Irradiation Conditions," *Synthetic Communications*, Vol. 32, No. 16, 2002, pp. 2467-2476.
<http://dx.doi.org/10.1081/SCC-120003394>
- [20] G. Penieres, I. Bonifas, G. Lopez, *et al.*, "Synthesis of Benzimidazoles in Dry Medium," *Synthetic Communications*, Vol. 30, No. 12, 2000, pp. 2191-2195.
<http://dx.doi.org/10.1080/00397910008087397>
- [21] K. Bougrin, A. Loupy, A. Petit, *et al.*, "Nouvelle Voie de Synthèse des 2-Trifluorométhylarylimidazoles sur Montmorillonite K10 en 'Milieu Sec' Sous Micro-Onde," *Tetrahedron*, Vol. 57, No. 1, 2001, pp. 163-168.
[http://dx.doi.org/10.1016/S0040-4020\(00\)00992-3](http://dx.doi.org/10.1016/S0040-4020(00)00992-3)
- [22] S. R. Kumar, Y. Venkateswarlu and P. Leelavathi, "Synthesis of 1,5-Benzodiazepines Catalyzed by Zinc Triflate in Solvent-Free Medium," *Asian Journal of Chemistry*, Vol. 23, No. 4, 2011, pp. 1611-1614.
- [23] X. Zhu and A. Ganesan, "Regioselective Synthesis of 3-Alkylindoles Mediated by Zinc Triflate," *The Journal of Organic Chemistry*, Vol. 67, No. 8, 2002, pp. 2705-2708.
<http://dx.doi.org/10.1021/jo010996b>
- [24] K. C. Lekhok, D. Bhuyan, D. Prajapati and R. C. Boruah, "Zinc Triflate: A Highly Efficient Reusable Catalyst in the Synthesis of Functionalized Quinolines via Friedlander Annulations," *Molecular Diversity*, Vol. 14, No. 4, 2010, pp. 841-846.
<http://dx.doi.org/10.1007/s11030-009-9214-0>
- [25] V. N. Kumar, G. Sridhar, E. Laxminarayana, B. Mallaiah and T. Giri, "Selective Deprotection of *t*-Butyldimethylsilyl (TBDMS) Ethers Using Zinc (II) Trifluoromethanesulfonate," *Journal of Drug Chemistry (IJPI)*, Vol. 2, No. 10, 2012, 4p.
- [26] D. Atukuri, R. K. Ravindra and P. K. Pramod, "Zinc Triflate Catalyzed Facile Synthesis of Novel 1,2,4-Triazolinone Derivatives Using 3-Arylsydnone as Synthons," *Letters in Organic Chemistry*, Vol. 10, No. 7, 2013, pp. 510-517.
<http://dx.doi.org/10.2174/1570178611310070009>
- [27] A. J. Blacker, M. M. Farah, M. I. Hall, S. P. Marsden, O. Saidi and J. M. J. Williams, "Synthesis of Benzazoles by Hydrogen-Transfer Catalysis," *Organic Letters*, Vol. 11, No. 9, 2009, pp. 2039-2042.
<http://dx.doi.org/10.1021/ol900557u>
- [28] S. B. Sapakal, K. F. Shelke, S. S. Sonar, B. B. Shingate and M. S. Shingare, "Acidic Ionic Liquid Catalyzed Environmentally Friendly Synthesis of Benzimidazole Derivatives," *Bulletin of the Catalysis Society of India*, Vol. 2, 2009, p. 78.
- [29] V. A. Sontakke, S. Ghosh, P. P. Lawande, B. A. Choa-pade and V. S. Shinde, "A Simple, Efficient Synthesis of 2-Aryl Benzimidazoles Using Silica Supported Periodic Acid Catalyst and Evaluation of Anticancer Activity," *ISRN Organic Chemistry*, Vol. 2013, 2013, Article ID: 453682.
- [30] J. Peng, M. Ye, C. Zong, F. Hu, L. Feng, X. Wang and C. Chen, "Copper-Catalyzed Intramolecular C-N Bond Formation: A Straightforward Synthesis of Benzimidazole Derivatives in Water," *The Journal of Organic Chemistry*, Vol. 76, No. 2, pp. 716-719.
<http://dx.doi.org/10.1021/jo1021426>