

**Facile Synthesis of Open Mesoporous Carbon Nanofibers with Tailored Nanostructure as a Highly Efficient Counter Electrode in CdSe Quantum-Dot-Sensitized Solar Cells†**

Baizeng Fang,<sup>a,b,c</sup> Minwoo Kim,<sup>a</sup> Sheng-Qiang Fan,<sup>a,d</sup> Jung Ho Kim,<sup>a</sup>

David P. Wilkinson,<sup>\*,b,c</sup> Jaejung Ko,<sup>\*,a</sup> and Jong-Sung Yu<sup>\*,a</sup>

<sup>a</sup> *Department of Advanced Materials Chemistry, BK21 Research Team, Korea University, 208 Seochang, Jochiwon, ChungNam 339-700, Republic of Korea,*

<sup>b</sup> *Department of Chemical & Biological Engineering, University of British Columbia,*

*2360 East Mall, Vancouver, B.C., Canada V6T 1Z3, <sup>c</sup> Clean Energy Research Centre,*

*2360 East Mall, Vancouver, B.C., Canada V6T 1Z3, and <sup>d</sup> Department of Applied*


*Chemistry, School of Science, Xi'an Jiaotong University, Xi'an, Shaanxi 710049,*

*People's Republic of China*


**Supporting Information**

---

\*To whom correspondence should be addressed. E-mail: [jsyu212@korea.ac.kr](mailto:jsyu212@korea.ac.kr) (J.-S. Yu); [jko@korea.ac.kr](mailto:jko@korea.ac.kr) (J. Ko); [dwilkinson@chbe.ubc.ca](mailto:dwilkinson@chbe.ubc.ca) (D. P. Wilkinson)



**Fig. S1.** Representative SEM images with various magnifications for the MCNFs.



**Fig. S2.** Representative SEM image for the activated carbon (Duksan Pharm. Co. Korea).




Fig. S3. Typical nitrogen adsorption-desorption isotherm at 77 K and the derived PSD for the activated carbon (Duksan Pharm. Co. Korea).