



Fallacia californica sp. nov. (Bacillariophyta), a new freshwater diatom species from streams in California, USA

ROSALINA STANCHEVA^{1*} & KALINA M. MANOYLOV²

¹Department of Biological Sciences, California State University San Marcos, San Marcos, California 92096, USA.

²Department of Biological and Environmental Sciences, Georgia College and State University, Milledgeville, GA 31061 USA

*Corresponding author (E-mail: rhrstov@csusm.edu)

Abstract

Small naviculoid diatoms remain largely taxonomically unexplored in the United States, due to the inability for clear differentiation of unique characteristics under light microscope magnification during routine identification. In a recent study of benthic stream diatoms from California and as a part of taxonomy data quality assurance and quality control process it became evident that taxonomic reevaluation of a common *Fallacia* species is needed. A new small-celled *Fallacia* species is described here and compared to similar taxa. Nomenclatural and ecological discussions, based on historical and current literature, are presented for each taxon. Type materials were consulted for taxonomic evaluation and comparison. The new taxon is described with light microscopy and scanning electron microscopy. *Fallacia californica* could be distinguished from other small *Fallacia* species with coarse areolae by the unique combination of following features: 1) distinct lyre-shaped sternum, 2) asymmetrical central area, and 3) variable central striae with one to four inner areolae absent, and unilateral gap between the central striae on the mantle. The new freshwater species was distributed in warm alkaline waters with medium conductivity and elevated nutrients content. The majority of populations have been recorded in coastal streams, where they could be abundant, but the species could be scattered in other inland running waters as well.

Keywords: *Fallacia*, Bacillariophyta, diatom, streams, California, new species, scanning electron microscopy

Introduction

Small naviculoid diatoms are widely distributed in periphyton, can vary in abundance, and provide important information towards the ecological assessment of each habitat (Van Dam *et al.* 1994, Wetzel *et al.* 2015). Taxonomic evaluation using high quality research microscopes, while estimating relative abundance of diatom taxa within site, remains a necessary step in bioassessment (Manoylov 2014). The taxonomic composition of the freshwater algae in California, particularly from streams and rivers has been reported as part of national programs and state surveys. In the last decade, the extensive bioassessment sampling by the Surface Water Ambient Monitoring Program (SWAMP) of the California Water Resource Control Board allowed a more complete picture of the local diatom flora from streams. Taxonomic analysis of the data led to the description of new-to-science species belonging to the diatom genera *Amphora* Ehr. ex Kütz., *Halamphora* (Cleve) Levkov, *Rhoicosphenia* Grunow, *Gomphonema* Grunow, and *Gomphoneis* Cleve (Stepanek & Kociolek 2013, Thomas & Kociolek 2015, Stancheva *et al.* 2016), with more in progress. A recent diatom quality assurance and control process, similar to the USEPA (2013) procedures, indicated that taxonomic reevaluation of a common, and often locally abundant *Fallacia* species is needed.

The diatom genus *Fallacia* Stickle & D.G. Mann was described in Round *et al.* (1990) with transfer of 59 species from *Navicula* Bory. The unique combination of characters combine H-shaped chloroplast, lateral depressed sterna interrupting the striae, and a finely porous conopeum (Li *et al.* 2014b; Spaulding *et al.* 2010). Most *Fallacia* taxa are characteristic of marine and brackish benthic habitats, where they often form rich epipsammic communities in coastal and estuarine intertidal areas (Witkowski 1993, Sabbe *et al.* 1999, Witkowski *et al.* 2000). Currently, there are 89 *Fallacia* species recognized (Guiry & Guiry 2018). Taxa ascribed to the genus *Fallacia* have been increasing recently with transfer and description of several new species (e.g., Sabbe *et al.* 1999, Witkowski 1991, Procopiak & Fernandes 2003, Metzeltin *et al.* 2005, Li *et al.* 2015, Van De Vijver & Cox 2015, Genkal & Yarushina 2017),

but the morphological variation and diversity of the freshwater species remain poorly understood. Taxonomic and autecological characteristics of intertidal marine and brackish species from this genus were analyzed in depth (i.e., Sabbe *et al.* 1999). The species diversity of *Fallacia* in freshwater habitats is much lower, for instance, 17 species were reported from central Europe (Cantonati *et al.* 2017), the same number of species from Uruguay (Metzeltin *et al.* 2005), 15 species presented from Baikal with 47% of those described as new to sciences (Kulikovskiy *et al.* 2012), and 23 species are known from streams in California (SWAMP Master List).

The most prominent morphological feature of the genus *Fallacia* is the lyre-shaped hyaline marking (sternum, canal), located on either side of the raphe. The markings represent hollow canals which are formed between the depressed sterna and the finely porous conopeum, which covers the valve surface (Liu *et al.* 2012). Striae are uniseriate, interrupted by lateral sterna (rib) internally and composed by round internally-closed areolae (Liu *et al.* 2012) (see Fig. 1). Sabbe *et al.* (1999) separated three complex morphological clusters within *Fallacia*: 1) species with predominantly marginal striae; 2) species with distinctly panduriform lyre, and 3) species complex around *Fallacia tenera* (Hust.) D.G. Mann, characterized by striae split to two coarse areolae by longitudinal hyaline rib.

Recently, Liu *et al.* (2012) separated several species into the new genus *Pseudofallacia* Y. Liu, J. P. Kociolek & Q. X. Wang, based on the presence of a longitudinal rib on either side of the axial area, which interrupts the striae internally, in contrast to the lyre-shaped canal in *Fallacia*. Also, the central area in *Pseudofallacia* is large, expanded transversely to the valve margins, in contrast to the small central area in *Fallacia*. The striae in *Pseudofallacia* have one elongated, externally closed areola.

The common freshwater species *F. tenera* was transferred to the new genus *Pseudofallacia* by Liu *et al.* (2012). This new taxonomic position was accepted in several publications (Ector *et al.* 2015, Genkal & Yarushina 2017), but some recent authors used the genus name *Fallacia* (i.e. Li *et al.* 2014, Miscoe *et al.* 2017). Li *et al.* (2014) provided ultrastructural evidence that *F. tenera* does possess the typical characteristics of *Fallacia*, the most important of which is the lyre-shaped canal, which does not have areolae in it, as illustrated for *Pseudofallacia oculata* Krasske (Liu *et al.* 2012, Fig. 15). The complicated taxonomy and nomenclature of *F. tenera* are discussed in detail by Li *et al.* (2014) and further in this paper, because it is closely related to the diatom species presented here. As part of the required taxonomic reconciliation process (USEPA, 2013) it became evident that all taxonomists recognized a potentially new taxon, which combines the coarse areolae, characteristic for *Fallacia* cf. *tenera* with distinct panduriform lyre-shaped markings and dense areolae, typical for *Fallacia* cf. *cryptolyra* (Brockmann) Stickle & D.G. Mann.

The goal of this study was to describe the most common, and often locally abundant *Fallacia* taxon in coastal streams in California. The new species is illustrated with light microscopy (LM) and scanning electron microscopy (SEM) and compared with closely related taxa. Distributional and ecological data from stream locations in California are also presented.

Material and methods

For this study we used diatom samples obtained between 2013 and 2017 as part of the SWAMP program in California. The diatom samples were collected along with environmental variables using the multihabitat sampling protocol (Fetscher *et al.* 2009). For LM observations, the preserved diatom samples were cleaned by the hydrogen peroxide method, mounted in Naphrax®, and 600 diatom valves were identified to species level and counted (for details see Stancheva *et al.* 2015). The original analysis of the samples was performed at the California State University San Marcos and EcoAnalysts Inc. For this study, diatom slides were scanned additionally after diatom enumeration, in order to obtain a size ranges a minimum of 50 *Fallacia* valves. LM imaging of the specimens was performed using a Leica DM2500 microscope with differential interference contrast optics and Leica DFC295 Camera attached to the microscope (Leica Microsystems, Wetzlar, Germany) at the Georgia College and State University.

SEM was performed with cleaned specimens air dried onto cover glasses, attached to aluminum stubs, sputter-coated with 5 nm of iridium or platinum. Initial observations of frustular ultrastructure were performed with JEOL SEM (JSM-IT100) at Georgia College and State University. The diatom material was examined and pictured in high vacuum mode using a Zeiss SIGMA 500 (Carl Zeiss Microscopy, Thornwood, NY, USA) with an accelerating voltage of 5 kV at the Nano3 Facility at the University of California, San Diego. Holotype slides and material are deposited at the Georgia College and State Natural History University Museum Algae Collection, diatom collection and isotype slides and type material are maintained by the author Rosalina Stancheva at the California State University San Marcos.

In addition, we observed three slides from Hustedt collection at the Friedrich Hustedt Zentrum für

Diatomeenforschung Institute for Polar and Marine Research: 1) Slide N8/50 *Navicula tenera* Hust., R. S. Java, Ranu Klindungan Lake 1, 1930. St., 2) Slide 230/70 R. M. Kamerun Lagune 1b, 1939. H.2. (for *Navicula auriculata* Hust.), and 3) Slide 397/94, R. M. Arcachon, 1960. H. (for *Navicula dissipata* Hust.). Terminology used followed Anonymous (1975), Ross *et al.* (1979), and Round *et al.* (1990).

Results

Fallacia californica Stancheva & Manoylov, sp. nov. (Figs. 1–38)





FIGURE 1. SEM image of *Fallacia californica* Stancheva & Manoylov, sp. nov. with indication of main structural elements of the valve as follow: stars—lyre-shaped sternum; dots—hyaline rib interrupting the striae; arrows—areolae adjacent to the raphe branches; diamonds—inner areolae; arrowheads—areolae near the mantle. Scale bar 1 μm .

Description


Valves are elliptical with rounded poles, 4–5.6 μm wide, 6–14.5 μm long (Figs 2–20). In girdle view (Fig. 9) ends look rounded too. The raphe is filiform and slightly arched toward the primary side of the valve. Terminal raphe fissures curved toward the secondary side of the valve. The striae are parallel in the center, radiate near the poles, 21–10 μm , composed of coarse areolae resolvable with LM. The lyre-shaped sternum distinctly panduriform, constricted in the center (Figs 3, 4, 10, 16, 17, 19), often clearly visible in the smallest valves (Fig. 15). Striae longitudinally interrupted by a hyaline rib which split the striae into two or three areolae of different sizes. The hyaline rib is often enlarged to fan-shaped near the valve center (Figs 2, 7, 8, 10, 15, 18). The areolae closer to the mantle are larger, transapically elongated, one per stria, uniform and equally spaced through the valve; occasionally the central two areolae are slightly distant and form unilateral gap (Figs 2–6, 8, 10, 12, 15–19). The inner areolae are smaller, one or two per stria, forming a longitudinal line which is curved inwards near the center (Figs 7, 10, 16, 18). The inner areolae are asymmetrical on both sides of the central area, smaller, variable in size and position, weakly developed, widely spaced or absent, or sometimes longer than the other areolae (Figs 10, 20). The central area is asymmetrical, small or transapically expanded to unilateral fascia in some specimens. Areolae adjacent to the raphe branches are well developed, single or double, forming equally long lines on both sides of the raphe (Figs 7, 10, 15, 16, 19).

SEM observations

Internally, small helictoglossae are visible at the polar terminals of the raphe, slightly curved to the primary side of the valve (Figs 21, 24). On both sides of the raphe, depressed lyre-shaped lateral sternum is interrupting the striae



FIGURES 2–20. *Fallacia californica* Stancheva & Manoylov, *sp. nov.* Figs. 2–15, type material from Aliso Creek, California, USA (site 901M14126), Fig. 9 girdle view. Figs 16–20 from Soquel Creek (site 304PS0338). Note the distinct panduriform lyre sternum (Figs 2–4, 15); the hyaline rib, which is enlarged to fan-shaped near the valve center (Figs 2, 7, 10, 18) and enlarged central area on the opposite side (Figs 2, 5, 10, 18). LM images, scale bar 10 μ m.




FIGURES 21–32. *Fallacia californica* Stancheva & Manoylov, *sp. nov.* type material from Aliso Creek, California, USA and from Soquel Creek (Figs 26–28). Figs 21–26 internal valve view. Figs 27–32 external valve view. Arrowheads show the gap between central striae. SEM images, scale bars 1 μm .

(Figs 21–25, 35). The striae outside the sternum are crossed by longitudinal hyaline rib, which does not appear to be depressed (Figs 22–26). A longitudinal row with transapically elongated large areolae is positioned on the valve mantle (Figs 21–25). These areolae are equidistant, except for the two central areolae on the primary side, which are separated by larger gap (Figs 24, 30–32). The inner areolae are positioned on the valve face. They are smaller and vary from elongated in a single row to circular in a single or double row near the center of the valve (Figs 21–25). One to four central striae on the primary side are absent (Figs 24, 25). The striae between the raphe and lyre-shaped sternums are transapically elongated, constricted, in a single row, or circular in pairs (Figs 22, 23, 35). The rows with areolae on both sides of the raphe branches are relatively symmetrical. Internally, the areolae are occluded by thin hymenes, which are often dissolved, but partially visible on Figs 24, 25, 33.

Externally, the areolae on the valve face are covered by a finely porous conopeum from the raphe sternum to the valve mantle (Figs 27–32). The edge of mantle possesses finger-like protrusions, each with two to four “pegs”, which fasten the connection of the conopeum and the mantle (Figs 30, 31, 36). The central 4–6 finger-like mantle protrusions

are longer and more pronounced than the other (Figs 29, 31). The images with broken conopeum (Figs 34, 37) show that the lyre-shaped lateral depressed sterna and the finely porous conopeum comprised two longitudinal canals that connect with the exterior by pores on both sides of terminal raphe fissure. The cingulum is composed of at least one wide open girdle band (Figs 28, 32, 37, 38).

Type: USA. California: Aliso Creek, Orange County, 33.54743949, -117.72230590, *Jim Mann, May 17, 2016* (holotype GCP4211 circled specimen on the slide; isotype RS! 015, circled specimen on slide and material, CSUSM, USA).



FIGURES 33–38. *Fallacia californica* Stancheva & Manoylov, *sp. nov.* type material from Aliso Creek, California, USA and from Soquel Creek (Figs 34, 37). Figs 33, 35 internal valve view. Figs 34, 36–38 external valve view. Fig. 33 shows areolae with remnants from thin hymens. Figs 34, 37 eroded valves with degraded conopeum show the canal of lyre-shaped sternum. Fig. 35 internal valve showing distinctly depressed lyre-shaped sternum. Fig. 36 finger-like protrusions, each with two to four “pegs”, which fasten the connection of the conopeum and the mantle. Fig. 38 shows girdle band. SEM images, scale bars 1 μm .

Etymology:—The epithet refers to the state of California, USA, where the species was first observed.

Distribution and ecological notes:—Distributed in twenty one coastal streams in central and southern California and in Rock Creek located at the foothills of the Sierra Nevada Mts. *F. californica* was locally abundant in seven coastal stream sites with relative abundance ranging from 6.2 to 16.6% in diatom counts. The salinity in *F. californica* habitats was below 0.5 ppt, except for the type locality with elevated salinity 1.8 ppt. Specific conductivity was 39–1567 $\mu\text{S cm}^{-1}$ and 3434 $\mu\text{S cm}^{-1}$ in the type locality, pH range was 7.1–8.6. Total nitrogen concentrations ranged from 0.1 to 2.2 mg L^{-1} , and total phosphorus from 0.06 to 0.16 mg L^{-1} . The distribution data and environmental variables are presented in Table 1.

TABLE 1. Stream locality data and environmental variables in the habitats of *Fallacia californica* Stancheva & Manoylov, sp. nov. NA – not available.

Site ID	901MI4126	304PS0338	304WDCAH1	204PS0358	201LAG140	201LAG150	201LAG165	201LAG210	201LAG241	201LAG280	201WLK098	201WLK120
Sapling date	5/17/2016	5/17/2016	5/17/2016	5/23/2016	09-Jun-16	01-Jun-16	14-Jun-16	07-Jun-16	06-Jun-16	31-May-16	24-May-16	23-May-16
<i>Fallacia californica</i> relative abundance (%)	12.2	16.7	6.3	9.5	4.5	0.2	3.2	1.2	2.3	9.3	4.0	0.3
Stream name	Aliso Creek	Soquel Creek	Waddell Creek	Crow Creek	Nicasio Creek	Halleck Creek	Lagunitas Creek	Lagunitas Creek	San Geronimo Creek	San Geronimo Creek	Walker Creek	Chileno Creek
Latitude	33.54743	36.99348236	37.113717	37.69543831	38.07034	38.06669	38.05839	38.01861	38.008307	38.013003	38.215233	38.20764
Longitude	-117.28263	-121.955562	-122.26978	-122.058428	-122.77051	-122.70264	-122.76522	-122.73306	-122.703405	-122.660755	-122.864597	-122.79456
Alkalinity as CaCO ₃ (mg L ⁻¹)	288	190	95	350	85.6	73.1	80.2	74.3	170	213	59.8	147
Ammonia as N (mg L ⁻¹)	2.18	0.0194	0.0098	0.0068	0.0166	0.0058	0.0086	0.0042	0.0064	0.0057	0.0153	0.0269
Ash Free Dry Mass (mg cm ⁻²)	NA	6.08	6.61	6.51	NA	NA	NA	NA	NA	NA	NA	NA
Chloride (mg L ⁻¹)	486.2	48	21.5	66.5	11.1	20.7	10.3	9.48	21.2	21.8	15.5	29.1
Chlorophyll a (µ cm ⁻²)	NA	20.3	17.3	21.3	1.03	0.49	0.11	0.36	0.31	1.22	0.93	1.78
Dissolved Organic Carbon (mg L ⁻¹)	NA	2.31	5.08	5.6	2.93	2.08	2.90	2.06	1.29	2.27	6.11	12.1
Hardness as CaCO ₃ (mg L ⁻¹)	1182.3	108	99.4	400	93.7	89.7	88.2	81.4	197	236	68.4	165
Nitrate + Nitrite as N (mg L ⁻¹)	1.39	0.0102	0.147	0.2	0.0248	0.0433	0.0382	0.0644	0.277	0.230	0.159	0.292
Nitrogen, Total (mg L ⁻¹)	NA	0.109	0.219	0.494	0.153	0.170	0.165	0.162	0.427	0.370	0.525	1.26
OrthoPhosphate as P (mg L ⁻¹)	0.418	0.112	0.0668	0.148	0.0239	0.0185	0.0233	0.0167	0.0454	0.0797	0.0609	0.158
Oxygen, Dissolved (mg L ⁻¹)	8.29	9	9.2	10.7	8.43	9.67	9.05	10.75	9.52	6.41	9.31	5.74
pH	7.85	8.4	7.9	8.6	7.47	8.07	7.82	8.81	7.94	7.38	7.65	7.21
Phosphorus as P (mg L ⁻¹)	0.349	0.0875	0.0623	0.156	0.0301	0.0204	0.0156	0.0160	0.0402	0.0700	0.0728	0.251
Salinity (ppt)	1.8	0.4	0.2	0.4	0.10	0.12	0.10	0.09	0.20	0.23	0.09	0.19
Specific Conductivity (µS cm ⁻¹)	3434	830	330	883	204.1	246.3	200.7	186.4	407.6	473.7	188.1	390.7
Sulfate (mg L ⁻¹)	1026.5	155	37.4	103	NA	NA	NA	NA	NA	NA	NA	NA
Suspended solids (mg L ⁻¹)	2.8	0.4	0.7	8.1	23.8	NA	NA	13.1	15.4	3.5	12.8	2.9
Temperature (°C)	19.4	14.7	14.9	15.3	14.969	16.95	14.014	12.819	15.290	14.810	13.842	14.57

NA - not available

...continued on the next page


TABLE 1. (Continued)

Site ID	201WLK160	201WLK178	201WLK230	206PET400	304GZR	308SBCAHI	404M07360	404M07361	403FCA022	517FCA222	535CR1051
Sapling date	18-May-16	16-May-16	25-May-16	10-May-16	6/5/2017	18/May/2016	6/8/2017	6/6/2017	5/29/2013	7/10/2013	6/25/2014
<i>Fallacia californica</i> relative abundance (%)	6.2	3.8	0.3	0.2	0.8	0.3	0.7	16.3	0.3	3.0	0
Stream name	Walker Creek	Walker Creek	Arroyo Sausal	Lichau Creek	Gazos Creek	Soberanes Creek	Medea Creek	Topanga Canyon	Santa Clara River	Rock Creek	Tuolumne River
Latitude	38.17545	38.163658	38.14342	38.29444	37.175211	36.455100	34.15478	34.10139	34.354720	39.300000	37.5929
Longitude	-122.82044	-122.783144	-122.72472	-122.66629	-122.353366	-121.922610	-118.75924	-118.62114	-119.022500	-120.968890	-121.061
Alkalinity as CaCO ₃ (mg L ⁻¹)	50.8	47.7	83.4	172	120	125	NA	NA	NA	NA	NA
Ammonia as N (mg L ⁻¹)	0.0160	0.0113	0.0050	0.0340	NA	0.009	NA	NA	0.12	NA	NA
Ash Free Dry Mass (mg cm ⁻²)	NA	NA	NA	NA	NA	18.4	NA	NA	7.49	2.27	NA
Chloride (mg L ⁻¹)	13.2	13.0	31.7	60.0	NA	40.4	NA	NA	80	1.4	NA
Chlorophyll a (µ cm ⁻²)	1.14	1.48	3.25	5.94	NA	5.2	NA	NA	40.7	5.5	NA
Dissolved Organic Carbon (mg L ⁻¹)	6.66	7.75	5.52	8.45	NA	2.8	NA	NA	1.9	NA	NA
Hardness as CaCO ₃ (mg L ⁻¹)	56.6	54.9	94.9	272	NA	102	NA	NA	764	12.3	NA
Nitrate + Nitrite as N (mg L ⁻¹)	0.265	0.289	0.0022	0.164	NA	0.05	NA	NA	2.1	0.05	NA
Nitrogen, Total (mg L ⁻¹)	0.717	0.781	0.326	0.624	NA	0.1	NA	NA	2.25	0.05	NA
OrthoPhosphate as P (mg L ⁻¹)	0.0871	0.105	0.0550	0.274	NA	0.08	NA	NA	0.06	0.04	NA
Oxygen, Dissolved (mg L ⁻¹)	9.51	9.87	9.25	5.77	10.22	10	NA	NA	NA	NA	NA
pH	7.91	7.82	8.66	7.67	7.91	8.4	NA	NA	8.1	7.13	NA
Phosphorus as P (mg L ⁻¹)	0.0989	0.122	0.0963	0.333	NA	0.045	NA	NA	0.027	0.011	NA
Salinity (ppt)	0.08	0.08	0.13	0.36	0.13	0.1	NA	NA	NA	NA	NA
Specific Conductivity (µS cm ⁻¹)	164.4	159.7	277.8	738.4	304.5	168	NA	NA	1567	39	NA
Sulfate (mg L ⁻¹)	NA	NA	NA	NA	NA	9.1	NA	NA	0.36	0.36	NA
Suspended solids (mg L ⁻¹)	NA	NA	21.1	6.7	NA	3.5	NA	NA	1.3	2.4	NA
Temperature (°C)	14.36	12.70	14.582	13.69	15.2	12.9	NA	NA	NA	NA	NA

NA - not available

Comments:—We used two large *Fallacia* populations from southern California (Aliso Creek) and the central coast (Soquel Creek discharging into Monterey Bay) for detailed LM and SEM observations to illustrate the morphological variation in this species. The silica thickness of diatom valves of this species varies significantly depending on different water chemistry, in addition to the valve diminution (see Figs 16–19). The LM appearance of the hyaline lyre in *F. californica* is similar to *F. cryptolyra*. However, the striae in *F. cryptolyra* are much finer and denser (26 to 36 in 10 μm according to Sabbe *et al.* 1999), and are not interrupted by hyaline rib longitudinally. *F. californica* could be distinguished from other small *Fallacia* species with course areolae by the unique combination of following features: 1) distinct panduriform lyre-shaped sternum; 2) asymmetrical central area; and 3) variable central striae with one to four inner areolae absent, and unilateral gap between the central striae on the mantle.

The species with most similar morphology, i.e. *F. tenera*, has distinctly different valve structure, besides the larger valves and the lower number of striae. It is easily distinguished by the uniform striae along the valve margins and particularly near the central area, narrower lanceolate lyre, and a single or incomplete double row of areolae along the raphe. In the studied data set, *F. tenera* was recorded with a single valve in Aliso Creek (Fig. 39) and with several valves in Tuolumne River in the Central Valley (Figs 40–42), showing considerable variation in valve outline and the pattern of areolae adjacent to the raphe branches. A comparison of morphological features of *F. californica* with similar species from *F. tenera* group is presented in Tables 2 and 3.



FIGURES 39–46. *Fallacia tenera* (Hust.) D. G. Mann. Fig. 39 specimen from Aliso Creek, Orange County, California. Figs 40–42 specimens from Tuolumne River, Central Valley, Stanislaus County, California. Figs 43–46 specimens from Ranu Klindungan Lake, Java, Hustedt's slide N8/50 *Navicula tenera*, R. S. Java, Ranu Klindungan Lake 1, 1930. LM images, scale bar 10 μm .

Navicula tenera Hust. 1937 (Figs 43–46)

Type locality: Ranu Klindungan Lake, Java, Indonesia.

Valve are 5–6.5 µm wide, 12–14 µm long with striae 16–18 in 10 µm according to Hustedt (1937). In Hustedt slide N8/50 we observed 17 valves, which corresponded to *N. tenera*. Valves are 4.9–6 µm wide, 12–14 µm long, with 18–20 striae in 10 µm. Marginal striae are uniform along the valve, in some specimens there was a shorter single central stria (Figs 44 and 46). There is a single row of circular areolae on concave side of the raphe branches, and occasionally few isolated areolae on the other side of the raphe (Figs 43, 45, 46).

In Hustedt's slides 230/70 and 397/94 we did not observe diatom valves corresponding to descriptions and size ranges of *Navicula auriculata* and *N. dissipata* (see Table 1).

Discussion

LM species identification within *Fallacia* is based on shape, size and structure of the valve, hyaline lyre, lateral hyaline areas and longitudinal ribs, along with striae structure and density (Sabbe *et al.* 1999). *F. californica* is closely related to species from *F. tenera* group, because of the presence of a longitudinal rib on the valve face outside the lyre, which is splitting each stria to 2–3 coarse areolae. The hyaline lyre in *F. californica* is well developed, depressed, panduriform and distinct under LM even in the smallest specimens. However, in some species of this group, for instance in *F. tenera* and *F. arenaria* Sabbe & Vyverman, there is a tendency for obligate or facultative absence of depression on the lyre-shaped sternum (Sabbe *et al.* 1999). According to Sabbe *et al.* (1999), the distinguishing of the species belonging to *F. tenera* complex is based mainly on the structure, number and position on the valve of the hyaline areas. *F. californica* is distinguished from all close species based on specific appearance of the distinct panduriform lyre, in combination with longitudinal rib typically enlarged towards the center, and unilaterally expanded central area.

The current understanding about the morphological variation of *F. tenera* is unclear (Li *et al.* 2014), since Schoeman & Archibald (1979) lumped freshwater *Navicula tenera* (syn. *N. uniseriata* Hust.) from Ranu Klindungan Lake in Indonesia (Hustedt 1937), with *Navicula insociabilis* var. *dissipatoides* Hust. from the Weser River in Germany (Hustedt 1957) and marine *Navicula auriculata* from Cameroon lagoon (syn. *N. dissipata*, Hustedt 1966). Besides the size and valve outline differences, the freshwater species have unilateral row of areolae near the raphe, in contrast to well developed two rows of areolae on both sides of the raphe in marine species in combination with weakly developed or absent inner puncta of the central striae (see Table 2). This broad concept has been accepted by Krammer & Lange-Bertalot (1986), Sabbe *et al.* (1999), Witkowski *et al.* (2000), Cantonati *et al.* (2017) which resulted in morphologically and ecologically heterogeneous *F. tenera* species complex. Reevaluation of the type materials of the freshwater species of *Fallacia* is needed, in conjunction with ecological, morphological and molecular data, in order to clarify the infrageneric concepts.

In North America, Patrick & Reimer (1966) illustrated *N. tenera* with single row of coarse puncta on concave side of the raphe and reported it for California. Specimens with two rows of puncta on both sides of the raphe were defined as *N. auriculata*. The newly described species *Fallacia californica* shows similarity in striae position and structure to the marine *N. auriculata* by having two rows of puncta bordering the raphe and variable differentiation of the central striae and inner areolae on both sides of the raphe (feature not recognized by Patrick & Reimer 1966), in contrast to the uniform striae in *F. tenera* from our data set (see Figs 39–42). However, *F. californica* differs from both taxa by its smaller valves, denser striae, distinct panduriform lyre and asymmetrical central area (Table 2 and 3). During the routine analysis of samples from California, *F. californica* should be carefully compared with other sporadically reported finely-striated taxa, e.g. *F. cryptolyra*, *F. insociabilis* (Krasske) D.G. Mann, *F. monoculata* (Hust.) D.G. Mann, *F. sublucidula* (Hust.) D.G. Mann, *F. omissa* (Hust.) D.G. Mann. *F. californica* shares some structural similarities also with a small-sized *F. cf. teneroides* (Hust.) D.G. Mann, recorded from sandy sediments in the mouth of the Westerschelde estuary, The Netherlands (Sabbe *et al.* 1999).

The new species can be found in running alkaline waters in California with medium conductivity, elevated nutrient concentrations and salinity below 0.5 ppt, occasionally inhabiting streams with higher electrolyte content. The majority of populations have been recorded in coastal streams, where they could be abundant, but the species could be scattered in inland running waters as well. The dominant taxa in the *F. californica* communities were *Nitzschia inconspicua* Grunow, *N. microcephala* Grunow, *Navicula gregaria* Donkin, *Amphora pediculus* (Kütz.) Grunow, and the epiphytic diatoms *Cocconeis placentula* Ehr., *Planothidium lanceolatum* (Bréb. ex Kütz.) Lange-Bert., *P. frequentissimum* (Lange-Betr.) Lange-Bert., *P. delicatulum* (Kütz.) Round & Bukh., and *Rhoicosphenia abbreviata* (C. Ag.) Lange-Bert.

TABLE 2. Comparison of size range of *Fallacia californica* Stancheva & Manoylov, *sp. nov.* with similar taxa belonging to *Fallacia tenera* group reported in the literature. NA—not available.

Taxon	Width (um)	Length (um)	Striae in 10 um	Locality	Reference	SEM Figure	LM figure
<i>Fallacia californica</i> Stancheva & Manoylov	4–5.6	6–14.5	20–23	Streams in California	This study	Figs. 20–37	Figs. 1–19
<i>Navicula tenera</i> Hust.	5–6.5	12–14	16–18	Ranu Klindungan Lake, Java, Indonesia	Hust. (1937)	NA	Fig. 11 in Shoeman & Archibald (1979)
<i>Navicula auriculata</i> Hust.	6	16	20	Cameroon lagoon, Cameroon, West Coast of Africa, marine	Hust. (1944)	NA	Hust. Diatom Collection EarthCape Database
<i>Navicula dissipata</i> Hust.	6.5	24	16	Nordasot, West Coast of Norway	Hust. (1944)	NA	Figs. 13–15 in Shoeman & Archibald (1979)
<i>Navicula dissipata</i> Hust.	5–7	9–27	14–17	Baltic Sea, Norway, euryhaline, marine	Hust. (1961–1966)	NA	Hust. Diatom Collection EarthCape Database
<i>Fallacia tenera</i> (Hust.) D.G. Mann (as <i>Navicula teneroides</i> Hust.)	6–6.5	11.5–14	17–20	Tidal flat near the entrance of the Yahagi-Furukawa River in Atsumi Bay, Japan	Takano (1990)	1–6	NA
<i>Fallacia tenera</i> (Hust.) D.G. Mann	5.4–8.6	8.5–24.1	17–20	River mouth on Ubara beach, Japan	Li <i>et al.</i> (2014)	Figs. 9–30	Figs. 1–8
<i>Fallacia tenera</i> (Hust.) D.G. Mann	5.7–6.3	11–14	18–20	Rivers in France	Ector <i>et al.</i> (2015)	NA	Figs. 1–25
<i>Fallacia tenera</i> (Hust.) D.G. Mann	5–5.4	9.7–10.6	19–21.5	Westerschelde estuary, The Netherlands	Sabbe <i>et al.</i> (1999)	Figs. 75, 78, 82	Figs. 1–4
<i>Fallacia tenera</i> (Hust.) D.G. Mann	5	13	13	Shallow pool in Makauwahi Cave, Kauai, Hawaii, USA	Miscoe <i>et al.</i> (2017)	NA	Fig. 104
<i>Navicula auriculata</i> Hust.	5–6	13–16	16–20	USA, fresh and brackish water	Patrick & Reimer (1966)	NA	NA
<i>Navicula tenera</i> Hust.	5–7	11–14	16–18	USA, California, fresh and brackish water	Patrick & Reimer (1966)	NA	NA
<i>Navicula tenera</i> Hust.	4–9	9–27	13–22	Rivers, streams and lagoons in South Africa	Shoeman & Archibald (1979)	Figs. 29–42	Figs. 16–28
<i>Navicula tenera</i> Hust.	4–9	9–27	13–22	Central Europe, freshwater, euryhaline	Krammer & Lange-Bertalot (1986)	NA	Figs. 19–23
<i>Fallacia tenera</i> (Hust.) D.G. Mann	4–9	9–27	13–22	Gulf of Gdansk and Mecklenburg Bay, Baltic Sea, brackish water, occasionally in freshwater	Witkowski <i>et al.</i> (2000)	NA	Pl. 71: 52–56
<i>Pseudofallacia tenera</i> (Hust.) Liu, Kociolek & Wang	4–9	9–27	13–22	Central Europe, in freshwater with high electrolyte content	Cantonati <i>et al.</i> (2017)	NA	Figs. 35–38
<i>Navicula teneroides</i> Hust.	4.5	9–12	24	Lago de Maracaibo, Venezuela, brackish water	Hust. (1956)	NA	Hust. Diatom Collection EarthCape Database
<i>Fallacia cf. teneroides</i> (Hust.) D.G. Mann	4.4	8.7–9.4	22–23	Westerschelde estuary, The Netherlands	Sabbe <i>et al.</i> (1999)	Fig. 81	Fig. 5, 6
<i>Fallacia cf. teneroides</i> (Hust.) D.G. Mann	3.5–5	9–12	24	Norwegian Sea, Narvik	Witkowski <i>et al.</i> (2000)	NA	Pl. 71: 58, 59
<i>Fallacia arenaria</i> Sabbe & Vyverman	3.9–5	6.4–9.8	26.5–27.5	Westerschelde estuary, The Netherlands; Blakeney Point (England); German Wadden Sea	Sabbe <i>et al.</i> (1999)	Figs. 76, 77, 79, 80	Figs. 11–18
<i>Fallacia meridionalis</i> Metzeltin, Lange-Bert. & García-Rodríguez	5.5–6.2	11.5–15.5	19–20	Several small rivers that flow into coastal lagoons in Uruguay	Metzeltin <i>et al.</i> (2005)	Pl. 61: 37	Pl. 61: 30–36

TABLE 3. Comparison of *Fallacia californica* Stancheva & Manoylov, *sp. nov.* with similar taxa from *Fallacia tenera* group, based on valve morphology reported in the references listed in Table 2.

Species	Lyre-shaped marking	Longitudinal hyaline rib crossing the striae	Areolae outside the hyaline rib	Areolae between the hyaline rib and the lyre	Central areolae between the hyaline rib and the lyre	Areolae along the raphe branches	Central area
<i>Fallacia californica</i> Stancheva & Manoylov	wide; panduriform	parallel to valve margin; slightly enlarged to fan-shaped near the valve center	one transapically elongated areola per stria; gap between central areolae on primary side	one transapically elongated or two small circular areolae per stria; striae variable; areolae irregularly spaced or absent near the center	areolae on primary side fewer or absent; areolae on secondary side small circular, single or double, irregular, situated closer to central area; rarely fewer	one transapically elongated or two small circular areolae per stria on both sides; rows equal in length and nearly symmetrical	small or broad; asymmetrical transapically expanded towards primary side
<i>Fallacia tenera</i> (Hust.) D.G. Mann	narrow; lanceolate	parallel to valve margin; uniform in size	one transapically elongated areola per stria; striae uniform, equally spaced	one circular or slightly elongated areola per stria; striae uniform, regularly spaced	not differentiated	circular areolae in single row on both sides; row on secondary side shorter with smaller, fewer or absent areolae	broad; asymmetrical
<i>Fallacia auriculata</i> (Hust.) D.G. Mann (Syn. <i>N. dissipata</i> Hust.)	narrow; lanceolate	parallel to valve margin; abruptly enlarged near the valve center on primary side	one transapically elongated areola per stria; striae uniform, equally spaced	one slightly elongated areola per stria; smaller circular weakly developed areolae near the center	areolae on primary side fewer or absent; areolae on secondary side weakly developed	circular areolae in single row on both sides; row on secondary side shorter with fewer areolae	broad; nearly symmetrical; slightly enlarged on primary side
<i>Fallacia teneroides</i> (Hust.) D.G. Mann	narrow; lanceolate	absent or LM indistinguishable	very small, LM indistinguishable	very small, LM indistinguishable	not differentiated	circular areolae in double row on primary side; single row on secondary side; rows equally long on both sides	broad; asymmetrical
<i>Fallacia cf. teneroides</i> (Hust.) D.G. Mann	narrow; lanceolate; slightly constricted	broad; fan-shaped	one transapically elongated areola per stria; striae uniform, equally spaced	one circular areola per stria; striae uniform, equally spaced	circular areolae in a single not interrupted row, slightly curved towards the valve center	circular areolae in single row on both sides; row on secondary side shorter with fewer areolae	absent
<i>Fallacia arenaria</i> Sabbe & Vyverman	not visible	parallel to valve margin; uniform in width	one transapically elongated or two small circular areolae per stria; striae uniform, equally spaced	one circular areola per stria; striae uniform, equally spaced	circular areolae in a single not interrupted row, slightly curved towards the valve center	few circular areolae in single unequal rows on both sides	small; symmetrical
<i>Fallacia meridionalis</i> Metzeltin, Lange-Bert. & García-Rodríguez	narrow; lanceolate	parallel to valve margin; uniform in width	one transapically elongated areola per stria; striae uniform, equally spaced	one slightly elongated areola per stria; striae uniform, equally spaced	not differentiated	elongated areolae in single row on both sides; rows equal in length and nearly symmetrical	broad; symmetrical

Acknowledgements

The California State Water Resources Control Board Surface Water Ambient Monitoring Program (SWAMP) provided funding and data. Dr. Lisa Underwood (EcoAnalysts Inc.) provided diatom distributional data. Dr. Robert Sheath contributed taxonomic discussion and edits of the manuscript. We are grateful to Sarah Olischläger from Alfred-Wegener-Institute Bremerhaven for sending slides from the Hustedt collection. The authors acknowledge the assistance of Ryan Anderson with electron microscopy imaging at the Nano3 Facility at the University of California San Diego. We thank the editor Dr. Evan Thomas and two anonymous reviewers for their valuable comments, which improved the presentation of this paper. KMM SEM work at Georgia College and State University was supported by National Science Foundation grant MIP-1531843.

References

- Anonymous (1975) Proposals for a standardization of diatom terminology and diagnoses. *Nova Hedwigia, Beihefte* 53: 323–354.
- Cantonati, M., Kelly, M.G. & Lange-Bertalot, H. (2017) *Freshwater Benthic Diatoms of Central Europe*. Koeltz Botanical Books, Schmittens-Oberreifenberg, Germany, 942 pp.
- Ector, L., Wetzel, C.E., Novais, M.H. & Guillard, D. (2015) *Atlas des diatomées des rivières des Pays de la Loire et de la Bretagne*. DREAL Pays de la Loire, Nantes.
- Fetscher, A.E., Busse, L. & Ode, P.R. (2009) *Standard Operating Procedures for Collecting Stream Algae Samples and Associated Physical Habitat and Chemical Data for Ambient Bioassessments in California*. California State Water Resources Control Board Surface Water Ambient Monitoring Program (SWAMP) Bioassessment SOP 002. (updated May 2010). Available from: http://www.waterboards.ca.gov/water_issues/programs/swamp/docs/sop_algae.pdf (accessed 19 February 2018)
- Genkal, S.I. & Yarushina, M.I. (2017) The Species of the genus *Fallacia* A.J. Stickle & D.G. Mann (Bacillariophyta) in Russia: morphology, taxonomy, ecology and distribution. *International Journal on Algae* 19: 99–110.
<https://doi.org/10.1615/InterJAlgae.v19.i2.10>
- Guiry, M.D. & Guiry, G.M. (2018) *AlgaeBase*. World-wide electronic publication, National University of Ireland, Galway. Available from: <http://www.algaebase.org> (accessed 19 February 2018)
- Hustedt Diatom Collection EarthCape Database (2006–2015) Version 15.2. EarthCape OY. Available from: <http://hustedt.awi.de/#ViewID=Home> (accessed 1 March 2018)
- Hustedt, F. (1934) Systematische und ökologische Untersuchungen über die Diatomeen-Flora von Java, Bali und Sumatra nach dem Material der Deutschen Limnologischen Sunda-Expedition. “Tropische Binnenge wässer, Band VII”. *Archiv für Hydrobiologie, Supplement* 15: 187–295, pls 13–20.
- Hustedt, F. (1937) Systematische und ökologische Untersuchungen über die Diatomeen-Flora von Java, Bali und Sumatra nach dem Material der Deutschen Limnologischen Sunda-Expedition. “Tropische Binnenge wässer, Band VII”. *Archiv für Hydrobiologie, Supplement* 15: 187–295, pls 13–20.
- Hustedt, F. (1944) Neue und wenigbekannte Diatomeen. *Bericht der Deutschen Botanischen Gesellschaft* 61: 271–290, 8 pls.
- Hustedt, F. (1956) Diatomeenausdem Lago de Maracaibo in Venezuela. In: *Ergebnisse der deutschen limnologischen Venezuela-Expedition 1952 (F. Gessner & V. Vareschi)*. Deutscher Verlag der Wissenschaften, Berlin, pp. 93–140.
- Hustedt, F. (1957) Die Diatomeenflora des Flußsystems der Weser im Gebiet der Hansestadt Bremen. *Abhandlungen herausgegeben vom Naturwissenschaftlichen Verein zu Bremen* 34: 181–440.
- Hustedt, F. (1961–1966) Die Kieselalgen Deutschlands, Österreichs und der Schweiz. In: Dr. L. Rabenhorsts (Ed.) *Kryptogamenflora von Deutschland, Österreich und der Schweiz, Band 7, Teil 3*. Akademische Verlagsgesellschaft, Leipzig, 816 pp.
- Krammer, K. & Lange-Bertalot, H. (1986) Bacillariophyceae 1. Teil: Naviculaceae. In: *Süßwasserflora von Mitteleuropa. Vol. 2 (1)*. VEB Gustav Fischer Verlag, Jena, pp. 1–876, 206 pls., 2976 figs.
- Kulikovskiy, M., Lange-Bertalot, H., Metzeltin, D. & Witkowski, A. (2012) Lake Baikal: hotspot of endemic diatoms I. In: Lange-Bertalot, H. (Ed.) *Iconographia Diatomologica. Annotated Diatom Micrographs*. Vol. 23. Diversity-Taxonomy-Identification. Koeltz Scientific Books, Königstein, Germany, 861 pp., 156 pls.
- Li, Y., Suzuki, H., Nagumo, T. & Tanaka, J. (2014) Auxosporulation, morphology of vegetative cells and perizonium of *Fallacia tenera* (Hust.) D.G. Mann (Bacillariophyceae). *Phytotaxa* 164: 239–254.
<https://doi.org/10.11646/phytotaxa.164.4.3>
- Li, Y., Suzuki, H., Nagumo, T., Tanaka, J., Sun, Z. & Xu, K. (2015) *Fallacia decussata*, sp. nov.: a new marine benthic diatom (Bacillariophyceae) from Northeast Asia. *Phytotaxa* 224: 258–266.

<https://doi.org/10.11646/phytotaxa.224.3.4>

- Liu, Y., Kociolek, J.P., Fan, Y. & Wang, Q. (2012) *Pseudofallacia* gen. nov., a new freshwater diatom (Bacillariophyceae) genus based on *Navicula occulta* Krasske. *Phycologia* 51: 620–626.
<https://doi.org/10.2216/11-098.1>
- Manoylov, K.M. (2014) Taxonomic identification of algae (morphological and molecular): species concepts, methodologies, and their implication for ecological bioassessment. *Journal of Phycology* 50: 409–424.
<https://doi.org/10.1111/jpy.12183>
- Metzeltin, D., Lange-Bertalot, H. & García-Rodríguez, F. (2005) Diatoms of Uruguay. Compared with other taxa from South America and elsewhere. *Iconographia Diatomologica* 15: 1–736.
- Miscoe, L.H., Johansen, J.R., Kociolek, J.P. & Lowe, R. (2016) Investigation of the cave diatom flora of Kauai, Hawaii: an emphasis on taxonomy and distribution. *Bibliotheca Phycologica* 120: 3–74.
- Patrick, R. & Reimer, C.W. (1966) *The Diatoms of the United States exclusive of Alaska and Hawaii, V. 1. Monographs of the Academy of Natural Sciences of Philadelphia 13*. Philadelphia PA., 688 pp.
- Procopiak, L.K. & Fernandes, L.F. (2003) Valve morphology of the benthic diatom *Fallacia marnieri* (Manguin) Witkowski (Sellaphoraceae-Bacillariophyta). *Brazilian Journal of Biology* 63: 113–119.
<https://doi.org/10.1590/S1519-69842003000100015>
- Ross, R., Cox, E.J., Karayeva, N.I., Mann, D.G., Paddock, T.B.B., Simonsen, R. & Sims, P.A. (1979) An amended terminology for the siliceous components of the diatom cell. *Nova Hedwigia, Beihefte* 64: 513–533.
- Round, F.E., Crawford, R.M. & Mann, D.G. (1990) *The Diatoms. Biology and Morphology of the Genera*. Cambridge University Press, UK, 747 pp.
- Sabbe, K., Vyverman, W. & Muylaert, K. (1999) New and little-known *Fallacia* species (Bacillariophyta) from brackish and marine intertidal sandy sediments in Northwest Europe and North America. *Phycologia* 38: 8–22.
<https://doi.org/10.2216/i0031-8884-38-1-8.1>
- Schmidt, A. (1934–1936) *Atlas der Diatomaceen-kunde*. Leipzig. O.R. Reiland Series VIII (Heft 101–102): pls. 401–408 [F. Hustedt].
- Schoeman, F.R. & Archibald, R.E.M. (1979) *The Diatom Flora of Southern Africa No.5*. CSIR Special Report WAT 50, Pretoria, 65 pp.
- Sonneman, J., Sincock, A., Fluin, J., Reid, M., Newall, P., Tibby, J. & Gell, P.A. (2000) *An Illustrated Guide to Common Stream Diatom Species from Temperate Australia. Identification Guide No. 33*. Murray—Darling Freshwater Research Centre: Albury.
- Spaulding, S.A., Lubinski, D.J. & Potapova, M. (2010) Diatoms of the United States. <http://westerndiatoms.colorado.edu> (accessed 11 November 2017)
- Stancheva, R., Busse, L., Kociolek, J.P. & Sheath, R.G. (2015) *Standard Operating Procedures for Laboratory Processing and Identification of Stream Algae*. California State Water Resources Control Board Surface Water Ambient Monitoring Program (SWAMP) Bioassessment SOP 0003. Available from: http://www.waterboards.ca.gov/water_issues/programs/swamp/bioassessment/docs/sop_algae_lab.pdf (accessed 19 February 2018)
- Stancheva, R., Sheath, R.G. & Kociolek, J.P. (2016) New freshwater gomphonemoid diatoms from streams in the Sierra Nevada mountains, California, USA. *Phytotaxa* 289: 118–134.
<https://doi.org/10.11646/phytotaxa.289.2.2>
- Stepanek, J.G. & Kociolek, J.P. (2013) Several new species of *Amphora* and *Halamphora* from the western USA. *Diatom Research* 28: 61–76.
<https://doi.org/10.1080/0269249X.2012.735205>
- Takano, H. (1990) Semiconical covers in valves of *Navicula teneroides* Hustedt. *Diatom* 5: 5–7.
- Thomas, E.W. & Kociolek, J.P. (2015) Taxonomy of three new *Rhoicosphenia* (Bacillariophyta) species from California, USA. *Phytotaxa* 204: 1–21.
<https://doi.org/10.11646/phytotaxa.204.1.1>
- Van Dam, H., Mertenés, A. & Sinkeldam, J. (1994) A coded checklist and ecological indicator values of freshwater diatoms from the Netherlands. *Netherlands Journal of Aquatic Ecology* 28: 117–33.
<https://doi.org/10.1007/BF02334251>
- Van De Vijver, B. & Cox, E.J. (2015) *Fallacia emmae* sp. nov., (Bacillariophyta) a new soil-inhabiting diatom species from the Sub-Antarctic Region. *Cryptogamie, Algologie* 36: 245–254.
<https://doi.org/10.7872/crya/v36.iss3.2015.245>
- USEPA (2013) National Rivers and Streams Assessment 2013-14: Quality Assurance Project Plan. EPA-841-B-12-007. U.S. Environmental Protection Agency, Office of Water, Washington, DC. Available from: https://www.epa.gov/sites/production/files/2016-04/documents/nrsa1314_qualityassuranceprojectplan_v2.1_final.pdf (accessed 19 February 2018)
- Wetzel, C.E., Ector, L., Van De Vijver, B., Compère, P. & Mann, D.G. (2015) Morphology, typification and critical analysis of some ecologically important small naviculoid species (Bacillariophyta). *Fottea* 15 (2): 203–234.

<https://doi.org/10.5507/fot.2015.020>

Witkowski, A., Lange-Bertalot, H. & Metzeltin, D. (2000) Diatom Flora of Marine Coasts I. *In*: Lange-Bertalot, H. (Ed.) *Iconographia Diatomologica. Vol. 7. Annotated Diatom Micrographs. Diversity-Taxonomy-Identification*. Koeltz Scientific Books, Königstein, Germany, 925 pp., 219 pls.

Witkowski, A. (1991) *Fallacia cassubiae* sp. nov., a new brackish-water diatom from the Puck Bay (Southern Baltic Sea), Poland. *Diatom Research* 6: 401–409.

<https://doi.org/10.1080/0269249X.1991.9705184>

Witkowski, A. (1993) *Fallacia florinae* (Moeller) comb. nov. a marine, epipsammic diatom. *Diatom Research* 8: 215–219.

<https://doi.org/10.1080/0269249X.1993.9705254>