

Falls in older adults with Major depressive disorder (MDD)

Stubbs, Brendon; Stubbs, Jean; Solomon, Gnanaraj; Soundy, Andrew

DOI:

[10.1017/S104161021500126X](https://doi.org/10.1017/S104161021500126X)

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Stubbs, B, Stubbs, J, Solomon, G & Soundy, A 2016, 'Falls in older adults with Major depressive disorder (MDD): a systematic review and exploratory meta-analysis of prospective studies', *International Psychogeriatrics*, vol. 28, no. 1, pp. 23-29. <https://doi.org/10.1017/S104161021500126X>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Copyright © International Psychogeriatric Association 2015

Checked Feb 2016

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



**Falls in older adults with Major depressive disorder (MDD):
A systematic review and exploratory meta-analysis of
prospective studies**

Journal:	<i>International Psychogeriatrics</i>
Manuscript ID:	IPG-04-15-123.R1
Manuscript Type:	Review Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Stubbs, Brendon; University of Greenwich, Health and Social Care Stubbs, Jean; Pharmacy Gnanaraj, Solomon; South London and Maudsley, Physiotherapy Soundy, Andy; University of Birmingham, Physiotherapy
Keywords:	Falls, Mood Disorders

SCHOLARONE™
Manuscripts

Only

Falls in older adults with Major depressive disorder (MDD): A systematic review and exploratory meta-analysis of prospective studies

Brendon Stubbs^{1, 2}, Jean Stubbs³, Solomon Donald Gnanaraj² Andrew Soundy⁴

Running title: major depression and falls

Submission to International Psychogeriatrics

1. School of Health and Social Care, University of Greenwich, Southwood Site Avery Hill Road, Eltham, London, UK
2. Physiotherapy department, South London and Maudsley NHS Foundation Trust, Denmark Hill, London, UK
3. Research Associate, St Andrews Healthcare, Northampton, NN1 5DG, UK
4. Department of Physiotherapy, School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, Birmingham, B15 2TT, UK

*Corresponding author: Brendon Stubbs, University of Greenwich, New Eltham, Avery Hill Road, London, SE9 2UG, United Kingdom. Telephone 004420 8331 8000, fax : 00441604696126 email brendonstubbs@hotmail.com

Abstract

Background

Depressive symptomology is now widely recognised as a key risk factor for falls. The evidence regarding the impact of Major depressive disorder (MDD) on falls is unclear. A systematic review and exploratory meta-analysis was undertaken to explore the relationship between MDD and falls.

Method

Major electronic database were searched from inception till April 2015. Studies that defined MDD and measured falls prospectively in older adults (≥ 60 years) were included. Studies relying on depressive symptomology were excluded. The methodological quality of included articles was assessed and study findings were synthesised using an exploratory meta-analysis.

Results

From a potential of 415 articles, only 3 studies met the inclusion criteria. This included 976 unique older adults with a range of mean age from ≥ 65 to 83 years. The methodological quality of included studies was satisfactory. None of the included studies' primary aim was to investigate the relationship between MDD and falls. The exploratory meta-analysis demonstrated older adults with MDD are at increased risk of falling compared to non-depressed older adults (odds ratio (OR) 4.0, 95% CI 2.0 to 8.1, $I^2=60\%$, $n=976$).

Conclusion

There is a paucity of research considering falls in older adults with MDD. Our results demonstrate that the odds of falling appear to be greater among people with MDD (OR 4.0) than in previous meta-analyses that have only considered subthreshold depressive symptoms. Given the distinct nature and challenges with MDD, more research is required to better understand the falls risk in this group.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Key words: falls, major depressive disorder, major depression

For Review Only

Introduction

Falls and depression are two of the most pervasive and burdensome issues among the expanding older adults population (Gillespie *et al.*, 2012; Murray *et al.*, 2012). Within the last decade there has been an increased focus on the impact of depressive symptoms on mobility and falls risk. Although the relationship is complex (Iaboni and Flint, 2013) the preponderance of research to date has demonstrated an increased risk. The evidence was summarised by (Kvelde *et al.*, 2013) who conducted a meta-analysis and found that depressive symptoms are associated with falls (odds ratio (OR) 1.46, 95% confidence interval (CI) 1.27–1.67). The results were similar to an earlier meta-analysis of depressive symptoms and falls (Deandrea *et al.*, 2010). One of the central components of treatment for depressive symptomology is antidepressant medication (Olfson Marcus 2009). However, a recent meta-analysis confirmed that older adults in receipt of selective serotonin reuptake inhibitors (SSRI) inhibitors are more likely to fall (OR 1.64, 95% CI 1.46-1.85,) and experience injurious falls including those resulting in fracture (OR 1.72, 95% CI 1.55-1.90; (Stubbs, In press)). Other antidepressant medications are also associated with an increased falls risk (Woolcott *et al.*, 2009).

To date, research regarding depression and falls has overwhelmingly focussed on depressive symptoms typically determined through screening tools which are notoriously inaccurate (Mitchell *et al.*, 2009). For instance, many rates of depressive symptoms reported in older age are grossly inflated (Fiske *et al.*, 2009) and false positives are a particular issue (Mitchell *et al.*, 2009). Little research appears to have addressed the relationship between major depressive disorders (MDD) and falls in older adults. However, MDD in older age is a unique and clear manifestation in its own right aside from subthreshold depressive symptoms with particular marked effects on an older adult's mental and physical health (Fiske *et al.*, 2009). Lifetime prevalence of MDD is approximately 16% (Kessler *et al.*, 2003), whilst around 1-5% of older adults are affected but frequencies as high as 10-12% and 14-42% are seen among medical inpatients and long term care facilities (LTCF) respectively

(Fiske et al., 2009). Coincidentally, these are two settings where falls are particularly problematic (Stubbs et al., 2015). MDD is diagnosed through a structured clinical interview and is associated with profound individual, healthcare and societal costs (Rodda *et al.*, 2011). Indeed, mortality and morbidity is particularly high from MDD in older age (Rodda *et al.*, 2011). Given the more severe and distinct nature of MDD in comparison to subthreshold depressive symptomology, one may assume that the relationship between MDD and falls is more marked.

Given this gap in the literature, this systematic review set out to address the current limitations in the literature by undertaking a systematic review and meta-analysis focussing on relationship between MDD and falls in older adults.

Methods

This systematic review was conducted in accordance with the meta-analysis of observational studies in epidemiology (MOOSE) guidelines (Stroup *et al.*, 2000) and reported in accordance with the PRISMA statement (Moher *et al.*, 2009) following a predetermined but unpublished protocol.

Inclusion and exclusion criteria

Articles were eligible that a) included older adults with a mean age over 60 years b) defined MDD according to a structured clinical assessment or confirmed MDD through a medical diagnosis. C) Measured falls prospectively. Articles in any setting were considered. Articles monitoring falls retrospectively were excluded due to concerns regarding reverse causality. Articles investigating depressive symptoms identified by a screening tool (e.g. Patient and Health Questionnaire, (Kroenke *et al.*, 2001) were excluded unless a clear diagnosis of MDD was made. Only articles written in English were considered. We did not attempt to identify ongoing or unpublished studies.

Search strategy

Two authors conducted searches on PubMed, EMBASE, PsycARTICLES, and CINAHL from inception until April, 2015 using the medical subject headings terms 'depression' OR 'depressive' or 'major depression' or 'major depressive disorder' and 'falls'. In addition, citation chasing of the included studies and recent systematic reviews investigating depressive symptoms, antidepressant medication and falls was undertaken (Gebara *et al.*, 2014; Iaboni and Flint, 2013; Kvelde *et al.*, 2013; Launay *et al.*, 2013).

Study selection

After removal of duplicates, two authors screened the titles and abstracts of all potentially eligible articles. A list of full text articles was developed and the full texts of these articles were reviewed. The eligibility criteria were applied and a final list of included articles was reached.

Methodological study appraisal

Two authors independently completed methodological quality assessment of included articles using the Newcastle Ottawa Scale (NOS; (Wells *et al.* 2010)). If any disagreement arose, a third author was available for mediation. The NOS provides an assessment of the quality of non-randomised controlled trials and its content validity and reliability have been established (Wells *et al.* 2010). Included studies are judged on 9 items across three key areas: selection of the participants, comparability of the participants and outcomes. Each study receives an overall score for methodological quality of up to 9 points (one for each item) and scores of 5 and above are considered of satisfactory quality (Wells *et al.* 2010).

Data extraction

Data extraction was undertaken using a predetermined database. Our primary outcome was any fall (1>) in the follow up period within each study. We defined a fall as ‘an unexpected event in which the participants come to rest on the ground, floor, or lower level’ (Lamb *et al.*, 2005). The data collected from each article included: study design, geographical region, setting, demographic details of both groups (mean age, % females), details regarding falls (how they were measured, duration and results), details regarding MDD, antidepressant medication and details regarding other medications.

Statistical analyses

Due to the anticipated heterogeneity, data were pooled with a random effects meta-analysis using Comprehensive meta-analysis (version 3). The OR and 95% CI were calculated for each study and pooled together to provide an overall result. Numerous subgroup analyses were planned including according to setting (community and long term care facilities/ hospitals) and gender. An anticipated subgroup analyses according to antidepressant medical class (selective serotonin reuptake inhibitors (SSRI) and tricyclic antidepressants (TCA)) and type of fall (any fall or recurrent fall) were also

1
2
3 scheduled. In order to investigate sources of heterogeneity, meta regression analyses were planned
4
5 when sufficient data were available ($N > 3$) including study setting, median year of data collection, age
6
7 (mean years), gender (percentage females), prevalence of those in receipt of antipsychotic
8
9 treatment, prevalence taking antidepressants and geographical region. Publication bias was assessed
10
11 with the Begg-Mazumdar Kendall's tau (Begg and Mazumdar, 1994) and Egger bias test (Egger *et al.*,
12
13 1997).
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

Results

Search results and included participants

After the removal of duplicates, the initial database searches yielded 415 potential studies. The most common reason for exclusion was because the study considered depressive symptoms and not clearly defined MDD (N=34). Following the application of the eligibility criteria, 3 prospective studies met the inclusion criteria and were included in the Meta analysis (Gaßmann *et al.*, 2009; Granek *et al.*, 1987; Rosendahl *et al.*, 2003). Full details of the search results including reasons for exclusion are summarized in Figure 1.

Place Figure 1 around here

None of the included studies primary aim was to investigate the relationship between MDD and falls. There were a total of 1,068 older adults represented across the included studies, but complete data on MDD and falls was only available from 530 older adults (85.2%) from one study (Gaßmann *et al.*, 2009). Thus, a total of 976 unique participants were included in the systematic review. The mean age of the sample ranged from 81 (fallers, Granek *et al.*, 1987) to over 65 years (Gaßmann *et al.*, 2009). Two studies' diagnosed MDD based on medical records (Gaßmann *et al.*, 2009, Granek *et al.*, 1987) whilst the other (Rosendahl *et al.*, 2003) specifically stated MDD was diagnosed according to the DSM III (APA) criteria. Two of the studies clearly defined a fall using similar definitions (Rosendahl *et al.* 2003, Gaßmann *et al.*, 2009) and two were based in LTCF (Gaßmann *et al.*, 2009, Granek *et al.*, 1987) whilst Gaßmann *et al.*, (2009) recruited a random population cohort. The length of follow up ranged from 6 months (Gaßmann *et al.*, 2009) to one year (Rosendahl *et al.*, 2003). All of the included articles scored 5 on the NOS. A summary of the included studies is presented in Table 1.

Place Table 1 around here

Meta-analysis investigating falls in those with MDD

1
2
3 It was possible to pool data on falls from all 3 studies' including a total of 976 older adults. The
4
5 overall pooled OR was 4.0 (95% CI 2.0-8.1, $p<0.0001$) and is displayed in Figure 2. There was a
6
7 moderate amount of heterogeneity ($I^2=60\%$) but there was no evidence of publication bias according
8
9 to the Egger test ($=4.1$, $p=0.4$) or the Begg test ($=.66$, $p=0.3$).
10

11
12 *Place Figure 2 around here*
13
14

15 In an exploratory subgroup analysis utilising the two studies in LTCF the pooled OR was 3.3 (95% CI
16
17 1.6 – 6.8, $I^2=63\%$).
18

19
20 There were insufficient data to conduct the other planned subgroup analyses or meta-regression
21
22 analyses.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion

To our knowledge, the current systematic review is the first to focus on the prospective relationship between MDD and falls in older adults. Moreover, the current study is the first to conduct a formal meta-analysis of this relationship establishing a greatly increased risk of falls in those with MDD with an OR of 4.0 (95% 2.0 to 8.1). In an exploratory subgroup analyses the odds of falling remain high in older adults with MDD in LTCF (OR 3.3, 95% CI 1.6 – 6.8). Given the marked burden of both MDD and falls in older adults, coupled with the established relationship between depressive symptoms and falls (Kvelde et al., 2013) it is surprising that we identified only 3 prospective studies investigating the relationship between MDD and falls. Whether or not the OR of 4.0 is an entirely accurate reflection of the risk of falls in people with MDD is unclear due to the limitations in the dataset, but our results indicate older adults with MDD are at a higher risk of falls compared to those with subthreshold depressive symptoms. It is also important to note our finding that none of the authors set out with the primary aim to specifically investigate falls among older people with MDD.

Although there are clearly a number of limitations to our results, the finding that people with MDD appear to be at even greater risk of falls than those with subthreshold depressive symptoms (Kvelde et al., 2013) are concerning. This is especially given the fact that this group have reduced femoral hip bone mineral density (Stubbs et al., 2015b) and that those in receipt of certain type of antidepressant medication such as SSRI are more likely to fall (Stubbs, 2015). Whilst the relationship between falls and SSRI medication is broadly unequivocal, one cannot assume that all those in receipt of SSRI have MDD. This is a point exemplified by (Quach et al., 2013) who found that although a relationship was evident between antidepressant medication and falls, 53% of those in receipt of antidepressant medication had no clinically significant depressive symptoms and the number with MDD is likely to be considerably less. Currently, SSRI medication forms the frontline treatment for MDD in older adults and since the data concerning falls risk and SSRI medication is overwhelmingly pertained through observation studies (Stubbs 2015), there is no concrete reason to

change this unless data can be derived from RCTs. As others have commented in the context of depressive symptoms and falls (Kvelde *et al.*, 2013, Iabona *et al.*, 2013) the relationship between MDD and falls is likely to be influenced by a number of other factors such as lower levels of physical activity (Eggermont *et al.*, 2012) accelerating sensorimotor deconditioning leading to reduced balance and poorer mobility (Leveille *et al.*, 2007). Moreover, the number of pain sites and pain severity are known to be linked to depression (Denkinger *et al.*, 2014) and also falls (Stubbs *et al.*, 2014) and this may play an account for this relationship. Further to this, heightened psychological concerns related to falling that may be present in those with depression (Hull *et al.*, 2012) such as fear of falling, which is also a key risk factor for falls (Deandrea *et al.*, 2010) it is clear to see why the relationship is complex.

Given that exercise is effective in reducing depressive symptomology in older adults (Bridle *et al.*, 2012) there is potential for exercise as an intervention in this group. The potential for exercise as an intervention is also particularly promising as a recent umbrella review of meta analyses of randomised controlled trial interventions found that exercise is the most consistently effective intervention to prevent falls (Stubbs *et al.*, 2015). When one considers that exercise is essential for overall health and wellbeing, prevents the onset of disability in older adults (Tak *et al.*, 2013) and is broadly as effective as pharmacological interventions in reducing cardiovascular disease mortality (Naci and Ioannidis, 2013) its potential role as a health promotion and fall prevention strategy are profound. Naturally people with MDD may experience a range of barriers to engaging in physical activity and motivation strategies may help (Vancampfort *et al.*, 2013) and physical therapists may devise appropriate adaptive interventions.

Although our preliminary results identify that older adults with MDD are more likely to fall and that this relationship appears stronger than in studies considering subthreshold depressive symptoms are unsurprising, there are clearly a number of limitations that need to be considered. First, although just under 1,000 participants were in our analysis, we only identified 3 studies that met our criteria.

However this is a finding in its own right shedding light on the lack of consideration of the risk of falls in older adults with MDD. Second, there was limited information on potential confounders such as psychotropic medication and comorbidities which will influence this relationship. Specifically, due to inadequate reporting in the original manuscripts, it was not possible to clearly illustrate details regarding other falls risk factors between the MDD and control cohorts such as comorbidities, polypharmacy and pain. Future prospective research is clearly needed to disentangle these relationships to better understand falls risk in older adults with MDD. Third, there was evidence of some heterogeneity in the analyses. Finally, although we quantitatively established there was no evidence of publication bias, these tests may be underpowered due to the small number of studies. Nevertheless, allowing for these caveats, our results are consistent indicating a higher risk of falls in older adults with MDD. Future work is required that sets out with the primary aim to investigate falls in older adults with MDD. This work should follow international falls prevention guidelines such as those proposed by PROFANE (Lamb *et al.*, 2005) and clearly investigate the influence of psychotropic medication. Future research should seek to clearly define a fall. Within our meta-analysis, two of the included studies defined a fall, whilst it is unclear if the definition of a fall influences the outcomes of a study, defining a fall is good practice (Lamb *et al.*, 2005). Most importantly, an RCT is required to investigate interventions to prevent falls in older adults with MDD. This could include both an exercise and SSRI arm which investigates the impact of the interventions on both falls and mood. Such an RCT may also assist in answering the lingering questions regarding the influence of SSRI medication on falls.

Conclusion

There is a paucity of research considering falls in older adults with MDD, exemplified by our review in which we only identified 3 prospective studies. However, our results demonstrate that the odds of falling appear to be greater among people with MDD (OR 4.0) than in previous meta-analyses that have only considered subthreshold depressive symptoms.

Conflict of interest

No authors have any to declare. This project received no funding.

Author's roles

Design of study – BS, JS, SG, AS. Searches, acquisition of data and synthesis BS, JS, AS. Meta analysis
BS. BS wrote the first draft of the manuscript; all authors provided critical comments and approved
the final version.

For Review Only

References

Stubbs B. (In press). A meta analysis in falls in older adults taking selective serotonin reuptake inhibitors confirms an association but by no means but by no means implies causation. *American Journal of Geriatric Psychiatry*.

Begg, C. B. and Mazumdar, M. (1994). Operating characteristics of a rank correlation test for publication bias. *Biometrics*, 50, 1088-1101.

Bridle, C., Spanjers, K., Patel, S., Atherton, N. M. and Lamb, S. E. (2012). Effect of exercise on depression severity in older people: systematic review and meta-analysis of randomised controlled trials. *The British Journal Of Psychiatry: The Journal Of Mental Science*, 201, 180-185.

Deandrea, S., Lucenteforte, E., Bravi, F., Foschi, R., La Vecchia, C. and Negri, E. (2010). Risk factors for falls in community-dwelling older people: a systematic review and meta-analysis. *Epidemiology (Cambridge, Mass.)*, 21, 658-668.

Denkinger, M. D., Lukas, A., Nikolaus, T., Peter, R. and Franke, S. (2014). Multisite pain, pain frequency and pain severity are associated with depression in older adults: results from the ActiFE Ulm study. *Age And Ageing*.

Egger, M., Davey Smith, G., Schneider, M. and Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *BMJ (Clinical Research Ed.)*, 315, 629-634.

Eggermont, L. H. P., Penninx, B. W. J. H., Jones, R. N. and Leveille, S. G. (2012). Depressive Symptoms, Chronic Pain, and Falls in Older Community-Dwelling Adults: The MOBILIZE Boston Study. *Journal of the American Geriatrics Society*, 60, 230-237.

Fiske, A., Wetherell, J. L. and Gatz, M. (2009). Depression in older adults. *Annual Review of Clinical Psychology*, 5, 363-389.

Gaßmann, K. G., Rupprecht, R. and Freiberger, E. (2009). Predictors for occasional and recurrent falls in community-dwelling older people. *Zeitschrift für Gerontologie und Geriatrie*, 42, 3-10.

Gebara, M. A., Lipsey, K. L., Karp, J. F., Nash, M. C., Iaboni, A. and Lenze, E. J. (2014). Cause or Effect? Selective Serotonin Reuptake Inhibitors and Falls in Older Adults: A Systematic Review. *The American Journal Of Geriatric Psychiatry: Official Journal Of The American Association For Geriatric Psychiatry*.

Gillespie, L. D., et al. (2012). Interventions for preventing falls in older people living in the community. *Cochrane Database of Systematic Reviews*.

Granek, E., et al. (1987). Medications and diagnoses in relation to falls in a long-term care facility. *Journal Of The American Geriatrics Society*, 35, 503-511.

Iaboni, A. and Flint, A. J. (2013). The complex interplay of depression and falls in older adults: A clinical review. *The American Journal of Geriatric Psychiatry*, 21, 484-492.

Kvelde, T., et al. (2013). Depressive Symptomatology as a Risk Factor for Falls in Older People: Systematic Review and Meta-Analysis. *Journal of the American Geriatrics Society*, 61, 694-706.

Lamb, S. E., Jørstad-Stein, E. C., Hauer, K. and Becker, C. (2005). Development of a common outcome data set for fall injury prevention trials: the prevention of Falls Network Europe consensus. *Journal of the American Geriatrics Society*, 53, 1618-1622.

Launay, C., De Decker, L., Annweiler, C., Kabeshova, A., Fantino, B. and Beauchet, O. (2013). Association of depressive symptoms with recurrent falls: a cross-sectional elderly population based study and a systematic review. *The Journal Of Nutrition, Health & Aging*, 17, 152-157.

Leveille, S. G., Bean, J., Ngo, L., McMullen, W. and Guralnik, J. M. (2007). The pathway from musculoskeletal pain to mobility difficulty in older disabled women. *Pain (03043959)*, 128, 69-77.

Mitchell, A. J., Vaze, A. and Rao, S. (2009). Clinical diagnosis of depression in primary care: a meta-analysis. *Lancet*, 374, 609-619.

Moher, D., Liberati, A., Tetzlaff, J. and Altman, D. G. (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Clinical Trials*, 6, 1-6.

Naci, H. and Ioannidis, J. P. A. (2013). Comparative effectiveness of exercise and drug interventions on mortality outcomes: metaepidemiological study. *BMJ (Clinical Research Ed.)*, 347, f5577-f5577.

- Quach, L., et al.** (2013). Depression, Antidepressants, and Falls Among Community-Dwelling Elderly People: The MOBILIZE Boston Study. *Journals of Gerontology Series A: Biological Sciences & Medical Sciences*, 68, 1575-1581.
- RC, K., et al.** (2003). The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *Jama*.
- Rodda, J., Walker, Z. and Carter, J.** (2011). Depression in older adults. *BMJ: British Medical Journal (Overseas & Retired Doctors Edition)*, 343, 683-687.
- Rosendahl, E., Lundin-Olsson, L., Kallin, K., Jensen, J., Gustafson, Y. and Nyberg, L.** (2003). Prediction of falls among older people in residential care facilities by the Downton index. *Aging Clinical And Experimental Research*, 15, 142-147.
- Stroup, D. F., et al.** (2000). Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA: The Journal Of The American Medical Association*, 283, 2008-2012.
- Stubbs, B., Binnekade, T., Eggermont, L., Sepehry, A. A., Patchay, S. and Schofield, P.** (2014). Pain and the Risk for Falls in Community-Dwelling Older Adults: Systematic Review and Meta-Analysis. *Archives of Physical Medicine & Rehabilitation*, 95, 175-187.e179.
- Stubbs, B., Brefka, S. and Denking, M.** (2015). What works to prevent falls in community dwelling older adults? An umbrella review of meta-analyses of randomised controlled trials. *Physical Therapy*.
- Vancampfort, D., et al.** (2013). The importance of self-determined motivation towards physical activity in patients with schizophrenia. *Psychiatry Research*, 210, 812-818.
- Wells, G., Shea, B., O'Connell, D. and Peterson, J. e. a.** The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses.
- Woolcott, J. C., et al.** (2009). Meta-analysis of the impact of 9 medication classes on falls in elderly persons. *Archives of Internal Medicine*, 169, 1952-1960.

Figure 1. PRISMA 2009 flow diagram for search strategy




Figure 2 - meta analysis of falls in older adults with MDD




Table 1 – summary of included studies

Study	Setting	Design	Participant details	MDD assessment	Define a fall?	Falls acquisition	NOS score
Gaßmann et al 2009	Community	Prospective population cohort	N=622 (530*) N=322 male, 297 female All 65≥ years 107 fell≥1	MDD diagnosis confirmed in medical records.	Yes	Prospective questionnaire on falls for 6 months.	5
Rosendahl et al 2003	LTCF	Prospective	N=78, 81.6 years, 72% female, 41% used walking aid	DSM-III-R criteria	Yes	Prospective follow up for 12 months.	5
Granek et al 1987	LTCF	Prospective	N=368, fallers 83 years and non-fallers 81 years	MDD diagnosis confirmed in medical records	No	Prospective follow up with recording of falls incidents.	5

Key: NOS= Newcastle Ottawa Scale, LTCF= long term care facility, MDD= major depressive disorder

*Some participants did not have complete falls and MDD data, so only 530 included in meta-analysis.