

# False teeth: conodont-vertebrate phylogenetic relationships revisited

## **Susan TURNER**

Monash University Geosciences, Box 28E, Victoria 3800,  
and Queensland Museum, Geosciences Annex,  
122 Gerler Road, Hendra, Queensland 4011 (Australia)  
sue.turner@qm.qld.gov.au

## **Carole J. BURROW**

Queensland Museum, Geosciences Annex,  
122 Gerler Road, Hendra, Queensland 4011 (Australia)  
carole.burrow@gmail.com

## **Hans-Peter SCHULTZE**

Natural History Museum, The University of Kansas,  
1345 Jayhawk Blvd., Lawrence, Kansas 66046-7561 (USA)  
hp1937@ku.edu

## **Alain BLIECK**

Université de Lille 1, Sciences de la Terre, FRE 3298 du CNRS Géosystèmes,  
F-59655 Villeneuve d'Ascq cedex (France)  
alain.blieck@univ-lille1.fr

## **Wolf-Ernst REIF†**

Eberhard-Karls-Universität, Institut für Geowissenschaften,  
Sigwartstraße 10, D-72076 Tübingen (Germany)

## **Carl B. REXROAD**

Indiana Geological Survey, 611 North Walnut Grove,  
Bloomington, Indiana 47405-2208 (USA)

## **Pierre BULTYNCK**

Department of Paleontology, Royal Belgian Institute of Natural Sciences,  
Vautier street 29, B-1000 Brussels (Belgium)

## **Godfrey S. NOWLAN**

Geological Survey of Canada,  
3303 – 33rd Street NW, Calgary, Alberta, T2L 2A7 (Canada)

---

Turner S., Burrow C. J., Schultze H.-P., Blieck A., Reif W.-E.†, Rexroad C. B., Bultynck P. & Nowlan G. S. 2010. — False teeth: conodont-vertebrate phylogenetic relationships revisited. *Geodiversitas* 32 (4): 545-594.

**ABSTRACT**

An evidence-based reassessment of the phylogenetic relationships of conodonts shows that they are not “stem” gnathostomes, nor vertebrates, and not even craniates. A significant group of conodont workers have proposed or accepted a craniate designation for the conodont animal, an interpretation that is increasingly becoming established as accepted “fact”. Against this prevailing trend, our conclusion is based on a revised analysis of traditional morphological features of both discrete conodont elements and apparatuses, histological investigation and a revised cladistic analysis modifying that used in the keystone publication promoted as proof of the hypothesis that conodonts are vertebrates. Our study suggests that conodonts possibly were not even chordates but demonstration of this is beyond the scope of this paper. To summarize, in conodonts there is low cephalization; presence of simple V-shaped trunk musculature and unique large-crystal albid material in the elements; lack of a dermal skeleton including characteristic vertebrate hard tissues of bone, dentine and enamel; lack of odontodes with bone of attachment and a unique pulp system; lack of segmentally-arranged paraxial elements and dermal elements in median fins, all of which supports neither a vertebrate nor a craniate relationship for conodonts.

**KEY WORDS**

Chordata,  
Craniata,  
conodont animals/  
elements,  
character coding,  
cladistic analysis,  
palaeo-histology.

**RÉSUMÉ**

*Des pseudo-dents : une réévaluation des relations phylogénétiques des conodontes et des vertébrés.*

Une réévaluation des relations phylogénétiques des conodontes est fondée sur de nouvelles preuves. Elle montre que les conodontes ne sont ni des gnathostomes-souches, ni des vertébrés, ni même des crâniates. Un groupe significatif de spécialistes des conodontes a proposé, ou accepté, que ces organismes soient considérés comme des crâniates, une interprétation qui est en train de s'installer en tant que fait avéré. Notre conclusion va à l'encontre de cette tendance; elle est fondée sur une révision des traits morphologiques traditionnels à la fois des éléments isolés et des assemblages de conodontes, sur les données histologiques et sur une analyse cladistique révisée, ce qui modifie les conclusions de la publication principale qui a promu l'hypothèse selon laquelle les conodontes seraient des vertébrés. Notre étude suggère même que les conodontes n'aient pas été des chordés, mais la démonstration de cette hypothèse va au-delà de l'objectif de cet article. En résumé, chez les conodontes, le degré de céphalisation est faible; la musculature du tronc a une forme simple en V; les éléments isolés montrent un tissu blanc avec des cristaux de grande taille, uniques pour ce tissu; il n'y a pas de squelette dermique incluant les tissus durs caractéristiques des vertébrés tels que l'os, la dentine et l'émail; il n'y a pas d'odontodes avec leur os et leur système pulpaire unique; il n'y a pas d'éléments paraxiaux disposés de façon segmentée sur le corps, ni d'éléments dermiques aux nageoires médianes. Tout cet ensemble de caractères ne permet pas d'argumenter des affinités entre conodontes et vertébrés ou crâniates.

**MOTS CLÉS**

Chordata,  
Craniata,  
animal-conodonte,  
codage des caractères,  
analyse cladistique,  
paléohistologie.

---

*This paper is dedicated to the memory of our colleague,  
Professor Dr Wolf-Ernst Reif (1945-2009),  
who died just after acceptance of this paper.*

## INTRODUCTION

The zoological affinity of conodonts, small exclusively marine animals of the Palaeozoic to early Mesozoic eras, has been vigorously debated ever since their phosphatic parts (calcium phosphate or apatitic conodont “elements”) were first described by Pander (1856). For example, Müller (1981) listed almost 50 publications for the period 1856–1975 that variously suggested affinity with ten different taxonomic entities, including such diverse groups as vertebrates, annelids and plants. Chinese terminology reflects this with 11 different words equivalent to higher taxa for conodonts (Wang Cheng-Yuan, pers. comm. to ST, 1984). Of the multitude of proposed conodont relationships, Sweet (1988: 170–184) easily dismissed most, including strong evidence against any affinity with myxinooids (hagfishes) and even questioned the wisdom of a chordate relationship. Sweet (1988: 172, 173) discussed but provided no refutation for Tillier & Cuif’s (1986) claim that conodonts might be related to aplacophoran molluscs (but see Briggs *et al.* 1987), where they noted similarities between the two, having discovered calcium phosphate in the teeth and mandibles of one aplacophoran taxon. Calcium phosphate of course is not limited to conodonts (or vertebrates) but is found in several “invertebrate” taxa, such as in phyllocarids, nemertean stylets, some brachiopod shells, and also likely in the radular teeth of some chitons (Watabe 1990), and so Tillier & Cuif’s (1986) conclusions should be regarded as based on rather simplistic comparisons. Sweet (1988) concluded his analysis of possible affinities of the conodont organism with the suggestion that they could probably best be assigned to a separate phylum and were the result of yet another experiment in evolution that eventually became extinct. Here, we wish to continue the debate because we do not accept the increasingly prevailing paradigm that conodonts are vertebrates.

In referring to “conodonts” here we concentrate on only the euconodonts (= conodonts *s.s.*) or “complex” conodonts, as did e.g., Donoghue & Aldridge (2001), Donoghue *et al.* (2000, 2008), and others, who claim that conodonts are vertebrates (see below) and restrict their hypothesis of

conodont interrelationships to euconodonts. The relationships of the latter to other conodont groups (protoconodonts, paraconodonts) are still controversial (see references in Reif 2006). However, all conodont groups need to be considered in the context of their possible relationship to vertebrates if the conodont groups are closely related. Szaniawski (1983) interpreted protoconodonts as close relatives of chaetognaths reiterating this view in 1987 but then stressing that the evolutionary link between proto- and para-conodonts remains to be conclusively established. In 2002 he provided evidence for the protoconodont origin of chaetognaths. On the other hand, on the basis of well-preserved material from the Upper Cambrian of Sweden, Szaniawski & Bengtson (1993) made a strong case for a close evolutionary link between para- and euconodonts. Müller & Hinz-Schallreuter (1998) considered all three groups together, and noted a diversity of histological structures: the earliest supposed euconodont *Cambropustula* Müller & Hinz, 1991 from the lower Upper Cambrian, for instance, lacks “white matter” (= albid tissue, an essentially opaque formless tissue “characterized by voids, which may be interlamellar spaces, or concentrations of small, densely packed, irregularly shaped cellulose” [Lindström & Ziegler 1971]), supposedly an evolutionary novelty within conodonts *s.s.* (Table 1); they considered the protoconodonts to be “ancestors” of paraconodonts. Others consider that paraconodonts and euconodonts is a totally artificial separation of a biologic continuum (Nicoll, pers. comm. 2009); the only real distinction between the two is the absence of white matter in the former. Both groups have complex apparatus structures that are similar in related species either side of the albid tissue divide. Nevertheless, Donoghue & Aldridge (2001) maintained the separation of the three groups when considering conodont relationships to vertebrates. The problem then is how to reconcile e.g., Szaniawski & Bengtson’s (1993) and Müller & Hinz-Schallreuter’s (1998) hypotheses of euconodont phylogenetic relationships with Donoghue & Aldridge’s (2001: fig. 6.4) and Smith *et al.*’s (2001: fig. 5.5) hypotheses. Even with these contradicting views, to test their methodology we follow here Donoghue & Aldridge’s (2001) claim

TABLE 1. — Comparison of terminologies (homologies) employed for different hard tissues of the conodont element by principal sources and a selection of papers cited. **1**, Gross 1954, 1957, 1960; Müller & Nogami 1971; Müller 1981; Schultze 1996; Reif 2006; **2**, Briggs 1992; Sansom *et al.* 1992, 1994; Aldridge *et al.* 1993; Aldridge & Purnell 1996; Smith *et al.* 1996, 2001; Samson 1996; Janvier 1997; Donoghue 1998, 2001; Donoghue *et al.* 1998, 2000, 2006; Donoghue & Sansom 2002; Aldridge & Briggs 2009; **In bold**: first use of terms.

Conodont author	Crown tissue	“White matter”	Basal filling
Pander 1856	konzentrische Lamellen (concentric lamellae)	kleine Zellen oder Höhlen (with small cells or bubbles)	hohl, Pulpa (hollow, pulp cavity)
Branson & Mehl 1933			bony (no structure of ordinary bone)
Hass 1941	lamellae	cellular or cancellate structure	hollow
Gross 1954, 1957 Gross 1960	Lamellen (lamellae) Lamellen (lamellae)	Scheinpulpa (pseudo pulp) durch Bläschenbildung getrübt Teil (part cloudy by formation of bubbles)	Basisfüllung (basal filling) dicke Lamellen in Richtung der Lamellen der Krone (thick lamellae in line with those of the crown)
Schmidt <i>in</i> Schmidt & Müller 1964	Schmelz (enamel)	–	Dentin (dentine)
Lindström 1964	lamellae	<b>white matter</b> with small irregular cells	lamellae
Lindström & Ziegler 1971		<b>albid</b>	
Barnes <i>et al.</i> 1973	<b>hyaline</b>	white matter	growth lamellae in continuation with those of the crown
Müller & Nogami 1971; Müller 1981	growth lamellae		lamellae in continuation with crown lamellae
Lindström & Ziegler 1981	concentric lamellae	recrystallized with holes	different tissues
“German” school <sup>1</sup> Barskov <i>et al.</i> 1982	lamellar tissue	white matter	spongy bone
Aldridge <i>et al.</i> 1986	hyaline lamellae, crystallites parallel to direction of growth		Lamellae not always in continuation with those of the crown
Sweet 1988	lamellae of hyaline	recrystallisation of hyaline	lamellae continuous with hyaline lamellae
Wright 1990b	lamellae with crystallites parallel to direction of growth	finely crystalline with holes of > 1.0 µm diameter	
Dzik 1986, 2000	enamel: large elongated crystallites of apatite	not bone	isometric apatite crystallites, dentine
Szaniawski 1987; Szaniawski & Bengtson 1993	lamellae		two layers of lamellae
Hall 1990	elongate, prismatic and short, platy crystallites		amorphous, cryptocrystalline masses
“British school” <sup>2</sup>	enamel	cellular bone or kind of enamel tissue	dentine (mesodentine, lamellar-to-spheritic tubular or atubular), or globular cartilage
Kemp & Nicoll 1995, 1996; Kemp 2002b	contains collagen therefore not enamel	absence of collagen therefore not bone	similarity to cartilage
Donoghue 1998; Donoghue <i>et al.</i> 2000; Donoghue & Aldridge 2001	enamel	distinct conodont tissue; developmentally homologous to enamel; cellular dermal bone	dentine; globular calcified cartilage or dentine
Kemp 2002a	large, flat, oblong crystals parallel to long axis of element		
Guo <i>et al.</i> 2005			tubular, atubular and spherulitic dentine

TABLE 1. — Continuation.

Conodont author	Crown tissue	“White matter”	Basal filling
Trotter <i>et al.</i> 2007 Aldridge & Briggs 2009	elongate, well-aligned crystals translucent hyaline of lamellae = enamel	extraordinarily large crystals white matter	opaque basal body – dentine with tubules + calci- spheres
Dzik 2009 this paper	lamellin hyaline not enamel	white matter not bone	not dentine

that euconodonts have to be considered separately when testing their relationship with vertebrates.

The discovery particularly of the Scottish (Early Carboniferous, Mississippian), South African (Late Ordovician) and also Early Silurian (one from Wisconsin) remains of whole and partial specimens showing soft tissues preserved with conodont elements towards the anterior end of an elongate animal (Fig. 1A, B), precipitated the interpretation of a chordate-like anatomy for conodonts (Smith *et al.* 1987; Aldridge & Briggs 2009). Indeed, Aldridge *et al.* (e.g., 1986, 1993), Briggs (1992) and Sansom *et al.* (1992) went further, arguing that conodonts are vertebrates (but see discussion in Reif 2006; Bliciek *et al.* 2009; Bultynck 2009 and here). This opinion became virtual dogma with the publication of an extensive evaluation of chordate and conodont characters by Donoghue *et al.* (2000), who provided a cladistic analysis where conodonts became “stem gnathostomes” (see also Donoghue *et al.* 2006). We disagree with their conclusion and find that, based on the physical evidence, it is doubtful that conodonts were craniates.

In the present paper we put forward our case stating the need for a refutation of Donoghue *et al.*'s (2000) hypothesis, listing evidence against a conodont-vertebrate relationship, incorporating this data in a cladistic analysis based on the matrix they used, discussing in more detail our reasons against vertebrate relationship, followed by our conclusions. We think equally that whether conodonts are or are not truly chordates is still an open question, but a demonstration of such would require a far larger cladistic analysis than is the object of this paper or recovery of conodont animals with more clearly preserved diagnostic structures.

#### SYSTEMATIC NOMENCLATURE NOTE

We make here a brief point about the use of terminology such as “crown group” (CG), “stem group” (SG), “total group” (TG). Using the definition of Jefferies (1979), a CG is the smallest monophyletic group, or clade, to contain the last common ancestor of all extant members, and all of that ancestor's descendants; all organisms that are more closely related to this CG than to any other living group are referable to its SG (Hennig 1969, 1983). As living taxa are by definition in the CG, it follows that all members of its SG are extinct, and thus that SGs only have fossil members. A CG plus its SG considered together then constitute the “total group”. Accepting these definitions presupposes that a SG is perforce paraphyletic (e.g., Jefferies 1979; Donoghue 2005). [But note that discrepancies can appear in the literature such as in Donoghue *et al.*'s (2006: fig. 1) paper where Chondrichthyes are included in the SG Gnathostomes whereas Acanthodii are not even considered.]

Another problem is the use of the same name for the “total group” as for the “crown group” (e.g., gnathostomes, tetrapods, etc.). In the case of tetrapods, the TG Tetrapodomorpha Ahlberg, 1991 includes the SG fossil piscine sarcopterygians down to the next extant sister group, the dipnoans. The SG includes piscine and tetrapod-like sarcopterygians. The content of the CG Tetrapoda depends on the position of the extant forms in a phylogenetic tree (Laurin & Anderson 2004). In contrast, in the case of gnathostomes, the TG called Gnathostomata by Donoghue *et al.* (2000) creates a problem, because the next extant taxon is the Petromyzontida; no name based on a phylogeny has been suggested.

This TG Gnathostomata should then include as the SG most fossil “agnathans”, and therefore forms that lack characteristic gnathostome jaws. The TG Gnathostomata is thus not different from what could be named “euvertebrates” in the following topology (myxinoid (lampreys + euvertebrates)) (see Fig. 6). We prefer to use an apomorphy-based definition of a TG Gnathostomata (Placodermi (Chondrichthyes (Acanthodii + Osteichthyes))), that is, vertebrates with jaws, and distinguish it from a CG Eugnathostomata. The CG Eugnathostomata (Chondrichthyes (Acanthodii + Osteichthyes)) would include all extant gnathostomes and their fossil relatives, but not their fossil sister group Placodermi.


#### “BRITISH SCHOOL” CONCEPTS OF CONODONTS AND VERTEBRATES

Discovery by British palaeontologists (Briggs *et al.* 1983) of reasonably complete and partial conodont-bearing animal specimens preserved with conodont elements distorted from life position but still in the head region (Fig. 1A, B), opened a new door on the interpretation of biological affinity because for the first time there was a “real conodont animal”. The ten specimens from the Lower Carboniferous Granton Shrimp-Bed, Edinburgh (Clarkson [1985: 5] thought that the original “looked like a small lamprey”), combined with Silurian specimens from Wisconsin (Smith *et al.* 1987) and the more than 100 complete apparatuses, some in partial bodies, from the Upper Ordovician Cedarberg Formation in South Africa (Aldridge & Briggs 2009), formed an early, and highly variable cohort of diverse biological information to assimilate into a single conodont animal model. As noted by others (e.g., Briggs *et al.* 1983: 3; Sweet 1988: 28-32; Conway

Morris 1989: 138; Bultynck 2009), the preservation is moderate, and interpretation of structures remains open to discussion (see e.g., Fig. 1C, E-H). The “British School” of conodont study that claims that conodonts are vertebrates mainly emanates from the work of R. A. Aldridge and his students at Leicester University (Aldridge & Briggs 2009); several within this group have begun to class the elements as “microvertebrate” remains (e.g., in the British Micropalaeontological Society literature), a designation with which we also disagree based on the characters discussed below.

Despite widespread scientific challenges (e.g., Kemp & Nicoll 1995, 1996; Schultze 1996; Kemp 2002a, b; Turner *et al.* 2004; Reif 2006; see e.g., Fig. 2) to such opinions and interpretations, conodonts continue to be touted as vertebrates, and even as stem gnathostomes, in scientific publications (e.g., Donoghue *et al.* 2000, 2008; Purnell 2001; Holland & Chen 2001; Sansom *et al.* 2001, 2005; Smith *et al.* 2001; Donoghue & Sansom 2002; Janvier 2003, 2007a, 2008; Albanesi & Bergström 2004; Donoghue & Purnell 2005; Dzik 2009; Sire *et al.* 2009), and increasingly now in text- and other books both scholarly and semi-popular (e.g., Prothero 1998; Liem *et al.* 2001; Benton 2005), and on the all-pervading Internet. There has been an increasing acceptance that conodonts are vertebrates in recent scientific and “informed” popular literature (e.g., Mallatt & Chen 2003; Hall 2005; Kuhn & Barnes 2005; Guo *et al.* 2005; Janvier 2006, 2007a, 2008; see discussion by Blicek *et al.* 2009) and even on websites (Janvier 1997, 2001), with little or no acknowledgement of contrary arguments (e.g., Aldridge & Briggs 2009). The mostly dogmatic promulgation and uninformed acceptance of the hypothesis that conodonts are vertebrates has also invaded the molecular biology/

FIG. 1. — Conodont animal fossils showing soft tissues preserved and a selection of hypothetical interpreted reconstructions (not to scale): **A**, conodont fossil, *Clydagnathus? cf. cavusformis* in lateral view (anterior to the left) from the Lower Carboniferous (Visean) Granton Shrimp Bed Lagerstätte, near Edinburgh, Scotland, after Aldridge *et al.* (1993: fig. 3; specimen RMS GY 1992.41.1); **B-B'**, interpretative drawing of conodont animal modified from Aldridge *et al.* (1993), with highlighted conodont elements above the animal, Lower Carboniferous from Scotland, redrawn by Aldridge *et al.* (1993) after Briggs *et al.* (1983: fig. 2); **B**, part of IGSE 13821; **B'**, assemblage from counterpart IGSE 13822; **B''**, head region of IGSE 13821; **C**, interpretation of whole conodont animal based on *Besselodus* Aldridge, 1982 elements and *Clydagnathus? cf. cavusformis* body modified from Dzik (1986: fig. 4); **D**, interpretation of conodont restoration with external conodont element array on front of head and added (unsubstantiated) branchial openings (modified from figure by David Baines in Aldridge & Briggs (2009: fig. 4.2)); **E**, interpretative restoration of whole conodont animal modified from Pridmore *et al.* (1997: fig. 4B; note that Janvier's [2009] reconstruction is virtually identical to this one); **F**, interpretation of whole conodont animal with large eyes and two hypophysial openings, modified from Donoghue *et al.* (2000: fig. 6D); **G**, interpretation of



conodont animal head (arrow: anterior) using *Polygnathus* Hinde, 1879 elements by Nicoll (1995: text-fig. 1) (reproduced with permission from the author); **H**, for comparison an interpretation of a “naked” lower vertebrate (mid-Palaeozoic agnathan) *Jamoytius* White, 1946, from a similar (Lower Silurian) Lagerstätte in Scotland, showing fin configuration, muscle blocks, and branchial openings (modified from a restoration by Colin Newman in Dixon *et al.* [1988: 26, figure]). IGSE, British Geological Survey, Murchison House Edinburgh, Scotland; RMS, now Museums of Scotland, Edinburgh, Scotland. Abbreviations: Pa, Pb, M, Sb, Sc: usual nomenclature for conodont elements. Scale bars: A, 2 mm; B', B'', 2 mm; B, C-G: 5 mm.

genomics literature (see e.g., Shimeld & Holland 2000; Kawasaki *et al.* 2004), and conodonts are now accepted by many as more highly evolved than lampreys and hagfishes.

However, as noted by Hall (2005), not everyone accepts this view and if the work of the majority of conodont element workers is consulted (e.g., Walliser 1994; Nicoll 1995; Belka 2004), the claims would not have been as extreme. Most conodont workers have not become deeply embroiled in the debate because most are busy utilising conodont elements to solve geological problems and several, including R. D. Norby (pers. comm. to ST, 1997) think that leaving conodonts (for the time being) as “protochordates” provides a solution (but see now the arguments of Raineri [2006] and Reif [2006]).

#### POSSIBLE FUNCTION OF SOME CONODONT APPARATUSES

The conodonts (eu + para) first appear to have developed mineralized tissues to support the feeding apparatus in the late or early Late Cambrian. A trend in conodonts for an increase in morphologic complexity from coniform single denticles in the Late Cambrian to ramiform elements with multiple denticles on one or more processes to complex pectiniform blades and plates that can have surficial denticles and ridges can be observed. From the smooth surfaces of Cambrian forms, there developed ridges, grooves and surface striations or patterning. Some surfaces of some elements show regular to irregular reticulated surfaces that are thought to have been a reflection of the cell pattern of the tissue that covered the element and formed the growth layers of apatite that were accreted on the surface of the element with growth. All, or almost all, have morphologically differentiated apparatuses. Experimentation occurred in conodont apparatus architecture in the Late Cambrian and through much of the Ordovician. Particularly, Ordovician conodont apparatuses exhibit complexity with the S vs P element split an early development. By the Silurian, apparatus architecture variety had become more limited and during the Devonian the pattern is almost stable remaining that way until the end-Triassic extinction of conodonts (e.g., Nicoll 1995).

So what are the possible functions of the conodont apparatus elements in the conodont animal? It is accepted by all sides of the discussion that the elements are located in the anterior or head region of the animal and that the apparatus elements were involved in the capture and ingestion of food. It is the mechanism of food capture and ingestion that is in question. It is also generally agreed that the morphological variation observed in the elements of any given apparatus suggests that the different parts served different functions.

Although there exists one notational location scheme for septimembrate apparatuses (Sweet 1981) that can be used for the major part of euconodont taxa, there is no unique typical conodont apparatus structure. The apparatus generally consists of 15 elements of 7 element types (septimembrate) but many geologically younger apparatuses, especially of Triassic age, have 15 elements of 8 element types (octimembrate). In the later example one of the element types, which in the early to mid Palaeozoic had been represented by two identical element pairs, have differentiated into two morphologically discrete element types. Some apparatuses consist of only 4 P elements (Polyplacognathidae, see Sweet 1988: 71).

Conodont apparatuses are generally composed of three basic types of elements that have three possible distinct functions. The anteriorly, and transversely oriented, M elements served to keep large material/particles out of the food stream and protect the more fragile S elements. The S elements followed and, presumably covered by ciliated tissues, collected the food particles. Lastly are the P elements and these could do quite different jobs depending on their morphology. If just S and P elements are considered, there are a number of different types of conodont animals that must have had very different feeding strategies: coniform – coniform apparatuses (*Terodontus*, *Drepanodus*, *Drepanoistodus*); coniform – ramiform apparatuses (no known taxa); coniform – pectiniform apparatuses (*Jumudontus*, *Pelekysnathus*, *Icriodus*); ramiform – ramiform apparatuses (*Erraticodon*, *Cordylodus*); ramiform – pectiniform apparatuses (*Polygnathus*, *Ozarkodina*) (for full tabulation see Nicoll 1992: Table 1). In those apparatuses with pectiniform P, such as *Polygnathus*






FIG. 2. — Histology of conodont element (showing continuous growth) vs vertebrate scale or tooth (odontode) formed in a papilla of mesenchymous cells: **A**, drawing of longitudinal section showing continuous growth with lateral additions and formation of serration of the conodont *Gnathodus texanus* Roundy, 1926 modified from Gross (1954: fig. 2, 1-5, subsequent addition of serrations) *contra* Donoghue (1998) Type 4 growth (reproduced in Reif 2006: fig. 4) where a basal body (or is it white matter?) apparently occurs in each serration (reproduced with permission from *Senckenbergiana lethaea*); **B**, **B'**, longitudinal section through conodont "*Ctenognathus*" Pander, 1856 showing interrupted growth = "healing" (after Gross 1957: fig. 2D, F); **C**, schematic cross section to show continuous growth (arrows) of conodont element and basal filling (after Gross 1957: fig. 4); **D**, **D'**, comparison of cross section of conodont element with osteichthyan (mammalian) tooth with pulp cavity (from Schultze 1996: fig. 2A, B; reproduced with permission from the author); **E**, **E'**, formation of vertebrate odontodes in comparison to conodont growth in A. The odontode is formed by a papillary organ (with enamel organ, odontoblasts and osteoblasts in a single morphogenetic step). It cannot grow, but is enlarged instead by addition of a new odontode, that is formed by a new papillary organ resulting from a new interaction between ectoderm and mesenchyme (modified from Reif 2006: fig. 3a): **thick line**, enamel or enameloid; **radiating lines**, dentine with pulp cavity underneath; **basal tissue**, attachment bone = cement with thin lines of deposition. Note the presence of pulp in vertebrate examples. Abbreviations: **al**, lamellar tissue; **b**, basis; **bb**, basal body; **bo**, bone; **b.gr**, basal groove; **bri**, interruption = break of growth in lamellar tissue; **bt**, basis cone; **ce**, cement; **de**, dentine; **e**, enamel; **f**, fine fibers; **la**, lamellar tissue; **tf**, inner basal filling; **t.gr**, boundary between the two kinds of filling of the basal groove; **w.m.**, albid tissue or white matter. a, b, from original author figures. Scale bars: 0.1 mm.

*linguiformis* or *Palmatolepis* or *Neogondolella*, the elements were oriented laterally, opposed across the axial plane of the food chain (Nicoll 1987). In those apparatuses with coniform or ramiform P elements, they would have been oriented with the cusp tips pointing in the same direction (Nicoll 1995). Analyses of this sort demonstrated that many different conodont animals evolved with different feeding strategies.

Conodont elements are complex and consisted of two parts, a crown and an attachment structure (Fig. 2C). The histology of each of these is distinct (see below). The crown was that part of the element that was in contact with the external environment and thus the interactive part of the element, be it analogous to a tooth or a tissue-support structure. The attachment structure (attachment cone or plate or basal body; Fig. 2D) was connected to the inner hollow of the crown (or to its lower surface in planate elements) and to the muscle or ligament tissue of the conodont animal that controlled its movement. In most cases, only the crown tissue is recovered from the rock that contained the collected specimens, but attachment structures are common in some localities and can be especially common in some genera and element types (Nicoll 1995).

Here we do not accept the “British” reconstructions of the conodont animal (see e.g., Fig. 1D, F), and interpretations of the function of conodont apparatuses, i.e. elements functioning as teeth. As for an alternative view, Nicoll (1995) commented that the apparatus structure could have functioned as part of a microphagous filter-feeding structure and put the apparatus in an amphioxus-like body (Fig. 1G) to explain and interpret the anatomical relationships, claiming that there were three major and different working arrangements of conodont elements and that none could have served as a cutting function but that the Pa elements might have served in something of a crushing capacity (but see also below and Fig. 3C). The elements were at least partially, if not completely, covered by ciliated tissue and did not have to go through the difficult-to-explain process of being “retracted” for new layers to be secreted. Nicoll (1995; and e.g., Kemp & Nicoll 1995) strongly supported a non-vertebrate affinity, with which we concur. Crown growth was

centrifugal (layers added increasing the size and complexity of the element with growth) and may or may not have occurred periodically (Zhang *et al.* 1997), which does not occur in vertebrate teeth.

Based on known specimens, we contend that a conodont apparatus should not be equated in any way with an array of biting, chewing, crushing/grinding vertebrate teeth (cf. Reif 2006). In the case of some specific genera or species (Purnell & von Bitter 1992; Purnell 1995b), such comparisons have been made but they should not be used for assigning conodonts to vertebrates. Others (e.g., Gedik & Çapkinoglu 1996) claim a parasitic mode of life for conodonts, where by attaching themselves to the soft-tissue of a host-animal and sucking its fluids, the elements could show wear-traces, but those traces could heal during the time of attachment to the host animal, with major breaks visible later but as yet the morphogenetic evidence has not been presented.

A different point of view is given herein on the basis of the seximembrate apparatus of *Polygnathus linguiformis linguiformis* Hinde, 1879 (Fig. 3C). Two groups of elements can be recognized in this species on the basis of their location and their morphology. The anterior part of the apparatus consists of a set of ramiform elements (S and M) with mostly fine, delicate denticles, which were located in the anterior part of the buccal cavity (Nicoll 1985: fig. 10), i.e. inside the mouth. They are generally considered, also in other conodont genera, as a food-grasping/filtering system (e.g., Nicoll 1985; Purnell 1993; Walliser 1994); again note the symmetry, which is opposed to any vertebrate array.

The second part of the apparatus consists of two pairs of pectiniform elements. A first pair of comb-shaped Pb elements, followed by a pair of platform elements (Pa) that most likely were located in the posterior part of the buccal cavity in the pharynx, presumably close to the opening of the gut (Nicoll 1985). Grinding, crushing or cutting activities have been proposed for them (Nicoll 1985; cf. Purnell 1995a for *Idiognathus* Gunnell, 1931). The upper surface of the platform of the Pa element in *Polygnathus linguiformis* is characterized by a median, longitudinal crest (the carina) flanked on both sides by a longitudinal depression (the adcarinal troughs).




FIG. 3. — Morphogenesis of euconodont elements: **A**, hypothetical euconodont morphogenesis of apparatus, based on *Clydagnathus cavusformis* Rhodes, Austin & Druce, 1969, Stage 1 – “shards” (cf. Reif 2006) of apparatus. There is no dental lamina, dental papilla, or tooth bud structure present (cf. Reif 1984); a pulp is absent; the tissues are not living but laid down like layers of inorganic crystals; repair is done by living tissue surrounding the element; **B**, Stage 2, continuous centrifugal growth with outer layers forming multi-cusps; whole apparatuses must form in one contemporaneous session to work, unlike vertebrate rotational dentitions (even in the most advanced mammals there are three separate cycles possible: milk, adult and wisdom teeth); **C**, Stage 3, the working complex in dorsal view, anterior to top; oesophagus to bottom: reconstructed seximembrate apparatus of *Polygnathus linguiformis linguiformis* Hinde, 1879 (assembled by PB, and see Sweet 1998: 99) based on elements from sample BT 18, S Morocco, Tafilalt, Lower Givetian Bou Tchrafine section (see Bultynck 1987). The location of the elements is based on the most generally accepted scheme. The elements are shown in such way that their outline and ornamentation can be clearly recognized; their orientation does not correspond to the original natural orientation in the apparatus. So, the **M** and **S** elements (except the **Sa**) and the **Pb** element should be turned upward over an angle of 45°. The anterior and posterior processes of the **Pb** element meet at an angle of about 130°, similar to the deflection of the posterior tongue of the **Pa** element. The orientation of the **P** elements is a matter of discussion. The elements of the apparatus are deposited at the Institut royal des Sciences naturelles de Belgique, Brussels, under catalogue number I.R.Sc.N.B. no. b5177. Scale bar: 500 µm.

The platform ends in a tongue-like structure, deflected outwards and downwards (see Nicoll 1987: plate 5.3, figs 8-12). During life the element was covered by (epidermal) tissue, the cells of which could leave an imprint on the surface of the element (see Weddige 1989: fig. 14). The course and the height

of the carina, the depth of the adcarinal troughs and the form and orientation of the posterior end of the platform are variable and diagnostic for species, not only in the genus *Polygnathus* Hinde, 1879. The morphology of this type of **Pa** element is not adapted for good occlusion between the right and

left Pa elements [Note that contrarily good occlusion is necessary for the functioning of vertebrate teeth.] An alternative interpretation might be that contractions in the epidermal tissue bring food particles via the Pb elements to the Pa elements and these are then guided via the adcarinal troughs to the gut opening. The tongue-like posterior part of the Pa element might have assisted a swallowing movement at the opening.

Finally, it should be stressed that the architecture of at least some evolved multimembrate conodont apparatuses show no convincing similarities with tooth arrangements in the buccal cavity either of agnathans or primitive Gnathostomata and functioned in a completely different way.

#### CHORDATE, CRANIATE, VERTEBRATE CHARACTERS

The major phyla of the deuterostomes are the Echinodermata, Hemichordata (including the Pterobranchia, the Enteropneusta and the Graptolithina), and the Chordata. The latter traditionally includes the subphyla Tunicata. The term “protochordates” has commonly been applied to all these taxa except the Echinodermata and Craniata. The Echinodermata, “protochordates” and Craniata supposedly share in common the deuterostomate condition (at least in recent taxa) whereby the gastropore of the embryo becomes the anus of the adult, and which shows a modified trisegmental body plan; and most possess gill-slits and a central axial structure, a notochord that provides some skeletal support. However, this “situation” exemplifies the difficulty of the problem of comparing an echinoderm, an enteropneust or graptolite, and a chordate, a real difficulty as these organisms exhibit very different morphologies. In other words, this difficulty deals with deep nodes of a cladogram, when the phylogeny is built for such a general or basic systematic question, where defining homologous features is fundamental.

The Echinodermata are deuterostomes, and the carpod echinoderms are considered by Jefferies (2001 and citations therein) and followers as closely related to craniates but few agree with his “calchordate” hypothesis. Amongst the hemichordates, both the

Enteropneusta and the Pterobranchia show a tripartite body plan and the latter possess a single pair of gill slits. The body plan of the long-extinct graptolites is unknown. None of these minor phyla possesses any significant resemblance to the conodonts; it is amongst the Chordata and especially Craniata that certain resemblances have been claimed.

The phylum Chordata has been diagnosed by the presence of characters such as a notochord, a dorsal hollow nerve cord, pharyngeal gill slits, segmented muscle blocks (myomeres) (Fig. 4B-E), and a post-anal tail. Of the modern members of the phylum, only the craniates possess a substantial fossil record because of the major preservational bias for apatitic hard tissues — bone, dentine and their possible precursors (for definitions of hard tissues and their development see Francillon-Vieillot *et al.* 1990). This lack of potentially basal chordate fossil material has proven a major obstacle in the search for chordate origins (cf. Blicek 1992). Indeed, Garstang (1928) based his theory of a paedomorphic origin solely on the embryology of extant chordates, and postulated that an organism similar to a tunicate or cephalochordate larva could have acquired sexual maturity without metamorphosing, thus providing a spring-board for the evolution of chordates and vertebrates.

The discovery of exceptionally preserved soft-bodied biotas in Konservat-Fossil-Lagerstätten has provided opportunities to examine and describe fossil lampreys and myxinoids from the mid-Palaeozoic (Janvier & Lund 1983; Bardack 1997; Poplin *et al.* 2001; Gess *et al.* 2006), and also purported fossil chordates from the Cambrian. Widely varying interpretations have been proposed for relationships of soft-bodied Cambrian taxa to living groups. *Emmonsaspis cambrensis* (Walcott, 1890) from the Lower Cambrian of Vermont has been allied with the graptolites, chordates, arthropods, and frond-like organisms since its initial description (Conway Morris 1993a, b). Even the most widely accepted earliest chordate, *Pikaia gracilens* Walcott, 1911, from the Middle Cambrian Burgess Shale, was originally interpreted as a polychaete annelid, and has since been allied with the cephalochordates based on synapomorphies such as chevron-shaped myomeres and an anteriorly extending notochord




FIG. 4. — Schematic frontal and cross sections of anterior end of a conodont animal compared with a cephalochordate and selected chordates. The left side of the animals shows dorsal surface features and muscle segmentation and the right side is cut away to show internal features at the level of the branchial region (if present): **A**, generalized conodont showing the position of conodont elements (association of structures based on Briggs *et al.* 1983, which is still highly imaginative with little reality beyond the general shape: note the diametrically opposed symmetry of apparatus to any vertebrate); **B**, amphioxus *Branchiostoma lanceolata* (Pallas, 1774); note that the notochord reaches into an anterior extension, but not the neural cord and not the dorsal “fin” (see 9) (after Bracegirdle & Miles 1978 and Parker & Haswell 1921); **C**, hagfish *Myxine glutinosa* Linnaeus, 1758 (after Dean 1899 and Marinelli & Strenger 1954); **D**, lamprey *Lampetra fluviatilis* (Linnaeus, 1758) (after Marinelli & Strenger 1956: figs 6, 10, 17, 36, 42); **E**, shark *Squalus* sp. (frontal section after Liem *et al.* 2001: fig. 18-2C). 1, muscular “cone”; 2, conodont apparatus (after Aldridge 1987: fig. 1.3 and Norby 1976); 3a, V-shaped myomeres; 3b, W-shaped myomeres; 4, medial tube; 5, notochord; 6, nerve cord; 7, velum (oral hood); 8, gill; 9, fin-ray box; 10, mouth; 11, pharynx; 12, branchial duct; 13a, nasal sacs; 13b, nasohypophysial opening (NB: not 13 for both as in Donoghue *et al.* 2000); 13c, nares; 14, hypophysial duct; 15, hypophysial sac; 16, gill arch; 17, eye; 18, mandibular arch with odontodes (teeth); 19, gill chamber; 20, external gill slit; 21, oesophagus; 22, spiracle; 23, brain; 24, anterior margin of pharynx; 25, hyoid arch.

(Conway Morris 1998), despite the presence of two anterior tentacles unknown in cephalochordates, but recently has again been excluded (Janvier 2003). With its two tentacles, it looks more like *Tullimonstrum gregarium* Richardson, 1966 (Pennsylvanian, Mazon Creek, Illinois), which has variously been compared to annelids, arthropods and molluscs (see Beall 1991; Conway Morris 1991, 1993a; Dzik 2000).

Discovery of older material from the Lower Cambrian Chengjiang Formation of Yunnan has led to finds such as *Yunnanozoon lividum* Chen *et al.* 1995 and *Cathaymyrus diadexus* Shu, Conway Morris and Zhang, 1996 (Shu *et al.* 1996a), with differently coloured features and impressions interpreted as a notochord, muscle blocks, and gill slits comparable with the preservation of soft tissues in fossil lampreys and myxinoids (and we would argue that the structures look more like gills than anything seen

in *Mayomyzon* Bardack & Zangerl, 1968 from the Pennsylvanian of Illinois, USA). *Yunnanozoon lividum* has also been considered to be a hemichordate (Shu *et al.* 1996b), or the most basal chordate (Chen *et al.* 1999), with *C. diadexus* as only a junior synonym of *Y. lividum* (Chen & Li, 1997). Still the search for chordate ancestors is anything but resolved. However, the deep dorsal body or fin of *Yunnanozoon* (M1 to M22 in Chen *et al.* 1995) does resemble the dorsal fin of the Permian cephalochordate from South Africa (Oelofsen & Loock 1981). In addition, with other possible early craniates from the same locality, such as *Myllokunmingia fengjiaoa* Shu, Zhang & Han, 1999 (= *Haikouichthys ercaicunensis* Luo, Hu & Shu, 1999; see Shimeld & Holland 2000: fig. 2) with gill pouches, the timing of chordate origins might be as early as Early Cambrian, or even Precambrian (Turner *et al.* 2004). As yet, however, there are no

Cambrian complete, mineralized, conodont or vertebrate body fossils, which contrasts with the “Cambrian explosion” when so many mineralized “invertebrate” taxa appeared (calcitic, phosphatic, siliceous, etc.).

Of course, we know that for phylogeneticists the problem of origin of any taxon cannot be linked absolutely to time. This is primarily concerned with systematics. However, when considering that the earliest claimed vertebrates (craniates) are possibly Early Cambrian in age [i.e. accepting the Chinese ones discussed above], this is solely based upon the known fossil record, and can therefore be used as a test of the phylogenetic scheme adopted.

Returning to modern chordate groups, all known chordates possess asymmetrically-organized internal organs, which are linked to control by proteins encoded by genes expressed on the left side of all known vertebrate embryos (Boormann & Shimeld 2002a, b). Raineri (2006), however, recently refuted the chordate affinities of the protochordates based on the development of their notochord and central nervous system on the ventral rather than dorsal side (indicating that they are gastroneurians, bilateralia with ventral neural cord), the muscular structure of their notochord, and lack of attachment of the axial musculature to the notochord. This is not a new idea (Arendt & Nübler-Jung 1994; Bergström 1996, 1997; Bergström *et al.* 1998) for the problem was already discussed in the 19th century. In any case, the definitive paper on chordate relationships based on whole-genome analyses of selected tunicates, a lancelet, and vertebrates (i.e. representatives of the three modern chordate groups) supported retention of cephalochordates in the deuterostomes, but basal to tunicates and chordates, with amphioxus (lancelets) as the most basal chordate (Putnam *et al.* 2008: fig. 2).

The craniates are characterized by presence of a neural crest, a notochord ventral to the neural cord (spinal cord) and additional characters connected with the brain (e.g., Janvier 2008). Reif (2006) reduced the chordates to craniates = hagfishes + vertebrates. Here we keep a conservative view of chordates with two basal groups, i.e. tunicates and cephalochordates, and a crown group, i.e. craniates including hagfishes and vertebrates.

## WHAT IS A VERTEBRATE?

What, then, are the characters that define a vertebrate? To quote Raineri (2006: 271) “the dawn of the vertebrates came into being when the dorsal ectoderm was turned into neural tissue on mesodermal induction”; or from Janvier (2003: 526), the diagnosis of the vertebrates is “based on two developmental characters [...] the presence of neural crests and epidermal placodes”. The resulting physical characters unique to vertebrates include “odontogenic tissues of the dermal skeleton and the branchial skeleton” and the “formation of the major vertebrate sensory organs, such as the olfactory, optic and otic capsules, and the lateral line system”. Following recent work on hagfish embryology by Ota *et al.* (2007), the neural crest character can be expanded to the presence of delaminating neural crest, previously thought to be a character of all vertebrates except hagfishes.

One of the sticking points in the debate for most people relates to what is a “true” vertebrate. Many conodont workers are using a very loose definition of what is a vertebrate compared to most vertebrate workers, backed up by the so-called “Total Group” Concept of Jefferies (Reif 2004 and see above). The debate can only continue when everyone involved agrees on the definition of a vertebrate (Janvier 2003; Reif 2006), and to promulgate understanding, this definition is communicated widely.

In general morphology, vertebrates possess a notochord, lying ventral to the central nervous system, at some stage in the life history, an extensive central nervous system, dermal placodes that develop into sense organs, the neural crest, and an exo- and endoskeleton. The primitive tripartite body plan is confined to the embryo and ontogeny provides a succession of new segments arising by subdivision of the second segment. Corresponding to this multiplication of segments are the muscular somites and the series of segmental gill slits. All possess a post-anal tail at some stage in their life history. In the development of their phosphatic hard tissues, vertebrates other than lampreys possess forms of dentine and bone (see further below).

## CONODONT ELEMENT HARD TISSUES VS VERTEBRATE TISSUES

Conodont elements exhibit exceptionally diverse histological structure (Müller 1981; Hall 1990), and variation, caused by intrinsic factors as well as the effects of diagenesis on individual elements can result in differing fine structure for elements of the same or closely related taxa. *Cordylodus* Pander, 1856 elements from Ordovician deposits at Sukrimagu in Tallinn, Estonia have no trace of the thin high-organic layers described in elements of this genus by Szaniawski & Bengtson (1993). The one *Cordylodus* specimen illustrated by Sansom *et al.* (1992) from the Lower Ordovician of Estonia reveals fewer details of the hard tissues in the element, although it does appear to have layers of hydroxyapatite arranged as radial crystals. Similar variation can be found in other genera of conodonts and in other parts of the conodont element. Elements of *Chirognathus* Branson & Mehl, 1933, also from the Harding Sandstone, are described as having scalloped growth lamellae, intersected by parallel tubules and occasional calcospherites in the basal body, which Smith *et al.* (1995: 310, 311, fig. 3A-C) considered to be indicative of vertebrate dentine. Elements of the same taxon from the same deposit have also been described as having growth lamellae with undulations of diagenetic origin, and the histological appearance of this specimen is given no special phylogenetic significance (Müller 1981: fig. 21.4).

Here we discuss some of the interpretations of conodont hard tissues, and give examples of those in and outside the vertebrate paradigm. Table 1 shows different interpretations of the tissues of the conodont elements. One major problem is lack of training in histology, which leads to misunderstanding of hard tissues. The intention of Gross (1954, 1957, 1960), a noted palaeohistologist, was to demonstrate how the conodont tissues differ from those of vertebrates (see Fig. 2A-C).

### Dentine

We consider that dentine is a prime hard tissue denoting a vertebrate (e.g., Turner *et al.* 2004). Donoghue *et al.* (2000, 2006) considered conodont basal bodies to be formed of dentine, but the globular tissue they

discuss cannot be homologous with vertebrate dentine, which grows centripetally (Gross 1954, 1957; Schultze 1996; Reif 2006; Fig. 2E), and is added basally not topically (Trotter *et al.* 2007). Even just considering the basal body structure, Carter & Lutz (1990: pl. 25, fig. D) illustrated the calcitic, lathic Regular Simple Prismatic (RSP) outer shell layer of the bivalve mollusc *Anomia simplex*, which looks more like dentine than anything in conodonts. Additionally, Dong *et al.* (2005) studied the basal tissue structure in “the earliest euconodonts, presumed to be the most plesiomorphic” (*Cambroistodus*, *Dasytodus*, *Granatodontus*, *Hirsutodontus*, *Proconodontus*, and *Teridontus*) showing a wide spectrum of variation in fabric from atubular lamellar, lamellar with “multiple point nucleation sites”, tubular to fibrous, none of which corresponds to dentine *per se* (see also Dzik 2009). Euconodont elements have no (pulp) cavity, hence not the slightest trace of blood and nerve supply (Fig. 2), nor any other sign that there was a living tissue such as dentine (Gross 1954; Schultze 1996; Reif 2006). This last point we consider most important. Interestingly, there seems to have been little research done on the development of the vertebrate odontode pulp system (pulp, pulp cavity) by molecular developmental biologists (e.g., Hall 2005).

Openings in the white matter of conodont elements are too small to have housed osteocytes (bone-forming cells) or odontoblasts (*contra* Sansom *et al.* 1992; Dzik 2009; see Schultze 1996). As noted above, the tissue material at the base (basal body) of the conodont element (when present, for generally the basal body is unknown, having been lost in most taxa in post-mortem taphonomic processes) does not have a structure consistent with any known (ortho)dentine (Kemp & Nicoll 1996; Fig. 2C-E).

### Enamel

Vertebrate enamel is a highly structured hard tissue, almost devoid of organic matter in the mature state, and containing small crystals of calcium hydroxyapatite (Carlson 1990; Warshawsky 1989). True enamel is found in sarcopterygians, but actinopterygians also possess enamel (ganoin) instead of the more usual collagen-based enameloid in chondrichthyans,

also a highly organized material (Carlson 1990) but formed below the basal membrane in teeth (Reif 2006: figs 4F, 5C, D, H). Dipnoans (lungfish: Kemp 1992, 2003; Satchell *et al.* 2000; Kemp & Barry 2006; Barry & Kemp 2007), like other sarcopterygians (Schultze 1969; Smith 1989), are among those “advanced” fish (i.e. basal to tetrapods) having true enamel in the dentition. Enamel (ganoine) of a similar protoprismatic form in developing stages is also found in the scales of primitive fossil and living actinopterygians like *Erpetoichthys calabaricus* (Smith, 1865) (Zylberberg *et al.* 1997). In these fish taxa, the early enamel resembles the initial stages of enamel formation in mammals (Satchell *et al.* 2000), although, at least in lungfish, it develops into protoprisms with a unique crystalline structure (Kemp & Barry 2006).

Thus, enamel in the dentition or scales of vertebrates (sarcopterygians and tetrapods), also arranged in layers but with a different morphogenesis (Reif 2006; see Fig. 2Db, E), has slender, elongate spicular crystals of calcium hydroxyapatite, perpendicular to and external to the basal membrane surface of the tooth or scale. The crystals are invariably arranged in specific ways, depending on the animal from which they came, and (despite Donoghue’s 2001 doubt) are oriented perpendicular to the surface of the tooth or scale (e.g., Sander 2000). The close association of the enamel with a dentine-enamel junction is also a distinctive character of vertebrate enamel (Fig. 2E). When a complete series of well-preserved sarcopterygian material covering developing and mature stages is examined, non-prismatic to prismatic enamel can be observed (Carlson 1990). In mammals, enamel prisms are highly ordered but patterns vary enormously among the different groups (e.g., Koenigswald 2000). In reptiles (Sander 1997, 2000) and amphibians, the enamel is less highly ordered.

Subsequent to the description of the hard tissue histology of conodont elements as exhibiting structures found in highly evolved vertebrates such as sharks and mammals (Sansom *et al.* 1992), conodont element fine structure was classified into three broad types, lamellar crown tissue, white matter, and basal tissue (Donoghue 1998; Donoghue & Chauffe 1998). These authors considered lamellar crown, or hyaline, tissue

to be homologous with vertebrate enamel, despite the large size of the component crystals (Donoghue 1998: 653), and the complete lack of any prismatic structure (Donoghue & Chauffe 1998). Variation in orientation of the crystals among different conodont genera was also considered unimportant. The lamellar crown tissue was described (Donoghue 1998: 655, 658) as distinct and separate from the white matter with which it interdigitates (forming the centre of serrations in conodont elements, see Gross 1954). However, these three characters, large crystal size, close association with the lateral hard tissue or white matter (Schultze 1996), and the lack of prismatic structure in the conodont hyaline tissue (Fig. 2B) indicate that the latter in conodont elements is not homologous to vertebrate enamel (Table 1). Lack of equivalence of the two tissues is emphasized when the numerous conodont elements with longitudinally or obliquely arranged mineral crystals, such as in *Panderodus* Ethington, 1959, are taken into consideration.

Notwithstanding the claims that conodont animals are vertebrates and the elements are true vertebrate teeth, transmission (TEM) and scanning electron microscopy (SEM) has shown that the mineralized component of the hyaline tissue of two Ordovician conodont taxa known only from elements, *Panderodus* and *Cordylodus*, consists of large, flat, oblong crystals, arranged in layers that run parallel to the long axis of the element. Within the layers in *Cordylodus*, crystals of hyaline tissue are positioned across the layer, perpendicular to the surface of the element. In *Panderodus*, the crystals are arranged obliquely or in line with the layer. The hydroxyapatite crystals in conodont hyaline tissue are exceptionally large, with no trace of prisms, unlike fish protoprismatic enamel, or the highly organized prismatic enamel of mammals (e.g., Kemp 2002a; Trotter & Eggins 2006; Trotter *et al.* 2007).

Light and scanning electron microscopy can provide conflicting evidence, even when the same taxon is used (Sansom *et al.* 1992; Szaniawski & Bengtson 1993). Some analyses indicate that the hyaline tissue of conodont elements cannot be enamel because it consists of bipartite layers (e.g., one Triassic conodont illustrated by Zhang *et al.* [1997] shows it clearly and they makes a point of it), not found in any



vertebrate enamel (Szaniawski & Bengtson 1993). The hyaline tissue is certainly high in organic matter, possibly remains of collagen (although doubted by Aldridge & Briggs 2009), also not a characteristic of vertebrate enamel (Kemp 1999, 2002a; Kemp & Nicoll 1996; Trotter & Eggers 2006).

#### *Bone, cartilage and other tissues*

Nevertheless, the hard tissues of conodont elements have been described as vertebrate, comparable to equivalent structures in sharks, other extinct fish and mammals (Barnes *et al.* 1973; Barskov *et al.* 1982; Sansom *et al.* 1992). The hyaline tissue of the element crown is described by them as radial crystallite enamel and the “albid tissue” (= white matter of others; Table 1) as bone with lamellae and osteocyte lacunae, the latter having canaliculi to house the cellular processes of the osteocytes. Depending on species, the single (rarely preserved but erroneously shown as multiple “growth cavities” in the histogenesis scheme of Donoghue 1998: fig. 9) basal body is alleged to include spheritic calcified cartilage, comparable to a similar tissue in sharks (Sansom *et al.* 1992) and in *Eriptychius* Walcott, 1892 from the Upper Ordovician Harding Sandstone (Smith *et al.* 1996), or mesodentine, as in certain younger fossil fish (Sansom *et al.* 1994; Donoghue 1998), or even lamellin (Dzik 2009). These comparisons based on superficial resemblances have been used to support the classification of conodonts among the vertebrates (Aldridge & Purnell 1996) although, as we here emphasize, this determination is not universally accepted (e.g., Kemp & Nicoll 1995; Kemp 2002a, b; Müller 1981; Schultze 1996; Walliser 1994; Table 1).

Similarly, the albid tissue cannot be bone because it contains no organic residues, and the spaces in the tissue in unaltered elements are too small to be osteocyte lacunae (Fähræus & Fähræus-von Ree 1994; Kemp & Nicoll 1995). True bone is mineralized and reacts in polarized light; Ca-phosphate is mineralized differently in enamel, dentine, bone and mineralized cartilage. The original work of Sansom *et al.* (1992) was based on material that can best be described as highly altered, so the identification of botryoidal mineralization in the basal body of *Cordylodus* elements as spheritic calcified cartilage

can be understood. The structures described in the *Cordylodus* elements bear no resemblance to spheritic calcified cartilage in Recent elasmobranch material (e.g., Francillion-Viellet *et al.* 1990; Dean & Summers 2006).

*Contra* the earlier work, Donoghue *et al.* (2006: 282) now disavow any bone in the conodont “oral skeleton”. Thus, removing the conodonts from the equation, even Donoghue *et al.* (2006) state that “true” enamel is only found in CG osteichthyans. Comparison of the ultrastructure of non-prismatic hyaline tissue in conodont elements and the organized enamel of vertebrates provides little support for a close phylogenetic relationship between vertebrates and conodonts. Trotter & Eggers (2006) and Trotter *et al.* (2007: 108) have recently shown that the large albid crystals of the crown tissue of a euconodont element are quite different from the fine crystalline tissue of dermal bone, dentine and enamel of vertebrates, contradicting specifically Sansom *et al.* (1992, 1994), Smith *et al.* (1996), Donoghue (1998), and Donoghue & Aldridge (2001). The crystals of hydroxyapatite in conodont hyaline tissue are large, with no trace of a prismatic arrangement, unlike the protoprismatic enamel of fish teeth and scales, or the highly organized prismatic enamel of mammal teeth. In addition, crystal arrangement in conodont hyaline tissue varies widely among conodont taxa (Wright 1990). Crystal arrangements similar to those of fish enamel are found in higher vertebrates, but none resembles, in any respect, any of the crystalline arrangements to be found in the hyaline tissue of conodont elements. All those findings on conodont vs vertebrate mineralized tissues support the arguments of Kemp (2002a) and Reif (2006) who refuted the arguments of Donoghue *et al.* (2000, 2003, 2006, 2008).

#### *Other tissue characters*

Conodont elements lack the colour range and lustre seen in vertebrate microfossils (e.g., Ørving 1973: fig. 3 and see cover of *The Australian Geologist Newsletter* 107, 1998). This consistent difference also refutes the presence of dentine with tubules in conodonts as these structures allow vertebrate fossil remains to infuse colour from the surrounding matrix. The conodont and microvertebrate colour indices are

therefore different although vertebrate material does respond to thermal changes, but with a different (as yet uncalibrated) set of gradations (Turner 1994b). Regarding co-occurring conodont elements and early vertebrate remains, another difference can be advocated. In her study of Silurian microvertebrate remains and conodonts from the Baltic Basin and from central Asia (Tuva, NW Mongolia, South Siberia), Z. Zigaite (pers. comm. to AB) found that thelodont tissues retain less  $^{18}\text{O}$  than conodonts, an outcome which gives a higher recalculated palaeotemperature of sea water based on dentinous thelodont scales than conodont elements, which is probably because of a major difference in the ultrastructure of the respective mineralized tissues. This trend is increased when a stronger diagenetic alteration occurred, as in Tuva where most vertebrate microremains (thelodonts, mongolepids, acanthodians) are whitish and give much too high palaeotemperatures (of *c.* 50°C!).

Turner & Blicek (1995) also considered conodont vs vertebrate micro-ornaments. Although the imprinting of external cells of overlying soft tissue on (dermal) hard parts is possible in many animals, the patterns seen in many conodont elements are different from those in vertebrate scales and teeth. These impressions were used as an argument for conodonts being vertebrates by Simonetta *et al.* (1999) but, as Reif (2006: 418) also showed, cell impressions are not exclusive to the surface of enamel nor can they be used as conclusive identification for enamel in conodonts. For comparison, Märss (2006) reviewed micro-ornaments in a wide variety of vertebrate scales with surficial enamel, enameloid, and dentine.

To summarize, there is so much evidence that conodont elements are not and have nothing to do with teeth that it is not even a question of whether they can be vertebrate. There is no pulp cavity in highly evolved conodonts, even if the basal body could be regarded as such in early conodonts (Müller & Hinz-Schallreuter [1998] reckoned that the latter appeared as an evolutionary novelty within the euconodonts). Schultze (1996), Reif (2006), and others cited here, have shown that the hard tissues in euconodont elements do not have the morphogenetic history or structure to be interpreted as vertebrate, let alone teeth.

## RELATIONSHIPS OF CONODONT'S BASED ON MORPHOLOGY OF CRANIATES AND "PROTOCHORDATES"

As noted above, cephalochordates and craniates share a well-developed system of somites; nevertheless, Raineri (2006) considered this a convergence and excluded cephalochordates from the deuterostomes. Thus, in considering cephalochordates, fossil possibilities include *Yunnanozoon* (= *Haikouella*), the Permian *Palaeobranchiostoma* Oelofsen & Loock, 1981, and the mid-Cambrian Burgess Shale *Pikaia* Walcott, 1911, which is still not fully accepted. This discussion of cephalochordate structure is therefore based mainly on a handful of Recent species belonging to two families (at most) and to a single order, exemplified by *Branchiostoma lanceolata* (Costa 1834) (Fig. 4B).

Craniates and *Branchiostoma* share a number of features, some of which have a functional origin in locomotion. Chief among these are the skeletal, muscular and nervous elements, specifically the notochord (*Branchiostoma* lives in the sediment with the notochord on the ventral side, therefore the discussion as to where "dorsal" and "ventral" is in *Branchiostoma*, see above), the hollow dorsal nerve cord, and the organisation of the musculature into folded segmental somitic blocks. In both groups during ontogeny each of the three gill-slits on each side folds into a U shape, with the fold dorsal in position (Fig. 4). Conodonts and craniates differ in the following major ways.

### *Cephalization and eyes*

Protochordates in general, including cephalochordates, show little sign of cephalization. Accepting the interpretation of the Cambrian *Yunnanozoon* from China as a cephalochordate indicates that the cephalization in early cephalochordates (Mallatt & Chen 2003) was further developed than in the extant *Branchiostoma*. By contrast, the higher craniates show perhaps the greatest level of cephalization of any animals, involving as many as nine segments (Balfour 1877). It should be noted, however, that cephalization is not restricted to a single monophyletic group. In any mobile animal, that part which meets the environment first – the

anterior end, is likely to develop a concentration of sense organs. The pressures towards cephalization are thus present in all mobile groups, viz., shrimps and worms, snails, frogs, dogs, etc. All Recent craniates, including hagfishes and lampreys (Fig. 4), have a “new head” comprising organs produced by delamination and migration of neural crest cells; this process has only recently been demonstrated in hagfishes (Ota *et al.* 2007).

Conodont body fossils show little sign of cephalization except for the possible anterior internal oral/branchial feeding apparatus (Fig. 4A; see discussion in Nicoll 1995) but no nasal or hypophysial openings are known, unlike vertebrates (cf. Fig. 1F, H). Again, as emphasized by others (e.g., Walliser 1994, and pers. comm. 2009), the symmetry and operational movement of the conodont elements within the apparatus is at 90° to that of any vertebrate tooth array or branchial system, thus again mitigating against their being either true vertebrate teeth or vertebrate *per se*. The level of cephalization in this regard is not higher than the oral hood of *Branchiostoma* (Fig. 4B). Briggs *et al.* (1983) identified the large paired dark stains at the anterior end of the Granton conodont *Clydagnathus?* cf. *cavusformis* (*C. windsorensis* (Globensky, 1967)) as eyes (Fig. 1A, D, F). Subsequently, Gabbott *et al.* (1995) and Purnell (1995b) interpreted muscle fibres in a similar position on the giant conodont *Promissum pulchrum* Kovács-Endrődy, 1987 from the Soom Shale (Cedarberg Formation, latest Ordovician [latest Hirnantian]-earliest Silurian [earliest Rhuddanian], Table Mountains, South Africa; Vandenbroucke *et al.* 2009) as extrinsic eye muscles. These structures, preserved as semicircular or somewhat rhombic bands/sheets (Donoghue *et al.* 2000: fig. 4G), are unlike those of any known vertebrate eye, appearing to be the remnants of muscular half-rings or cones rather than discrete eye muscles (see also Reif 2006). Trunk and tail myomeres (muscle blocks) of conodonts are V-shaped (Fig. 1A) and not W-shaped as in all craniates including hagfish and lamprey (see e.g., Pridmore *et al.* 1997: fig. 4). The V-shape of the myomeres in the South African conodonts are highlighted by post-mortem shrinkage, but by comparison the purported eye muscles appear to be broad sheets. Even the shape of the

“eye” stains of *Clydagnathus?* are unusual, being preserved as flattened cones with the apices meeting medially (Aldridge 1987: fig. 1.9B), and thus not comparable to vertebrate sclerotic capsules or eyes and eye muscles. Are these paired structures actually lateral, or could they have been dorsal and ventral? When Briggs (2003) noted that “The toothlike elements [...] are consistently preserved to one side of the head”, he presumably inferred the orientation from the position of the “eyes”. Even if, unlikely as it seems, these structures were eyes or eyespots, the tunicate Larvacea and larval Ascidiacea also possess eyespots.

This feature, then, is inadequate for distinguishing between craniate and protochordate affinities for conodonts. For comparison, Janvier & Arsenault (2007) distinguished lateral and median stains without identifying the lateral stains as eyes at the correct position even in the indubitable craniate *Euphanerops* Woodward, 1900.

#### *Anterior end of notochord*

Adult *Branchiostoma* have a notochord that extends into the preoral region, to the anterior tip of the animal (Fig. 4B). In all other protochordates and in all craniates the notochord (when present) is confined to that posterior portion of the body behind the hypophysis or its homologue (Huxley 1858; Carlisle 1953). This is also the anterior termination of the notochord in embryonic *Branchiostoma*, and the anterior extension develops only later in ontogeny (Berrill 1987). It thus appears to be a secondary development, which may indeed be confined to the single order Branchiostomatoidea, the order that includes all Recent species, whereas the supposed Early Cambrian cephalochordate *Yunnanozoon* (= *Haikouella*) is comparable to craniates in cephalization and a notochord reaching to the hypophysial region (Mallatt & Chen 2003). In ontogeny the notochord develops by pinching off the dorsal region of the gut (the archenteron), and is thus confined to the post-oral region behind the hypophysis. From a functional point of view the anterior extension, which follows in *Branchiostoma*, provides a stiffening, which might originally have been an important adaptation to burrowing. It might also have impeded cephalization in branchi-

ostomatoids (and also in *Palaeobranchiostoma*), and in any cephalochordates possessing this adaptation, but not in Early Cambrian *Yunnanozoon*. Raineri (2006), however, has countered homology of the notochord in *Branchiostoma* and craniates.

There is no certain evidence for a notochord in conodont body fossils. Briggs (2003: 277) stated that “Paired axial lines that run the length of the trunk might represent the gut or the notochord.” If this structure is a notochord, then its extent conforms to the general condition found in Tunicata and Craniata, not the adult branchiostomatoidean condition. We have argued above that forward growth of the notochord is a specialized adaptation to a burrowing habit, and may not be of more than ordinal value as a distinguishing feature. There is every reason to think that it was not characteristic of early cephalochordates (*Yunnanozoon*; Mallatt & Chen 2003), any more than it is found in any other group of protochordates.

#### *Skeletal elements*

Vertebrates possess skeletal elements in addition to the notochord. Apart from (internal) viscerocrania, these comprise first, the segmentally-arranged paraxial elements (of bone or cartilage), which later give rise to such structures as the vertebrae and ribs; and, second, dermal elements, consisting primarily of bone and dentine, and forming scales, teeth and fin rays (thus excluding hagfishes). In nearly all known early vertebrates, these odontodes form an exoskeleton but can also be found lining the internal surface of the mouth to the pharynx, a feature retained in many living fishes (e.g., Reif 2002; Märss *et al.* 2007). Reif (2006 and references therein: e.g., Fig. 2E) has discussed the errors of Donoghue’s (1998) interpretation of conodont morphogenesis providing clear morphogenetic diagrams for vertebrate odontodes and showing how the structure and growth of conodont elements does not match the Odontode Regulation Theory (Reif 2002) in any way. As noted above, Trotter *et al.* (2007) also showed that conodont element tissues are clearly distinct in crystal size from any vertebrate tissue. There is also no mineralized keratin in conodont elements, and so affinities with rasping teeth of lampreys and hagfishes are also very unlikely.

Despite a few older claimed records, definite “fish” scales *per se* (i.e. non-conodont elements) first appear in the fossil record in the Early Ordovician (Turner *et al.* 2004; Young 1997, 2009). Lampreys possess the paraxial elements but lack dermal elements (see also Fin-rays section). Their possession of endoskeletal fin rays is not evidence of an ancestor possessing a dermal skeleton. Hagfishes possess a caudal cartilage with cartilaginous rays (Retzius 1892; homology to radial or fin rays uncertain). *Branchiostoma* lacks both paraxial dermal and endoskeletal elements.

Conodont body fossils also lack any trace of paraxial, dermal or endoskeletal elements (see conclusions). Nevertheless, elements in the caudal region were compared with those in hagfish by Janvier (1998). Samples of disjunct conodont elements from throughout the stratigraphic range of the group show no evidence of skeletal elements in the organism other than those of an anterior feeding apparatus. *Conopiscius* Briggs & Clarkson, 1987, found in the Carboniferous Granton Shrimp Beds with *Clydagnathus?*, possibly had mineralized scales associated with its V-shaped myomeres, and was claimed as an agnathan; Dzik (2009) recently asserted a conodont affinity for *Conopiscius*, but its relationships, and the presence of scales, are still uncertain.

#### *Folding of the muscular somites*

Each muscle segment of *Branchiostoma* is folded into a V shape, with the angle of the V directed forward. All craniates, including hagfishes, lampreys and gnathostomes, and even the earliest fossil fish, in which the structure can be distinguished, show a more elaborate folding, into a W shape. In effect, the dorsal and ventral wings of the W-shaped muscle block provide a separately controllable musculature for the compressed dorsal and ventral body margins and for the median fins, where present. The V-shaped pattern could be interpreted as another indication of a convergent evolution of free-moving animals compared to craniates (Raineri 2006).

Conodont body fossils show V-shaped folding with the angle of the V directed forward (Fig. 1A). This, indeed, forms the basis of one of the argu-

ments for their chordate nature (Donoghue *et al.* 2000). They lack any anteriorly-directed refolding of the wings of the V to form a W shape. In this they are compatible with a cephalochordate condition, not with craniates. However, whereas young vertebrate embryos have V-shaped myotomes (many examples in Moser *et al.* 1984), this may be interpreted as a plesiomorphous state and does not help with conodont relationships.

#### *Fin rays*

With the exception of hagfishes, the dermal median fins of craniates are supported by endoskeletal fin radials, which are articulated at the base and supplied with a musculature derived from the forwardly-reflected wings of the W-shaped somites. The fins of *Branchiostoma*, in contrast, are supported by passive non-segmental box-like structures, which lack musculature or basal articulation and provide merely a stiffening (Fig. 4B).

The median fins are clearly supported by some kind of “fin rays” in conodont body fossils. These “fin rays”, however, seem not to correspond to the muscular somites and lack any trace of basal articulation or musculature. Indeed, with simple V-folding of the somites the basis for any fin-ray musculature is lacking. Accordingly, the conodont fin rays are more like the *Branchiostoma* box rays than the fin radials of craniates.

#### *Gill slits*

Cephalochordates possess U-shaped gill slits, which form two slits or openings that are homologous to one primary gill slit of a craniate. Despite the many conodont animals having been studied, only one is reported to show structures that have been interpreted, very tentatively, as four possible gill pouches (Briggs *et al.* 1983: fig. 3A; Donoghue *et al.* 2000: fig. 3C). However, we cannot identify these structures in those published figures. Considering that eyes, eye muscles, myomeres, notochord and caudal fin rays have supposedly been identified, it seems unlikely that gill structures would not also have been preserved in conodont specimens, if they were actually present. By comparison, they are present in the Chinese Cambrian chordates (e.g., Shu *et al.* 2003) and fossil hagfishes and lampreys.

Sansom *et al.* (2010) proposed a new approach in experimental taphonomy of basal and early chordates in order to constrain the interpretations of their soft-bodied fossil representatives, and consequently to improve the analysis of the phylogenetic relationships. They thus focussed on individual character changes dependent on decomposition stages rather than on features of whole organisms. This way of analysing fossils is certainly interesting and promising. However, Sansom *et al.* (2010) published only a limited decay study based on only three specimens each of *Branchiostoma* and *Ammocoetus* (= *Lampetra larva*). [Three may be considered as statistically weak, but we can surmise that this is only a preliminary study.] They let the specimens rot without sediment cover, which would have protected the decaying specimen in most natural cases. Soft tissue preservation (except impregnated soft tissue) requires immediate cover and in addition special conditions within the sediment. Even with such conditions, each organism reacts differently, e.g., fat content is different from group to group, etc. (e.g., Schäfer 1972). Sansom *et al.* (2010: fig. 3) presented a sequence of resulting decay events on a simple tree for both species. They established five decay stages for the two specimens, from the complete specimen to a stage with notochord and some indication of muscle myomeres. The gill basket resists decay to a late stage – in *Branchiostoma* to stage 4 (their “stem chordate”) and in *Lampetra* to stage 3 (their “stem vertebrate”). They transferred their interpretations onto a deuterostome phylogeny (Sansom *et al.* 2010: fig. 1a) to show the possible position of several chordate and vertebrate fossils showing exceptional (soft tissue) preservation, which have been subjected to varying interpretations and placement in phylogenies.

One has to distinguish here between different interpretations of structures and the decay process. The yunnanozoans are extremely well preserved even with buccal tentacles, and thus show no decay comparable with Sansom *et al.*'s (2010: fig. 2) decay sequence and so their different placement in phylogenies are actually differences in interpretation (in contrast to Briggs 2010). In

reality, one can only be certain of the original form of a decay or fossilization process if the original form is found, as in the case of scaumenallization (Béland & Arsenault 1985). Sansom *et al.* (2010) omitted conodonts. Conodont fossils have no gills although supposed eyes are well preserved (decay stage 2 of larval *Lampetra*). Based on the decay schedules given by Sansom *et al.* (2010), conodonts would not even fit their “stem chordate” stage. That supports our conclusion on other reasoning presented here that conodonts were not even chordates.

### Pathology

Pathological factors often illuminate morphogenesis. Hass (1941: pl. 13; figs 4, 5), writing on conodont element morphology, considered rejuvenation of injured parts; he noted that it is mostly thinner extremities that are broken away, and that the element can be rejuvenated or rebuilt, although the new growth axes do not always align with the old, and that there can be several restorations. Hass (1941) also referred to an observation by Furnish (1938: pl. 41 fig. 31) on a partly regenerated Early Ordovician specimen of *Drepanodus subarcuatus*: “Since the cusp is thin and blade-like, most specimens are broken and many individuals show evidence of replacement in the apical portion”; and also discussed “Suppression of parts” [presumably cf. “suppressed denticles” per various glossaries] where he indicated that growth axes are suppressed during growth mostly by lack of room. These elements do show mode of growth, but it is quite different from that of vertebrate odontodes (cf. Reif 1982, 2002). Lindström (1964: fig. 3C) gave a good example of a thin section through a regenerated break, and he also discussed the process of regrowth at length (see also Gross 1954; Fig. 2B). Weddige (1990) documented numerous abnormalities from developmental and traumatic causes, and gives sufficient detail to show that conodonts have no pathologies that relate them to equivalent anomalies in the dentition of lower vertebrate hard tissues (e.g., Reif 1984); consequences of trauma and disease in vertebrate hard parts differ significantly from equivalent accidents in conodonts, in that they are generally not repaired.

## THE NEW CLADISTIC ANALYSIS

It has now been accepted for some time that what is still informally called “agnathan fishes” corresponds to a paraphyletic grouping, incorporating both extant (cyclostomes) and extinct (ostracoderms) taxa. The phylogenetic relationships of the parts of this group (myxinioids, lampreys, pteraspidomorphs, anaspids, osteostracans, galeaspids, pituriaspids, thelodonts) are still currently discussed, and cannot be considered as resolved (see e.g., the various proposals by Janvier 1981, 1997, 1998, 2001, 2003, 2006, 2007a, b, 2008, 2009; Märss *et al.* 2007). However, this paraphyletic group is at the crux of vertebrate evolution, especially regarding the origin of the head and neural crest-derived tissue (Northcutt 1996). In contrast to the living *Branchiostoma* (Holland & Holland 2001), extant and extinct lampreys (e.g., Hardisty & Potter 1972) and all other vertebrates (e.g., Janvier 2008) possess a complex brain and placodes that contribute to well-developed eyes, as well as auditory and olfactory systems, i.e. they are craniates. These sensory systems were arguably a trigger to subsequent vertebrate diversifications. However, although these systems are known from skeletal forms and other impressions in agnathans (e.g., Märss *et al.* 2007; Janvier 2008), the vertebrate structures identified in the Early Cambrian *Myllokunmingia* (= *Haikouichthys*) from the Chengjiang Fossil-Lagerstätte are doubted, despite Shu *et al.*'s (1999, 2003) and Zhang *et al.*'s (2001) claims to the contrary. Although *Myllokunmingia* resembles somewhat the ammocoete larva of modern lampreys, there is no evidence of vertebrate hard tissues nor of a brain; nevertheless, a purported branchial system with gill pouches is present. Chengjiang fossils are preserved as coloured stains, indentations and impressions on the rock matrix, as is the case with fossil soft-bodied hagfishes and lampreys. Here we include them in two of our analyses, coded in accordance with these structures being correctly identified by the original authors, but have also processed the data with only the taxa used in Donoghue *et al.*'s (2000) original matrix. One change was made to their taxa, by our nominating the species for the thelodont genus *Loganellia*, viz. *L. scotica*, as some

characters code differently for other thelodont taxa (cf. Märss *et al.* 2007).

Donoghue *et al.*'s (2000) analysis is based on their *a priori* assumption that euconodonts are chordates. As noted by Janvier (2003: 526), "The position of euconodonts [...] in most current phylogenies is largely imposed by assumptions [our emphasis] about the presence of certain characters such as extrinsic [*sic*] eye muscles or gills". Despite being unconvinced that gills are present in conodonts, we have kept the gill-character codings used by Donoghue *et al.* (2000), even though we regard the "?" state as being nonapplicable rather than unknown for conodonts.

Despite the various pros and cons of different approaches to taxon sampling and character coding, as discussed by Donoghue *et al.* (2000), we have followed their methods, taxa and characters as closely as possible. The main revision to the data matrix of Donoghue *et al.* (2000) is our deletion of the physiological and other characters (6, 9-13, 24, 33, 36, 39, 42, 53, 56, 57, 83-103) that are not known for any fossil taxa (euconodont, agnathan or gnathostome; see Appendix 1). Otherwise, we have minimized changes to the original characters, altering their character 27, pouch-shaped gills, to our character 20, gill opening shape: 0-, simple slit; 1-, pore; 2-, slit opening to chamber, so that tunicates and jawed fishes code differently, and altering their character 41, large lateral head vein, to our character 31, lateral head vein. In this new analysis, character codings have been revised as detailed in Appendix 2. The character matrix (Table 2) incorporates revisions based on recent publications on several taxa; in particular the new description of *Euphanerops* (Janvier & Arsénault 2007) led to changes in codings for characters 6 (7), 21 (28), 25 (32), 26 (34), 29 (38), 33 (44), 36 (47), 37 (48), 39 (50), 42 (54), 43 (55), 44 (58), 48-60 (62-74), 66 (80), and 67 (81). Donoghue *et al.* (2000) used PAUP 3.1.1 for their parsimony analysis, whereas we used the updated PAUP 4.0b10 program for Windows (Swofford 2002), while also using equal-weight, unordered multistate characters, and branch-and-bound tree-building routine (i.e. heuristic search), but with a data matrix of just 68 of the original 103 characters.

The article by Donoghue *et al.* (2000) included cladistic analyses of chordates (including conodonts, with the *a priori* assumption that they are chordates) that incorporated physiological as well as morphological and histological characters. In their cladogram and even in their preferred trees (Donoghue *et al.* 2000: fig. 14a: ACCTAN [= accelerated character state transformation], fig. 14b: DELTRAN [= delayed character state transformation]), the position of conodonts is poorly supported as they sit above a node that is characterized by 45 synapomorphies in their fig. 14a (39 in their fig. 14b) above the node Craniata with Cephalochordata, of which conodonts have only seven codable characters (1, 2, 19, 28, 46, 51 and 65), i.e. only 16% (*ibid.*, fig. 14b: 18%); thus 84% (*ibid.*, fig. 14b: 82%) are missing or inapplicable. In addition, only two of the 20 homoplasies are present in conodonts, i.e. only 10% (Donoghue *et al.* 2000: fig. 14b: 14%) present; a third homoplasy (character 50 = preanal median fold) below the node is even coded 0 for conodonts. This is both an illogical and unexplainable position for conodonts. Nevertheless, these authors dismissed all other contemporary views of conodont relationships as virtually unscientific because they did not include a "numerical cladistic analysis". Their approach, we contend, was actually based on a preferential data set, with a near-complete data matrix only being possible for extant rather than fossil taxa. Their main analysis resulted in the conclusion that conodonts "are the most plesiomorphic member of the total group [our emphasis] Gnathostomata" (Donoghue *et al.* 2000: 191; also Gess *et al.* 2006). This leads to a semantic inconsistency (see discussion above) because "gnathostomes" are defined as jawed vertebrates with teeth; as noted above, a "stem" gnathostome without teeth therefore cannot be a gnathostome and be within the "total group" and conodont elements are therefore not homologous with vertebrate teeth (e.g., Gross 1954; Schultze 1996; Kemp 2002a; Reif 2006). Reif (2004) also discussed the misunderstanding of Hennig's usage of "stem group" and the "Total Group Concept" of Jefferies (1979) showing that Hennig never intended his method to be extended back in time.

TABLE 2. — Data matrix for the 17 original taxa used by Donoghue *et al.* (2000: table 1) plus *Myllokunmingia* and *Yunnanozoon*. “?” applies to both inapplicable and unknown codings. Multiple state character codings are unordered, the default “MSTaxa = uncertain” (rather than “polymorph” or “variable”) was used.

Our character nos.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25
Donoghue <i>et al.</i> character nos.	1	2	3	4	5	7	8	14	15	16	17	18	19	20	21	22	23	25	26	27	28	29	30	31	32
Tunicata	0	?	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	?	?	?	?	0
Cephalochordata	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Myxinoidea	1	1	0	0	1	1	0	2	1	1	1	1	0	1	0	0	0	1	1	1	0	0	0\1	0	0\1
Petromyzontida	1	1	0	2	1	1	1	2	1	1	2	1	1	2	0	0	1	2	1	1	1	1	1	1	0
Heterostraci	1	1	1	1	?	1	1	1	0	0	?	1	?	2	1	0	1	2	2	1	1	1	0	0	0
Astraspis	1	1	?	?	?	?	?	?	?	?	?	?	?	?	?	?	0	1	1	1	1	1	1	1	0
Eriptychius	1	1	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Arandaspida	1	1	?	2	?	1	?	?	?	?	?	?	?	?	?	?	?	?	2	1	1	1	0	1	0
Anaspida	1	1	?	2	?	1	?	2	1	1	2	?	?	?	?	?	?	?	1	2	1	1	1	0	1
Jamoytius	1	1	?	?	?	?	?	2	1	1	1	?	?	?	?	?	?	?	?	?	1	1	0	1	0
Euphanerops	1	1	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Osteostraci	1	1	0	2	1	?	1	2	1	1	2	2	1	2	1	1	1	2	2	1	1	1	1	0	1
Galeaspida	1	1	1	2	?	1	1	1	0	1	2	1	1	2	1	1	1	2	2	1	1	1	0	1	0
Loganellia scotica	1	1	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	1	2	1\2	1	1	1	1	0
Pituriaspida	1	1	?	?	?	?	1	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Jawed vertebrates	1	1	1	1\2	1	1	1	1	0	0	0	1	1	3	1	1	1	2	1\2	2	1	1	1	0	0
Conodonta	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
Yunnanozoon/Haikouella	0	1	?	?	?	?	0	?	0	0	0	?	?	?	?	?	?	?	?	?	0	1	1	0\1	0
Myllokunmingia/Haikouichthys	1	1	?	?	?	?	0	2	?	1	1	?	?	?	?	?	?	?	?	?	1	1	1	0\1	0

Our character nos.	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48
Donoghue <i>et al.</i> character nos.	34	35	37	38	40	41	43	44	45	46	47	48	49	50	51	52	54	55	58	59	60	61	62
Tunicata	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Cephalochordata	0	0	0	0	1	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0
Myxinoidea	0	0	0	0	1	1	1	0\1	0	0	0	0	1	1	1	1	0	0	0	0	0	0	0
Petromyzontida	1	0	1	1	0	1	0	1	1	0	0	1	0	1	1	1	1	1	1	0	0	1	0
Heterostraci	1	0	?	?	?	?	?	?	0	0	?	0	0	0	1	1	?	?	1	?	?	?	0
Astraspis	?	?	?	?	?	?	1	0	0	?	0	0	?	?	?	?	?	?	?	?	?	?	1
Eriptychius	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	1
Arandaspida	?	?	0	?	?	?	?	?	0	0	?	0	0	1	0	1	?	?	?	?	?	?	0
Anaspida	?	0	?	?	?	?	?	0	0	1	1	1	0	1	0	1	?	?	?	?	?	?	1
Jamoytius	?	1	?	?	?	?	?	?	?	?	?	0	0	?	0	1	?	1	?	?	?	?	0
Euphanerops	1	0	?	1	?	?	?	?	0	1	1	1	0	1	0	1	?	0	1	0	?	?	1
Osteostraci	1	1	1	1	0	1	1	1	0	1	0	1	2	0	1	1	?	?	1	1	1	1	0
Galeaspida	1	1	?	?	?	?	1	1	0	0	?	?	0	1	1	?	?	1	1	1	1	1	0
Loganellia scotica	?	0	?	?	?	?	?	0	1	1	1	1	0	1	0	1	?	?	?	?	?	?	1
Pituriaspida	?	1	?	?	?	?	?	?	?	?	?	?	1	?	?	?	?	?	?	1	1	1	?
Jawed vertebrates	1	0	1	1	1	1	0\1	1	1	1	0\1	1	2	0	1	0	0	1	1	1	1	1	0
Conodonta	?	?	?	?	?	?	?	?	0	0	?	0	0	1	0	1	0	0	0	?	?	?	0
Yunnanozoon/Haikouella	1	0	?	?	?	?	?	?	0	0	?	0	0	0	0	0	0	0	0	0	0	?	?
Myllokunmingia/Haikouichthys	1	0	?	?	?	?	?	?	1	?	1	1	0	?	0	?	0	?	1	0	0	?	?

Donoghue *et al.* (2000) included numerous variations of their main analysis, which they used to illustrate “worst case” scenarios by leaving out all characters for which fossil taxa coded “?”, and changing the conodont coding to 0 for several contentious characters (eye muscles, histology). Their “worst” result showed conodonts as more

derived (crownward) than hagfishes (Donoghue *et al.* 2000: fig. 11D). We present here a review of the morphological and histological characters, which leads us to consider that conodonts are neither vertebrates nor craniates. Using these characters, we present a new cladistic analysis that counters the conclusions of the Donoghue *et al.*'s



TABLE 2. — Continuation.

Our character nos.	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68
Donoghue <i>et al.</i> character nos.	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82
Tunicata	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Cephalochordata	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Myxinoidea	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Petromyzontida	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Heterostraci	0	0	1	1	1	0	2	0	1	1	2	1 $\frac{1}{2}$	1	1	0	2	1	0	0	0
Astraspis	0	0	1	1	1	0	2	2	1	0	2	1	1	?	?	1	0	?	?	0
Eriptychius	?	1	1	1	1	0	2	2	1	0	2	1	0	?	?	1	0	?	?	?
Arandaspida	?	?	1	1	1	0	0	0	1	1	2	2	1	0	2	1	?	?	?	1
Anaspida	0	0	1	0	1	0	0	0	0	0	2	2	0	1	0	1	0	0	0	0
Jamoytius	?	?	1	?	?	?	?	?	?	?	?	2	0	0	0	0	0	0	0	0
Euphanerops	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0
Osteostraci	1	1	1	0	0	1	1	0	1	0	2	1	0	1	0	2	0	1	1	1
Galeaspida	1	1	1	0	1	0	0	0	0	0	1	1	0	1	0	2	0	1	1	0
Loganellia scotica	0	0	1	0	?	0	1	0	0	0	1	1	0	0	1	1	0	0	?	0
Pituriaspida	1	?	1	?	?	?	?	?	?	?	?	?	?	?	?	2	1	1	?	?
Jawed vertebrates	1	1	1	0	0	1	1	1	1	0	2	1	0	0	1	1	0	0	1	1
Conodonta	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Yunnanozoon/Haikouella	0	0	0	0	0	0	0	0	0	0	0	?	0	0	0	0	0	0	0	0
Myllokunmingia/Haikouichthys	0	?	0	0	0	0	0	0	0	0	0	?	0	0	0	0	0	0	0	?

(2000) analysis. However, our main criticisms of the claims of the “British School” are based on sound morphological and histological arguments that really do not rely on a computer.

Before giving the results of our new phylogenetic analysis, we want to mention three recent analyses that resulted in differing conodont-vertebrate relationships: 1) conodonts with “?” in polytomy with lampreys, etc.: (myllokunmingiids (hagfishes, lampreys, ? *euconodonts*, *Euphanerops* (*anaspids* ((*arandaspids*, *astraspids*, *heterostracans*) (*thelodonts* (*galeaspids* (*pteraspids*, *osteostracans*, jawed vertebrates)))))) (Janvier 2007a: fig. 1.13). In this analysis, Janvier (2007a: 32) considered that euconodonts are “regarded by many paleontologists as the basalmost stem gnathostomes”, he cited only Schultze’s (1996) paper with opposing opinion, and continued: “The phylogenetic position of euconodonts as stem gnathostomes remains tenuously supported, and they may turn out to be either more closely related to hagfishes or lampreys (or cyclostomes as a whole), or even stem vertebrates.” (for a further opinion, see Janvier 2008); 2) conodonts with “?” in polytomy with anaspids etc.: (cephalochordates (*Myllokunmingiida* (hagfishes, lam-

preys, ? *Euphaneropidae* (? *Euconodonts*, *Anaspida*, (*Arandaspida*, *Astraspida*, *Heterostraci*), *Thelodonti* (*Galeaspida* (*Pituriaspida*, *Osteostraci* (*Placodermi* (*Chondrichthyes* (*Acanthodii*, *Osteichthyes*)))))) (Janvier 2007b: fig. 2.3), with the relationships of euconodonts to or within vertebrates still debated by this author (Janvier 2007b: 65): “Euconodonts share with crown-group vertebrates the presence of median fin radials and are best placed as stem gnathostomes, notably on the basis of their ability to develop mineralized skeletal elements made of apatite”; and Janvier (2008) who places the conodonts in his cladogram with a question mark, and a polytomy; and 3) conodonts below (more stem-ward of) craniates (our position): (*Cephalochordata* (*Haikouella* (*Conodonts* (*Myxinoidea* (*Petromyzontida* (*Heterostraci* (*Anaspida* (*Thelodontida* (*Osteostraci* (*Placodermi* (*Chondrichthyes* (*Acanthodii*, *Osteichthyes*)))))))))) (Wilson *et al.* 2007: fig. 3.1).

Since our paper was proposed for publication, a series of papers appeared in various books and journals of biology and/or palaeontology (Janvier 2008, 2009; Donoghue *et al.* 2008; Koentges 2008; Paris *et al.* 2008; Aldridge & Briggs 2009; Huysseune *et al.* 2009; Sire *et al.* 2009). These authors




FIG. 5. — Cladograms generated by PAUP 4.01b10 for Windows (Swofford 2002), data (see Table 2) compiled with NEXUS Data Editor (Page 1999), using 68 unordered, equal weight characters, heuristic search, starting trees obtained by stepwise addition with random addition sequence, tree-bisection-reconnection branch swapping; trees generated using Treeview X (Page 1996): **A-C**, using the original 17 taxa, characters of Donoghue *et al.* (2000) recoded (see Appendices 1; 2); Consistency index (CI) = 0.5971, Homoplasy index (HI) = 0.4029, and Retention index (RI) = 0.6940 (see Appendix 4 for CI, HI and RI values for the single characters); 27 shortest trees of equal length = 139 steps, with numbered nodes (cf. apomorphy lists, Appendix 3) for the trees illustrated and 50% majority rule bootstrap values given in italics (under node number) at nodes supported by the bootstrap analyses; **A**, ACCTRAN character-state optimisation (accelerated appearance of character states), tree 6 of 27 equal length trees; **B**, DELTRAN character-state optimisation (delayed appearance of character states), tree 4 of 27 equal length trees; **C**, strict consensus of 27 shortest trees for the original 17 taxa; **D**, strict consensus of 207 shortest trees (DELTRAN) or 212 shortest trees (ACCTRAN) of equal length = 145 steps for the original 17 taxa plus *Myllokunmingia* and *Yunnanozoon*.




Fig. 6. — DELTRAN character-optimisation tree with presentation of character changes. See Appendix 1 for character list and character states. CRANIATA, synapomorphies: 1, neural crest present; 2, brain present; 5, divided pituitary present; 6, optic tectum present; 12, paired olfactory organ; 19, sensory lines in grooves; 20, gill openings pouch-shaped; 31, lateral head vein present; homoplasies: 8, single nasal opening; 9, nasopharyngeal duct present; 10, single nasopharyngeal opening; 41, visceral arches fused to neurocranium; VERTEBRATA, synapomorphies: 4, pineal organ present and uncovered; 7, cerebellum present; 13, extrinsic eye muscles present; 14, two semicircular canals; 17, sensory line system with neuromasts; 21, symmetrical gill position; 26, gill lamellae with filaments; 28, close position of atrium and ventricle; 29, closed pericardium; 43, arcualia present; homoplasies: 11, dorsal position of nasohypophysial opening; 18, sensory line grooves or canals present on head and body; 23, slanting row of gill openings; EUVERTEBRATA, synapomorphies: 48, dermal trunk skeleton; 51, calcified dermal skeleton; homoplasies: 60, rod-shaped scales; GNATHOSTOMATA, synapomorphies: 14, three semicircular canals; 20, gill opening as slits to chambers; 56, enamel present; homoplasies: 30, paired dorsal aortae; 61, oakleaf-shaped tubercles; 63, denticles in pharynx; reversals: 10, no single nasohypophysial opening; 11, no nasohypophysial opening; 41, visceral arches not fused to neurocranium; 62, no oral plates; 64, small micromeric dermal head covering in adult state; CONODONTA: no synapomorphies in our data matrix, these would be hyaline and albid (white matter) tissue and their connection, base not in unit with crown and conodont apparatus.

accepted in general the conclusions of Donoghue *et al.* (2000, 2003, 2006, 2008) that conodonts are 1) vertebrates, and 2) “stem gnathostomes”. We do not want to comment on these papers in detail here because we focus on the origin of the discussion, that is, the phylogenetic analysis of Donoghue *et al.* (2000). We emphasize that those papers do not critically evaluate the arguments of Donoghue *et al.*, and that, in some cases, they are contradictory in their own developments (see e.g., Janvier 2009), who says in the English version [p. 211] that “Fossils thus show that bone and teeth have preceded jaws”, although in the French version [p. 214] he says that “l’os et la dentine ont précédé l’apparition des mâchoires” – bone and dentine have preceded the occurrence of jaws, but without any mention of teeth.

## RESULTS OF THE NEW CLADISTIC ANALYSIS

Our analysis of the revised data matrix (Table 2) results in cladograms for the original 17 taxa (Figs 5A, B; 6) where (Tunicata + Cephalochordata) appear as sister taxa of (Conodonts + Craniata). Vertebrates (node 31) are characterized by nearly the same synapomorphies and homoplasies as in Donoghue *et al.* (2000), but do not include the conodonts.

The list of characters for craniates in Donoghue *et al.*'s (2000: fig. 14a, ACCTTRAN) cladogram includes nine synapomorphies and 15 homoplasies. Of these, seven synapomorphies [1<sup>1</sup> (1<sup>1</sup>) neural crest present, 2<sup>1</sup> (2<sup>1</sup>) brain present, 5<sup>1</sup> (5<sup>1</sup>) pituitary divided, 6<sup>1</sup> (7<sup>1</sup>) optic tectum present, 12<sup>1</sup> (18<sup>1</sup>) paired

olfactory organ, 14<sup>1</sup> (20<sup>1</sup>) one semicircular canal, and 40<sup>1</sup> (51<sup>1</sup>) ability to synthesise creatine phosphatase] and five homoplasies [8<sup>2</sup> (14<sup>2</sup>) single median nasal opening, 10<sup>1</sup> (16<sup>1</sup>) single nasohypophysal opening, 11<sup>1</sup> (17<sup>1</sup>) terminal nasohypophysal opening, 18<sup>1</sup> (25<sup>1</sup>) sensory line grooves on head, and 19<sup>1</sup> (26<sup>1</sup>) sensory lines in grooves] appear in our ACCTRAN analysis at node 33 (Craniata + Conodonta), even though all these characters are unknown (coded “?”, see Table 2) in conodonts. In contrast our ACCTRAN analysis (Appendix 3) shows only one of Donoghue *et al.*'s (2000) synapomorphies for craniates (40<sup>1</sup> (51<sup>1</sup>)) and one additional homoplasy (38<sup>1</sup> (49<sup>1</sup>) hypocercal tail) at node 33 (Craniata + Conodonta). This and the following results from our DELTRAN analysis (Figs 5B; 6) are significant differences. Donoghue *et al.*'s (2000) synapomorphies [1<sup>1</sup> (1<sup>1</sup>), 2<sup>1</sup> (2<sup>1</sup>), 5<sup>1</sup> (5<sup>1</sup>), 6<sup>1</sup> (7<sup>1</sup>), 12<sup>1</sup> (18<sup>1</sup>) and 20<sup>1</sup> (27<sup>1</sup>) pouch-shaped gills] and homoplasies [8<sup>2</sup> (14<sup>2</sup>), 10<sup>1</sup> (16<sup>1</sup>), 19<sup>1</sup> (26<sup>1</sup>) and 41<sup>1</sup> (52<sup>1</sup>) visceral arches fused to neurocranium] appear at node 32 (Craniata) of our DELTRAN analysis with all of these characters except 41<sup>1</sup> (52<sup>1</sup>) being synapomorphies, thus placing the conodonts outside the craniates. Characters 31<sup>1</sup> (41<sup>1</sup>) lateral head vein and 9<sup>1</sup> (15<sup>1</sup>) nasopharyngeal duct present are additional synapomorphies at this node in our DELTRAN analysis.

The Vertebrata (Fig. 6; Appendix 3) are characterized by nine synapomorphies [7<sup>1</sup> (8<sup>1</sup>) cerebellum present, 13<sup>1</sup> (19<sup>1</sup>) extrinsic eye musculature, 14<sup>2</sup> (20<sup>2</sup>) two semicircular canals, 17<sup>1</sup> (23<sup>1</sup>) sensory line system with neuromasts, 21<sup>1</sup> (28<sup>1</sup>) symmetrical gill position, 26<sup>1</sup> (34<sup>1</sup>) gill lamellae with filaments, 28<sup>1</sup> (37<sup>1</sup>) close position of atrium to ventricle, 29<sup>1</sup> (38<sup>1</sup>) closed pericardium, and 43<sup>1</sup> (55<sup>1</sup>) arcualia present], and two homoplasies [4<sup>2</sup> (4<sup>2</sup>) uncovered pineal organ, and 18<sup>2</sup> (25<sup>2</sup>) sensory grooves and canals on head and body] in our analyses (ACCTRAN and DELTRAN; also both transformations produce the same strict consensus tree, Fig. 5C); two additional different homoplasies appear in ACCTRAN [34<sup>1</sup> (45<sup>1</sup>) anal fin separate, and 50<sup>1</sup> (64<sup>1</sup>) calcified cartilage present] and DELTRAN [11<sup>2</sup> (17<sup>2</sup>) nasohypophysal opening dorsal, and 23<sup>1</sup> (30<sup>1</sup>) lateral gill openings in slanting row]. Donoghue *et al.*'s (2000) cladogram shows 24(!) additional synapomorphies [(6<sup>2</sup>), (12<sup>1</sup>), (13<sup>1</sup>), (24<sup>1</sup>), (39<sup>1</sup>), (42<sup>1</sup>), (57<sup>1</sup>), (83<sup>1</sup>-

94<sup>1</sup>), (96<sup>1</sup>-99<sup>1</sup>), (102<sup>1</sup>)] for vertebrates, which are all characters unknown in fossil taxa and therefore eliminated from our analysis, as noted earlier. Character 7<sup>1</sup> [(8<sup>1</sup>) cerebellum present] appears as an additional synapomorphy for vertebrates in our analyses relative to the synapomorphies in Donoghue *et al.*'s (2000). Two [4<sup>2</sup> (4<sup>2</sup>), 34<sup>1</sup> (45<sup>1</sup>)] of the four homoplasies for vertebrates in our ACCTRAN analysis (Fig. 5A; Appendix 3) are not present in Donoghue *et al.*'s (2000) cladogram, which shows four homoplasies [(29<sup>1</sup>) branchial series with fewer than 10 openings, (30<sup>1</sup>) slanting row of gill openings, (49<sup>1</sup>) hypocercal tail, (50<sup>0</sup>) no preanal median fold] and two reversals for vertebrates [(40<sup>0</sup>) no paired dorsal aortae, (95<sup>1</sup>) eliminated in our analysis] not present in our analysis. These characters [23<sup>1</sup> (30<sup>1</sup>), 30<sup>0</sup> (40<sup>0</sup>), 38<sup>1</sup> (49<sup>1</sup>), 39<sup>0</sup> (50<sup>0</sup>)] appear more basal in our analyses with the exception of character 22<sup>1</sup> (29<sup>1</sup>) (elongate branchial series), which occurs within lower vertebrates.

In addition, six synapomorphies [(8<sup>1</sup>) cerebellum present, (9<sup>1</sup>) pretrematic branches in branchial nerves, (21<sup>1</sup>) vertical semicircular canals forming loops, (41<sup>1</sup>) lateral head vein present, (60<sup>1</sup>) occiput enclosing vagus and glossopharyngeal, and (65<sup>1</sup>) calcified dermal skeleton present] place the conodonts “above” (more crownward) the petromyzontids in Donoghue *et al.*'s (2000) cladogram. Of these characters, number (9) was eliminated, and numbers 7 (8), 15 (21), 31 (41) and 46 (60) are coded as “?” (unknown) and 51 (65) as “0” (absent) for conodonts in our data matrix.

The position of conodonts in our cladograms (Figs 5; 6) is the result of changing the coding of 12 characters (see Appendix 2) for conodonts (1 (1), 2 (2) and 13 (19) from 1 to ?; 20 (27) from 0 to ?; 41 (52), 42 (54), 43 (55), 55 (69) from ? to 0; 51 (65), 56 (70) and 63 (77) from 1 to 0; 59 (73) from 2 to 0). These recodings change the synapomorphies (eight) and homoplasies (three) between conodonts and craniates/vertebrates in Donoghue *et al.*'s (2000) cladogram. In our data matrix 50% of our 68 characters are unknown for conodonts (Table 2). The result of our analysis is an improved version of the data set of Donoghue *et al.* (2000); it demonstrates that conodonts are neither vertebrates nor craniates.

We also ran the analysis leaving pituriaspids and *Eriptychius* out of the matrix as most data for these taxa are unknown; the consensus tree did show a little better resolution. Concerning *Eriptychius*, the phylogenetic position given by Donoghue *et al.* (2000; also Donoghue & Aldridge 2001) as the sister-group of gnathostomes appears unusual. After their detailed auto-critical evaluation of the character analysis of *Eriptychius*, Donoghue *et al.* (2000: 217) did recognize that they “cannot claim that the evidence for the association of *Eriptychius* with gnathostomes [...] is well supported”. *Eriptychius* actually has 82% of missing data in Donoghue *et al.*'s (2000) paper, and some characters are miscoded. For instance, character 78 (state 1: dermal head covering in adult state small micromeric) does not apply to the dermal head cover of *Eriptychius*, which is more likely to be meso/macromeric (Denison 1967). Most characters that link *Eriptychius* to gnathostomes are homoplastic (Donoghue *et al.* 2000: fig. 14a, b), and the only one that is resolved as a synapomorphy to both taxa (character 20: number of semi-circular canals in labyrinth) is coded “?” for *Eriptychius* by Donoghue *et al.* (2000: table 1). So, their result was not strongly supported and has been abandoned in more recent papers (Donoghue & Sansom 2002; Donoghue *et al.* 2003).

In order to test relationships of the purported Chinese Cambrian chordates, we included in our matrix *Myllokunmingia*, synonymous with *Haikouichthys* (Hou *et al.* 2002; Janvier 2003) as noted above, and *Yunnanozoon*, now considered synonymous with *Haikouella*. Shimeld & Holland (2000: fig. 1) showed a hypothesis of phylogenetic relationship between living members of the phylum Chordata plus *Myllokunmingia* (as a claimed craniate, Holland & Chen 2001) and *Yunnanozoon* (= *Haikouella*, a claimed basal chordate, Mallatt & Chen 2003), but they left out other claimed fossil chordates, including *Pikaia* and *Cathamyrius* (see above) that they claimed were possibly related to *Branchiostoma* and the euconodonts (the claimed possible vertebrates). Our analysis shows some support for that of Holland & Chen (2001) with *Myllokunmingia* as a vertebrate stemward of Petromyzontida, and *Yunnanozoon* as a chordate between Tunicata and

Cephalochordata in some of the 200+ best trees. The consensus tree (Fig. 5D) places Tunicata, Cephalochordata, *Yunnanozoon* and Craniata in a polytomy, and *Myllokunmingia* with Vertebrata, Myxinoidea, *Jamoytius* (Fig. 1H), *Euphanerops* and Conodonta in another polytomy.

However, our consensus trees (Fig. 5C, D) do not help to resolve interrelationships within vertebrates, indicating that the data matrix is not adequate for this purpose, although elimination of poorly known taxa greatly increases the resolution of the analysis.

#### SUMMARY: WHY CONODONTS ARE NOT VERTEBRATES

Perhaps development of the (buccal-pharyngeal) bipartite conodont apparatus in advanced para- and euconodonts reflects an alternative pathway to allow a soft-bodied worm-shaped animal to increase in size, support a filter-feeding lifestyle, and even possibly indulge in macrophagy. It is interesting that even the more extreme “conodonts are vertebrates” people seem to accept that protoconodonts are related to chaetognaths (see e.g., in Reif 2006) or even that chaetognaths descend from protoconodonts (Szaniawski 2002). These organisms are sometimes classified among the protostomes (Lecointre & Le Guyader 2001), sometimes among the deuterostomes (references in Janvier 1998); they are barely known as fossils, and then in Palaeozoic times, from Cambrian and Carboniferous Fossil-Lagerstätten (Benton 1993; Vannier *et al.* 2007). However, if it is confirmed that protoconodonts are phylogenetically related to chaetognaths, and also confirmed that a phylogenetic relationship does exist between protoconodonts, paraconodonts and euconodonts (Vannier *et al.* 2007, and references therein), the latter would also be related to chaetognaths, rather than to chordates, a hypothesis already advocated on the basis of molecular analysis (Kasatkina & Buryi 1996). Moreover, if conodonts are related to chaetognaths, and if the latter are the sister-group of craniates as indicated by some molecular analyses (Christofferson & Araújo-de-Almeida 1994), this would also place conodonts in the sister-group of

craniates (see e.g., Peterson 1994; Pridmore *et al.* 1997). However, the most recent molecular analysis by Putnam *et al.* (2008: fig. 1) placed the “hemichordate” acorn worms (i.e. Enteropneusta) + sea urchins (Echinodermata) as sister group to chordates.

Instead, we consider that conodonts s.l. represent one or more “invertebrate animals” (a paraphyletic grouping) with phosphatic elements that, unlike vertebrates, disappeared by the end of Triassic (cf. Gross 1954). A major part of the argument for excluding all conodonts from craniates (and thus from vertebrates) and not interpreting their elements as teeth rests with the differences in hard tissues (Schultze 1996; Kemp & Nicoll 1996; Kemp e.g., 2002a, b; Reif 2006; Trotter *et al.* 2007).

We summarize our lines of evidence as follows: – cephalization is low in conodonts, possibly at the level of extant cephalochordates, whereas it is much higher in craniates where cephalization reaches an advanced state of development;

– conodonts lack segmentally-arranged paraxial elements, a feature of vertebrates;

– conodont trunk musculature is simple V-shaped as opposed to a W-shaped in craniates;

– conodonts lack dermal elements in median fins, whereas median fins of vertebrates possess dermal fin rays that are articulated at the base with supporting cartilaginous elements;

– conodonts lack a dermal skeleton including bony plates whereas all vertebrates have odontodes with bone of attachment and a unique pulp system (see Fig. 2 for comparative morphogenesis);

– conodont element ultrastructure, revealed by TEM and SEM, has a crystalline structure very different in crystal size and arrangement than it is in vertebrates. The albid material of conodonts is formed by extraordinarily large crystals in contrast to the fine small crystals of bone, dentine and enamel in vertebrates;

– conodont hyaline material shows a histochemical reaction for collagen, which is not present in vertebrate enamel;

– conodont element albid material (white matter) does not react for collagen, which is a major element in vertebrate bone;

– the lacunae in the albid material are too small for eukaryote cells such as osteocytes;

– a cladistic analysis based on the recorded data set of Donoghue *et al.* (2000) supports neither a vertebrate nor a craniate relationship for conodonts.

Above, we have summarized from the work of several of us and in addition cite new evidence. There is no evidence that conodonts share any of the attributes of vertebrate hard tissues (Trotter *et al.* 2007) and the morphogenetic system of formation of vertebrate hard parts (Reif 2006). Sansom (2006), in his review of Hall’s (2005) book, corrected his own earlier work by stating “Conodonts did not possess cellular bone and probably did not possess cartilage within their oral-pharyngeal feeding elements, despite the interpretation of Sansom *et al.* (1992), a point that has become clear from later papers and a deeper understanding of the constraints of tissue topology within conodont elements”.

A further paper on patterning of hair in mammals (Sick *et al.* 2006) reiterated the basic difference between conodonts and vertebrates — that the former have no external dermal armour whatsoever, be it isolated odontodes (“scales”), even phosphatic nubs such as in *Hadimopanella* (once thought to be vertebrate and now referred to palaeoscolecid worms: see e.g., Hinz *et al.* 1990) or bony plates. In this respect they share nothing with the earliest definite vertebrates (*sensu* Turner *et al.* 2004), either plated such as *Arandaspis* or scaled such as *Sacabambaspis* and *Areyonga* (Young 1997, 2009). In vertebrates, odontodes (scales, teeth, denticles, etc.; Reif 1982, 2002, 2006) grow and are shed in particular ways; only rarely is there a “continuous” growth, e.g., fin spines, rat’s teeth, dicynodont and elephant tusks, but even these have a finite existence. In conodonts the elements grow outward continuously and thus are covered in soft tissue; they do not erupt and they are not shed. Because they are covered in tissue they must have been everted during feeding to function like vertebrate teeth. But, just covered by a thin tissue they could have been used for food grasping-filtering and transportation and sometimes as grinding plates (the platforms) during passage of food from the buccal cavity to the gut. This is true of the myxinoid and certain annelid/polychaete apparatuses. In myxinoids, there is the single palatal tooth attached to a cartilaginous plate. The lingual teeth also attach to a

cartilaginous dental plate with a longitudinal keel fitting into the grooved upper surface of the immovable cartilaginous basal plate making up the floor of the mouth (Retzius 1892). The described conodont impressions lack any evidence of necessary (appropriate) cartilaginous plates, yet traces of presumed “cartilaginous” fin rays are preserved, thus arguing against similar apparatuses in conodonts and myxinoids. Of course, the limited number of possible feeding mechanisms means that any superficial similarities might as well be a matter of function as of a genetic relationship. Further, there are obvious major differences between the internal structure of phosphatic conodont elements and the horny “teeth” of myxinoids.

## CONCLUSIONS

This paper constitutes a further refutation of the hypothesis that conodonts are “stem gnathostomes” or vertebrates. Conodonts are not recognized as an early vertebrate group that experimented with skeletal biomineralization. At most they might represent a cephalochordate grade of evolution, similar to the amphioxus *Branchiostoma*. *Contra*, e.g., Donoghue *et al.* (2000, 2003, 2008), Kuhn & Barnes (2005), and Janvier (2008), the plesion Conodonta Eichenberg, 1930 is in no way a sister group or a member, stem or otherwise of the phylum Craniata Linnaeus, 1758, its subphylum Vertebrata Linnaeus, 1758, nor of the superclass Gnathostomata Cope, 1889. Placing higher taxa Conodontophorida or Conodontia in the Chordata Bateson, 1886 or Craniata (e.g., Farrell 2004; Nelson 2006; King *et al.* 2009) is not accepted for the reasons outlined above. Instead, based on the evidence provided here, we support the hypotheses that:

- the phylogenetic status of conodonts *s.l.*, including proto-, para- and euconodonts is not resolved; at the moment the three groups are informal, with protoconodonts probably not monophyletic. Müller & Hinz-Schallreuter (1998) considered the three groups as related. Szaniawski (2002) favoured the idea that protoconodonts belonging to the evolutionary lineage of *Phakelodus* are probably the stem group of chaetognaths. Szaniawski & Bengtson

(1993) described a well documented evolutionary lineage from a paraconodont species to a euconodont species. Authors who relate euconodonts to craniates consider that the three groups have different phylogenetic relationships;

- conodont elements are not odontodes and do not possess vertebrate hard tissues, including globular cartilage, bone, lamellin, dentine, enameloid or enamel, and do not exhibit vertebrate morphogenesis (see Figs 2; 3); broken conodont elements could be repaired during the animal’s life, and therefore, unlike odontodes (scales, teeth, etc.) in vertebrates, the elements had to be infolded in tissue at least at times;

- conodont elements are not homologous with vertebrate teeth and do not represent the first vertebrate experiment in skeletonisation or mineralized feeding apparatus (see Figs 2; 3);

- conodont (euconodont) animals are not craniates, nor vertebrates, nor “stem gnathostomes” (see Figs 5; 6).

## Acknowledgements

We thank Dr Anne Kemp (Centre for Microscopy and Microanalysis, University of Queensland, St Lucia, Australia), and the late Dr David Carlisle (Canada), and our co-author Pr Dr Wolf-Ernst Reif (Institut für Geowissenschaften, Eberhard-Karls Universität Tübingen, IGTU), who sadly passed away in 2002 and June 2009, respectively, and Dr Robert S. Nicoll (Canberra) for their inspiration and discussion; Dr Julie Trotter (Department of Earth and Marine Sciences, Australian National University, Canberra) for sharing data whilst in press; Pr Dr Otto H. Walliser (Zentrum für Geowissenschaften, Abteilung Geobiologie, Göttingen University), Dr Rodney D. Norby (Illinois State Geological Survey, Champaign, Illinois), Dr Bruce Lieberman (University of Kansas, Department of Geology, Lawrence, Kansas), and Dr Zivile Zigaite (University of Vilnius, Lithuania) for comments on our work. Judy Bracefield (Brisbane), Verena Benz and Peter Brandscheid (IGTU) provided assistance. ST acknowledges an Australian Academy of Sciences grant, which enabled insightful discussions with and kind hospitality of Prs Drs

W.-E. Reif and J. Nebelsick in 2006; and to the latter for support at IGTU in 2007-2009. ST and CJB acknowledge basic facilities at the Queensland Museum. We thank Drs A. Kemp, H. Lelièvre and M. Ginter, two unnamed referees, Gaël Clément and the editor-in-chief, Dr D. Merle, for their constructive reviews.

## REFERENCES

- ALBANESI G. L. & BERGSTRÖM S. M. 2004. — Conodonts: Lower to Middle Ordovician Record, in WEBBY B. D., PARIS F., DROSER M. L. & PERCIVAL I. G. (eds), *The Great Ordovician Biodiversity Event*. Columbia University Press, New York: 312-326 + 395-466.
- ALDRIDGE R. J. 1987. — Conodont palaeobiology: a historical review, in ALDRIDGE R. J. (ed.), *Palaeobiology of conodonts*. British Micropalaeontological series, Ellis Horwood: 11-34.
- ALDRIDGE R. J. & PURNELL M. A. 1996. — The conodont controversy. *Trends in Ecology & Evolution* 11 (11): 463-468.
- ALDRIDGE R. J. & BRIGGS D. E. G. 2009. — The discovery of the conodont anatomy and its importance for understanding the early history of vertebrates, in SEPKOSKI D. & RUSE M. (eds), *The Paleobiological Revolution. Essays on the Growth of Modern Paleontology*. University of Chicago Press, Chicago & London: 73-88.
- ALDRIDGE R. J., BRIGGS D. E. G., CLARKSON E. N. K. & SMITH M. P. 1986. — The affinities of conodonts – new evidence from the Carboniferous of Edinburgh, Scotland. *Lethaia* 19: 279-291.
- ALDRIDGE R. J., BRIGGS D. E. G., SMITH M. P., CLARKSON E. N. K. & CLARK N. D. L. 1993. — The anatomy of conodonts. *Philosophical Transactions of the Royal Society, London*, Ser. B, 340: 405-421.
- ANDRES D. 1988. — Strukturen, Apparate und Phylogenie primitiver Conodonten. *Palaeontographica A* 200: 105-152.
- ARENDT D. & NÜBLER-JUNG K. 1994. — Inversion of dorsoventral axis. *Nature* 371: 26.
- BALFOUR F. M. 1877. — The development of elasmobranch fishes. *Journal of Anatomy and Physiology* 11 (3): 406-490.
- BARDACK D. 1997. — Fishes: "Agnatha", Acanthodii, and Osteichthyes, in SHABICA C. W. & HAY A. A. (eds), *Richardson's Guide to The Fossil Fauna of Mazon Creek*. Northeastern Illinois University, Chicago, Illinois: 226-243.
- BARNES C. R., SASS D. B. & MONROE E. A. 1973. — Ultrastructure of some Ordovician conodonts, in RHODES F. H. T. (ed.), *Conodont Paleozoology. Geological Society of America Special Paper* 141: 1-30.
- BARRY J. C. & KEMP A. 2007. — High resolution transmission electron microscopy of developing enamel in the Australian lungfish, *Neoceratodus forsteri* (Osteichthyes: Dipnoi). *Tissue and Cell* 39: 387-398.
- BARSKOV I. S., MOSKALENKO T. A. & SAROSTINA L. P. 1982. — New evidence for the vertebrate nature of conodontophorids. *Paleontological Journal* 1982 (1): 82-90. [translation from *Paleontologicheskii Zhurnal* 1982 (1): 80-86].
- BEALL B. S. 1991. — The Tully Monster and a new approach to analyzing problematica, in SIMONETTA A. M. & CONWAY MORRIS S. (eds), *The Early Evolution of Metazoa and the Significance of Problematic Taxa*. Cambridge University Press, Cambridge: 271-285.
- BÉLAND P. & ARSENAULT M. 1985. — Scauménellisation de l'Acanthodii *Triazeugacanthus affinis* (Whiteaves) de la formation d'Escuminac (Dévonien supérieur de Miguasha, Québec): révision du *Scaumenella mesacanthi* Graham-Smith. *Canadian Journal of Earth Sciences* 22 (4): 514-524.
- BELKA Z. 2004. — †Conodonta, in WESTHEIDE W. & RIEGER R. (eds), *Spezielle Zoologie. Teil: Wirbel- oder Schädeltiere*. Spektrum, Heidelberg & Berlin: 173.
- BENTON M. J. (ed.) 1993. — *The Fossil Record* 2. Chapman & Hall, London: XVII + 845 p.
- BENTON M. J. 2005. — *Vertebrate Paleontology*. 3rd edition. Blackwell Publishing Ltd., Oxford: XII + 474 p.
- BERGSTRÖM J. 1996. — Upside-down evolution? *Geologiska Föreningens i Stockholm Förhandlingar* 118, Jubilee Issue: A58-59.
- BERGSTRÖM J. 1997. — Origin of high-rank groups of organisms. *Paleontological Research* 1: 1-14.
- BERGSTRÖM J., NAUMANN W. W., VIEWEG J. & MARTÍMUS M. 1998. — Conodonts, calcichordates and the origin of vertebrates. *Mitteilungen aus dem Museum für Naturkunde Berlin, Geowissenschaftliche Reihe* 1: 81-92.
- BERRILL N. J. 1987. — Early chordate evolution. *International Journal of Invertebrate Reproduction and Development* 11: 1-27.
- BLIECK A. 1992. — At the origin of chordates. *Geobios* 25: 101-113.
- BLIECK A. & GOUJET D. 1983. — *Zascinaspis laticephala* nov. sp. (Agnatha, Heterostraci) du Dévonien inférieur du Spitsberg. *Annales de Paléontologie (Vert.-Invert.)* 69: 43-56.
- BLIECK A., TURNER S., BURROW C. J., SCHULTZE H.-P. & REXROAD C. B. 2009. — Organismal biology, phylogeny and strategy of publication: why conodonts are not vertebrates, in 69th Annual Meeting Society of Vertebrate Paleontology & 57th Symposium on Vertebrate Paleontology and Comparative Anatomy (SVP/PCA) (Univ. Bristol, UK, Sept. 23-26, 2009). Program and Abstracts. *Journal of Vertebrate Paleontology* 29 (Suppl. to no. 3): 65A.



- BOORMANN C. J. & SHIMELD S. M. 2002a. — The evolution of left-right asymmetry in chordates. *BioEssays* 24: 1004-1011.
- BOORMANN C. J. & SHIMELD S. M. 2002b. — Pitx homeobox genes in *Ciona* and amphioxus show left-right asymmetry is a conserved chordate character and define the ascidian adenohipophysis. *Evolution and Development* 4: 354-365.
- BRACEGIRDLE B. & MILES P. H. 1978. — *An Atlas of Chordate Structure*. Heinemann Educational Books, London, VII + 119 p.
- BRANSON E. B. & MEHL M. G. 1933. — Conodont studies. *University of Missouri Studies* 8: 349 p.
- BRIGGS D. E. G. 1992. — Conodonts: A major extinct group added to the vertebrates. *Science* 256: 1285-1286.
- BRIGGS D. E. G. 2003. — The role of decay and mineralization in the preservation of soft-bodied fossils. *Annual Review of Earth and Planetary Sciences* 31: 275-301.
- BRIGGS D. E. G. 2010. — Decay distorts ancestry. *Nature* 463: 741-743.
- BRIGGS D. E. G., CLARKSON E. N. K. & ALDRIDGE R. J. 1983. — The conodont animal. *Lethaia* 16: 1-14.
- BRIGGS D. E. G., ALDRIDGE R. J. & SMITH M. P. 1987. — Conodonts are not aplacophoran molluscs. *Lethaia* 20: 381-382.
- BULTYNCK P. 1987. — Pelagic and neritic conodont successions from the Givetian of pre-Sahara Morocco and the Ardennes. *Bulletin de l'Institut royal des Sciences naturelles de Belgique, Sciences de la Terre* 57: 149-181.
- BULTYNCK P. 2009. — To be or not to be: can conodonts be vertebrates?, in P. GODEFROIT & O. LAMBERT (eds), *EAVP Extraordinary Meeting – Tribute to Charles Darwin and Bernissart Iguanodons* (Brussels, 9-14 Febr. 2009). Royal Institute of Natural Sciences of Belgium, Brussels: 24.
- CARLISLE D. B. 1953. — On the origin of the pituitary body of chordates. *Nature* 172: 1098.
- CARLSON S. J. 1990. — Vertebrate dental structures (chapter 21), in CARTER J. G. (ed.), *Skeletal Biomineralization: Patterns, Processes and Evolutionary Trends*. Volume I. Van Nostrand Reinhold, New York: 531-556.
- CARTER J. G. & LUTZ R. A. 1990. — Bivalvia (Mollusca), in CARTER J. G. (ed.), *Skeletal Biomineralization: Patterns, Processes and Evolutionary Trends*. Volume II. Part 2 Atlas and Index. Van Nostrand Reinhold, New York: 5-28, pls 1-121.
- CHEN J.-Y., DZIK J., EDGEcombe G. D., RAMSKÖLD L. & ZHOU G.-Q. 1995. — A possible Early Cambrian chordate. *Nature* 377: 720-722.
- CHEN J.-Y., HUANG D.-Y. & LI C.-W. 1999. — An early Cambrian craniate-like chordate. *Nature* 402: 518-522.
- CHRISTOFFERSON M. L. & ARAÚJO-DE-ALMEIDA E. A. 1994. — A phylogenetic framework of the enterocoela (Metamerie: Coelomata). *Revista Norestina de Biologia (Brazil)* 9: 172-208.
- CLARKSON E. N. K. 1985. — 1. The Granton Shrimp-Bed, Edinburgh: crustaceans, conodonts and conservation. *Earth Science Conservation* 22: 3-8.
- CONWAY MORRIS S. 1989. — Conodont palaeobiology: recent progress and unsolved problems. *Terra Nova* 1: 135-150.
- CONWAY MORRIS S. 1991. — Problematic taxa: a problem for biology or biologists?, in SIMONETTA A. M. & CONWAY MORRIS S. (eds), *The Early Evolution of Metazoa and the Significance of Problematic Taxa*. Cambridge University Press: 19-24.
- CONWAY MORRIS S. 1993a. — The fossil record and the early evolution of the Metazoa. *Nature* 361 (6409): 219-225.
- CONWAY MORRIS S. 1993b. — Ediacaran-like fossils in Cambrian Burgess Shale-type faunas of North America. *Palaeontology* 36 (3): 593-635.
- CONWAY MORRIS S. 1998. — *The Crucible of Creation. The Burgess Shale and the Rise of Animals*. Oxford University Press, Oxford: XXIII + 242 p.
- COSTA O. G. 1834. — *Annuario zoologico. Cenni zoologici, ossia descrizione sommaria delle specie nuove di animali scoperti in diverse contrade del regno nell'anno 1834*. Napoli, 99 p.
- DEAN B. 1899. — On the embryology of *Bdellostoma stouti*, in *Festschrift zum siebenzigsten Geburtstag von Carl F. v. Kupffer*. Gustav Fischer, Jena: 221-276.
- DEAN M. N. & SUMMERS A. P. 2006. — Mineralized cartilage in the skeleton of chondrichthyan fishes. *Zoology* 109: 164-168.
- DENISON R. H. 1967. — Ordovician vertebrates from Western United States. *Fieldiana: Geology* 16 (6): 131-192.
- DIXON A. D., HOYTE D. A. N. & RÖNNING O. 1988. — *Fundamentals of Craniofacial Growth*. CRC Press, New York, 487 p.
- DONG X.-P., DONOGHUE P. C. J. & REPETSKI J. E. 2005. — Basal tissue structure in the earliest euconodonts: Testing hypotheses of developmental plasticity in euconodont phylogeny. *Palaeontology* 48 (2): 411-421.
- DONOGHUE P. C. J. 1998. — Growth and patterning in the conodont skeleton. *Philosophical Transactions of the Royal Society of London, Series B*, 353: 633-666.
- DONOGHUE P. C. J. 2001. — Microstructural variation in conodont enamel is a functional adaptation. *Proceedings of the Royal Society of London, Series B*, 268: 1691-1698.
- DONOGHUE P. C. J. 2005. — Saving the stem group – a contradiction in terms? *Paleobiology* 31 (4): 553-558.
- DONOGHUE P. C. J. & CHAUFFE K. 1998. — *Concho-*

- dontus*, *Mitrellataxis* and *Fungulodus*: conodonts, fish, or both? *Lethaia* 31: 283-292.
- DONOGHUE P. C. J. & ALDRIDGE R. J. 2001. — Origin of a mineralized skeleton, in AHLBERG P. E. (ed.), *Major Events in Early Vertebrate Evolution – Palaeontology, phylogeny, genetics and development*. Systematics Association, Special Volume Series 61. Taylor & Francis, London & New York: 85-105.
- DONOGHUE P. C. J. & SANSOM I. J. 2002. — Origin and Early Evolution of Vertebrate Skeletonization. *Microscopy Research and Technique* 59: 352-372.
- DONOGHUE P. C. J. & PURNELL M. A. 2005. — Genome duplication, extinction and vertebrate evolution. *Trends in Ecology and Evolution* 20 (6): 312-319.
- DONOGHUE P. C. J., PURNELL M. A. & ALDRIDGE R. J. 1998. — Conodont anatomy, chordate phylogeny and vertebrate classification. *Lethaia* 31 (3): 211-219.
- DONOGHUE P. C. J., FOREY P. L. & ALDRIDGE R. J. 2000. — Conodont affinity and chordate phylogeny. *Biological Reviews* 75: 191-251.
- DONOGHUE P. C. J., SMITH M. P. & SANSOM I. J. 2003. — The origin and early evolution of chordates: molecular clocks and the fossil record, in DONOGHUE P. C. J. & SMITH M. P. (eds), *Telling the Evolutionary Time: Molecular Clocks and the Fossil Record*. Taylor & Francis, London & New York: 190-223.
- DONOGHUE P. C. J., SANSOM I. J. & DOWNS J. P. 2006. — Early evolution of vertebrate skeletal tissues and cellular interactions, and the canalization of skeletal development. *Journal of Experimental Zoology* 306B: 278-294.
- DONOGHUE P. C. J., PURNELL M. A., ALDRIDGE R. J. & ZHANG S.-X. 2008. — The interrelationships of “complex” conodonts (Vertebrata). *Journal of Systematic Palaeontology* 6 (2): 119-153.
- DZIK J. 1986. — Chordate affinities of the conodonts, in HOFFMANN A. & NITECKI M. (eds), *Problematic Fossil Taxa*. Oxford Monographs on Geology and Geophysics no. 5. Oxford University Press, Oxford, New York: 240-254.
- DZIK J. 2000. — The origin of the mineral skeleton in chordates, in HECHT M. K., MACINTYRE R. J. & CLEGG M. T. (eds), *Evolutionary Biology*. volume 31. Kluwer Academic; Plenum Publisher, New York: 105-154.
- DZIK J. 2009. — Conodont affinity of the enigmatic Carboniferous chordate *Conopiscius*. *Lethaia* 41: 31-38.
- FÄHRAEUS L. E. & FÄHRAEUS-VON REE G. E. 1994. — Histomorphology of sectioned and stained 415 Ma old soft-tissue matrix from internal fluorapatite skeletal elements of an extinct taxon, Conodontophorida (Conodontia), in KOBAYASHI I., MUTVEI H. & SAHNI A. (eds), *Structure, Formation and Evolution of Fossil Hard Tissues*. Tokyo University Press, Tokyo: 107-112.
- FARRELL J. R. 2004. — Siluro-Devonian conodonts from the Camelford Limestone, Wellington, New South Wales. *Palaeontology* 47: 937-982.
- FRANCILLON-VIEILLOT H., BUFFRENIL V. DE, CASTANET J., GERAUDIE J., MEUNIER F. J., SIRE J.-Y., ZYLBERBERG L. & RICQLES A. DE 1990. — Microstructure and mineralization of vertebrate skeletal tissues, in CARTER J. G. (ed.), *Skeletal Biomineralization*. Van Nostrand, New York: 471-530.
- FURNISH W. M. 1938. — Conodonts from the Prairie du Chien (Lower Ordovician) beds of the upper Mississippi valley. *Journal of Paleontology* 12: 318-340.
- GABBOTT S. E., ALDRIDGE R. J. & THERON J. N. 1995. — A giant conodont with preserved muscle tissue from the Upper Ordovician of South Africa. *Nature* 374 (6525): 800-803.
- GARTANG W. 1928. — The origin and evolution of larval forms. *Nature* 211: 366.
- GEDIK I. V. & ÇAPKINOĞLU S. 1996. — Conodonts, were they (the first?) parasitic animals?, in *Sixth International Conodont Symposium (ECOS VI, July 4-6, 1996, Warszawa)*. Abstracts: 19.
- GESS R. W., COATES M. I. & RUBIDGE B. S. 2006. — A lamprey from the Devonian of South Africa. *Nature* 443: 981-984.
- GROSS W. 1954. — Zur Conodonten-Frage. *Senckenbergiana lethaea* 35 (1/2): 73-85.
- GROSS W. 1957. — Über die Basis der Conodonten. *Paläontologische Zeitschrift* 31: 78-91.
- GROSS W. 1960. — Über die Basis bei den Gattungen *Palmatolepis* und *Polygnathus* (Conodontida). *Paläontologische Zeitschrift* 34: 40-58.
- GUO W., DONG X.-P., ZENG X.-C. & ZHAO L. 2005. — Histological observation of some primitive euconodonts of Late Cambrian from Liaoning. *Progress in Natural Science* 15 (12): 1084-1088.
- HALL B. K. 2005. — *Bones and Cartilage: Developmental and Evolutionary Skeletal Biology*. Elsevier Academic Press, San Diego: XXXVII + 760 p.
- HALL J. C. 1990. — Conodontophorida, in CARTER J. G. (ed.), *Skeletal Mineralization: Patterns, Processes and Evolutionary Trends*. Volume II. Part 10 Atlas and Index. Van Nostrand Reinhold, New York: 81-85.
- HARDISTY A. & POTTER I. (eds) 1972. — *The Biology of Lampreys*. Volume 2. Academic Press, New York, XIV + 466 p.
- HASS W. H. 1941. — The morphology of conodonts. *Journal of Paleontology* 15: 71-81.
- HENNIG W. 1969. — *Die Stammesgeschichte der Insekten*. W. Kramer, Frankfurt/Main, *Senckenberg-Buch* 40, 436 p.
- HENNIG W. 1983. — Stammesgeschichte der Chordaten. *Fortschritte in der zoologischen Systematik und Evolutionsforschung* 2. P. Parey, Hamburg, Berlin, 208 p.
- HINDE G. J. 1879. — On conodonts from the Chazy and Cincinnati Group of the Cambro-Silurian, and

- from the Hamilton and Genesee-Shale division of the Devonian, in Canada and the United States. *Quarterly Journal of the Geological Society, London* 35: 351-369.
- HINZ, I., KRAFT, P., MERGL, M. & MÜLLER, K. J. 1990. — The problematic *Hadimopanella*, *Kaimenella*, *Milaculum* and *Utahphospha* identified as sclerites of Palaeoscolecida. *Lethaia* 23: 217-221.
- HOLLAND L. Z. & HOLLAND N. D. 2001. — Evolution of neural crest and placodes: amphioxus as a model for the ancestral vertebrate? *Journal of Anatomy* 199: 85-98.
- HOLLAND N. D. & CHEN J.-Y. 2001. — Origin and early evolution of the vertebrates: new insights from advances in molecular biology, anatomy, and palaeontology. *BioEssays* 23: 142-151.
- HOU X.-G., ALDRIDGE R. J., SIVETER D. J., SIVETER D. J. & FENG X.-H. 2002. — New evidence of the anatomy and phylogeny of the earliest vertebrates. *Proceedings of the Royal Society, London B* 269: 1865-1869.
- HUXLEY T. H. 1858. — On the theory of the vertebrate skull. Croonian Lecture delivered 18 November, 1858. *Proceedings of the Royal Society, London* 9: 381-457 [imprint 1859].
- HUYSSSEUNE A., SIRE J.-Y. & WITTEN P. E. 2009. — Evolutionary and developmental origins of the vertebrate dentition, in *The Integument Story: Origins, Evolution and Current Knowledge* (Paris, July 2007). *Journal of Anatomy* 214 (4): 465-476.
- JANVIER P. 1981. — The phylogeny of the Craniata, with particular reference to the significance of fossil "agnathans". *Journal of Vertebrate Paleontology* 1(2): 121-159.
- JANVIER P. 1997. — Euconodonts, in *Tree of Life*. World Wide Web address: <http://www.tolweb.org/Euconodonts/14832> (accessed 2009).
- JANVIER P. 1998. — Les Vertébrés avant le Silurien. *Geobios* 30 (7): 931-950.
- JANVIER P. 2001. — Euconodonts†, in *Tree of Life*. World Wide Web address: <http://phylogeny.Arizona.edu/tree/eukaryotes/animals/chordata/euconodonts/> (accessed 2009).
- JANVIER P. 2003. — Vertebrate characters and the Cambrian vertebrates, in NÉRAUDEAU D. (ed.), *Les chemins de l'évolution: sur les pas de Steven Jay Gould*. *Comptes-Rendus Palevol* 2 (6-7): 523-531.
- JANVIER P. 2006. — Modern look for ancient lamprey. *Nature* 443: 921-924.
- JANVIER P. 2007a. — Living primitive fishes and fishes from deep time, in MCKENZIE D. J., FARREL A. P. & BRAUNER C. J. (eds), *Primitive Fishes*. Fish Physiology 26. Elsevier, Amsterdam: 1-51.
- JANVIER P. 2007b. — Homologies and evolutionary transitions in early vertebrate history, in ANDERSON J. S. & SUES H.-D. (eds), *Major Transitions in Vertebrate Evolution*. Indiana University Press, Bloomington and Indianapolis: 57-121.
- JANVIER P. 2008. — The brain in the early jawless vertebrates: evolutionary information from an empty nutshell. *Brain Research Bulletin* 75: 314-318.
- JANVIER P. 2009. — Les premiers Vertébrés et les premières étapes de l'évolution du crâne, in COPPENS Y., PADIAN K., DE RICQLÈS A. & TAQUET P. (eds), *Histoire évolutive de la vie*. *Comptes-Rendus Palevol* 8 (2-3): 209-219.
- JANVIER P. & LUND R. 1983. — *Hardistiella montanensis* n. gen. et sp. (Petromyzontida) from the Lower Carboniferous of Montana with remarks on the affinities of the lampreys. *Journal of Vertebrate Paleontology* 2: 407-413.
- JANVIER P. & ARSENAULT M. 2007. — The anatomy of *Euphanerops longaevis* Woodward, 1900, an anaspid-like jawless vertebrate from the Upper Devonian of Miguasha, Quebec, Canada. *Geodiversitas* 29 (1): 143-216.
- JEFFERIES R. P. S. 1979. — The origin of chordates – a methodological essay, in HOUSE M. R. (ed.), *The Origin of Major Invertebrate Groups*. Systematics Association, Special Volume 12. Academic Press, London: 443-477.
- JEFFERIES R. P. S. 2001. — The origin and early fossil history of the chordate acustico-lateralis system, with remarks on the reality of the echinoderm-hemichordate clade, in AHLBERG P. E. (ed.), *Major Events in Early Vertebrate Evolution – Palaeontology, Phylogeny, Genetics and Development*. Systematics Association, Special Volume Series 61. Taylor & Francis, London & New York: 40-66.
- KASATKINA A. P. & BURYI G. I. 1996. — On the relation of chaetognaths and conodonts. *Albertiana* 18: 21-23.
- KAWASAKI K., SUZUKI T. & WEISS K. M. 2004. — Genetic basis for the evolution of vertebrate mineralized tissue. *Proceedings of the National Academy of Sciences* 101 (31): 11356-11361.
- KEMP A. 1992. — Ultrastructure of the developing dentition in the Australian lungfish, *Neoceratodus forsteri*, in SMITH P. & TCHERNOV E. (eds), *Structure, Function and Evolution of Teeth*. Freund Publisher, Tel Aviv: 11-33.
- KEMP A. 1999. — A refined method for the staining of organic remnants in conodont elements. *Ichthyolith Issues* 20: 34.
- KEMP A. 2002a. — Hyaline tissue of thermally unaltered conodont elements and the enamel of vertebrates. *Alcheringa* 26: 23-36.
- KEMP A. 2002b. — Amino-acid residues in conodont elements. *Journal of Paleontology* 76: 518-528.
- KEMP A. 2003. — Ultrastructure of developing tooth plates in the Australian lungfish, *Neoceratodus forsteri* (Osteichthyes: Dipnoi). *Tissue and Cell* 35:

- 401-426.
- KEMP A. & NICOLL R. S. 1995. — Protochordate affinities of conodonts. *Courier Forschungsinstitut Senckenberg* 182: 235-245.
- KEMP A. & NICOLL R. S. 1996. — A histochemical analysis of biological residues in conodont elements, in TURNER S. & BLIECK A. (eds), *Gross Symposium, Volume 1. Modern Geology* 20 (3-4): 287-302 (re-published in TURNER S. & BLIECK A. [eds], *Gross Symposium, Volume 2. Modern Geology* 21 [1-2]: 197-213 with 2 colour plates [1997]).
- KEMP A. & BARRY J. C. 2006. — Prismatic dentine in the Australian lungfish, *Neoceratodus forsteri* (Osteichthyes: Dipnoi). *Tissue and Cell* 38: 127-140.
- KING C. M., ROBERTS C. D., BELL B. D., FORDYCE R. E., NICOLL R. S., WORTHY T. H., PAULIN C. D., HITCHMOUGH R. A., KEYES I. W., BAKER A. N., STEWART A. L., HILLER N., MCDOWALL R. M., HOLDAWAY R. N., MCPHEE R. P., SCHWARZHANS W. W., TENNYSON A. J. D., RUST S. & MACADIE I. 2009. — Chapter 24: Phylum Chordata – lancelets, fishes, amphibians, reptiles, birds, mammals, in GORDON D. P. (ed.), *New Zealand Inventory of Biodiversity. Vol. One: Kingdom Animalia*. Canterbury University Press, Christchurch: 433-553.
- KOENIGSWALD W. v. 2000. — Two different strategies in enamel differentiation: Marsupialia versus Eutheria, in TEAFORD M. F., SMITH M. M. & FERGUSON M. W. J. (eds), *Development, Function and Evolution of Teeth*. Cambridge University Press, Cambridge, UK: 107-118.
- KOENTGES G. 2008. — Teeth in double trouble. *Nature* 455: 747-748.
- KUHN T. S. & BARNES C. R. 2005. — Ordovician conodonts from the Mithaka Formation (Georgina Basin, Australia): regional and paleobiogeographical implications. *Geologica Acta* 3 (4): 317-337.
- LAURIN M. & ANDERSON J. S. 2004. — Meaning of the name Tetrapoda in the scientific literature: an exchange. *Systematic Biology* 53: 68-80.
- LECOINTRE G. & LE GUYADER H. 2001. — *Classification phylogénétique du Vivant*. Belin, Paris, 543 p. (translated to English: *The Tree of Life: A Phylogenetic Classification*. Harvard University Press, Cambridge, Massachusetts, 2006)
- LIEM K. F., BEMIS W. E., WALKER W. F. J. & GRANDE L. 2001. — *Functional Anatomy of the Vertebrates: an Evolutionary Perspective*. Third edition. Harcourt College Publishers, Orlando, Florida: XVII + 703 p.
- LINDSTRÖM M. 1964. — *Conodonts*. Elsevier, Amsterdam, VI + 196 p.
- LINDSTRÖM M. & ZIEGLER W. 1971. — Feinstrukturelle Untersuchungen an Conodonten. 1. Die Überfamilie Panderodontacea. *Geologica et Palaeontologica* 5: 9-33.
- LINDSTRÖM M. & ZIEGLER W. 1981. — Surface micro-ornamentation and observations on internal composition, in ROBISON R. A. (ed.), *Treatise on Invertebrate Paleontology, Part W: Miscellanea, Supplement 2, Conodonta*. Geological Society of America and University of Kansas, Lawrence: W41-W52.
- MALLATT J. & CHEN J.-Y. 2003. — Fossil sister group of craniates: predicted and found. *Journal of Morphology* 258: 1-31.
- MARINELLI W. & STRENGER A. 1954. — *Vergleichende Anatomie und Morphologie der Wirbeltiere*. Band 1. II. Lieferung: *Lampetra fluviatilis* (L.). Franz Deuticke, Wien: 1-80.
- MARINELLI W. & STRENGER A. 1956. — *Vergleichende Anatomie und Morphologie der Wirbeltiere*. Band 1. II. Lieferung: *Myxine glutinosa* (L.). Franz Deuticke, Wien: 81-172.
- MÄRSS T. 2006. — Exoskeletal ultrasculpture of early vertebrates. *Journal of Vertebrate Paleontology* 26 (2): 235-252.
- MÄRSS T., TURNER, S. & KARATAJUTE-TALIMAA, V. N. 2007. — “Agnatha” II, Thelodonti, in SCHULTZE H.-P. (ed.), *Handbook of Paleichthyology*, Vol. 1B. Verlag Dr Friedrich Pfeil, Munich, 143 p.
- MOSER H. G., RICHARDS W. J., COHEN D. M., FAHAY M. P., KENDALL A. W. JR & RICHARDSON S. L. (eds) 1984. — *Ontogeny and Systematics of Fishes*. American Society of Ichthyologists and Herpetologists, Special Publication Number 1: IX + 760 p.
- MÜLLER K. J. 1981. — Zoological affinities of conodonts, in ROBISON, R. A. (ed.), *Treatise on Invertebrate Palaeontology, Part W: Miscellanea, Supplement 2, Conodonta*. Geological Society of America, University of Kansas, Lawrence: W78-W82.
- MÜLLER K. J. & NOGAMI Y. 1971. — Über den Feinbau der Conodonten. *Memoirs of the Faculty of Sciences Kyoto University, Series of Geology and Mineralogy* 38: 1-88.
- MÜLLER K. J. & HINZ I. 1991. — Upper Cambrian conodonts from Sweden. *Fossils and Strata* 28: 153 p.
- MÜLLER K. J. & HINZ-SCHALLREUTER I. 1998. — Internal structure of Cambrian conodonts. *Journal of Paleontology* 72 (1): 91-112.
- NELSON J. S. 2006. — *Fishes of the World*. 4th Edition. John Wiley & Sons Inc., Hoboken, XIX + 601 p.
- NICOLL R. S. 1985. — Multielement composition of the conodont species *Polygnathus xylus xylus* Stauffer, 1940 and *Ozarkodina brevis* (Bischoff and Ziegler, 1957) from the Upper Devonian of the Canning Basin Western Australia. *BMR Journal of Australian Geology and Geophysics* 9: 133-147.
- NICOLL R. S. 1987. — Form and function of the Pa element in the conodont animal, in ALDRIDGE R. J. (ed.), *Palaeobiology of Conodonts*. Ellis Horwood, Chichester: 77-90.
- NICOLL R. S. 1992. — Analysis of conodont apparatus organisation and the genus *Jumudontus* (Conodonta),

- a coniform-pectiniform apparatus structure from the Early Ordovician. *BMR Journal of Australian Geology and Geophysics* 13: 213-228.
- NICOLL R. S. 1995. — Conodont element morphology, apparatus reconstruction and element function: a new interpretation of conodont biology with taxonomic implications. *Courier Forschungsinstitut Senckenberg* 182: 247-262.
- NORBY R. D. 1976. — Conodont apparatuses from Chesterian (Mississippian) strata of Montana and Illinois. Unpublished PhD Thesis, University of Illinois at Champaign-Urbana, USA, 203 p.
- NORTHCUTT R. G. 1996. — The origin of craniates: neural crest, neurogenic placodes, and homeobox genes. *Israel Journal of Zoology* 42: 273-313.
- OELOFSEN B. W. & LOOCK J. C. 1981. — A first record of a fossil cephalochordate. *Palaeontographia Africana* 24: 17.
- ØRVIG T. 1973. — Fossila fisktänder i svepelektronmikroskopet: gamla frågeställningar i ny belysning [Fossil fish teeth in the scanning electron microscope: old problems in new light]. *Flora och Fauna* 68(4): 166-173 (in Swedish).
- OTA K. G., KURAKU S. & KURATANI S. 2007. — Hagfish embryology with reference to the evolution of the neural crest. *Nature* 446: 672-675.
- PAGE R. D. M. 1996. — Treeview: an application to display phylogenetic trees on personal computers. *Computer Applications in the Biosciences* 12: 357-358. Also World Wide Web address: <http://taxonomy.zoology.gla.ac.uk/rod/treeview.html> (accessed 2009).
- PAGE R. D. M. 1999. — *Nexus Data Editor for Windows*. World Wide Web address: <http://taxonomy.zoology.gla.ac.uk/rod/NDE/nde.html> (accessed 2009).
- PANDER C. H. 1856. — *Monographie der fossilen Fische des silurischen Systems der russisch-baltischen Gouvernements*. Kaiserliche Akademie der Wissenschaften, St. Petersburg, X + 91 p.
- PARIS M., ESCRIVA H., SCHUBERT M., BRUNET F., BRTKO J., CIESIELSKI F., ROECKLIN D., VIVAT-HANNAH V., JAMIN E. L., CRAVEDI J.-P., SCANLAN T. S., RENAUD J.-P., HOLLAND N. D. & LAUDET V. 2008. — *Amphioxus* postembryonic development reveals the homology of chordate metamorphosis. *Current Biology* 18: 825-830.
- PARKER T. J. & HASWELL W. A. 1921. — *A Text-Book of Zoology*. Vol. II. Macmillan and Co., London, XX + 714 p.
- PERNÈGRE V. & GOUJET D. 2007. — The genus *Gigantaspis* Heintz, 1962 (Vertebrata, Heterostraci) from the Lower Devonian of Spitsbergen. *Palaeontology* 50: 323-346.
- PETERSON K. J. 1994. — The origin and early evolution of the Craniata, in PROTHERO D. R. & SCHOCH R. M. (eds), Major features of vertebrate evolution. *The Paleontological Society, Short Courses in Paleontology* 7: 14-37.
- POPLIN C., SOTTY D. & JANVIER P. 2001. — Un Myxinoïde (Craniata, Hypertreti) dans le *Konservat-Lagerstätte* Carbonifère supérieur de Montceau-les-Mines (Allier, France). *Comptes-rendus de l'Académie des Sciences, Paris, Sciences de la Terre et des planètes* 332: 345-350.
- PRIDMORE P. A., BARWICK R. E. & NICOLL R. S. 1997. — Soft anatomy and the affinities of conodonts. *Lethaia* 29 (4): 317-328.
- PROTHERO D. R. 1998. — *Bringing Fossils to Life. An Introduction to Paleobiology*. WCB McGraw-Hill, Boston, X + 457 p.
- PURNELL M. A. 1993. — Feeding mechanisms in conodonts and the function of the earliest vertebrate hard tissues. *Geology* 21: 375-377.
- PURNELL M. A. 1995a. — Microwear on conodont elements and macrophagy in the first vertebrates. *Nature* 374 (6525): 798-800.
- PURNELL M. A. 1995b. — Large eyes and vision in conodonts. *Lethaia* 28: 187-188.
- PURNELL M. A. 2001. — Scenarios, selection, and the ecology of early vertebrates, in AHLBERG P. E. (ed.), *Major Events in Early Vertebrate Evolution. Palaeontology, Phylogeny, Genetics and Development*. Systematics Association Special Volume Series 61. Taylor & Francis, London and New York: 188-208.
- PURNELL M. A. & VON BITTER P. 1992. — Blade-shaped conodont elements functioned as cutting teeth. *Nature* 359: 629-630.
- PUTNAM N. H., BUTTS T., FERRIER D. E. K., FURLONG R. F., HELLSTEN U., KAWASHIMA T., ROBINSON-RECHAVI M., SHOGUCHI E., TERRY A., YU J.-K., BENITO-GUTIÉRREZ E., DUBCHAK I., GARCIA-FERNÁNDEZ J., GIBSON-BROWN J. J., GRIGORIEV I. V., HORTON A. C., JONG P. J. DE, JURKA J., KAPITONOV V. V., KOHARA Y., KUROKI Y., LINDQUIST E., LUCAS S., OSOEGAWA K., PENNACCHIO L. A., SALAMOV A. A., SATOU Y., SAUKA-SPENGLER T., SCHMUTZ J., SHIN-I T., TOYODA A., BRONNER-FRASER M., FUJIYAMA A., HOLLAND L. Z., HOLLAND P. W. H., SATOH N. & ROKHSAR D. S. 2008. — The amphioxus genome and the evolution of the chordate karyotype. *Nature* 453: 1064-1071.
- RAINERI M. 2006. — Are protochordates chordates? *Biological Journal of the Linnean Society, London*, 87 (2): 261-284 (with Erratum: *Biological Journal of the Linnean Society, London*, 87 [4]: 637 [2006]).
- REIF W.-E. 1982. — Evolution of dermal skeleton and dentition in vertebrates. The Odontode Regulation Theory, in HECHT M., WALLACE B. & PRANCE G. T. (eds), *Evolutionary Biology* 15. Plenum Press, New York: 287-368.
- REIF W.-E. 1984. — Pattern regulation in shark dentition, in MALACHINSKI G. M. (ed.), *Pattern Formation, a Primer in Developmental Biology*. Macmillan

- Publishing Co., New York, 603-621.
- REIF W.-E. 2002. — Evolution of the dermal skeleton of vertebrates: Concepts and methods. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 223 (1): 53-78.
- REIF W.-E. 2004. — Problematic issues of cladistics: 5. Flexible phylogenetic classification (FPC) of higher taxa of the Vertebrata. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 233 (3): 369-395.
- REIF W.-E. 2006. — Conodonts, odontodes, stem-groups, and the ancestry of enamel genes. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 241 (3): 405-439.
- RETZIUS M. G. 1892. — Das Caudalskelet der *Myxine glutinosa*. *Biologiska Föreningen i Förhandlingar, Stockholm* 3: 81-84.
- SANDER P. M. 1997. — Non-mammalian synapsid enamel and the origin of mammalian enamel prisms: The bottom-up perspective, in KOENIGSWALD W. VON & SANDER P. M. (eds), *Tooth Enamel Microstructure*. AA Balkema, Rotterdam: 41-62.
- SANDER P. M. 2000. — Prismless enamel in amniotes: terminology, functions, and evolution, in TEAFORD M. F., SMITH M. M. & FERGUSON M. W. J. (eds), *Development, Function and Evolution of Teeth*. Cambridge University Press, Cambridge, UK: 92-106.
- SANSOM I. J. 2006. — Book Review. *Bones and Cartilage: Developmental and Evolutionary Skeletal Biology*. Brian K. Hall. 2005. Elsevier Academic Press, San Diego. xxxviii+760 pp., with illustrations. *Evolution & Development* 8 (4): 389-390.
- SANSOM I. J., SMITH M. P., ARMSTRONG H. A. & SMITH M. M. 1992. — Presence of the earliest vertebrate hard tissues in conodonts. *Science* 256: 1308-1311.
- SANSOM I. J., SMITH M. M. & SMITH M. P. 1994. — Dentine in conodonts. *Nature* 368: 591.
- SANSOM I. J., SMITH M. M. & SMITH M. P. 2001. — The Ordovician radiation of vertebrates, in AHLBERG P. E. (ed.), *Major Events in Early Vertebrate Evolution. Palaeontology, Phylogeny, Genetics and Development*. Systematics Association, Special Volume Series 61. Taylor & Francis, London and New York: 156-171.
- SANSOM I. J., DONOGHUE P. C. J. & ALBANESI G. 2005. — Histology and affinity of the earliest armoured vertebrate. *Biology Letters* 1 (4): 446-449.
- SANSOM R. S., GABBOTT S. E. & PURNELL M. A. 2010. — Non-random decay of chordate characters causes bias in fossil interpretation. *Nature* 463: 797-800.
- SATCHELL P. G., SHULER C. F. & DIEKWISCH T. G. H. 2000. — True enamel coverings in teeth of the Australian lungfish *Neoceratodus forsteri*. *Cell and Tissue Research* 299: 27-37.
- SCHÄFER W. 1972. — *Ecology and Paleoecology of the Marine Environment*. Oliver and Boyd, Edinburgh, 568 p.
- SCHMIDT H. & MÜLLER K. J. 1964. — Weitere Funde von Conodonten-Gruppen aus dem oberen Karbon des Sauerlandes. *Paläontologische Zeitschrift* 38: 105-135.
- SCHULTZE H.-P. 1969. — Die Faltenzähne der rhipidistiiden Crossopterygier, der Tetrapoden und der Actinopterygier-Gattung *Lepisosteus*; nebst einer Beschreibung der Zahnstruktur von *Onychodus* (struniformer Crossopterygier). *Palaeontographica Italica* 65 (n. ser. 35): 63-137.
- SCHULTZE H.-P. 1996. — Conodont histology: an indicator of vertebrate relationship?, in TURNER S. & BLIECK A. (eds), *Gross Symposium, Volume 2. Modern Geology* 20 (3-4): 275-285.
- SHIMELD S. M. & HOLLAND P. W. H. 2000. — Vertebrate innovations. *Proceedings of the National Academy of Sciences of the United States of America* 97 (9): 4449-4452.
- SHU D.-G., CONWAY MORRIS S., HAN J., ZHANG Z.-F., YASUI K., JANVIER P., CHEN L., ZHANG X.-L., LIU J.-N., LI Y. & LIU H.-Q. 2003. — Head and backbone of the Early Cambrian vertebrate *Haikouichthys*. *Nature* 421 (6922): 526-529.
- SHU D.-G., CONWAY MORRIS S. & ZHANG X.-L. 1996a. — A *Pikaia*-like chordate from the Lower Cambrian of China. *Nature* 384: 156-157.
- SHU D.-G., ZHANG X.-L. & CHEN L.-Z. 1996b. — Reinterpretation of *Yunnanzoon* as the earliest known hemichordate. *Nature* 380: 428-430.
- SHU D.-G., LUO H.-L., CONWAY MORRIS S., ZHANG X.-L., HU S.-X., CHEN L., HAN J., ZHU M., LI Y. & CHEN L.-Z. 1999. — Lower Cambrian vertebrates from South China. *Nature* 402 (6757): 42-46.
- SICK S., REINKER S., TIMMER J. & SCHLAKE T. 2006. — WNT and DKK determine hair follicle spacing through a reaction-diffusion mechanism. *Science* 31: 1447-1450.
- SIMONETTA A. M., PUCCI A. & DZIK J. 1999. — Hypotheses on the origin and early evolution of chordates in the light of recent zoological and paleontological evidence. *Italian Journal of Zoology* 66: 99-109.
- SIRE J.-Y., DONOGHUE P. C. J. & VICKARYOUS M. K. 2009. — Origin and evolution of the integumentary skeleton in non-tetrapod vertebrates. *Journal of Anatomy* 214 (4): 409-440.
- SMITH M. M. 1989. — Distribution and variation in enamel structure in the oral teeth of sarcopterygians: its significance for the evolution of a protoprismatic enamel. *Historical Biology* 3: 97-126.
- SMITH M. P., BRIGGS D. E. G. & ALDRIDGE R. J. 1987. — A conodont animal from the lower Silurian of Wisconsin, USA, and the apparatus architecture of panderodontid conodonts, in ALDRIDGE R. J. (ed.), *Palaeobiology of conodonts*. British Micropalaeontological Society Series, Ellis Horwood, Chichester: 91-104.
- SMITH M. M., SANSOM I. J. & SMITH M. P. [quoted as Smith P.] 1995. — Diversity of the dermal skeleton

- in Ordovician to Silurian vertebrate taxa from North America: histology, skeletogenesis and relationships, in LELIÈVRE H., WENZ S., BLIECK A. & CLOUTIER R. (eds), *Premiers Vertébrés et Vertébrés Inférieurs. Geobios, Mémoire Spécial* 19: 65-70.
- SMITH M. M., SANSOM I. J. & SMITH M. P. 1996. — “Teeth” before armour: the earliest vertebrate mineralized tissues, in TURNER S. & BLIECK A. (eds), *Gross Symposium, Volume I. Modern Geology* 20 (3-4): 303-319.
- SMITH M. P., SANSOM I. J. & COCHRANE K. D. 2001. — The Cambrian origin of vertebrates, in AHLBERG P. E. (ed.), *Major Events in Early Vertebrate Evolution. Palaeontology, Phylogeny, Genetics and Development*. Systematics Association, Special Volume Series 61. Taylor & Francis, London and New York: 67-84.
- SWEET W. C. 1981. — Morphology and composition of elements- Macromorphology of elements and apparatuses, in ROBISON R. A. (ed.), *Treatise on Invertebrate Paleontology, Part W, Miscellanea*, Supplement 2, *Conodonts*. Geological Society of America and University of Kansas, Lawrence: W5-W20.
- SWEET W. C. 1988. — *The Conodonts: Morphology, Taxonomy, Paleoecology, and Evolutionary History of a Long-extinct Animal Phylum*. Oxford Monographs on Geology and Geophysics 10. Clarendon Press, Oxford, x + 212 p.
- SWOFFORD D. 2002. — PAUP 4.0b10 for Windows. Sinauer Associates, Sunderland, Massachusetts. World Wide Web address: <http://paup.csit.fsu.edu/win.html>.
- SZANIAWSKI H. 1983. — Structure of protoconodont elements. *Fossils and Strata* 15: 21-27.
- SZANIAWSKI H. 1987. — Preliminary structural comparisons of protoconodont, paraconodont, and euconodont elements, in ALDRIDGE R. J. (ed.), *Palaeobiology of Conodonts*. Ellis Horwood, Chichester: 35-47.
- SZANIAWSKI H. 2002. — New evidence for the protoconodont origin of chaetognaths. *Acta Palaeontologica Polonica* 47 (3): 405-419.
- SZANIAWSKI H. & BENGTON S. 1993. — Origin of euconodont elements. *Journal of Paleontology* 67 (4): 640-654.
- TILLIER S. & CUIF J.-P. 1986. — L'animal-conodonte est-il un Mollusque Aplacophore? *Comptes-rendus de l'Académie des Sciences, Paris* 303 (II): 627-632.
- TROTTER J. A. & EGGINS S. M. 2006. — Chemical systematics of conodont apatite determined by laser ablation ICPMS. *Chemical Geology* 233: 196-216.
- TROTTER J. A., GERALD J. D. F., KOKKONEN H. & BARNES C. R. 2007. — New insights into the ultrastructure, permeability, and integrity of conodont apatite determined by transmission electron microscopy. *Lethaia* 40 (2): 97-110.
- TURNER S. 1985. — Remarks on the early history of chondrichthyans, thelodonts, and some “higher elasmobranchs”, in COOPER R. (ed.), *Hornibrook Symposium* (Christchurch, 1985). *New Zealand Geological Survey Record* 9: 93-95.
- TURNER S. 1994a. — Thelodont squamation. *Ichthyolith Issues* 13: 12-15.
- TURNER S. 1994b. — Thermal alteration of microscopic fish remains. *Ichthyolith Issues* 13: 25-26.
- TURNER S. & BLIECK A. 1995. — Conodont-vertebrate comparison and significance of micro-ornament, in BROCK G. A. (ed.), *The AUSCOS-1/BOUCOT Symposium (Macquarie University, Sydney, July 18-21, 1995)*. Abstracts: 105.
- TURNER S., BLIECK A. & NOWLAN G. S. 2004. — Vertebrates (Agnathans and Gnathostomes), in WEBBY B. D., PARIS F., DROSER M. L. & PERCIVAL I. G. (eds), *The Great Ordovician Biodiversity Event*. Columbia University Press, New York: 327-335 + 395-466.
- VANDENBROUCKE T. R. A., GABBOTT S. E., PARIS F., ALDRIDGE R. J. & THERON J. N. 2009. — Chitinozoans and the age of the Soom Shale, an Ordovician black shale Lagerstätte, South Africa. *Journal of Micropalaeontology* 28: 53-66.
- VANNIER J., STEINER M., RENVOISE E., HU S.-X. & CASANOVA J.-P. 2007. — Early Cambrian origin of modern food webs: evidence from predator arrow worms. *Proceedings of the Royal Society B* 274: 627-633.
- WALLISER O. H. 1994. — Architecture of the polygnathid conodont apparatus, in KÖNIGSHOF P. & WERNER R. (eds), *Willi Ziegler-Festschrift I. Courier Forschungsinstitut Senckenberg* 168: 31-36.
- WARSHAWSKY H. 1989. — Organization of crystals in enamel. *The Anatomical Record* 224: 242-262.
- WATABE N. 1990. — Calcium phosphate structures in invertebrates and protozoans, in CARTER J. G. (ed.), *Skeletal Biomineralization: Patterns, Processes and Evolutionary Trends*. Volume I, Chapter 4. Van Nostrand Reinhold, New York: 35-44.
- WEDDIGE K. 1989. — Conodonten – problematische Fossilien. *Natur und Museum* 119 (3): 67-82.
- WEDDIGE K. 1990. — Pathological conodonts. *Courier Forschungsinstitut Senckenberg* 118: 563-589.
- WILSON M. V. H., HANKE G. F. & MÄRSS T. 2007. — Paired fins of jawless vertebrates and their homologues across the “agnathan”-gnathostome transition, in ANDERSON J. S. & SUES H.-D. (eds), *Major Transitions in Vertebrate Evolution*. Indiana University Press, Bloomington and Indianapolis: 122-149.
- WRIGHT J. 1990. — Conodont geochemistry: a key to the Palaeozoic. *Courier Forschungsinstitut Senckenberg* 118: 227-305.
- YOUNG G. C. 1997. — Ordovician microvertebrate remains from the Amadeus Basin, central Australia. *Journal of Vertebrate Paleontology* 17: 1-25.
- YOUNG G. C. 2009. — An Ordovician vertebrate from western New South Wales, with comments on Cam-

- bro-Ordovician vertebrate distribution patterns. *Alcheringa* 33: 79-89.
- ZHANG S., ALDRIDGE R. J. & DONOGHUE P. C. J. 1997. — An Early Triassic conodont with periodic growth? *Journal of Micropalaeontology* 16: 65-72.
- ZHANG X.-L., SHU D.-G., LI Y. & HAN J. 2001. — New sites of Chengjiang fossils: crucial windows on the Cambrian explosion. *Journal of the Geological Society of London* 158: 211-218.
- ZYLBERBERG L., SIRE J.-Y. & NANJI A. 1997. — Immunodetection of amelogenin-like proteins in the ganoine of experimentally regenerating scales of *Calamoichthys calabaricus*, a primitive actinopterygian fish. *The Anatomical Record* 249: 86-95.

*Submitted on 4 August 2008;  
accepted on 26 February 2010.*



## APPENDIX 1

List of characters used in the data matrix. The first number (in bold) applies to our matrix; the second number (in italics and parentheses) corresponds to that used in the Donoghue *et al.*'s (2000: table 1) matrix. Characters 20 and 31 have been changed as noted in the text.

- |                |  |                |  |
|----------------|--|----------------|--|
| <b>1</b> (1)   | neural crest: 0. absent; 1. present  | <b>32</b> (43) | subaponeurotic vascular plexus: 0. absent; 1. present  |
| <b>2</b> (2)   | brain: 0. absent; 1. present   | <b>33</b> (44) | separate dorsal fin: 0. absent; 1. present   |
| <b>3</b> (3)   | olfactory peduncles: 0. absent; 1. present   | <b>34</b> (45) | anal fin separate: 0. absent; 1. present   |
| <b>4</b> (4)   | pineal organ: 0. absent; 1. present and covered;<br>2. present and uncovered                             | <b>35</b> (46) | unpaired fin ray supports closeset: 0. absent;<br>1. present   |
| <b>5</b> (5)   | divided pituitary: 0. absent; 1. present   | <b>36</b> (47) | paired lateral fin folds: 0. absent; 1. present  |
| <b>6</b> (7)   | optic tectum: 0. absent; 1. present  | <b>37</b> (48) | constricted pectorals: 0. absent; 1. present   |
| <b>7</b> (8)   | cerebellum: 0. absent; 1. present  | <b>38</b> (49) | tail shape: 0. isocercal; 1. hypocercal; 2. epicercal  |
| <b>8</b> (14)  | number of nasal openings: 0. none; 1. paired; 2. single<br>median  | <b>39</b> (50) | preanal median fold: 0. absent; 1. present   |
| <b>9</b> (15)  | nasopharyngeal duct: 0. absent; 1. present   | <b>40</b> (51) | ability to synthesise creatine phosphatase: 0. absent;<br>1. present                                 |
| <b>10</b> (16) | single nasohypophyseal opening: 0. absent;<br>1. present   | <b>41</b> (52) | visceral arches fused to neurocranium: 0. absent;<br>1. present                                      |
| <b>11</b> (17) | position of nasohypophyseal opening: 0. none; 1. ter-<br>minal; 2. dorsal                                | <b>42</b> (54) | trematic rings: 0. absent; 1. present  |
| <b>12</b> (18) | olfactory organ: 0. absent; 1. paired; 2. unpaired   | <b>43</b> (55) | arcualia: 0. absent; 1. present  |
| <b>13</b> (19) | extrinsic eye musculature: 0. absent; 1. present   | <b>44</b> (58) | braincase with lateral walls: 0. absent; 1. present  |
| <b>14</b> (20) | semicircular canals in labyrinth: 0. absent; 1. 1; 2. 2;<br>3. 3   | <b>45</b> (59) | neurocranium entirely closed dorsally, covering brain:<br>0. absent; 1. present                      |
| <b>15</b> (21) | vertical semicircular canal looped: 0. absent; 1. present  | <b>46</b> (60) | occiput enclosing vagus and glossopharyngeal: 0. ab-<br>sent; 1. present                             |
| <b>16</b> (22) | endolymphatic ducts open externally: 0. absent;<br>1. present  | <b>47</b> (61) | annular cartilage: 0. absent; 1. present   |
| <b>17</b> (23) | sensory line system with neuromasts: 0. absent;<br>1. present  | <b>48</b> (62) | trunk dermal skeleton: 0. absent; 1. present   |
| <b>18</b> (25) | sensory line grooves or canals: 0. absent; 1. present<br>on head only; 2. on head plus body              | <b>49</b> (63) | perichondral bone: 0. absent; 1. present   |
| <b>19</b> (26) | sensory line: 0. absent; 1. in grooves; 2. in canals   | <b>50</b> (64) | calcified cartilage: 0. absent; 1. present   |
| <b>20</b> (27) | gill opening shape: 0. present, simple slits; 1. present,<br>pouch-shaped; 2. present, slits to chambers | <b>51</b> (65) | calcified dermal skeleton: 0. absent; 1. present   |
| <b>21</b> (28) | gill relative position: 0. alternate; 1. symmetrical   | <b>52</b> (66) | spongy aspidin: 0. absent; 1. present  |
| <b>22</b> (29) | branchial series: 0. more than 10; 1. fewer than 10  | <b>53</b> (67) | lamellar aspidin: 0. absent; 1. present  |
| <b>23</b> (30) | gill openings lateral, slanting row: 0. absent;<br>1. present  | <b>54</b> (68) | cellular bone: 0. absent; 1. present   |
| <b>24</b> (31) | gills opening: 0. laterally; 1. ventrally  | <b>55</b> (69) | dentine: 0. absent; 1. present, mesodentine; 2. present,<br>orthodentine                             |
| <b>25</b> (32) | opercular flaps: 0. absent; 1. present   | <b>56</b> (70) | enamel/oid: 0. absent; 1. present, monotypic enamel;<br>2. present, enameloid                        |
| <b>26</b> (34) | gill lamellae with filaments: 0. absent; 1. present  | <b>57</b> (71) | three-layered exoskeleton: 0. absent; 1. present   |
| <b>27</b> (35) | mouth position: 0. terminal; 1. ventral  | <b>58</b> (72) | cancellar layer in exoskeleton with honeycomb-shaped<br>cavities: 0. absent; 1. present              |
| <b>28</b> (37) | relative position of atrium and ventricle: 0. well<br>separated; 1. close                                | <b>59</b> (73) | composition of scales/denticles: 0. absent; 1. present,<br>single odontode; 2. present, polyodontode |
| <b>29</b> (38) | closed pericardium: 0. absent; 1. present  | <b>60</b> (74) | scales: 0. absent; 1. present, diamond-shaped; 2. present,<br>rod-shaped                             |
| <b>30</b> (40) | paired dorsal aortae: 0. absent; 1. present  | <b>61</b> (75) | oakleaf-shaped tubercles: 0. absent; 1. present  |
| <b>31</b> (41) | lateral head vein: 0. absent; 1. present   | <b>62</b> (76) | oral plates: 0. absent; 1. present   |
|                |  | <b>63</b> (77) | denticles in pharynx: 0. absent; 1. present  |

## APPENDIX 1 — Continuation.

- 64 (78)** dermal head covering in adult state: 0. absent; 1. small micromeric; 2. large dermal plates or shield
- 65 (79)** large unpaired ventral and dorsal dermal plates on head: 0. absent; 1. present
- 66 (80)** massive endoskeletal head shield covering gills dorsally: 0. absent; 1. present
- 67 (81)** sclerotic ossicles: 0. absent; 1. present
- 68 (82)** ossified scleral capsule: 0. absent; 1. present

## APPENDIX 2

Changes of coding from Donoghue *et al.* (2000); their character numbers are listed in italics after our new numbering as in Appendix 1.

## TUNICATA

- 2 (2)** *Brain*: 0 to ?: larval ascidians have an anterior enlargement of the dorsal nerve chord.
- 22 (29)** *Elongate branchial series*: 0 to ?: the number of gill slits is very variable within tunicates (1-10), and it is unclear if the sessile forms with many slits are close to the ancestral form.
- 39 (50)** *Preanal median fold*: ? to 0: these folds could only occur in larval forms, but none are known.

## CEPHALOCHORDATA

- 4 (4)** *Pineal organ*: ? to 0: we do not recognize homology between their “frontal eye” and the pineal organ in vertebrates.
- 36 (47)** *Paired lateral fin folds*: 0 to 1: if “This refers to any lateral or ventrolateral fin-like fold” (Donoghue *et al.* 2000: 212), then the “metapleural” folds in *Branchiostoma* are included.

## MYXINOIDEA

- 4 (4)** *Pineal organ*: ? to 0: the pineal organ is absent, as noted by Donoghue *et al.* (2000).
- 12 (18)** *Olfactory organ*: 2 to 1: only the nasopharyngeal opening is unpaired, the olfactory organ is paired.
- 23 (30)** *Slanting gill openings lateral*: 0 to 0/1: different in different species and genera; one could even argue that they are partly ventral, promoting a recoding for character 24 (31).
- 25 (32)** *Opercular flaps*: 0 to 0/1: *Myxine garmani* has one common opening with slits covered by flap, in *Paramyxine atami* slits are crowded with flaps.
- 31 (41)** *Large lateral head vein*: 0 to 1: lateral head vein = anterior part of anterior cardinal or anterior vena cava is present.
- 33 (44)** *Dorsal fin*: 0 to 0/1: the anterior fin rays are free in contrast to rays of the caudal fin (Retzius 1892 refers to a dorsal fin).

- 38 (49)** *Tail shape*: 0 to 1: Retzius (1892: pl. 3, fig. 10) shows a hypocercal structure (also Marinelli & Strenger 1956: figs 108, 109).
- 42 (54)** *Trematic rings*: 1 to 0.
- 51 (65)** *Calcified dermal skeleton*: ? to 0: there is no skeleton and no record that they are able to synthesise creatine phosphatase.

## PETROMYZONTIDA

- 7 (8)** *Cerebellum*: 0 to 1: a small cerebellum lies below the posterior choroid plexus in the anterior part of the medulla oblongata.
- 9 (15)** *Nasopharyngeal duct*: 0 to 1: nasohypophyseal tract is present.
- 31 (41)** *Large lateral head vein*: 0 to 1: lateral head vein = anterior part of anterior cardinal or anterior vena cava present.

## HETEROSTRACI

- 4 (4)** *Pineal organ*: 2 to 1: pineal is covered, at least for the primitive forms. It is always covered by a “pineal macula” which is a zone of the dorsal carapace (dorsal shield) with a peculiar structure (thinner bone). Blicek & Goujet (1983) have described a pteraspid, called at that time *Zascinaspis laticephala* (now *Gigantaspis laticephala* after Pernègre & Goujet 2007), with a hole in place of the pineal macula, that they thought to be natural. This is exceptional and might be in fact an artefact, that is, a broken zone because of thinness of the bone at that place. So, must be coded 1.
- 6 (7)** *Optic tectum*: ? to 1: they had eyes, therefore an optic tectum should have been present; this assumption is less of a leap of faith than, say, Donoghue *et al.* (2000) inferring that having a pore-like gill opening indicates pouch-shaped gills.

## APPENDIX 2 — Continuation.

## 9-10 (15-16)

*Nasopharyngeal duct & single nasohypophyseal opening* respectively : must be coded 0 as there is no single nasohypophysial duct.

11 (17) *Position of nasohypophyseal opening* is not applicable, so remains ?.

32 (43) *Subaponeurotic vascular plexus*: 1 to ?: one should not use one single form (*Torpedaspis*), if all others do not show the structure.

35 (46) *Closely set unpaired fin ray supports*: 1 to ?: this character is non applicable. There is one unpaired fin, the caudal fin, but apparently no fin ray support.

47 (61) *Annular cartilage*: ? to 0: the structure of the mouth parts does not permit an annular cartilage.

60 (74) *Scale shape*: 1 to 1/2: both rod- and diamond-shaped scales occur, the former might be the primitive condition. In fact the patterns of scales are more diverse. They are “rod-shaped” in Ordovician taxa such as the arandaspids, which are not heterostracans, but basal pteraspidomorphs. They are rhombic (diamond-shaped) in *Eriptychius* (Ordovician, N America), and the APP group of Blicek *et al.* (1991), that is, anchipteraspids-protopteraspids-pteraspids in which are also included the psammosteids. They are more rectangular and thin on the flank of the cyathaspids (Blicek *et al.*'s 1991 CA group for cyathaspids-amphiaspids). However, rhombic-like scales also occur in this group: the smaller, lateral, lower scales of the trunk of cyathaspids, and a few isolated scales attributed to amphiaspids.

## ASTRASPIS

6 (7) *Optic tectum*: ? to 1: optic tectum should have been present (there are eyes).

21 (28) *Gills*: ? to 1: we would argue that the gill position is symmetrical.

33 (44) *Dorsal fin*: ? to 0: no separate dorsal fin.

34 (45) *Anal fin separate*: ? to 0: no separate ventral fin.

## ERIPTYCHIUS

6 (7) *Optic tectum*: ? to 1: optic tectum should have been present (there are eyes).

32 (43) *Subaponeurotic vascular plexus*: 1 to ?: We do not accept the homology between vascular canals in *Eriptychius* and the vascular plexus.

48 (62) *Trunk dermal skeleton*: ? to 1: trunk dermal skeleton must be present, if character 60 (74) is coded 1.

56 (70) *Enamellenameloid*: 1 to 2: the prismatic tissue is different to enamel in polarized light (Denison 1967).

## ARANDASPIDA

6 (7) *Optic tectum*: ? to 1: optic tectum should have been present (there are eyes).

17 (23) *Sensory line system with neuromasts*: ? to 1: we would argue that the lateral line system – even as grooves – has neuromasts.

36 (47) *Paired lateral fin folds*: ? to 0: no paired lateral fin folds.

38 (49) *Tail shape*: ? to 1: isocercal tail with median extension.

47 (61) *Annular cartilage*: ? to 0: the structure of the mouth parts does not permit an annular cartilage.

52 (66) *Spongy aspidin*: ? to 1: aspidin present.

53 (67) *Lamellar aspidin*: ? to 1: aspidin present.

54 (68) *Cellular bone*: ? to 0: Sansom *et al.* (2005) showed that arandaspids have a cellular bone.

## ANASPIDA

6 (7) *Optic tectum*: ? to 1: optic tectum should have been present (there are eyes).

9 (15) *Nasopharyngeal duct*: ? to 1: nasohypophysial opening is present.

17 (23) *Sensory line with neuromasts*: ? to 1: we would argue, if one accepts a lateral line system – even as grooves – it has neuromasts.

21 (28) *Gills*: ? to 1: we would argue that the gill position is symmetrical.

32 (43) *Subaponeurotic vascular plexus*: ? to 0: histology of scales points to subcutaneous vascularisation.

## JAMOYTIUS

6 (7) *Optic tectum*: ? to 1: optic tectum should have been present (there are eyes).

9 (15) *Nasopharyngeal duct*: ? to 1: if 11(17) (one nasohypophysial opening) is coded as 1, then this character (nasohypophysial duct) is also 1.

21 (28) *Gill relative position*: ? to 1: We would argue that the gill position is symmetrical.

47 (61) *Annular cartilage*: 1 to 0 (Janvier & Arsenault 2007).

## APPENDIX 2 — Continuation.

## EUPHANEROPS

Recoding follows Janvier & Arsenaault (2007); nevertheless here we interpret “lateral head stains” as eyes.

- 6 (7) *Optic tectum*: ? to 1: optic tectum should have been present (there are eyes).
- 21 (28) *Gill relative position*: ? to 1: symmetrical.
- 25 (32) *Opercular flaps*: ? to 0.
- 26 (34) *Gill lamellae with filaments*: ? to 1.
- 29 (38) *Closed pericardium*: ? to 1.
- 33 (44) *Dorsal fin*: ? to 0.
- 36 (47) *Paired lateral fin folds*: ? to 1.
- 37 (48) *Constricted pectorals*: ? to 0.
- 39 (50) *Preanal median fold*: ? to 0.
- 42 (54) *Trematic rings*: 1 to 0.
- 43 (55) *Arcualia*: ? to 1.
- 44 (58) *Braincase with lateral walls*: ? to 0.
- 48 (62) *Trunk dermal skeleton*: 1 to 0.
- 49 (63) *Perichondral bone*: ? to 0.
- 50 (64) *Calcified cartilage*: ? to 0.
- 51 (65) *Calcified dermal skeleton*: 1 to 0.
- 52-59 (66-73) ? to 0.
- 60 (74) *Scales*: 2 to 0.
- 66 (80) *Massive endoskeletal head shield over gills*: 0 to 1: “massive” endoskeleton covering gills dorsally.
- 67 (81) *Sclerotic ossicles*: ? to 0.

## OSTEOSTRACI

- 9 (15) *Nasopharyngeal duct*: 0 to 1: we would argue that there is a (very short) nasohypophysial duct.
- 12 (18) *Olfactory organ*: 1 to 2: olfactory organ is unpaired.

## GALEASPIDA

- 8 (14) *Number of nasal openings*: 2 to 1.
- 9 (15) *Nasopharyngeal duct*: 1 to 0: both nasal sacs lie on opposite sides of the median opening, and the unpaired hypophysial duct is in between.
- 33 (44) *Dorsal fin*: ? to 0: dorsal fin absent.
- 34 (45) *Anal fin separate*: 1 to 0: anal fin absent.
- 35 (46) *Closely set unpaired fin ray supports*: 1 to ? : not applicable.

## THELODONTI

*Loganellia scotica*

- 19 (26) *Sensory line*: 2 to 2/1: grooves could be possible too.
- 25 (32) *Opercular flaps*: 1 to 0: no opercular flaps known.

- 32 (43) *Subaponeurotic vascular plexus*: ? to 0: no indication of vascular plexus.

## PITURIASPIDA

Although there is no proof of bone or other calcified/phosphatic tissue, we accept consensus interpretation as a vertebrate rather than an arthropod.

- 65 (79) *Large head plates*: 0 to 1: if it is interpreted as a vertebrate, there are large plates.

## JAWED VERTEBRATES

- 4 (4) *Pineal organ*: 1 to 1/2: we do not know what is primitive (covered or uncovered pineal organ) in gnathostomes.
- 19 (26) *Sensory line*: 2 to 2/1: we do not know what is primitive (in grooves or canals) in gnathostomes.
- 20 (27) *Gill openings*: 0 to 2 : following new character codings.
- 32 (43) *Subaponeurotic vascular plexus*: 0 to 0/1: this type of system is present in the snout of sharks and primitive osteichthyans.
- 36 (47) *Paired lateral fin folds*: 0 to 0/1: possible stem gnathostomes from Northwest Territories, Canada (Wilson *et al.* 2007) as well as Lochkovian acanthothoracids from southeastern Australia (CJB pers. obs.) had ventrolateral rows of spines or spinelets.
- 55 (69) *Dentine*: 2 to 1: mesodentine is found in the oldest putative gnathostome (Ordovician *Skiichthys* Smith & Sansom 1997) and placoderms, regarded as the least derived gnathostome group.

## CONODONTA

- 1 (1) *Neural crest*: 1 to ? : the presence of dentine and extrinsic eye musculature on which the authors’ scoring for neural crest was based, are questioned.
- 2 (2) *Brain*: 1 to ? : presence of paired sensory organs and brain structure in conodonts is questionable.
- 13 (19) *Extrinsic eye musculature*: 1 to ? (linked to character 1): we are unconvinced that the paired anterior structures are eyes, let alone that the muscles associated with them are eye muscles.
- 21 (28) *Gill relative position*: 1 to ? : presence of gills is questionable, hence position in unknown or inapplicable.
- 35 (46) *Unpaired fin-ray supports closetset*: 1 to ? : we do not consider the fin-ray supports homologous with those of vertebrates.

## APPENDIX 2 — Continuation.

- 41 (52) *Visceral arches fused to neurocranium*: ? to 0: no visceral arches present.
- 42 (54) *Trematic ring*: ? to 0: no gills, so no trematic rings.
- 43 (55) *Arcualia*: ? to 0: arcualia never recorded.
- 51 (65) *Calcified dermal skeleton*: 1 to 0: many invertebrates have a calcified dermal skeleton, which is not homologous with that of vertebrates. Conodont elements are similarly not homologous with the dermal skeleton of any other group.
- 55 (69) *Dentine*: ? to 0: we do not accept that dentine is found in conodont elements.
- 56 (70) *Enamel/enameloid*: 1 to 0: there is no tissue with the optical properties of enamel.
- 59 (73) *Composition of the scales/denticles/teeth made up by several odontodes*: 2 to 0: conodont elements are “regarded as an independent experimentation, as a convergence to the dermal skeleton of vertebrates” by Reif (2002: 64), and cannot thus be homologous to odontodes.
- 63 (77) *Denticles in the pharynx*: 1 to 0: if one accepts Donoghue *et al.*'s (2000: 238) statement that “Whether conodont elements are homologous to true teeth is a moot point”, by their criteria, phosphatic tooth-like elements such as scolecodonts and phyllocarids would also “count” as teeth/denticles. Conodont elements are not homologous to thelodont and shark pharyngeal denticles (e.g., Turner 1985, 1994a; Reif 2006; Märss *et al.* 2007).

## APPENDIX 3

Apomorphy lists for character changes at nodes in examples of trees showing the same topology, generated by ACCTRAN (tree #6 of 27) and DELTRAN (tree #4 of 27) PAUP 4.0b10 analyses (see Fig. 5A, B). Character numbers (**Char.**) correspond to those described in Appendix 1; the black arrow “▶” under the change column represents unambiguous changes and the white arrow “▷” represents ambiguous changes; all changes are one step. Abbreviation: **CI**, consistency index. Note: 1, number in italics and parentheses corresponds to that used in the Donoghue *et al.*'s (2000; table 1) matrix.

ACCTRAN tree #6				DELTRAN tree #4			
Branch	Char. # 1	CI	Change	Branch	Char. # 1	CI	Change
node_34	30 (40)	0.333	0 ▶ 1	node_34	30 (40)	0.333	0 ▶ 1
▷ Cephalochordata				▷ Cephalochordata			
	36 (47)	0.25	0 ▶ 1		36 (47)	0.25	0 ▶ 1
	39 (50)	0.333	0 ▶ 1		39 (50)	0.333	0 ▶ 1
node_34 ▷ node_33	1 (1)	1	0 ▶ 1	node_34 ▷ node_33	38 (49)	0.667	0 ▶ 1
	2 (2)	1	0 ▶ 1		40 (51)	1	0 ▶ 1
	5 (5)	1	0 ▶ 1	node_33 ▷ node_32	1 (1)	1	0 ▶ 1
	6 (7)	1	0 ▶ 1	(Craniata)	2 (2)	1	0 ▶ 1
	8 (14)	0.667	0 ▶ 2		5 (5)	1	0 ▶ 1
	9 (15)	0.333	0 ▶ 1		6 (7)	1	0 ▶ 1
	10 (16)	0.333	0 ▶ 1		8 (14)	0.667	0 ▶ 2
	11 (17)	0.5	0 ▶ 1		9 (15)	0.333	0 ▶ 1
	12 (18)	1	0 ▶ 1		10 (16)	0.333	0 ▶ 1
	14 (20)	1	0 ▶ 1		12 (18)	1	0 ▶ 1
	18 (25)	0.667	0 ▶ 1		19 (26)	0.667	0 ▶ 1
	19 (26)	0.667	0 ▶ 1		20 (27)	1	0 ▶ 1
	20 (27)	1	0 ▶ 1		31 (41)	1	0 ▶ 1
	23 (30)	0.333	0 ▶ 1		41 (52)	0.5	0 ▶ 1
	31 (41)	1	0 ▶ 1	node_32 ▷ Myxinoidea	11 (17)	0.5	0 ▶ 1
	38 (49)	0.667	0 ▶ 1		14 (20)	1	0 ▶ 1
	40 (51)	1	0 ▶ 1		18 (25)	0.667	0 ▶ 1
node_33 ▷ node_32	41 (52)	0.5	0 ▶ 1		30 (40)	0.333	0 ▶ 1
node_32 ▷ Myxinoidea	30 (40)	0.333	0 ▶ 1		32 (43)	0.5	0 ▶ 1
	32 (43)	0.5	0 ▶ 1		39 (50)	0.333	0 ▶ 1
	39 (50)	0.333	0 ▶ 1	node_32	4 (4)	1	0 ▶ 2
				▷ node_31			
node_32 ▷ node_31	4 (4)	1	0 ▶ 2	(Vertebrata)	7 (8)	1	0 ▶ 1
	7 (8)	1	0 ▶ 1		11 (17)	0.5	0 ▶ 2
	11 (17)	0.5	1 ▶ 2		13 (19)	1	0 ▶ 1
	13 (19)	1	0 ▶ 1		14 (20)	1	0 ▶ 2
	14 (20)	1	1 ▶ 2		17 (23)	1	0 ▶ 1
	17 (23)	1	0 ▶ 1		18 (25)	0.667	0 ▶ 2
	18 (25)	0.667	1 ▶ 2		21 (28)	1	0 ▶ 1
	21 (28)	1	0 ▶ 1		23 (30)	0.333	0 ▶ 1
	26 (34)	1	0 ▶ 1		26 (34)	1	0 ▶ 1
	28 (37)	1	0 ▶ 1		28 (37)	1	0 ▶ 1
	29 (38)	1	0 ▶ 1		29 (38)	1	0 ▶ 1
	34 (45)	0.333	0 ▶ 1		43 (55)	1	0 ▶ 1
	43 (55)	1	0 ▶ 1	node_31 ▷ node_20	34 (45)	0.333	0 ▶ 1
	44 (58)	0.5	0 ▶ 1		47 (61)	1	0 ▶ 1
node_31 ▷ node_20	47 (61)	1	0 ▶ 1		50 (64)	0.333	0 ▶ 1
	50 (64)	0.333	0 ▶ 1	node_20	22 (29)	0.333	0 ▶ 1
				▷ Petromyzontida			
node_20	22 (29)	0.333	0 ▶ 1		33 (44)	0.333	0 ▶ 1
▷ Petromyzontida							
	33 (44)	0.333	0 ▶ 1		42 (54)	0.5	0 ▶ 1
	42 (54)	0.5	0 ▶ 1		44 (58)	0.5	0 ▶ 1
node_20	35 (46)	0.333	0 ▶ 1	node_20	35 (46)	0.333	0 ▶ 1
▷ Euphanerops				▷ Euphanerops			

## APPENDIX 3 — Continuation.

ACCTRAN tree #6				DELTRAN tree #4			
Branch	Char. # 1	CI	Change	Branch	Char. # 1	CI	Change
	36 (47)	0.25	0 > 1		36 (47)	0.25	0 > 1
	44 (58)	0.5	1 > 0		66 (80)	0.333	0 > 1
node_31 > node_30	66 (80)	0.333	0 > 1	node_31 > node_30	48 (62)	1	0 > 1
	3 (3)	0.5	0 > 1	(Euvertebrata)	51 (65)	1	0 > 1
	15 (21)	1	0 > 1		60 (74)	0.667	0 > 2
	45 (59)	1	0 > 1	node_30 > node_29	53 (67)	0.5	0 > 1
	46 (60)	1	0 > 1		59 (73)	0.667	0 > 2
	48 (62)	1	0 > 1		64 (78)	0.5	0 > 1
	51 (65)	1	0 > 1	node_29 > node_28	22 (29)	0.333	0 > 1
	53 (67)	0.5	0 > 1		60 (74)	0.667	2 > 1
	59 (73)	0.667	0 > 2	node_28 > node_27	3 (3)	0.5	1 > 0
node_30 > node_29	60 (74)	0.667	0 > 2		8 (14)	0.667	2 > 1
	36 (47)	0.25	0 > 1		9 (15)	0.333	1 > 0
	62 (76)	0.333	0 > 1		15 (21)	1	0 > 1
node_29 > node_28	64 (78)	0.5	0 > 1		32 (43)	0.5	0 > 1
	8 (14)	0.667	2 > 1		62 (76)	0.333	0 > 1
	9 (15)	0.333	1 > 0	node_27 > node_23	64 (78)	0.5	1 > 2
	22 (29)	0.333	0 > 1		52 (66)	1	0 > 1
	32 (43)	0.5	0 > 1		57 (71)	0.5	0 > 1
	60 (74)	0.667	2 > 1	node_23 > node_22	61 (75)	0.333	0 > 1
node_28 > node_27	67 (81)	0.5	0 > 1		4 (4)	1	2 > 1
	36 (47)	0.25	1 > 0		55 (69)	0.667	0 > 2
	57 (71)	0.5	0 > 1	node_22 > Heterostraci	10 (16)	0.333	1 > 0
	64 (78)	0.5	1 > 2		19 (26)	0.667	1 > 2
node_27 > node_23	10 (16)	0.333	1 > 0		23 (30)	0.333	1 > 0
	52 (66)	1	0 > 1		34 (45)	0.333	0 > 1
	58 (72)	0.5	0 > 1		38 (49)	0.667	1 > 0
	61 (75)	0.333	0 > 1		39 (50)	0.333	0 > 1
node_23 > node_22	65 (79)	0.333	0 > 1		58 (72)	0.5	0 > 1
	4 (4)	1	2 > 1		65 (79)	0.333	0 > 1
	38 (49)	0.667	1 > 0	node_22 > node_21	56 (70)	1	0 > 2
	39 (50)	0.333	0 > 1		64 (78)	0.5	2 > 1
	55 (69)	0.667	0 > 2	node_21 > <i>Astraspis</i>	18 (25)	0.667	2 > 1
node_22 > Heterostraci	67 (81)	0.5	1 > 0	node_21 > <i>Eriptychius</i>	50 (64)	0.333	0 > 1
	19 (26)	0.667	1 > 2		61 (75)	0.333	1 > 0
	23 (30)	0.333	1 > 0	node_23	22 (29)	0.333	1 > 0
node_22 > node_21	18 (25)	0.667	2 > 1	> Arandaspida			
	34 (45)	0.333	1 > 0		58 (72)	0.5	0 > 1
	56 (70)	1	0 > 2		60 (74)	0.667	1 > 2
	58 (72)	0.5	1 > 0		65 (79)	0.333	0 > 1
	64 (78)	0.5	2 > 1		67 (81)	0.5	0 > 1
	65 (79)	0.333	1 > 0	node_27 > node_26	68 (82)	0.5	0 > 1
node_21 > <i>Eriptychius</i>	50 (64)	0.333	0 > 1		16 (22)	1	0 > 1
	61 (75)	0.333	1 > 0		19 (26)	0.667	1 > 2
node_23 > Arandaspida	22 (29)	0.333	1 > 0		23 (30)	0.333	1 > 0
	60 (74)	0.667	1 > 2		44 (58)	0.5	0 > 1
	68 (82)	0.5	0 > 1		45 (59)	1	0 > 1
node_27 > node_26	16 (22)	1	0 > 1		46 (60)	1	0 > 1
	19 (26)	0.667	1 > 2		49 (63)	1	0 > 1
	23 (30)	0.333	1 > 0		50 (64)	0.333	0 > 1
	24 (31)	0.5	0 > 1	node_26 > node_25	67 (81)	0.5	0 > 1
	27 (35)	0.333	0 > 1		25 (32)	1	0 > 1
	35 (46)	0.333	0 > 1		33 (44)	0.333	0 > 1
	38 (49)	0.667	1 > 2		34 (45)	0.333	0 > 1
					37 (48)	1	0 > 1

## APPENDIX 3 — Continuation.

ACCTRAN tree #6				DELTRAN tree #4			
Branch	Char. # 1	CI	Change	Branch	Char. # 1	CI	Change
	49 (63)	1	0 > 1		38 (49)	0.667	1 > 2
	50 (64)	0.333	0 > 1		53 (67)	0.5	1 > 0
	66 (80)	0.333	0 > 1		54 (68)	1	0 > 1
node_26 > node_25	25 (32)	1	0 > 1		55 (69)	0.667	0 > 1
	33 (44)	0.333	0 > 1		57 (71)	0.5	0 > 1
	37 (48)	1	0 > 1	node_25 > node_24	68 (82)	0.5	0 > 1
	53 (67)	0.5	1 > 0		27 (35)	0.333	0 > 1
	54 (68)	1	0 > 1		35 (46)	0.333	0 > 1
	55 (69)	0.667	0 > 1	node_24 > Osteostraci	66 (80)	0.333	0 > 1
node_25 > node_24	68 (82)	0.5	0 > 1		3 (3)	0.5	1 > 0
	3 (3)	0.5	1 > 0		8 (14)	0.667	1 > 2
	8 (14)	0.667	1 > 2		9 (15)	0.333	0 > 1
	9 (15)	0.333	0 > 1		12 (18)	1	1 > 2
	12 (18)	1	1 > 2	node_24 > Pituriaspida	24 (31)	0.5	0 > 1
node_24 > Pituriaspida	65 (79)	0.333	0 > 1	node_24 > Pituriaspida	65 (79)	0.333	0 > 1
node_25	10 (16)	0.333	1 > 0	node_25	10 (16)	0.333	1 > 0
> Jawed vertebrate				> Jawed vertebrates			
	11 (17)	0.5	2 > 0	(Gnathostomata)	11 (17)	0.5	2 > 0
	14 (20)	1	2 > 3		14 (20)	1	2 > 3
	20 (27)	1	1 > 2		20 (27)	1	1 > 2
	24 (31)	0.5	1 > 0		30 (40)	0.333	0 > 1
	27 (35)	0.333	1 > 0		41 (52)	0.5	1 > 0
	30 (40)	0.333	0 > 1		56 (70)	1	0 > 1
	41 (52)	0.5	1 > 0		61 (75)	0.333	0 > 1
	56 (70)	1	0 > 1		62 (76)	0.333	1 > 0
	61 (75)	0.333	0 > 1		63 (77)	0.5	0 > 1
	62 (76)	0.333	1 > 0	node_26 > Galeaspida	64 (78)	0.5	2 > 1
	63 (77)	0.5	0 > 1		24 (31)	0.5	0 > 1
	64 (78)	0.5	2 > 1		27 (35)	0.333	0 > 1
node_26 > Galeaspida	66 (80)	0.333	1 > 0		59 (17)	0.667	2 > 1
	34 (45)	0.333	1 > 0		66 (80)	0.333	0 > 1
	57 (71)	0.5	1 > 0	node_28	33 (44)	0.333	0 > 1
				> <i>Loganellia scotica</i>			
	59 (73)	0.667	2 > 1		36 (47)	0.25	0 > 1
node_28	33 (76)	0.333	0 > 1		55 (69)	0.667	0 > 1
> <i>Loganellia scotica</i>							
	55 (77)	0.667	0 > 1		59 (17)	0.667	2 > 1
	59 (17)	0.667	2 > 1		63 (77)	0.5	0 > 1
	62 (76)	0.333	1 > 0	node_29 > Anaspida	36 (47)	0.25	0 > 1
	63 (77)	0.5	0 > 1		62 (76)	0.333	0 > 1
node_30 > <i>Jamoytius</i>	11 (17)	0.5	2 > 1	node_30 > <i>Jamoytius</i>	11 (17)	0.5	2 > 1
	27 (35)	0.333	0 > 1		27 (35)	0.333	0 > 1
	35 (46)	0.333	0 > 1		35 (46)	0.333	0 > 1
	42 (54)	0.5	0 > 1		42 (54)	0.5	0 > 1



## APPENDIX 4

Comparative character diagnostics – **CI** (consistency index), **HI** (homoplasy index), **RI** (retention index) values – for the 68 characters used in our analyses (Fig. 5A, B) and for the same 68 characters in tree 1 of three equal-length trees based on the codings in Donoghue *et al.* (2000), for the 17 base taxa (the strict consensus tree generated by their data for the 68 characters yields the same topology as their figure 7A tree).

Character	Donoghue <i>et al.</i> character number, with character description	Our analysis			Donoghue <i>et al.</i> analysis		
		CI	HI	RI	CI	HI	RI
1	(1 neural crest)	1	0	1	1	0	1
2	(2 brain)	1	0	0/0	1	0	1
3	(3 olfactory peduncles)	0.5	0.5	0.5	0.5	0.5	0.5
4	(4 pineal organ)	1	0	1	0.667	0.333	0
5	(5 divided pituitary)	1	0	1	1	0	1
6	(7 optic tectum)	1	0	1	1	0	1
7	(8 cerebellum)	1	0	1	1	0	1
8	(14 number of nasal openings)	0.667	0.333	0.667	0.667	0.333	0.5
9	(15 nasopharyngeal duct)	0.333	0.667	0.5	0.5	0.5	0
10	(16 single nasohyophyseal opening)	0.333	0.667	0.333	0.5	0.5	0.5
11	(17 position of nasohyophyseal opening)	0.5	0.5	0.333	0.5	0.5	0.333
12	(18 olfactory organ)	1	0	1	1	0	1
13	(19 extrinsic eye musculature)	1	0	1	1	0	1
14	(20 semicircular canals in labyrinth)	1	0	1	1	0	1
15	(21 vertical semicircular canal looped)	1	0	1	1	0	1
16	(22 endolymphatic ducts open externally)	1	0	1	1	0	1
17	(23 sensory line system with neuromasts)	1	0	1	1	0	1
18	(25 sensory line grooves or canals)	0.667	0.333	0.5	0.667	0.333	0.5
19	(26 sensory line)	0.667	0.333	0.667	0.667	0.333	0.8
20	(27 gill openings)	1	0	1	0.5	0.5	0.5
21	(28 gill relative position)	1	0	1	1	0	1
22	(29 elongate branchial series)	0.333	0.667	0.6	0.333	0.667	0.667
23	(30 gill openings lateral, slanting row)	0.333	0.667	0.5	0.333	0.667	0.6
24	(31 position of gill openings)	0.5	0.5	0	1	0	1
25	(32 opercular flaps)	1	0	1	0.5	0.5	0.5
26	(34 gill lamellae with filaments)	1	0	1	1	0	1
27	(35 mouth position)	0.333	0.667	0.333	0.5	0.5	0.667
28	(37 relative position atrium & ventricle)	1	0	1	1	0	1
29	(38 closed pericardium)	1	0	1	1	0	1
30	(40 paired dorsal aortae)	0.333	0.667	0	0.333	0.667	0
31	(41 lateral head vein)	1	0	1	1	0	1
32	(43 subaponeurotic vascular plexus)	0.5	0.5	0.667	0.333	0.667	0.333
33	(44 separate dorsal fin)	0.333	0.667	0.333	0.5	0.5	0.667
34	(45 anal fin separate)	0.333	0.667	0.5	0.333	0.667	0.6
35	(46 unpaired fin-ray supports closely set)	0.333	0.667	0.333	1	0	1
36	(47 paired lateral fin folds)	0.25	0.75	0	0.5	0.5	0.5
37	(48 constricted pectorals)	1	0	1	0.5	0.5	0.5
38	(49 tail shape)	0.667	0.333	0.667	0.667	0.333	0.75
39	(50 preanal median fold)	0.333	0.667	0	0.5	0.5	0.5
40	(51 ability to synthesise creatine phosphatase)	1	0	1	1	0	1
41	(52 visceral arches fused to neurocranium)	0.5	0.5	0.667	0.5	0.5	0.5
42	(54 trematic rings)	0.5	0.5	0	0.5	0.5	0.5
43	(55 arcualia)	1	0	1	1	0	1
44	(58 braincase with lateral walls)	0.5	0.5	0.667	1	0	1
45	(59 neurocranium entirely closed dorsally, covering brain)	1	0	1	1	0	1
46	(60 occiput enclosing vagus and glossopharyngeal)	1	0	1	1	0	1
47	(61 annular cartilage)	1	0	1	0.5	0.5	0.5

## APPENDIX 4 — Continuation.

Character	Donoghue <i>et al.</i> character number, with character description	Our analysis			Donoghue <i>et al.</i> analysis		
		CI	HI	RI	CI	HI	RI
48	(62 trunk dermal skeleton)	1	0	1	1	0	1
49	(63 perichondral bone)	1	0	1	1	0	1
50	(64 calcified cartilage)	0.333	0.667	0.6	0.5	0.5	0.75
51	(65 calcified dermal skeleton)	1	0	1	1	0	1
52	(66 spongy aspidin)	1	0	1	0.5	0.5	0.5
53	(67 lamellar aspidin)	0.5	0.5	0.8	0.333	0.667	0.6
54	(68 cellular bone)	1	0	1	0.5	0.5	0
55	(69 dentine)	0.667	0.333	0.75	0.4	0.6	0.25
56	(70 enamel/oid)	1	0	1	0.667	0.333	0.5
57	(71 three-layered exoskeleton)	0.5	0.5	0.8	0.333	0.667	0.6
58	(72 cancellar layer in exoskeleton with honeycomb-shaped cavities)	0.5	0.5	0	1	0	1
59	(73 composition of scales/denticles)	0.667	0.333	0.833	0.667	0.333	0.75
60	(74 scales)	0.667	0.333	0.857	0.667	0.333	0.857
61	(75 oakleaf-shaped tubercles)	0.333	0.667	0.333	0.5	0.5	0.667
62	(76 oral plates)	0.333	0.667	0.5	0.333	0.667	0.5
63	(77 denticles in pharynx)	0.5	0.5	0	0.333	0.667	0
64	(78 dermal head covering in adult state)	0.5	0.5	0.75	0.5	0.5	0.75
65	(79 large unpaired ventral and dorsal dermal plates on head)	0.333	0.667	0	1	0	1
66	(80 massive endoskeletal head shield covering gills dorsally)	0.333	0.667	0.333	1	0	1
67	(81 sclerotic ossicles)	0.5	0.5	0.667	0.5	0.5	0.667
68	(82 ossified scleral capsule)	0.5	0.5	0.5	0.333	0.667	0