

FAMILIES OF NEGATIVELY CURVED HERMITIAN MANIFOLDS

MICHAEL J. COWEN

ABSTRACT. A complex analytic family of compact hermitian manifolds has negative holomorphic sectional curvature in a neighborhood of any fibre having negative holomorphic sectional curvature.

1. Introduction. In [1, Hilfssatz 4, p. 120], Grauert and Reckziegel state:

If (Y, π, X) is an analytic family of compact Riemann surfaces of genus ≥ 2 , over a Riemann surface X , then for each point x_0 in X there is a neighborhood $V \subset X$ of x_0 and a differential metric on $Y|V$ such that $Y|V$ is strongly negatively curved.

Their construction of the metric is clear, but the proof of strong negative curvature involves a computation which is not altogether complete. (The proof, however, can be completed easily using the formula for the Gaussian curvature of the sum of two hermitian metrics [1, Aussage 1, p. 111].) The purpose of this note is to give a simpler computation which shows that the metric actually has holomorphic sectional curvature $\leq c < 0$ and hence *a fortiori* is strongly negatively curved [2, p. 39]. Indeed we will show that if the fibres of Y are n -dimensional compact manifolds each having holomorphic sectional curvature less than a negative constant, then $Y|V$ has negative holomorphic sectional curvature.

2. Definitions and statement of results. Let ds^2 be a hermitian metric on an n -dimensional complex manifold M , with $ds^2 = \sum g_{ij} dz_i d\bar{z}_j$ in local coordinates. Define the curvature tensor by

$$K_{ijkm} = \frac{\partial^2 g_{ij}}{\partial z_k \partial \bar{z}_m} - \sum_{p,q} \frac{\partial g_{ip}}{\partial z_k} g^{pq} \frac{\partial g_{qj}}{\partial \bar{z}_m} \quad \text{for } 1 \leq i, j, k, m \leq n.$$

M has holomorphic sectional curvature (which will be denoted by h.s.c.) less than a constant c if $-\sum k_{ijkm} s_i \bar{s}_j s_k \bar{s}_m < c$ for all holomorphic unit tangent vectors $s = \sum s_i \partial / \partial z_i$.

Received by the editors June 5, 1972.

AMS (MOS) subject classifications (1970). Primary 53C55; Secondary 32G05, 32H20.

Key words and phrases. Holomorphic sectional curvature, hyperbolic manifold, analytic family, differential metric.

© American Mathematical Society 1973

THEOREM. *Let (Y, π, X) be an analytic family of compact complex n -dimensional manifolds over a Riemann surface X , such that for each $x_0 \in X$, the fibre $Y_{x_0} = \pi^{-1}(x_0)$ has a hermitian metric of h.s.c. $\langle c \rangle < 0$, then there exists a neighborhood V of x_0 in X such that $Y|V$ has a hermitian metric of h.s.c. $\langle c' \rangle < 0$, with c, c' constants.*

REMARK. This generalizes Grauert and Reckziegel's result since a compact Riemann surface of genus ≥ 2 has a hermitian metric of Gaussian curvature (equals h.s.c. on a Riemann surface) less than a negative constant [2, Theorem 5.1, p. 12].

COROLLARY. *If σ is a holomorphic section of Y with isolated singularities in X , then σ extends as a holomorphic section to all of X .*

3. Construction of the metric. The construction is the obvious generalization of that in [1].

Since (Y, π, X) is locally trivial we can find a neighborhood V with coordinate z_{n+1} centered at x_0 , and neighborhoods U_1, \dots, U_r in Y such that $Y|V = \bigcup U_m$, each U_m has coordinates z_1, \dots, z_{n+1} with

$$\pi(z_1, \dots, z_{n+1}) = z_{n+1},$$

and z_1, \dots, z_n give coordinates in $U_m \cap Y_{z_{n+1}}$ for all z_{n+1} in V . The hermitian metric on $Y_0 = Y_{x_0}$ is of the form $\sum g_{ij}(z_1, \dots, z_n) dz_i d\bar{z}_j$ on $U_m \cap Y_0$ and thus can be extended to a pseudo-hermitian metric $\sum h_{ij} dz_i d\bar{z}_j$ on U_m , $1 \leq i, j \leq n$, by setting $h_{ij}(z_1, \dots, z_{n+1}) = g_{ij}(z_1, \dots, z_n)$. These pseudo-hermitian metrics can then be patched together by a partition of unity to give a pseudo-hermitian metric α on $Y|V$, such that $\alpha|Y_0$ is the original hermitian metric on Y_0 . That is, $\alpha = \sum k_{ij} dz_i d\bar{z}_j$ ($1 \leq i, j \leq n+1$) on U_m and $k_{ij}(z_1, \dots, z_n, 0) = g_{ij}(z_1, \dots, z_n)$ for $1 \leq i, j \leq n$. Since $\alpha|Y_0$ has h.s.c. $\langle c \rangle < 0$, it is clear that for z_{n+1} close enough to 0, $\alpha|Y_{z_{n+1}}$ will have h.s.c. $\langle c \rangle < 0$. By shrinking V we can assume (1) $V = \{|z_{n+1}| < t\}$, (2) $\alpha|Y_{z_{n+1}}$ has h.s.c. $\langle c \rangle < 0$ for all $z_{n+1} \in V$, and (3) k_{ij} , its first, and second partial derivatives are bounded on each U_m for $1 \leq i, j \leq n+1$. Since a disc in \mathbb{C} has a hermitian metric of Gaussian curvature -1 , we can put V in a larger disc and obtain a metric $h(z_{n+1}) dz_{n+1} d\bar{z}_{n+1}$ on V of Gaussian curvature $= -1$ such that h , its first, and second partials are bounded on V . Define a metric ds^2 on $Y|V$ for each $\lambda > 0$ by $ds^2 = \alpha + \lambda h(z_{n+1}) dz_{n+1} d\bar{z}_{n+1}$, i.e. on U_m , $ds^2 = \sum k_{ij} dz_i d\bar{z}_j + \lambda h(z_{n+1}) dz_{n+1} d\bar{z}_{n+1}$. Note that for large λ , ds^2 has negative h.s.c. in both the fibre and base directions. We wish to choose λ_0 so that for all $\lambda \geq \lambda_0$, ds^2 will have h.s.c. $\leq c'' < 0$. Clearly it suffices to do this on each U_m and then take the maximum of the λ_0 's so obtained.

4. **Proof of negative sectional curvature.** Assume we have shown the following:

(i) $K_{ijkm} \rightarrow \tilde{K}_{ijkm}$ for $1 \leq i, j, k, m \leq n$, uniformly on U_m as $\lambda \rightarrow \infty$, where $\tilde{K}_{ijkm}(z_1, \dots, z_{n+1})$ is the curvature of ds^2 restricted to $Y_{z_{n+1}} \cap U_m$.

(ii) $|K_{ijkm}| \leq M$ on U_m for all $1 \leq i, j, k, m \leq n+1$ except when $i=j=k=m=n+1$, and M is a constant.

$$(iii) \quad K_{ijkm} = \lambda \left(\frac{\partial^2 h}{\partial z_{n+1} \partial \bar{z}_{n+1}} - \frac{1}{h} \frac{\partial h}{\partial z_{n+1}} \frac{\partial h}{\partial \bar{z}_{n+1}} \right) + O(1),$$

when $i=j=k=m=n+1$, where $O(1)$ means a term which is uniformly bounded on U_m .

Since the Gaussian curvature of h is

$$\frac{1}{h} \left(\frac{\partial^2 h}{\partial z_{n+1} \partial \bar{z}_{n+1}} - \frac{1}{h} \frac{\partial h}{\partial z_{n+1}} \frac{\partial h}{\partial \bar{z}_{n+1}} \right) \leq -1$$

and h is bounded on U_m , we have:

(iii)' $K_{ijkm} \leq \lambda c'$ when $i=j=k=m=n+1$, where $c' < 0$ is a constant, for $\lambda \geq \lambda_0$.

Fix $z = (z_1, \dots, z_{n+1})$. If $s = \sum_{i=1}^n s_i (\partial/\partial z_i)$ is a holomorphic unit tangent vector to the fibre $Y_{z_{n+1}}$ then by (i) we have

$$-\sum K_{ijkm} s_i \bar{s}_j s_k \bar{s}_m \rightarrow -\sum \tilde{K}_{ijkm} s_i \bar{s}_j s_k \bar{s}_m < c < 0 \quad \text{as } \lambda \rightarrow \infty.$$

Hence by compactness of the unit sphere, we can choose λ_0 large enough so that for $\lambda \geq \lambda_0$ we have $-\sum K_{ijkm} s_i \bar{s}_j s_k \bar{s}_m < c$ for s tangent to the fibre. But if $s = \sum_{i=1}^{n+1} s_i \partial/\partial z_i$ is any holomorphic unit tangent vector, then by (ii) and (iii)' we have:

$$(*) \quad -\sum K_{ijkm} s_i \bar{s}_j s_k \bar{s}_m \leq -\sum_{i,j,k,m=1}^n K_{ijkm} s_i \bar{s}_j s_k \bar{s}_m + M \sum' |s_i| |s_j| |s_k| |s_m| + \lambda c' |s_{n+1}|^4,$$

where \sum' is the sum of the terms where at least one, but not all, of the i, j, k, m equals $n+1$. Thus if s is not tangent to the fibre, i.e., $s_{n+1} \neq 0$, then by taking λ_0 large enough we can insure that the h.s.c. is less than c_s in a neighborhood of s on the unit sphere, for all $\lambda \geq \lambda_0$. But from (*) it is also clear that if s is tangent to the fibre, then the h.s.c. is less than c_s in a neighborhood of s for all $\lambda \geq \lambda_0$. Therefore for each fixed z the h.s.c. at z is less than c_z for $\lambda > \lambda_0$ and hence by the relative compactness of U_m , the h.s.c. $< c < 0$ on U_m for $\lambda \geq \lambda_0$, which proves the Theorem.

Let $ds^2|_{Y_{z_{n+1}}} = \sum \tilde{k}_{ij} dz_i d\bar{z}_j$ be the metric restricted to the fibre, where $\tilde{k}_{ij} = k_{ij}$ for $1 \leq i, j \leq n$. Since

$$ds^2 = \sum k_{ij} dz_i d\bar{z}_j + \lambda h(z_{n+1}) dz_{n+1} d\bar{z}_{n+1} \equiv \sum g_{ij} dz_i d\bar{z}_j$$

where $1 \leq i, j \leq n+1$, it is easy to check that:

(a) $g^{ij} = \tilde{k}^{ij} + O(\lambda^{-1})$, $\partial g_{ip} / \partial z_k = \partial \tilde{k}_{ip} / \partial z_k$, $\partial^2 g_{ij} / \partial z_k \partial \bar{z}_m = \partial^2 \tilde{k}_{ij} / \partial z_k \partial \bar{z}_m$ for $1 \leq i, j, k, m, p \leq n$.

$$(b) \quad g^{ij} = \lambda^{-1} h(z_{n+1})^{-1} + O(\lambda^{-2}), \quad \frac{\partial g_{ip}}{\partial z_{n+1}} = O(1) + \frac{\partial h}{\partial z_{n+1}},$$

$$\frac{\partial^2 g_{ij}}{\partial z_{n+1} \partial \bar{z}_{n+1}} = O(1) + \lambda \frac{\partial^2 h}{\partial z_{n+1} \partial \bar{z}_{n+1}} \quad \text{for } i = j = p = n + 1.$$

(c) $g^{ij} = O(\lambda^{-1})$, $\partial g_{ip} / \partial z_k = O(1)$, $\partial^2 g_{ij} / \partial z_k \partial \bar{z}_m = O(1)$ otherwise. (Note. Since h is a function only of z_{n+1} , terms such as $\partial g_{ip} / \partial z_k$, for $i=p=n+1$ but $k \neq n+1$, do not involve λ or the derivatives of h .)

If $1 \leq i, j, k, m \leq n$ then

$$K_{ijkm} = \frac{\partial^2 \tilde{k}_{ij}}{\partial z_k \partial \bar{z}_m} - \sum_{p,q=1}^n \frac{\partial \tilde{k}_{ip}}{\partial z_k} (\tilde{k}^{pq} + O(\lambda^{-1})) \frac{\partial \tilde{k}_{qj}}{\partial \bar{z}_k} + O(\lambda^{-1})$$

$$= \tilde{K}_{ijkm} + O(\lambda^{-1}),$$

which proves (i). If $i=j=k=m=n+1$, then

$$K_{ijkm} = \lambda \frac{\partial^2 h}{\partial z_{n+1} \partial \bar{z}_{n+1}} + O(1) - \sum_{p,q=1}^n \frac{\partial g_{ip}}{\partial z_k} O(\lambda^{-1}) \frac{\partial g_{qj}}{\partial \bar{z}_m}$$

$$- \sum_{p=1}^n \frac{\partial g_{ip}}{\partial z_k} O(\lambda^{-1}) O(1) - \sum_{q=1}^n O(1) O(\lambda^{-1}) \frac{\partial g_{qj}}{\partial \bar{z}_k}$$

$$- (O(1) + \lambda \partial h / \partial z_{n+1})(\lambda^{-1} h^{-1} + O(\lambda^{-2}))(O(1) + \lambda(\partial h / \partial \bar{z}_{n+1}))$$

$$= \lambda(\partial^2 h / \partial z_{n+1} \partial \bar{z}_{n+1}) - h^{-1}(\partial h / \partial z_{n+1})(\partial h / \partial \bar{z}_{n+1})$$

$$+ O(1) + O(\lambda^{-1}) + O(\lambda^{-2})$$

which proves (iii). The proof of (ii) is obvious, since the only terms which are not $O(1)$ or $O(\lambda^{-1})$ are those appearing only when $i=j=k=m=n+1$.

5. Proof of Corollary. Assume σ has an isolated singularity at $x_0 \in H$. By the Theorem, there is a neighborhood $V = \{|z| < 1\}$ of x_0 such that $Y|V$ has a metric of h.s.c. $< c < 0$. Thus by [2, Theorem 4.11, p. 61], $Y|V$ is hyperbolic and, by a theorem of Mrs. Kwack [2, Theorem 3.1, p. 83], $\sigma: V - \{0\} \rightarrow Y|V$ has a holomorphic extension to $\sigma': V \rightarrow Y|V$ if there exists a suitable sequence of points $x_n \rightarrow x_0$ such that $\sigma(x_n) \rightarrow p_0 \in Y|V$. Since $Y|V$ is relatively compact in Y , the result follows.

6. Remarks. That X is a Riemann surface was not crucial to the proof of the Theorem and the proof goes through with obvious modifications when X is an arbitrary complex manifold. Then in the Corollary, σ

need only have singularities contained in an analytic set of codimension ≥ 1 in X , for σ to extend to all of X . The proof of the Corollary then follows from a result of Mrs. Kwack [2, Theorem 4.1, p. 86].

REFERENCES

1. H. Grauert and H. Reckziegel, *Hermiteische Metriken und normale Familien holomorpher Abbildungen*, Math. Z. **89** (1965), 108–125. MR **33** #2827.
2. S. Kobayashi, *Hyperbolic manifolds and holomorphic mappings*, Pure and Appl. Math., no. 2, Marcel Dekker, New York, 1970. MR **43** #3503.

DEPARTMENT OF MATHEMATICS, TULANE UNIVERSITY, NEW ORLEANS, LOUISIANA 70118

Current address: Department of Mathematics, Princeton University, Princeton, New Jersey 08540