

Fast and slow responses of Southern Ocean sea surface temperature to SAM in coupled climate models

Yavor Kostov · John Marshall · Ute Hausmann · Kyle C. Armour · David Ferreira · Marika Holland

Received: date / Accepted: date

Abstract We investigate how sea surface temperatures (SSTs) around Antarctica respond to the Southern Annular Mode (SAM) on multiple timescales. To that end we examine the relationship between SAM and SST within unforced preindustrial control simulations of coupled general circulation models (GCMs) included in the Climate Modeling Intercomparison Project phase 5 (CMIP5). We develop a technique to extract the response of the Southern Ocean SST to a hypothetical step increase in the SAM index. We demonstrate that in many GCMs, the expected SST step response function is nonmonotonic in time. Following a shift to a positive SAM anomaly, an initial cooling regime can transition into surface warming around Antarctica. However, there are large differences across the CMIP5 ensemble. In some mod-

els the step response function never changes sign and cooling persists, while in other GCMs the SST anomaly crosses over from negative to positive values only three years after a step increase in the SAM. This intermodel diversity can be related to differences in the models' climatological thermal ocean stratification in the region of seasonal sea ice around Antarctica. Exploiting this relationship, we use observational data for the time-mean meridional and vertical temperature gradients to constrain the real Southern Ocean response to SAM on fast and slow timescales.

Keywords Southern Ocean · Southern Annular Mode · surface westerlies · Atmosphere-ocean interaction · CMIP5

Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA 02139, USA
Tel.: +617-324-4569
E-mail: yavor@mit.edu

J. Marshall
Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

U. Hausmann
Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

K. C. Armour
University of Washington School of Oceanography, Box 357940, Seattle, WA 98195-7940, USA

D. Ferreira
Department of Meteorology, University of Reading, P.O. Box 243, Reading RG6 6BB, United Kingdom

M. M. Holland
National Center for Atmospheric Research, Boulder, CO 3090, USA

1 Introduction

In contrast to the strong global warming trend, the Southern Ocean (SO) has exhibited a gradual decrease in sea surface temperatures (SSTs) over recent decades (Figure 1, [Fan *et al.*, 2014; Armour *et al.*, 2015]). The large-scale geographic patterns of delayed and accelerated warming are related to the climatological background ocean circulation [Marshall *et al.*, 2014; Marshall *et al.*, 2015, Armour *et al.*, 2015; Hutchinson *et al.*, 2013; and Hutchinson *et al.*, 2015]. The high latitudes of the SO constitute an open channel with zonal flow, where the Antarctic Circumpolar Current (ACC) acts to thermally isolate the Antarctic region and limit poleward heat transport into the SO [Hutchinson *et al.*, 2013; Hutchinson *et al.*, 2015]. Moreover, in this region deep waters, unmodified by greenhouse gas forcing, are upwelled at the surface where they take up heat as the mean wind-driven circulation – partially compensated by the eddy circulation – transports them northward [Marshall *et al.*, 2015; Armour *et al.*, 2015]. The background circulation can therefore slow down the rate of surface warming in the SO relative to the

rest of the World Ocean. However, this mechanism of passive heat transport is not sufficient to explain the persistent cooling trends around Antarctica.

Some studies interpret the pattern of observed Southern Hemisphere SST trends as a response to a poleward shift and strengthening of the westerlies. These recent tendencies in the atmospheric circulation resemble the positive phase of the Southern Annular Mode (SAM) of natural variability, but they may in fact be a forced response [Thomas *et al.*, 2015], the result of ozone depletion [Thompson and Solomon, 2002; Gillett and Thompson, 2003; Sigmond *et al.*, 2011; Thompson *et al.*, 2011; Wang *et al.*, 2014]. Figure 1 illustrates the synchronous evolution of observed SST and SAM anomalies over the SO. The SST averaged between 55°S to 70°S is negatively correlated to the SAM index with $R = -0.65$ at a lag of 1 year. Multiple mechanisms have been proposed to explain the relationship between SST trends around Antarctica and poleward intensification of the westerlies.

Many studies conclude that a poleward intensification of the westerlies impacts SO SSTs by changing the ocean circulation [e.g., Hall and Visbeck, 2002; Oke and England, 2004; Russell *et al.*, 2006; Fyfe *et al.*, 2007; Ciasto and Thompson, 2008; Bitz and Polvani, 2012; Marshall *et al.*, 2014; Purich *et al.*, 2016]. The recent circulation changes have been confirmed by measurements of dissolved passive tracers [Waugh *et al.*, 2013; Waugh, 2014]. A positive SAM induces anomalous northward Ekman transport in the high latitude region of the Southern Hemisphere [Hall and Visbeck, 2002]. This gives rise to surface cooling poleward of 50°S. Ciasto and Thompson [2008] and Sen Gupta and England [2006] propose that the aforementioned oceanic mechanism complements SAM induced changes in the surface heat fluxes, and that both processes act in concert to set the spatial distribution of temperature anomalies around Antarctica.

Unlike Ciasto and Thompson [2008], Bitz and Polvani [2012] demonstrate that in the coupled CCSM3.5 GCM, an ozone-driven poleward intensification of the westerlies leads to an increase in SSTs throughout the SO. This result implies that changes in the winds cannot account for the observed cooling around Antarctica and may even have the opposite effect. Bitz and Polvani [2012] explain that poleward intensification by itself can lead to a positive SST response via anomalous Ekman upwelling of warmer water in the salinity-stratified circumpolar region. This highlights an apparent divergence in literature about the sign of the SO SST anomalies associated with a SAM-like pattern. A similar lack of consensus also carries over to studies which explore the connection between the westerly winds and SO sea ice. Hall and Visbeck [2002] suggest that a positive SAM causes sea ice expansion, while Sigmond and Fyfe [2014]

demonstrate that poleward intensification (forced by ozone depletion) is associated with a decrease in marine ice extent.

Ferreira *et al.* [2015] propose a theoretical framework that can resolve this ostensible disagreement about the sign of the SST anomaly associated with a poleward intensification of the westerlies. They use two different coupled GCMs to demonstrate that the SO response to winds in forced ozone depletion simulations is timescale-dependent. An atmospheric pattern similar to a positive SAM triggers short-term cooling followed by slow warming around Antarctica. The fast response is dominated by horizontal Ekman drift advecting colder water northward, while the slow response is sustained by Ekman upwelling of warmer water. Ferreira *et al.* [2015] show that the transition between the cooling and warming regime differs between two coupled GCMs.

In our work we examine how the SO responds to a poleward intensification of the westerlies in 23 state-of-the-art CMIP5 coupled models [Taylor *et al.*, 2012]. By analyzing the GCMs' control simulations, we are able to study the relationship between SAM and SO SST anomalies even in models which have not performed wind override experiments or targeted ozone depletion simulations. In agreement with Ferreira *et al.* [2015], our findings suggest that anomalous Ekman transport governs the SO response to SAM on interannual and decadal timescales. Furthermore, we interpret the diversity in the fast and slow responses across the CMIP5 ensemble in terms of the models' time-mean SO stratification. Finally, we use observational data for the ocean temperature climatology to constrain the SST step response function of the real SO.

2 Data and methods

The GCMs used in this study have made their experimental results publicly available through the CMIP5 initiative [Taylor *et al.*, 2012]. In our ensemble we include 23 models that have archived their output of ocean potential temperature, SST, and sea level pressure (SLP). We examine data from the CMIP5 preindustrial control simulations (pi-Control), which do not have any sources of external forcing. Thus all climate anomalies that we observe in these experiments can be attributed to internal variability. Moreover, the control simulations are hundreds of years long allowing us to perform statistical analysis with large samples of data. Table 1 provides additional information about the length of individual CMIP5 simulations. In order to conduct our analysis consistently across the ensemble, we convert all model output fields to the same regular latitude-longitude grid ($0.5^\circ \times 1^\circ$).

We define an annual-mean index for the SAM in each model as the first principal component of variability in SLP south of 20°S. Positive values of this index correspond to a poleward intensification of the westerly winds. In order

to remove the secular drift, we linearly detrend the SAM timeseries.

We calculate an area-weighted average of the annual-mean SST anomalies between 55°S and 70°S. As with the SAM index, we detrend the SST timeseries to eliminate the long-term drift. A comparison of the SO SST anomalies against the SAM index in CMIP5 models shows negative correlations at short lags (Figure 2). This is reminiscent of the synchronous evolution of westerly winds and SO SST seen in observations (Figure 1).

For each GCM, we estimate the impulse response function G (Green's function) of SO SST (55°S to 70°S) with respect to the SAM index. Following *Hasselmann et al.* [1993], we assume that the temperature timeseries can be represented as a convolution of G with a previous history of the SAM forcing:

$$\begin{aligned} SST(t) &= \int_0^{+\infty} G(\tau)SAM(t-\tau)d\tau + \varepsilon \\ &\approx \int_0^{\tau_{max}} G(\tau)SAM(t-\tau)d\tau + \varepsilon, \end{aligned} \quad (1)$$

where $SAM(t)$ is the SAM index normalized by its standard deviation σ_{SAM} , τ is the time lag in years, τ_{max} is an imposed maximum cutoff lag, and ε is residual noise. We discretize equation (1) to obtain

$$SST(t) \approx \sum_{i=0}^{\tau_{max}} G(i)SAM(t-i) + \varepsilon, \quad (2)$$

where coefficients $G(i)$ represent the response at different time steps after an impulse perturbation of magnitude σ_{SAM} . We then use a linear least-squares regression of the SO SST signal against the lagged SAM index to estimate $G(i)$ for $i = 0, \dots, \tau_{max}$.

In our intercomparison we take into account differences in the magnitude of SAM variability across the set of 23 models. We calculate $\overline{\sigma_{SAM}^{Ens}}$, the ensemble mean of the index standard deviations σ_{SAM} . We then rescale the estimated impulse response functions for each GCM, where we multiply $G(i)$ by the corresponding nondimensional ratio $\overline{\sigma_{SAM}^{Ens}}/\sigma_{SAM}$.

By selecting multiple shorter SST and SAM timeseries from the full control simulation and by varying the cutoff lag τ_{max} , we obtain a spread of estimates for the impulse response function $G(\tau)$ in a given model. Table 2 lists our fitting parameters and their values. We use the residuals ε to quantify the uncertainty $\sigma_{GreensFit}(t)$ on each of these least squares regressions. Figure 3a shows examples of impulse response estimates for three CMIP5 models, rescaled by $\overline{\sigma_{SAM}^{Ens}}$. Multiple fits span envelopes of uncertainty, while vertical bars denote the error margins $\sigma_{GreensFit}(t)$ on each fit. Note that in our analysis we use annual-mean SST. Hence the estimated Year 0 response is not zero, as it represents an average of the SST anomaly over the first months after a positive SAM impulse.

We integrate the impulse response function fits to obtain a spread of estimates for the SO step response function:

$$\begin{aligned} SST_{Step}(t) &= \int_0^t G(\tau)d\tau \\ &\approx \sum_{i=0}^t G(i), t \leq \tau_{max}. \end{aligned} \quad (3)$$

Each of the estimates corresponds to a different combination of start and end times for the timeseries, as well as different choices of τ_{max} . We calculate the mean $SST_{Step}(t)$ and the standard deviation $\sigma_{Spread}(t)$ which characterize our envelope of step response functions for a given model. We furthermore use the $\sigma_{GreensFit}(t)$ values to constrain the margin of error $\sigma_{StepFit}(t)$ on each individual estimate in our spread. We then combine $\sigma_{StepFit}(t)$ and $\sigma_{Spread}(t)$ in quadrature in order to quantify the total uncertainty $\sigma_{SST_{Step}}(t)$ on the mean $SST_{Step}(t)$ for a given GCM.

Figure 3b shows example step response functions calculated for the three models presented in Figure 3a. For each of the GCMs, we have shown only 100 different estimates, illustrating envelopes of uncertainty. In comparison, in our full analysis we perform more than 350 fits per model by selecting shorter timeseries from the control simulation and by specifying different values for τ_{max} . The vertical bars in Figure 3 indicate the error margins $\sigma_{StepFit}(t)$ based on the residuals of each regression.

The step response results are integral quantities, and hence they are smoother than the corresponding impulse response functions. However, a drawback is that the integrated errors grow larger in time. Nevertheless, Figure 3b demonstrates that even with generous envelopes of uncertainty and large error bars on the individual fits, we can still distinguish the estimated step response functions of different CMIP5 models.

We use synthetic noisy signals and artificially constructed systems with known step responses in order to test our methodology. The verification procedure is described and illustrated in detail in Appendix A. Multiple tests confirm the validity of our approach for estimating the SO response functions.

3 Results

Our estimated step response functions suggest notable inter-model differences in the SO SST response to SAM across the CMIP5 ensemble (Figure 4). Although all GCMs show initial cooling, many of them transition into a regime of gradual warming. If forced with a positive step increase in the SAM, a number of CMIP5 models – such as CanESM2, CCSM4, and CESM-CAM5 – are expected to show positive SST anomalies in the SO within a few years. In contrast, other ensemble members, including CNRM-CM5 and GFDL-ESM2M, do not exhibit such nonmonotonic response to a poleward intensification of the westerlies and instead

maintain negative temperature anomalies persisting for longer than a decade. What sets this intermodel diversity in the way the SO reacts to SAM on short and long timescales?

Following *Ferreira et al.* [2015], we examine whether the fast cooling regime is governed by northward wind-driven transport, advecting colder water up the climatological SO SST gradient. The anomalous horizontal heat flux Q_{hor} [W/m^2] in the SO Ekman layer of depth Z_{Ek} scales as

$$Q_{hor} \propto C_p \frac{[\tau'_x]}{f Z_{Ek}} \partial_y [\overline{SST}], \quad (4)$$

where $C_p \approx 4 \times 10^3$ J/kg/K is the specific heat of water, $[\tau'_x]$ is the zonally averaged zonal component of the anomalous surface wind-stress associated with SAM, f is the Coriolis parameter, and $\partial_y [\overline{SST}]$ is the meridional gradient of the zonally averaged climatological SST. As in *Ferreira et al.* [2015], we have assumed that eddy compensation in the thin Ekman layer is much smaller than the anomalous northward wind-driven transport. Since we have rescaled each SST response function by $\overline{\sigma_{SAM}^{Ens}}/\sigma_{SAM}$, we have eliminated some of the intermodel differences due to $[\tau'_x]$.

When we regress the estimated Year 1 cooling anomalies from our step responses against $\partial_y [\overline{SST}]$ averaged between 55° and 70° S, we see a strong anticorrelation with a Pearson's $R = -0.72$ (Figure 5a). This result is significant at the 5% level with $p < 0.01$ and highlights the importance of horizontal Ekman transport for governing the fast cooling regime during a positive phase of the SAM.

We also consider the role of Ekman upwelling for setting the long-term response to a step increase in the SAM index. We assume that the anomalous vertical heat flux Q_{vert} [W/m^2] in a subsurface layer of thickness Z_{sub} can be approximated as

$$Q_{vert} \propto -\delta \gamma C_p \left(\frac{\partial}{\partial y} \left[\frac{\tau'_x}{f} \right] \right) \frac{\Delta_z [\overline{\theta}]}{Z_{sub}}, \quad (5)$$

where $\Delta_z [\overline{\theta}]$ in $^\circ$ C is the inversion (i.e., the maximum vertical contrast) in the time-mean ocean potential temperature within the layer, and γ (unitless) denotes the efficiency with which the subsurface warming signal is communicated into the mixed layer. Parameter δ is a nondimensional factor $0 \leq \delta \leq 1$ that indicates whether we have full ($\delta = 0$), partial ($0 < \delta < 1$), or no ($\delta = 1$) compensation of the anomalous Ekman upwelling by the eddy-induced circulation.

If the slow response is indeed governed by upwelling of warmer water below the mixed layer, the bolus circulation cannot be neglected [*Ferreira et al.*, 2015]. Moreover, the fraction of eddy compensation ($1 - \delta$) is model dependent. The representation of mixed layer entrainment processes also differs across the CMIP5 ensemble. We therefore expect that both δ and γ may contribute to the intermodel spread in the slow SST response, along with the climatological SO temperature inversion $\Delta_z [\overline{\theta}]$.

Using Equation 5 as an Ansatz, we test the importance of the background thermal stratification $\Delta_z [\overline{\theta}]$ for setting differences in the slow response among CMIP5 GCMs. We calculate the average slope Λ [$^\circ$ C/year] of the step response functions between Year 1 and Year 7 after a step increase in the SAM. In many models this slope is predicted to be positive, corresponding to a slow warming. We compare Λ against the vertical temperature inversion $\Delta_z [\overline{\theta}]$ for the area-averaged water column between 55° and 70° S and between depths of 65 m and 550 m. We find that the slow response rates Λ across models are positively correlated with $\Delta_z [\overline{\theta}]$, with $R = +0.45$ (Figure 5b). This result is statistically significant with $p < 0.05$. It emphasizes that Ekman upwelling acting on the background temperature gradients contributes substantially to the intermodel spread in the slow SST responses to SAM.

The correlation between the rate Λ and the vertical temperature inversion $\Delta_z [\overline{\theta}]$ is not as strong as our result linking the rapid cooling response to the meridional SST gradients. We propose that the slow regime is more complicated than the fast one due in part to air-sea heat exchange [*Ferreira et al.*, 2015] but also due to multiple diverse processes within the ocean domain such as eddy compensation and mixed layer entrainment represented by coefficients δ and γ in Equation 5.

We acknowledge that the data points in our intermodel correlation analysis of the fast and slow response (Figures 5a and 5b) do not necessarily represent independent samples. Some CMIP5 ensemble members are in fact multiple versions of the same GCM with a different horizontal resolution (e.g., MPI-ESM-LR and MPI-ESM-MR). Other ensemble members are developed by the same institution (e.g., GFDL-CM3, GFDL-ESM2G, and GFDL-ESM2M) or belong to the same family of models and hence share common code or parameterizations [*Knutti et al.*, 2013]. Thus it is possible that we are inflating our sample size by redundantly including interdependent GCMs. On the other hand, we cannot know a priori which models may exhibit similarities or differences solely on the basis of their common genealogy. For instance, models MIROC-ESM and MIROC5 are related, but their predicted fast SST responses to SAM are statistically different (Figure 5a).

While acknowledging the limitations of this regression analysis, we attempt to extend our CMIP5 results to the real SO and place an observational constraint on the SST response to SAM. We calculate the climatological meridional SST gradients using data from the Reynolds Optimum Interpolation [*Reynolds et al.*, 2002] and compute a metric for time-mean vertical contrast in potential temperature using the Hadley Centre EN4 product [*Good et al.*, 2013]. We use these observationally based climatological SO temperature gradients and the linear relationships found among CMIP5 models (Figure 5) to estimate the fast and slow responses

in the real SO (denoted with stars in Figures 5a and 5b). Our results suggest an expected cooling of -0.13°C , with a standard error of 0.01°C , one year after a step increase in the SAM index. This is likely to be followed by a gradual reduction in the negative SST anomaly at a rate of $0.014^{\circ}\text{C}/\text{year}$, with a standard error of $0.003^{\circ}\text{C}/\text{year}$. The schematic Figure 6 illustrates our estimate for the SST step response function of the real SO.

4 Conclusions and discussion

In this study we have analyzed CMIP5 preindustrial control simulations and examined how SAM forces SO SSTs. In many GCMs the SST exhibits a two-timescale response to SAM: initial cooling followed by slow warming. As in *Ferreira et al.* [2015], we interpret the evolution of these temperature anomalies in terms of the wind-driven circulation redistributing the background heat reservoir. We show evidence that anomalous equatorward transport of colder water is responsible for the fast cooling response south of 50°S . Our results also suggest that the slow warming regime found in many GCMs is sustained by Ekman upwelling of warmer water in the haline stratified SO.

Across the CMIP5 ensemble, we find a notable intermodel spread in the SO SST response to poleward intensification of the westerlies. We relate the diversity of step response functions to differences in the background thermal stratification among the models. GCMs that have small meridional and large vertical temperature gradients in their SO climatology transition faster between the initial cooling and the slow warming regime. Our results imply that in order to better simulate the SST response to SAM, models need a realistic ocean climatology around Antarctica.

The model-specific results of our analysis have implications for attribution studies which evaluate the effects of greenhouse gas forcing and ozone depletion on the SO. For example, *Sigmond and Fyfe* [2014] analyze CMIP3 and CMIP5 output to determine the impact of the ozone hole on SO sea ice. Similarly, *Solomon et al.* [2015] design and conduct numerical experiments with CESM1(WACCM) to study how ozone depletion affects the circulation and sea water properties of the SO. Such in-depth attribution studies often employ a limited set of GCMs – for instance, only a few CMIP5 modeling groups provide output from ozone-only simulations [*Sigmond and Fyfe*, 2014]. However, individual GCMs have various biases in their mean ocean climatology [e.g., *Meijers et al.*, 2014; *Salleé et al.*, 2013]. Thus, we emphasize that the outcome of attribution experiments can be sensitive to the choice of models used. Realistic background temperature gradients are a prerequisite for simulating successfully the response of the SO to a poleward intensification of the westerlies, as the one seen in numerical experiments with ozone depletion.

Our results also identify criteria for constraining and critically assessing future projections of the Southern Hemisphere SST anomalies. Under scenarios with extended greenhouse gas emissions and gradual ozone recovery, CMIP5 models predict a significant and lasting poleward intensification of the westerlies throughout the 21st century [*Wang et al.*, 2014]. Based on our analysis, we suggest that those models which have smaller biases in their climatological stratification provide better estimates of future SST anomalies in the SO.

We point out that in our analysis we have neglected seasonal variations in ocean stratification and their impact on the SO SST response to wind changes. *Purich et al.* [2016] emphasize that in the summer a warm surface lens caps the colder subsurface winter water. Therefore, during this season, anomalous Ekman upwelling may complement rather than counteract the cooling effect of northward Ekman transport.

Our study has further limitations in its ability to account for the multiple diverse processes that take place in the SO. For example, *de Lavergne et al.* [2014] show that there are large differences among the CMIP5 models in their representation of deep convection around Antarctica. It is possible that certain GCMs which do not have strong SO convection, such as BCC-CSM1 and CNRM-CM5, may not be able to efficiently communicate a subsurface temperature signal into the mixed layer. This in turn can delay or preclude entirely the slow warming response to SAM in these models. The recurrence of convective and nonconvective periods in GCMs can also modify the variability of SO stratification about its mean climatology and affect the transition between the fast and slow SST responses [*Seviour et al.*, 2016].

Another potential deficiency in our work pertains to our treatment of atmosphere-ocean coupling. We have not explored any possible intermodel differences in the response of SO surface heat fluxes to SAM. In our linear response function analysis, we have also assumed that the SAM wind pattern forces the SST but not vice versa. However, *Sen Gupta and England* [2007] suggest that SO SST anomalies may feed back on the atmospheric circulation and increase the persistence of SAM. We treat such mechanisms as a source of error contributing to the uncertainty on our estimates of the step response functions.

It is also important to note that the CMIP5 ensemble members used in our analysis do not resolve eddies and rely on parameterizations. Therefore, these GCMs may be missing an important element of the ocean's response to winds. *Böning et al.* [2008] present observational evidence indicating that isopycnal slopes in the SO have not changed over the last few decades despite trends in the SAM. The *Böning et al.* [2008] results are consistent with the eddy-compensation phenomenon and support the possibility that unresolved eddy processes can strongly modulate anomalies

in the wind-driven circulation. Despite this shortcoming of our study, we reiterate that it is important to understand how poleward intensifying westerlies impact the SO in the very same models that are widely used to analyze historical climate change and make authoritative future projections.

Finally, our analysis can be used to make a qualitative estimate for the SST response to SAM in the real SO. Our results suggest that during a sustained positive phase of the SAM, SO SSTs can exhibit a non-monotonic evolution. A strong and rapid transient cooling may be followed by a gradual recovery. However, our results do not suggest a large warming rate during the slow response to SAM.

Our results have implications for surface heat uptake in the real SO and for the persistent expansion of the sea ice cover around Antarctica. The positive SAM trend over the last decades may have allowed a cooler SO to absorb more excess heat from the atmosphere in a warming world. Furthermore, SAM-induced negative SST anomalies may have contributed to the observed increase in SO sea ice extent [Holland et al., 2015; Kostov et al., 2016]. However, if the real SO exhibits a two-timescale response to SAM, the observed SST trends may reverse sign. Hence a sustained poleward intensification of the westerly winds – due to ozone and greenhouse gas forcing – can eventually contribute to a surface warming of the SO, a decreased rate of heat uptake, and a reduction in sea ice concentration. It is therefore important to constrain both the short-term and the long-term SO SST response to SAM.

Appendix A. Verification of the Methodology

We test our methodology from Section 2 in order to ascertain its reliability. Our verification procedure involves applying the regression algorithm to systems with a known prescribed step response function. The latter is convolved with a randomly generated order 1 autoregressive timeseries (AR(1)) that is 1000 years long and resembles a SAM forcing. The result of the convolution is our synthetic SST response, which is strongly diluted with a different AR(1) process characterized by longer memory. We choose parameters for the AR(1) models such that their autocorrelations resemble those of SAM and SO SST timeseries in the CMIP5 GCMs (for instance, Figure 7a and c). We conduct multiple verification tests with different choices of AR(1) parameters. We also vary the signal to noise ratio in our synthetic SST. Figure 7b and d show examples from two different tests.

Within every test we generate an ensemble of multiple synthetic SAM and SST signals with the same statistical properties but different random values. We apply our algorithm separately to each realization in the same fashion as our analysis of CMIP5 control simulations. The verification tests confirm the validity of our method for estimating step response functions.

Table 1 List of CMIP5 Control Simulations

Model Name	Control Run Length [Years]
ACCESS1-0	500
ACCESS1-3	500
BCC-CSM1	500
CanESM2	996
CCSM4	1051
CESM-CAM5	319
CMCC-CM	330
CNRM-CM5	850
GFDL CM3	500
GFDL-ESM2G	500
GFDL-ESM2M	500
GISS-E2-H	540
GISS-E2-R	550
IPSL-CM5A-LR	1000
IPSL-CM5A-MR	300
IPSL-CM5B-LR	300
MIROC5	670
MIROC-ESM	630
MPI-ESM-LR	1000
MPI-ESM-MR	1000
MRI-CGCM3	500
NorESM1-M	501
NorESM1-ME	252

Table 2 Fitting Parameters. We vary the maximum cutoff lag τ_{max} [Years]. Note that we use only $\tau_{max} = 50$ years and $\tau_{max} = 75$ years for models whose control simulation is shorter than 350 years. We use four different values of $\tau_{max} = 50$ where longer simulations are available. We also select shorter SST timeseries from the full control simulations by removing a certain percent of time steps from the beginning and the end of each model run.

Fitting Parameter	Parameter Space
τ_{max} [Years]	50, 75, 100, 150
Offset from the beginning of the full timeseries [% of simulation length]	0, 2.5, 5, 7.5, 10, 15, 20, 25, 30, 35, 40
Offset from the end of the full timeseries [% of simulation length]	0, 2.5, 5, 7.5, 10, 15, 20, 25, 30

Acknowledgements The CMIP5 data for this study is available at the Earth System Grid Federation (ESGF) Portal (<https://pcmdi9.llnl.gov/projects/esgf-llnl/>). Y.K. received support from an NSF MOBY grant, award #1048926. J.M., U.H., D.F., and M.H. were funded by the NSF FESD program, grant award #1338814. K.C.A. was supported by a James McDonnell Foundation Postdoctoral Fellowship and NSF grant OCE-1523641. We would like to thank the World Climate Research Programme and the Working Group on Coupled Modelling, which is in charge of CMIP5. We extend our appreciation to the organizations that support and de-

velop the CMIP infrastructure: the U.S. Department of Energy through its Program for Climate Model Diagnosis and Intercomparison and the Global Organization for Earth System Science Portals. We thank the CMIP5 climate modeling groups for providing their numerical output.

References

- Armour K, J Marshall, A Donohoe, J Scott, E Newsom (2015) Southern Ocean warming delayed by circumpolar upwelling and equatorward transport. (submitted)
- Bitz CM, Polvani LM (2012) Antarctic climate response to stratospheric ozone depletion in a fine resolution ocean climate model. *Geophysical Research Letters* 39:n/an/a. doi: 10.1029/2012GL053393
- Böning CW, A Disper, M Visbeck, SR Rintoul, FU Schwarzkopf (2008) The response of the Antarctic Circumpolar Current to recent climate change. *Nature Geoscience* 1:864-869. doi: 10.1038/ngeo362
- Ciasto LM, Thompson DWJ (2008) Observations of Large Scale Ocean Atmosphere Interaction in the Southern Hemisphere. *J. Climate*, 21, 1244-1259. doi: <http://dx.doi.org/10.1175/2007JCLI1809.1>
- Compo, G.P., J.S. Whitaker, and P.D. Sardeshmukh (2006) Feasibility of a 100 year reanalysis using only surface pressure data. *Bull. Amer. Met. Soc.*, 87, 175-190, doi: <http://dx.doi.org/10.1175/BAMS-87-2-175>
- de Lavergne C, JB Palter, ED Galbraith, R Bernardello, I Marinov (2014) Cessation of deep convection in the open Southern Ocean under anthropogenic climate change. *Nature Climate Change* 4, 278282, doi:10.1038/nclimate2132
- Dee DP, Uppala SM, Simmons AJ, et al (2011) The ERA-Interim reanalysis: configuration and performance of the data assimilation system. *Quarterly Journal of the Royal Meteorological Society* 137:553597. doi: 10.1002/qj.828
- Fan, T, C Deser, and DP Schneider (2014), Recent Antarctic sea ice trends in the context of Southern Ocean surface climate variations since 1950, *Geophys. Res. Lett.*, 41, 24192426, doi:10.1002/2014GL059239.
- Ferreira D, J Marshall, CM Bitz, S Solomon, and A Plumb (2015) Antarctic Ocean and Sea Ice Response to Ozone Depletion: A Two-Time-Scale Problem. *Journal of Climate* 28:12061226. doi: 10.1175/JCLI-D-14-00313.1
- Fyfe JC, Saenko OA, Zickfeld K, et al (2007) The Role of Poleward-Intensifying Winds on Southern Ocean Warming. *Journal of Climate* 20:53915400. doi: 10.1175/2007JCLI1764.1
- Gillett NP, Thompson DWJ (2003) Simulation of recent southern hemisphere climate change. *Science*, 302:2735. doi: 10.1126/science.1087440
- Good SA, Martin MJ, Rayner NA (2013) EN4: Quality controlled ocean temperature and salinity profiles and monthly objective analyses with uncertainty estimates. *Journal of Geophysical Research: Oceans* 118:67046716. doi: 10.1002/2013JC009067
- Hall A, Visbeck M (2002) Synchronous variability in the Southern Hemisphere atmosphere, sea ice, and ocean resulting from the Annular Mode. *Journal of Climate*, 15, 3043-3057. doi: [http://dx.doi.org/10.1175/1520-0442\(2002\)015;3043:SVITSH;2.0.CO;2](http://dx.doi.org/10.1175/1520-0442(2002)015;3043:SVITSH;2.0.CO;2)
- Hasselmann, K, R Sausen, E Maier-Reimer, and R Voss (1993), On the cold start problem in transient simulations with coupled atmosphere-ocean models. *Climate Dynamics*, 9(2), 53-61. doi: 10.1007/BF00210008
- Holland M, et al (2015) Sensitivity of Antarctic sea ice to SAM-associated wind anomalies in coupled climate models (in prep.).
- Hutchinson DK, England MH, Santoso A, Hogg AM (2013) Interhemispheric asymmetry in transient global warming: The role of Drake Passage. *Geophysical Research Letters* 40:15871593. doi: 10.1002/grl.50341
- Hutchinson, DK, MH England, AMcC Hogg and K Snow (2015) Interhemispheric Asymmetry of Warming in an Eddy Permitting Coupled Sector Model, *J. Climate*, 28, 73857406. doi: 10.1175/JCLI-D-15-0014.1
- Knutti, R, D Masson, and A Gettelman (2013), Climate model genealogy: Generation CMIP5 and how we got there, *Geophys. Res. Lett.*, 40, 11941199, doi:10.1002/grl.50256.
- Kostov Y, et al (2016) Southern Ocean cooling in a warming world: reassessing the role of westerly winds. (in prep.)
- Marshall J, KC Armour, JR Scott, Y Kostov, U Hausmann, D Ferreira, TG Shepherd, CM Bitz (2014) The ocean's role in polar climate change: asymmetric Arctic and Antarctic responses to greenhouse gas and ozone forcing. *Phil. Trans. R. Soc. A* 2014 372 20130040; DOI: 10.1098/rsta.2013.0040
- Marshall J, JR Scott, KC Armour, J-M Campin, M Kelley, and A Romanou (2015) The ocean's role in the transient response of climate to abrupt greenhouse gas forcing. *Climate Dynamics*, Vol. 44, Issue 7, pp 2287-2299. doi: 10.1007/s00382-014-2308-0.
- Meijers AJS (2014) The Southern Ocean in the Coupled Model Intercomparison Project phase 5. *Phil.Trans.R. Soc. A* 372: 20130296. doi: <http://dx.doi.org/10.1098/rsta.2013.0296>
- Oke P, England M (2004) Oceanic response to changes in the latitude of the Southern Hemisphere sub-polar westerly winds. *Journal of Climate*, 17, 10401054. doi: [http://dx.doi.org/10.1175/1520-0442\(2004\)017;1040:ORTCIT;2.0.CO;2](http://dx.doi.org/10.1175/1520-0442(2004)017;1040:ORTCIT;2.0.CO;2)
- Purich A, W Caj, MH England, and T Cowan (2016) Evidence for link between modelled trends in Antarctic sea ice and underestimated westerly wind changes. *Nature Communications*, in press.

- Reynolds RW, NA. Rayner, TM Smith, DC Stokes, and W Wang (2002) An Improved In Situ and Satellite SST Analysis for Climate. *J. Climate*, 15, 16091625. doi: [http://dx.doi.org/10.1175/1520-0442\(2002\)015;1609:AIISAS;2.0.CO;2](http://dx.doi.org/10.1175/1520-0442(2002)015;1609:AIISAS;2.0.CO;2)
- Russell JL, KW Dixon, A Gnanadesikan, RJ Stouffer, and JR Toggweiler (2006) The Southern Hemisphere Westerlies in a Warming World: Propping Open the Door to the Deep Ocean. *J. Climate*, 19, 63826390. doi: <http://dx.doi.org/10.1175/JCLI3984.1>
- Salleé, J-B, E Shuckburgh, N Bruneau, AJS Meijers, TJ Bracegirdle, and Z Wang (2013), Assessment of Southern Ocean mixed layer depths in CMIP5 models: Historical bias and forcing response, *J. Geophys. Res. Oceans*, 118, 18451862, doi:10.1002/jgrc.20157
- Sen Gupta A, England M (2006) Coupled OceanAtmosphereIce Response to Variations in the Southern Annular Mode. *J. Climate*, 19, 44574486. doi: <http://dx.doi.org/10.1175/JCLI3843.1>
- Sen Gupta A, England MH (2007) Coupled OceanAtmosphere Feedback in the Southern Annular Mode. *Journal of Climate* 20:36773692. doi: 10.1175/JCLI4200.1
- Seviour, WJM., Waugh, DW and Gnanadesikan, A (2016) Challenges in determining the transient response of the Southern Ocean to ozone depletion (in prep.)
- Sigmond M, Fyfe JC (2014) The Antarctic Sea Ice Response to the Ozone Hole in Climate Models. *Journal of Climate* 27:13361342. doi: 10.1175/JCLI-D-13-00590.1
- Sigmond M, Reader MC, Fyfe JC, Gillett NP (2011) Drivers of past and future Southern Ocean change: Stratospheric ozone versus greenhouse gas impacts. *Geophysical Research Letters* 38:n/an/a. doi: 10.1029/2011GL047120
- Solomon A, LM Polvani, KL Smith, and RP Abernathey (2015), The impact of ozone depleting substances on the circulation, temperature, and salinity of the Southern Ocean: An attribution study with CESM1(WACCM), *Geophys. Res. Lett.*, 42, doi:10.1002/2015GL064744.
- Taylor KE, Stouffer RJ, Meehl GA (2012) An overview of CMIP5 and the experiment design. *Bull Am Meteorol Soc* 93:485498. doi: <http://dx.doi.org/10.1175/BAMS-D-11-00094.1>
- Thomas JL, DW Waugh, A Gnanadesikan, 2015 Southern Hemisphere extratropical circulation: Recent trends and natural variability . *Geophys. Res. Lett.*, (submitted).
- Thompson D, Solomon S (2002) Interpretation of recent Southern Hemisphere climate change. *Science* 296 (5569), 895-899. [DOI:10.1126/science.1069270]
- Thompson DWJ, Solomon S, Kushner PJ, et al (2011) Signatures of the Antarctic ozone hole in Southern Hemisphere surface climate change. *Nature Geoscience* 4:741749. doi: 10.1038/ngeo1296
- Wang G, Cai W, Purich A (2014) Trends in Southern Hemisphere wind-driven circulation in CMIP5 models over the 21st century: Ozone recovery versus greenhouse forcing. *J. Geophys. Res. Oceans*, 119, 29742986, doi:10.1002/2013JC009589.
- Waugh DW, F Primeau, T DeVries, and M Holzer. Recent Changes in the Ventilation of the Southern Oceans. *Science*, 339 (6119), 568-570. [DOI:10.1126/science.1225411]
- Waugh DW (2014) Changes in the ventilation of the southern oceans. *Phil. Trans. R. Soc. A* 2014 372 20130269; DOI: 10.1098/rsta.2013.0269.




Fig. 1 Shown in black is the 1982-2014 timeseries of SST [$^{\circ}\text{C}$] averaged between 55°S and 70°S based on the NOAA Reynolds Optimum Interpolation [Reynolds *et al.*, 2002]. The 1980-2014 timeseries of the annual-mean SAM index based on the ERA Interim reanalysis [Dee *et al.*, 2011] is superimposed in gray. The index is defined as the first principal component of SLP variability south of 20°S and is normalized by its standard deviation. Solid lines indicate linear trends fitted to each timeseries. Note the reversed scale for the SAM timeseries shown on the right.




Fig. 2 Timeseries from the control simulation of model CCSM4: the SAM index in gray and the Southern Ocean (SO) SST anomaly averaged between 55°S to 70°S in black. Each index is detrended and rescaled by its standard deviation. The SST scale is shown on the left vertical axis, and the reversed scale for the SAM index is shown on the right. The SO SST is negatively correlated to the SAM index with $R = -0.37$ at a lag of 1 year.




Fig. 3 Annual-mean response of the Southern Ocean SST anomaly [°C] to: a) a positive impulse perturbation in the SAM index of magnitude equal to $\overline{\sigma}_{SAM}^{Ens}$; b) a positive step increase in the SAM index of magnitude equal to $\overline{\sigma}_{SAM}^{Ens}$. Different colors are used to distinguish the response functions in the three CMIP5 models shown: CCSM4, MPI-ESM-MR, and CNRM-CM5. For each model we show 100 fits that outline an envelope of uncertainty. Vertical error bars denote the margin of error for each fit.




Fig. 4 Annual-mean responses of the Southern Ocean SST [°C] to a step increase in the SAM index of magnitude $\overline{\sigma}_{SAM}^{Ens}$ – comparison across the CMIP5 ensemble. For each model we have shown only the mean estimate $SST_{Step}(t)$.




Fig. 5 a) Relationship between the models' climatological meridional SST gradients $\partial_y[\overline{SST}] [^{\circ}C / 100 km]$ in the Southern Ocean (55° - 70° S) and the Year 1 SST response $SST_{Step}(t=1) [^{\circ}C]$ to a step perturbation in the SAM index. The vertical error bars correspond to $\sigma_{SST_{step}(t=1)}$. b) Relationship between the climatological temperature inversion $\Delta_z[\overline{\theta}] [^{\circ}C]$ in the Southern Ocean (depth levels 65 m to 550 m) and the SST warming rate $\Lambda [^{\circ}C / year]$ which characterizes the slow response to a step increase in the SAM index. Legend: both a) and b) use the same color code and alphabetical order as in Figure 4 to distinguish the CMIP5 models analyzed. Straight lines indicate linear fits to the scatter where each data point is weighed by its uncertainty. The yellow stars denote estimates for the response of the real Southern Ocean based on observed climatological meridional SST gradients between 55° S and 70° S (NOAA Reynolds Optimum Interpolation *Reynolds et al.* [2002]) and the climatological $\Delta_z[\overline{\theta}]$ inversion (Hadley Centre EN4 dataset, *Good et al.* [2013]).




Fig. 6 Thick solid black lines: schematic for the estimated response of the real SO SST [°C] based on results from Figure 5. Thick dashed black lines denote two projections beyond Year 7: a linear extrapolation and a constant equilibrium. Thin gray lines indicate the estimated SO SST step response functions [°C] for the ensemble of 23 CMIP5 models shown with a different color scheme in Figure 4




Fig. 7 Application of the regression algorithm to systems with a known prescribed step response function. On the top row we show a test case where we assume long memory in our SAM and SST signals. The SST signal is diluted such that 60% of the variance is noise. In panel a) on the left, we show the lagged autocorrelations of SAM and SST in CCSM4 (gray dashed curves) and our synthetic artificially generated signals (solid black curves). In panel b) we show applications of the regression algorithm. The thick black curve is the true prescribed step response function. The thin gray curves and the vertical bars denote the estimated step response function $SST_{Step}(t)$ and the uncertainties $\sigma_{SST_{step}}(t)$ produced by applying our regression algorithm. The two gray curves in panel b) result from analyzing separate realizations in which we use the same prescribed step response and AR timeseries with the same statistical properties (illustrated in a)) but different random values. On the bottom row we show a test case where we assume shorter memory in the SAM and SST signals, but the SST signal is diluted with more noise, such that the forced response contributes only 20% of the total variance. Panels c) and d) are analogous to panels a) and b).