

Feedback regulation of cholesterol synthesis: sterol-accelerated ubiquitination and degradation of HMG CoA reductase

Russell A DeBose-Boyd¹

¹Department of Molecular Genetics, University of Texas Southwestern Medical Center, Dallas, TX 75390-9046, USA

3-Hydroxy-3-methylglutaryl coenzyme A (HMG CoA) reductase produces mevalonate, an important intermediate in the synthesis of cholesterol and essential nonsterol isoprenoids. The reductase is subject to an exorbitant amount of feedback control through multiple mechanisms that are mediated by sterol and nonsterol end-products of mevalonate metabolism. Here, I will discuss recent advances that shed light on one mechanism for control of reductase, which involves rapid degradation of the enzyme. Accumulation of certain sterols triggers binding of reductase to endoplasmic reticulum (ER) membrane proteins called Insig-1 and Insig-2. Reductase-Insig binding results in recruitment of a membrane-associated ubiquitin ligase called gp78, which initiates ubiquitination of reductase. This ubiquitination is an obligatory reaction for recognition and degradation of reductase from ER membranes by cytosolic 26S proteasomes. Thus, sterol-accelerated degradation of reductase represents an example of how a general cellular process (ER-associated degradation) is used to control an important metabolic pathway (cholesterol synthesis).

Keywords: ubiquitination, cholesterol synthesis, feedback regulation, HMG CoA reductase, statin, Scap-SREBP, Insig, ER-associated degradation

Cell Research (2008) 18:609-621. doi: 10.1038/cr.2008.61; published online 27 May 2008

Introduction

3-Hydroxy-3-methylglutaryl coenzyme A (HMG CoA) reductase catalyzes conversion of HMG CoA to mevalonate (Figure 1), the precursor of isoprenoid groups that are incorporated into many end-products including cholesterol, ubiquinone, heme, dolichol, and the farnesyl and geranylgeranyl groups that can become attached to many cellular proteins [1]. HMG CoA reductase has been long recognized as the rate-limiting enzyme in synthesis of cholesterol and as such is a primary focus of regulation. This is underscored by a multivalent system mediated by sterol and nonsterol isoprenoids that exerts stringent feedback control on reductase through multiple mechanisms [2]. The complexity of this regulatory system was first revealed in the late 1970s through the use of compactin, a member of the statin family of drugs that are potent competitive inhibitors of reductase [3, 4]. Treatment of cultured cells with compactin blocks production of mevalonate, thereby reducing levels of sterol

and nonsterol isoprenoids that normally govern feedback regulation of reductase. Cells respond to the inhibition of reductase by developing a drastic increase in reductase protein (~200-fold), owing to the combined effects of enhanced transcription of the reductase gene, efficient translation of reductase mRNA, and extended half-life of reductase protein. Complete reversal of this compensatory increase in reductase requires regulatory actions of both sterol and nonsterol end-products of mevalonate metabolism [2, 5]. Sterols inhibit the activity of sterol regulatory element-binding proteins (SREBPs), a family of membrane-anchored transcription factors that enhance cholesterol synthesis and uptake by modulating genes encoding cholesterol biosynthetic enzymes (including reductase) and the low density lipoprotein (LDL)-receptor [6]. An unknown nonsterol mevalonate-derived product controls the translational effects through a poorly understood mechanism that may be mediated by the complex 5'-untranslated region of the reductase mRNA [5]. Both sterol and nonsterol end-products of mevalonate metabolism combine to accelerate degradation of reductase protein through a mechanism mediated by the ubiquitin-proteasome pathway [7-9]. Through these mechanisms, the multivalent regulation of reductase

Correspondence: Russell A DeBose-Boyd

E-mail: Russell.DeBose-Boyd@utsouthwestern.edu




Figure 1 Schematic representation of the cholesterol synthetic pathway in animal cells. Reactions that require molecular oxygen are indicated, and specific inhibitors of various enzymes in the pathway are highlighted in red.

coordinates mevalonate metabolism such that essential nonsterol isoprenoids can be constantly supplied without risking the potentially toxic overproduction of cholesterol or one of its sterol precursors.

In all mammalian species that have been studied to date (i.e., human, hamster, rat, and mouse), reductase localizes to membranes of the endoplasmic reticulum (ER) and consists of 887 or 888 amino acids that can be separated into two contiguous domains (Figure 2A) [10-13]. The N-terminal domain of reductase is comprised of 339 amino acids and is integrated into membranes by virtue of eight membrane-spanning segments that are separated by short loops (Figure 2B) [14]. The 548-amino acid C-terminal domain of reductase projects into the cytosol and exerts all of the enzymatic activity [13]. The amino acid sequence of the membrane domain of reductase is strikingly conserved among mammalian species [15, 16], which suggested early on that the region may be important for more than just membrane anchorage. Indeed, two key observations have disclosed an important role for the membrane domain of reductase in sterol-accelerated degradation. (1) Expression of the truncated, cytosolic C-terminal domain of reductase produced a stable, catalytically active protein whose degradation was not influenced by sterols [17]. (2) A chimeric protein consisting of a fusion between the membrane domain of reductase and soluble β -galactosidase exhibited sterol-accelerated degradation similar to the wild-type full-length reductase [18]. These observations led to the hypothesis that the membrane domain of reductase somehow senses levels of membrane-embedded sterols, which triggers reactions that render the enzyme susceptible to proteolytic degradation [17]. This degradation occurs from ER membranes and can be blocked by inhibitors of the 26S proteasome, which leads to the accumulation of ubiquitinated forms of reductase [9, 19].

Ubiquitin and proteasomes have been implicated in degradation of reductase in the yeast *Saccharomyces cerevisiae* [20]. HMG2p, one of two reductase isozymes in yeast, is rapidly degraded when flux through the mevalonate pathway is high. Degradation of the other isozyme, HMG1p, is not regulated. Although the catalytic domains of yeast HMG2p and mammalian reductase show strong similarity (>50% identity over 540 amino acids), the membrane domains bear limited resemblances (<20% identity over 340 amino acids). Considering that the membrane domains of yeast and mammalian reductase are necessary and sufficient for accelerated degradation [18, 21], limited conservation between these regions provides an explanation for the observation that degradation is not triggered by sterols in yeast, but rather by nonsterol isoprenoids [22]. Despite these differences, regulated ubiquitination and degradation of reductase is employed by yeast and mam-




Figure 2 Domain structure of hamster HMG CoA reductase. **(A)** HMG CoA reductase consists of two distinct domains: a hydrophobic N-terminal domain with eight membrane-spanning segments that anchor the protein to ER membranes, and a hydrophilic C-terminal domain that projects into the cytosol and exhibits all of the enzyme's catalytic activity. **(B)** Amino acid sequence and topology of the membrane domain of hamster HMG CoA reductase. The lysine residues implicated as sites of Insig-dependent, sterol-regulated ubiquitination are highlighted in red and denoted by arrows. The YIYF sequence in the second membrane-spanning helix that mediates Insig binding is highlighted in yellow.

imals to modulate flux through the mevalonate pathway. For further insight into the pathway for degradation of HMG2p in yeast, readers are referred to several excellent reviews [20, 22, 23].

Insigs, polytopic proteins of the ER that mediate sterol-accelerated degradation of HMG CoA reductase

Crucial insights into the mechanism for sterol-accelerated degradation of reductase have emerged from comparisons made between reductase and Scap (the SREBP cleavage-activating protein). Similar to reductase, Scap contains two distinct domains: a hydrophobic N-terminal domain that spans the membrane eight times and a hydrophilic C-terminal domain that projects into the cytosol [24]. The C-terminal domain of Scap mediates a constitutive association with SREBPs; this interaction is required for Scap-dependent translocation of SREBPs from the ER to Golgi in sterol-depleted cells (Figure 3). Upon arrival in the Golgi, SREBPs encounter a pair of proteases (designated site-1 and site-2 proteases) that act successively to release soluble fragments from the membrane into the cytosol [25-29]. These processed forms of SREBPs then migrate from the cytosol into the nucleus and stimulate target gene expression, which results in increased synthesis and uptake of sterols [6]. The subsequent accumulation of sterols in ER membranes prevents proteolytic activation of SREBPs by blocking exit of Scap-SREBP complexes from the ER; transcription of SREBP target genes declines and cholesterol synthesis and uptake are suppressed. Inhibition of ER to Golgi transport of SREBPs results from sterol-induced binding of Scap to ER retention proteins called Insig-1 and Insig-2 [30, 31]. Insig binding occludes a cytosolic binding site in Scap recognized by COPII proteins, which incorporate cargo molecules into vesicles that deliver ER-derived proteins to the Golgi [32]. Scap-Insig binding is mediated by a segment of Scap's membrane domain that includes transmembrane helices 2-6 [25, 30]. A similar stretch of transmembrane helices is found in at least four other polytopic membrane proteins (including the Niemann Pick C1 protein, Patched, Dispatched, and reductase), all of which have been postulated to interact with sterols. Thus, the region has become known as the sterol-sensing domain [33]. The importance of the sterol-sensing domain in regulation of Scap is illustrated by findings that point mutations within the region disrupt Insig binding, thereby relieving mutant Scap-SREBP complexes from sterol-mediated ER retention [30, 31, 34-36].

The recognition of sequence resemblances between the sterol-sensing domains of Scap and reductase stimulated an appraisal of a role for Insigs in degradation of reduc-




Figure 3 Model for sterol-regulated Scap-SREBP pathway. SCAP is a sensor of sterols and an escort of SREBPs. In sterol-depleted cells, Scap facilitates export of SREBPs from the ER to the Golgi apparatus, where two proteases, Site-1 protease (S1P) and Site-2 protease (S2P), act to release the transcriptionally active, N-terminal bHLH-Zip domain of SREBPs from the membrane. The released bHLH-Zip domain migrates into the nucleus and binds to a sterol response element (SRE) in the enhancer/promoter region of target genes, activating their transcription. Accumulation of sterols in ER membranes triggers binding of Scap to one of two retention proteins called Insigs, which blocks incorporation of Scap-SREBP complexes into ER transport vesicles. As a result, SREBPs no longer translocate to the Golgi apparatus, the bHLH-Zip domain cannot be released from the membrane, and transcription of all target genes declines.

tase. This effort led to the following observations, which considered together divulge the action of at least one of the Insig proteins in sterol-accelerated degradation of reductase. First, when overexpressed by transfection in Chinese hamster ovary (CHO) cells, reductase cannot be degraded when the cells are treated with sterols [37]. Co-expression of Insig-1 restores sterol-accelerated degradation of reductase, suggesting the saturation of endogenous Insigs by the overexpressed reductase. Second, reduction of both Insig-1 and Insig-2 by RNA interference (RNAi) abolishes sterol-accelerated degradation of endogenous reductase [38]. Third, mutant CHO cells lacking both Insigs are impervious to sterol-stimulated degradation of reductase as well as sterol-mediated inhibition of SREBP processing [39].

Degradation of reductase coincides with sterol-induced binding of its membrane domain to Insigs [37], an action that requires a tetrapeptide sequence, YIYF, located in the second transmembrane segment of reductase (see Figure 2B) [38]. A mutant form of reductase in which the YIYF sequence is mutated to alanine residues no longer binds to Insigs and the enzyme is not subject to rapid degradation. The YIYF sequence is also present in the second

transmembrane domain of Scap, where it mediates sterol-dependent formation of Scap-Insig complexes [30, 31]. In fact, overexpressing the sterol-sensing domain of Scap in cells blocks Insig-mediated, sterol-accelerated degradation of reductase. Mutation of the YIYF sequence in the Scap sterol-sensing domain ablates this inhibition. This indicates that Scap and reductase bind to the same site on Insigs and the two proteins compete for limiting the amount of Insigs when intracellular sterol levels rise.

Insig-mediated ubiquitination of HMG CoA reductase

Evidence supporting a major role for the ubiquitin-proteasome pathway in sterol-accelerated degradation of reductase was first provided by the observation that proteasome inhibition blocks the process [19], leading to the accumulation of ubiquitinated forms of reductase on ER membranes [9]. This ubiquitination is obligatory for degradation of reductase and exhibits an absolute requirement for the presence of Insigs [38, 39]. Reduction of Insig-1 and Insig-2 mRNA by genetic mutation or RNAi-mediated knockdown abrogates sterol-dependent ubiquitination of endogenous reductase, rendering the enzyme refractory to accelerated degradation. Moreover, sterol-induced ubiquitination of reductase exhibits an absolute Insig requirement in transient transfection assays. Mutation of the YIYF sequence in reductase, which blocks Insig binding, prevents regulated ubiquitination and slows the enzyme's degradation. In contrast, conservative substitutions of arginine for lysines 89 and 248 in the membrane domain of reductase (Figure 2B) do not block Insig binding, but the substitutions rather abolish ubiquitination and subsequent degradation of reductase. Thus, lysines 89 and 248 in reductase are implicated as sites for Insig-mediated, sterol-induced ubiquitination. It is important to note that mutation of lysines 89 and 248 blocks ubiquitination and degradation of reductase in the context of the full-length enzyme, suggesting that the catalytic domain does not contribute to ubiquitination. This is consistent with the observation that the soluble catalytic domain is dispensable for sterol-regulated degradation [17].

How might Insig binding impart recognition of reductase by the ubiquitinating machinery? This question was addressed by examining reductase ubiquitination in a permeabilized cell system [40]. Sterol-depleted cells were permeabilized with low concentrations of the mild detergent digitonin such that nearly all of the cytosolic proteins were released into the supernatant upon centrifugation, whereas membrane proteins such as reductase remained associated with the pellet fraction. The pellet of permeabilized cells supports Insig-dependent ubiquitination of reductase that

is stimulated by additions of ATP, sterols, and rat liver cytosol *in vitro*. Surprisingly, reductase ubiquitination is potently stimulated by oxygenated derivatives of cholesterol, including 24-, 25-, and 27-hydroxycholesterol, but not by cholesterol itself. The significance of this finding will be discussed in more detail below.

Ubiquitination of proteins is a multistep process, involving the action of at least three types of enzymes [41]. In the first step, ubiquitin is activated by the ubiquitin-activating enzyme (E1), which forms a thiol ester between a reactive cysteine residue in E1 and the C-terminus of ubiquitin. Next, ubiquitin is transferred from E1 to a catalytic cysteine of the ubiquitin-conjugating enzyme (E2). The third type of enzyme, ubiquitin ligase (E3), facilitates transfer of activated ubiquitin from E2 to a lysine residue in the substrate (or a previously attached ubiquitin). Once a poly-ubiquitin chain of sufficient size is built, the substrate is recognized and subsequently degraded by proteasomes. Only two E1 enzymes exist, and both are cytosolic proteins. In contrast, a variety of E2s and E3s, both soluble and membrane-bound, have been described [42]. The exquisite sensitivity of substrate ubiquitination is ultimately determined by the E3, either alone or in combination with its cognate E2.

Fractionated S100 from HeLa cells was utilized to determine which component of the reductase ubiquitinating machinery (E1, E2 and E3) is provided by rat liver cytosol in the permeabilized cell system [40]. These fractions were first described by Hershko and co-workers [43, 44] and were generated by separating HeLa cell S100 into fractions that bind (Fraction II) or do not bind (Fraction I) an anion exchange resin. It was subsequently determined that Fraction I contained ubiquitin, whereas Fraction II contains E1 [45]. Fraction II effectively replaces rat liver cytosol for regulated ubiquitination of reductase in permeabilized cells, but Fraction I does not. Two observations indicate that Fraction II provides a source of ubiquitin activation in the permeabilized cell system. First, purified E1 replaces rat liver cytosol for sterol-regulated ubiquitination of reductase in permeabilized cells. Second, immunodepletion of E1 eliminates the reductase ubiquitinating activity of rat liver cytosol. These results demonstrate that E1 is the only cytosolic protein required for reductase ubiquitination, which indicates the reductase E2 and E3 are membrane-associated proteins. This notion is consistent with the localization of apparent sites of reductase ubiquitination, lysines 89 and 248, which are cytosolically exposed and are predicted to lie immediately adjacent to transmembrane helices three and seven (Figure 2B).

Results from the analysis of reductase ubiquitination in permeabilized cells indicated that Insig binding results in recruitment of enzymes that ubiquitinate reductase. Co-immunoprecipitation experiments, coupled with tandem

mass spectroscopy, were utilized to identify membrane proteins that associate with the sterol-dependent reductase-Insig complex. These studies revealed that Insig-1 binds to a known membrane-anchored ubiquitin ligase called gp78 [46]. The cDNA for gp78 predicts a 643-amino acid protein that can be divided into four domains. The N-terminal domain of 298 amino acids contains five to seven membrane-spanning helices that anchor the protein to ER membranes and mediate association with Insig-1. The membrane attachment region of gp78 is followed by a 43-amino acid region with a RING finger consensus sequence that confers ubiquitin ligase activity [47]. Following the RING domain is a 42-amino acid region homologous to Cue1p, an ER membrane protein in yeast that serves as a membrane anchor for Ubc7p, a cytosolic ubiquitin-conjugating enzyme [48]. Recently, this region of gp78 has been shown to directly bind to Ufd1, a cytosolic protein that modulates gp78 ubiquitin ligase activity, thereby enhancing ubiquitination and degradation of the enzyme's substrates [49]. Finally, the extreme C-terminus of gp78 (48 amino acids) mediates an interaction with VCP (Valosin-containing protein, also known as p97), an ATPase that has been implicated in the post-ubiquitination steps of ER-associated degradation (ERAD) [50].

At least three lines of evidence indicate that gp78, through its binding to Insig-1, initiates sterol-accelerated degradation of reductase. (1) Overexpression of the membrane domain of gp78 blocks Insig-mediated, sterol-accelerated degradation of reductase, suggesting that the membrane domain of gp78 competes with full-length gp78 for binding to Insig-1, thereby abolishing reductase ubiquitination. (2) Sterols trigger binding of gp78 to reductase in an Insig-dependent, sterol-regulated manner. The specificity of this binding is illustrated by the inability of gp78 to bind Scap, regardless of the presence or absence of sterols and/or Insigs. (3) RNAi-mediated knockdown of gp78 prevents sterol-regulated ubiquitination and degradation of endogenous reductase. Importantly, the effect of gp78 knockdown is specific inasmuch as knockdown of a related membrane-bound ubiquitin ligase, Hrd1, does not affect reductase ubiquitination. Another function of gp78, besides its role as a ubiquitin ligase, is to couple ubiquitination of reductase to degradation through its association with VCP. Indeed, coimmunoprecipitation experiments show that gp78 is an intermediary in association of VCP and Insig-1. Moreover, knockdown of VCP by RNAi prevents sterol-accelerated degradation of endogenous reductase and a dominant-negative ATPase-deficient mutant of VCP blocks sterol-regulated degradation of transfected reductase.

The identification of gp78 as an E3 ubiquitin ligase that mediates reductase ubiquitination has important implications for yet another mode of sterol regulation. The

regulation of Insig-1 contrasts that of reductase in that Insig-1 becomes ubiquitinated and is rapidly degraded by proteasomes in *sterol-depleted* cells [51]. Ubiquitination of Insig-1 is mediated by gp78 [52]. When sterols induce reductase to bind Insig-1, ubiquitination is diverted toward reductase and the enzyme becomes rapidly degraded. However, when sterols cause Scap to bind Insig-1, gp78 is displaced and no longer ubiquitinates Insig-1, thereby stabilizing the protein. This reaction helps to explain why reductase is degraded when it binds to Insig-1, whereas Scap binding to Insig-1 leads to retention in the ER. In addition, gp78-mediated ubiquitination and degradation of Insig-1 provide a mechanism for a recently appreciated process termed "convergent feedback inhibition" [51]. In sterol-depleted cells, Scap-SREBP complexes no longer bind Insig-1, which in turn becomes ubiquitinated and degraded. Thus, Scap-SREBP complexes are free to exit the ER and translocate to the Golgi, where the SREBPs are processed to the nuclear form that stimulates transcription of target genes, including the Insig-1 gene. Increased transcription of the Insig-1 gene leads to increased synthesis of Insig-1 protein, but the protein is ubiquitinated and degraded until sterols build up to levels sufficient to trigger Scap binding. Thus, inhibition of SREBP processing requires convergence of newly synthesized Insig-1 and newly acquired sterols.

The HMG CoA reductase sterol-sensing reaction

Oxysterols are derivatives of cholesterol that contain hydroxyl groups at various positions in the iso-octyl side chain [53, 54]. These compounds are synthesized in many tissues by specific enzymes called hydroxylases; oxysterols play key roles in cholesterol export and they are also intermediates in the synthesis of bile acids [55]. Oxysterols are significantly more soluble than cholesterol in aqueous solution, and thus can readily pass across the plasma membrane and enter cells. This property renders oxysterols such as 24-, 25-, and 27-hydroxycholesterol extremely potent in inhibiting cholesterol synthesis by stimulating binding of both reductase and Scap to Insigs. Oxysterols are present at very low concentrations (10^4 - to 10^6 -fold less than cholesterol) in tissues and blood, which raises questions as to whether they act through a similar mechanism as LDL-derived cholesterol to block cholesterol synthesis. In the case of Scap, the mode of action of these two classes of sterols is becoming clear. Cholesterol directly binds to the membrane domain of Scap in a specific and saturable fashion [56]. The interaction causes a conformational change in Scap that promotes Insig binding [57]. The addition of cholesterol *in vitro* to membranes isolated from sterol-depleted cells causes exposure of a cryptic trypsin

cleavage site, thereby altering the tryptic digestion pattern of Scap that can be monitored by immunoblot analysis [58]. Co-expression of Insig lowers the amount of cholesterol required to induce the conformational change in Scap. Oxysterols neither alter Scap's conformation *in vitro* nor bind to the protein's membrane domain, leading to the postulation of the existence of a membrane-bound oxysterol binding protein. Remarkably, Insig-2 has been recently defined as a membrane-bound oxysterol binding protein with binding specificity that correlates with the ability of oxysterols to inhibit SREBP processing [32, 59]. Thus, formation of the Scap-Insig complex can be initiated by either binding of cholesterol to the membrane domain of Scap or by binding of oxysterols to Insig. Both events prevent incorporation of Scap-SREBP into vesicles that bud from the ER en route to the Golgi. By analogy, the likely mechanism by which oxysterols stimulate degradation of reductase is through their binding to Insig.

In striking contrast to results obtained with Scap, the analysis of reductase ubiquitination in permeabilized cells revealed that the reaction was potently stimulated by oxysterols, but not by cholesterol [40]. These results led to a search for endogenous sterol regulators of reductase ubiquitination and degradation. Previous indirect studies implicated that lanosterol, the first sterol produced in the cholesterol biosynthetic pathway (Figure 1), or one of its metabolites participates in feedback inhibition of reductase. For example, genetic mutation or pharmacologic inhibition of lanosterol 14 α -demethylase, which catalyzes the first step in conversion of lanosterol to cholesterol (Figure 1), markedly reduces the amount of reductase activity in cells [60, 61]. These observations led to the evaluation of lanosterol and its metabolite 24,25-dihydrolanosterol as endogenous regulators of reductase ubiquitination and degradation [62]. When added to intact cells, lanosterol and 24,25-dihydrolanosterol potently stimulate ubiquitination and degradation of reductase through a reaction that requires the presence of Insig. The activity of both sterols is specific inasmuch as they do not inhibit processing of SREBPs. This is consistent with the inability of lanosterol to directly bind to Scap and Insig or alter Scap's conformation *in vitro* [58]. The action of lanosterol and 24,25-dihydrolanosterol is direct and does not require their conversion into an active metabolite as indicated by the reconstitution of reductase ubiquitination by simply incubating isolated membranes with the sterols and purified E1. Using this *in vitro* assay, the action of lanosterol and 24,25-dihydrolanosterol in stimulating ubiquitination and degradation of reductase was traced to methyl groups present in the 4 α , 4 β , and 14 α positions of the sterol ring.

Insig-mediated regulation of reductase is controlled by three classes of sterols: oxysterols, cholesterol, and methyl-

ated sterols such as lanosterol and 24,25-dihydrolanosterol. Oxysterols, which are derived from cholesterol, have dual actions in that they accelerate degradation of reductase and block ER to Golgi transport of Scap-SREBP through their direct binding to Insig. Cholesterol does not regulate reductase stability directly, but binds to Scap and triggers Insig binding, thereby preventing escape of Scap-SREBP from the ER. On the other hand, lanosterol selectively accelerates degradation of reductase without an effect on ER to Golgi transport of Scap-SREBP. Notably, the demethylation of lanosterol has been implicated as a rate-limiting step in the post-squalene portion of cholesterol synthesis, suggesting the reaction as a potential focal point in sterol regulation [63, 64]. Considering that lanosterol is the first sterol produced in cholesterol synthesis, it seems reasonable that it controls early steps in the pathway (i.e., the nonsterol branch) by stimulating reductase degradation. The accumulation of lanosterol is avoided, owing to its inability to block SREBP processing through Scap. This assures that mRNAs encoding enzymes catalyzing reactions subsequent to lanosterol remain elevated and lanosterol is metabolized to cholesterol. The importance of this conversion is highlighted by the observation that lanosterol cannot support cell growth in the absence of cholesterol and may be toxic [65]. This toxicity is likely due to the inability to optimize certain physiologic properties of cell membranes with regard to biological functions.

Oxygen sensing in the cholesterol biosynthetic pathway

The physiologic relevance of lanosterol as an endogenous regulator of reductase ubiquitination and degradation was deduced by the recognition that cholesterol synthesis is a highly oxygen-consumptive process. The synthesis of one molecule of cholesterol from acetyl-CoA requires eleven molecules of dioxygen, nine of which are consumed during the removal of the 4 α , 4 β , and 14 α methyl groups in lanosterol and its metabolite 24,25-dihydrolanosterol by the successive actions of lanosterol 14 α -demethylase and C4-methyl sterol oxidase (Figure 1). This led to speculation that oxygen deprivation (hypoxia) might block demethylation of lanosterol and 24,25-dihydrolanosterol and thereby stimulate degradation of reductase. Indeed, a recent study shows that hypoxia blunts cholesterol synthesis by inhibiting lanosterol and 24,25-dihydrolanosterol demethylation, causing both sterols to accumulate in cells [66]. Rapid degradation of reductase parallels hypoxia-induced accumulation of lanosterol and 24,25-dihydrolanosterol. This Insig-mediated degradation requires *de novo* sterol synthesis as indicated by its inhibition by compactin and the squalene monooxygenase inhibitor NB-598, but not the

lanosterol 14 α -demethylase inhibitor RS-21607 (see Figure 1). Although hypoxia accelerates degradation of reductase, processing of SREBPs remains unaffected. This finding is consistent with the observation described above that exogenous lanosterol stimulates degradation of reductase without inhibiting SREBP processing.

In addition to the accumulation of methylated sterols, the degradation of reductase in hypoxic cells also requires the action of the oxygen-sensitive transcription factor, HIF-1 α . In oxygenated cells, HIF-1 α is rapidly degraded owing to hydroxylation of specific proline residues in the protein [67]. Prolyl hydroxylation enhances binding of HIF-1 α to the von Hippel Lindau tumor suppressor protein (pVHL), which is the recognition component of a ubiquitin ligase that targets HIF-1 α for proteasomal degradation. Prolyl hydroxylation of HIF-1 α is catalyzed by a family of dioxygenases that use 2-oxoglutarate as a co-substrate and exhibit strict dependence for molecular oxygen [68-70]. When cells are deprived of oxygen, prolyl hydroxylation is inhibited, allowing HIF-1 α to escape degradation and accumulate to high levels. The stabilized HIF-1 α subunits associate with the constitutive HIF-1 β subunit, forming a heterodimeric transcription factor (HIF) that modulates

expression of more than 70 genes involved in both systemic and cellular responses to oxygen deprivation [71].

Evidence implicating a major role for HIF-1 α in the hypoxia-induced degradation of reductase is provided by both pharmacologic and genetic data. Treatment of oxygenated cells with dimethylxalylglycine (DMOG), a non-specific inhibitor of 2-oxoglutarate-dependent dioxygenases, not only stabilizes HIF-1 α [72] but also triggers rapid degradation of reductase through an Insig-dependent mechanism that requires de novo sterol synthesis. Genetic evidence for a role of HIF-1 α in reductase degradation is provided by the observation that the enzyme is refractory to hypoxia- and DMOG-induced degradation in mutant cells that are deficient in HIF-1 α . While these observations establish the importance of the action of HIF-1 α in oxygen-regulated degradation of reductase, they raise questions as to the HIF-target genes that mediate the response. In several DNA microarray analyses, Insig-1 and Insig-2 transcripts have been identified among those increased by either DMOG or hypoxia treatment [73-75]. This observation led to the subsequent discovery that DMOG and hypoxia enhance expression of both Insigs through a HIF-dependent mechanism. Considered together, these observations establish a




Figure 4 Mechanism for oxygen sensing in the cholesterol synthetic pathway. The link between synthesis of cholesterol and oxygen sensing in animal cells is provided by hypoxia-induced accumulation of lanosterol and 24,25-dihydrolanosterol and HIF-1 α -mediated induction of Insig-1 and Insig-2. Convergence of these responses leads to rapid degradation of HMG CoA reductase, thereby limiting synthesis of cholesterol.

connection between cholesterol synthesis and oxygen sensing in animal cells (Figure 4). These metabolic pathways are linked by two regulatory actions: (1) hypoxia-induced accumulation of the cholesterol biosynthetic intermediates lanosterol and 24,25-dihydrolanosterol; and (2) HIF-1 α mediated induction of Insigs. Convergence of these signals triggers rapid degradation of reductase, which ultimately limits synthesis of cholesterol and helps to guard against the wasting of cellular oxygen in the face of hypoxia.

Unanswered questions and future directions

Despite the recent advances in the understanding of molecular mechanisms underlying sterol-accelerated degradation of reductase, much remains to be determined. For instance, what is the mechanism by which lanosterol and 24,25-dihydrolanosterol trigger binding of reductase to Insigs? Do these methylated sterols directly bind the membrane domain of reductase in a reaction analogous to that of cholesterol and Scap? Unfortunately, attempts to demonstrate direct binding of methylated sterols to the membrane domain of reductase have been unsuccessful. Moreover, addition of lanosterol or 24,25-dihydrolanosterol to reductase-containing membranes *in vitro* fails to alter the tryptic pattern of the enzyme. Thus, the possibility exists that a distinct ER membrane protein binds to methylated sterols and, in turn, triggers binding of reductase to Insigs, thereby initiating reductase ubiquitination. Reductase is the target of statins, which are the most widely prescribed cholesterol-lowering drugs in humans. Interest in developing additional strategies that inhibit reductase has led to the discovery of nonsterol compounds, such as vitamin E (tocotrienols) and the bisphosphonate SR-12813, that mimic sterols in accelerating reductase degradation [76, 77]. The availability of such reagents may prove useful in the ongoing quest to define the molecular mechanisms for the reductase sterol-sensing reaction.

Another unresolved question in reductase degradation is the mechanism for delivery of ubiquitinated forms of the enzyme from the membrane to the cytosol for proteasomal degradation. Unlike model ERAD substrates that are either completely luminal or contain one transmembrane domain, proteasome inhibition leads to accumulation of ubiquitinated reductase on membranes, rather than in the cytosol [37]. This suggests that degradation of reductase is coupled to its ubiquitination and proceeds through a membrane-bound intermediate. However, reductase must be degraded as a unit without releasing the catalytic domain into the cytosol, which would defeat the purpose of regulated degradation.

Efficient degradation of reductase requires nonsterol isoprenoids derived from mevalonate in addition to sterols.

This was borne out of experiments showing that in compactin-treated cells sterols can trigger binding of reductase to Insigs and subsequent ubiquitination of the enzyme. However, the ubiquitinated reductase is not efficiently degraded unless the cells are also treated with mevalonate. This mevalonate requirement can be bypassed by the addition of geranylgeraniol (GG-OH), a 20-carbon isoprenoid, but not by the 15-carbon farnesol [38]. GGOH does not appear to trigger reductase ubiquitination, even though it augments sterol-accelerated degradation of the enzyme. This suggests the action of nonsterol isoprenoids in a post-ubiquitination step of reductase degradation.

The current view of sterol-accelerated degradation of reductase is illustrated in the model shown in Figure 5. The reaction is initiated by sensing of membrane-embedded sterols through direct or indirect interactions with the membrane domain of reductase. This interaction causes reductase to bind to a subset of Insigs that are associated with gp78, which mediates transfer of ubiquitin from the E2 Ubc7 to lysines 89 and 248 of reductase. Ubiquitination targets reductase for recognition by gp78-associated VCP, which, together with its cofactors, somehow extract ubiquitinated reductase from membranes and deliver it to proteasomes for degradation. The extraction step appears to be augmented by GG-OH. It seems likely that GG-OH, after its conversion to metabolically active geranylgeranylpyrophosphate (GG-PP), is incorporated into a protein that enhances the effect of sterols on reductase degradation. Possible candidates include geranylgeranylated Rab proteins, which are known to play key roles in various aspects of vesicular transport [78]. Thus, the possibility exists that a vesicle-mediated transport event delivers ubiquitinated reductase to a specific organelle or subdomain of the ER in which the protein is degraded. Notably, Ufd1 appears to play a key role in this pathway by enhancing gp78 ubiquitin ligase activity and modulating a post-ubiquitination step in reductase degradation. In addition, Ufd1 seems to bind to gp78 in a sterol-regulated fashion [49], but the significance of this is presently unknown. Complete elucidation of reductase degradation will likely require the reconstitution of post-ubiquitination steps of reductase degradation in a cell-free system.

What is the contribution of reductase degradation to overall cholesterol homeostasis in whole animals? Insigs appear to play a major role in regulation of reductase in the mouse liver. Genetic deletion of Insigs results in the accumulation of reductase to a level approximately 20-fold higher than that in wild type mice [79]. This accumulation is presumably attributable to the combination of both transcriptional and post-transcriptional regulation of reductase, but the extent to which each level of regulation contributed to the massive increase in reductase is unknown. Thus,




Figure 5 Pathway for sterol-accelerated degradation of HMG CoA reductase. Accumulation of 25-hydroxycholesterol, lanosterol, or 24,25-dihydrolanosterol in ER membranes triggers binding of the reductase to Insigs. A subset of Insigs is associated with the membrane-anchored ubiquitin ligase, gp78, which binds the E2 Ubc7 and VCP, an ATPase that plays a role in extraction of ubiquitinated proteins from ER membranes. Through the action of gp78 and Ubc7, reductase becomes ubiquitinated, which triggers its extraction from the membrane by VCP, and subsequent delivery to proteasomes for degradation. The post-ubiquitination step is postulated to be enhanced by geranylgeraniol through an undefined mechanism that may involve a geranylgeranylated protein, such as one of the Rab proteins.

studies that directly focus on reductase degradation are required in order to determine the contribution of protein stability to overall regulation of reductase in mice *in vivo* under various physiologic conditions, such as hypoxia.

The significance of Insig-mediated regulation of reductase in maintenance of cholesterol homeostasis is highlighted by the effectiveness of reductase inhibition in lowering plasma LDL-cholesterol in humans [80]. However, the inhibition of reductase disrupts normal feedback inhibition of the enzyme, and animals respond by developing a compensatory increase in reductase levels in the liver [81, 82]. Remarkably, a similar response has been observed in livers of statin-treated humans as well [83]. Knowledge of the mechanisms for this compensatory increase, particularly the contribution of degradation, may facilitate development of novel drugs that improve the effectiveness of statins, or in some cases provide alternative treatments. Such a drug would be modeled after lanosterol and 24,25-dihydrolanosterol, which selectively stimulate reductase degradation without affecting the Scap-SREBP pathway or LDL-receptor activity. In addition, elucidation of the

underlying mechanisms for sterol-accelerated, ERAD of reductase may have implications for degradation of other clinically important proteins such as the cystic fibrosis transmembrane conductance regulator (CFTR). Thus, further excitement will undoubtedly ensue once questions posed in this review begin to become clear.

Acknowledgments

Work in the DeBose-Boyd laboratory is supported by grants from the National Institutes of Health (HL20948), the Perot Family Foundation, the American Heart Association (0540128N), and the W.M. Keck Foundation.

References

- 1 Goldstein JL, Brown MS. Regulation of the mevalonate pathway. *Nature* 1990; **343**:425-430.
- 2 Brown MS, Goldstein JL. Multivalent feedback regulation of HMG CoA reductase, a control mechanism coordinating isoprenoid synthesis and cell growth. *J Lipid Res* 1980; **21**:505-517.
- 3 Endo A, Kuroda M, Tanzawa K. Competitive inhibition of 3-

- hydroxy-3-methylglutaryl coenzyme A reductase by ML-236A and ML-236B fungal metabolites, having hypocholesterolemic activity. *FEBS Lett* 1976; **72**:323-326.
- 4 Brown MS, Faust JR, Goldstein JL. Induction of 3-hydroxy-3-methylglutaryl coenzyme A reductase activity in human fibroblasts incubated with compactin (ML-236B), a competitive inhibitor of the reductase. *J Biol Chem* 1978; **253**:1121-1128.
 - 5 Nakanishi M, Goldstein JL, Brown MS. Multivalent control of 3-hydroxy-3-methylglutaryl coenzyme A reductase. Mevalonate-derived product inhibits translation of mRNA and accelerates degradation of enzyme. *J Biol Chem* 1988; **263**:8929-8937.
 - 6 Horton JD, Goldstein JL, Brown MS. SREBPs: activators of the complete program of cholesterol and fatty acid synthesis in the liver. *J Clin Invest* 2002; **109**:1125-1131.
 - 7 Roitelman J, Simoni RD. Distinct sterol and nonsterol signals for the regulated degradation of 3-hydroxy-3-methylglutaryl-CoA reductase. *J Biol Chem* 1992; **267**:25264-25273.
 - 8 McGee TP, Cheng HH, Kumagai H, Omura S, Simoni RD. Degradation of 3-hydroxy-3-methylglutaryl-CoA reductase in endoplasmic reticulum membranes is accelerated as a result of increased susceptibility to proteolysis. *J Biol Chem* 1996; **271**:25630-25638.
 - 9 Ravid T, Doolman R, Avner R, Harats D, Roitelman J. The ubiquitin-proteasome pathway mediates the regulated degradation of mammalian 3-hydroxy-3-methylglutaryl-coenzyme A reductase. *J Biol Chem* 2000; **275**:35840-35847.
 - 10 Ness GC, Spindler CD, Moffler MH. Purification of 3-hydroxy-3-methylglutaryl coenzyme A reductase from rat liver. *Arch Biochem Biophys* 1979; **197**:493-499.
 - 11 Edwards PA, Lemongello D, Fogelman AM. Purification and properties of rat liver 3-hydroxy-3-methylglutaryl coenzyme A reductase. *Biochim Biophys Acta* 1979; **574**:123-135.
 - 12 Brown MS, Dana SE, Dietschy JM, Siperstein MD. 3-Hydroxy-3-methylglutaryl coenzyme A reductase. Solubilization and purification of a cold-sensitive microsomal enzyme. *J Biol Chem* 1973; **248**:4731-4738.
 - 13 Liscum L, Finer-Moore J, Stroud RM, *et al*. Domain structure of 3-hydroxy-3-methylglutaryl coenzyme A reductase, a glycoprotein of the endoplasmic reticulum. *J Biol Chem* 1985; **260**:522-530.
 - 14 Roitelman J, Olender EH, Bar-Nun S, Dunn Jr WA, Simoni RD. Immunological evidence for eight spans in the membrane domain of 3-hydroxy-3-methylglutaryl coenzyme A reductase: implications for enzyme degradation in the endoplasmic reticulum. *J Cell Biol* 1992; **117**:959-973.
 - 15 Luskey KL, Stevens B. Human 3-hydroxy-3-methylglutaryl coenzyme A reductase. Conserved domains responsible for catalytic activity and sterol-regulated degradation. *J Biol Chem* 1985; **260**:10271-10277.
 - 16 Gertler FB, Chiu CY, Richter-Mann L, Chin DJ. Developmental and metabolic regulation of the *Drosophila melanogaster* 3-hydroxy-3-methylglutaryl coenzyme A reductase. *Mol Cell Biol* 1988; **8**:2713-2721.
 - 17 Gil G, Faust JR, Chin DJ, Goldstein JL, Brown MS. Membrane-bound domain of HMG CoA reductase is required for sterol-enhanced degradation of the enzyme. *Cell* 1985; **41**:249-258.
 - 18 Skalnik DG, Narita H, Kent C, Simoni RD. The membrane domain of 3-hydroxy-3-methylglutaryl-coenzyme A reductase confers endoplasmic reticulum localization and sterol-regulated degradation onto beta-galactosidase. *J Biol Chem* 1988; **263**:6836-6841.
 - 19 Inoue S, Bar-Nun S, Roitelman J, Simoni RD. Inhibition of degradation of 3-hydroxy-3-methylglutaryl-coenzyme A reductase *in vivo* by cysteine protease inhibitors. *J Biol Chem* 1991; **266**:13311-13317.
 - 20 Hampton RY. Genetic analysis of hydroxymethylglutaryl-coenzyme A reductase regulated degradation. *Curr Opin Lipidol* 1998; **9**:93-97.
 - 21 Hampton RY, Koning A, Wright R, Rine J. *In vivo* examination of membrane protein localization and degradation with green fluorescent protein. *Proc Natl Acad Sci USA* 1996; **93**:828-833.
 - 22 Hampton RY. Proteolysis and sterol regulation. *Annu Rev Cell Dev Biol* 2002; **18**:345-378.
 - 23 Hampton RY. ER-associated degradation in protein quality control and cellular regulation. *Curr Opin Cell Biol* 2002; **14**:476-482.
 - 24 Nohturfft A, Brown MS, Goldstein JL. Topology of SREBP cleavage-activating protein, a polytopic membrane protein with a sterol-sensing domain. *J Biol Chem* 1998; **273**:17243-17250.
 - 25 Hua X, Nohturfft A, Goldstein JL, Brown MS. Sterol resistance in CHO cells traced to point mutation in SREBP cleavage-activating protein. *Cell* 1996; **87**:415-426.
 - 26 Rawson RB, DeBose-Boyd R, Goldstein JL, Brown MS. Failure to cleave sterol regulatory element-binding proteins (SREBPs) causes cholesterol auxotrophy in Chinese hamster ovary cells with genetic absence of SREBP cleavage-activating protein. *J Biol Chem* 1999; **274**:28549-28556.
 - 27 DeBose-Boyd RA, Brown MS, Li WP, *et al*. Transport-dependent proteolysis of SREBP: relocation of site-1 protease from Golgi to ER obviates the need for SREBP transport to Golgi. *Cell* 1999; **99**:703-712.
 - 28 Nohturfft A, Yabe D, Goldstein JL, Brown MS, Espenshade PJ. Regulated step in cholesterol feedback localized to budding of SCAP from ER membranes. *Cell* 2000; **102**:315-323.
 - 29 Goldstein JL, DeBose-Boyd RA, Brown MS. Protein sensors for membrane sterols. *Cell* 2006; **124**:35-46.
 - 30 Yang T, Espenshade PJ, Wright ME, *et al*. Crucial step in cholesterol homeostasis: sterols promote binding of SCAP to INSIG-1, a membrane protein that facilitates retention of SREBPs in ER. *Cell* 2002; **110**:489-500.
 - 31 Yabe D, Brown MS, Goldstein JL. Insig-2, a second endoplasmic reticulum protein that binds SCAP and blocks export of sterol regulatory element-binding proteins. *Proc Natl Acad Sci USA* 2002; **99**:12753-12758.
 - 32 Sun LP, Seemann J, Goldstein JL, Brown MS. From the cover: sterol-regulated transport of SREBPs from endoplasmic reticulum to Golgi: Insig renders sorting signal in Scap inaccessible to COPII proteins. *Proc Natl Acad Sci* 2007; **104**:6519-6526.
 - 33 Kuwabara PE, Labouesse M. The sterol-sensing domain: multiple families, a unique role? *Trends Genet* 2002; **18**:193-201.
 - 34 Yabe D, Xia ZP, Adams CM, Rawson RB. Three mutations in sterol-sensing domain of SCAP block interaction with insig and render SREBP cleavage insensitive to sterols. *Proc Natl Acad Sci USA* 2002; **99**:16672-16677.
 - 35 Nohturfft A, Hua X, Brown MS, Goldstein JL. Recurrent G-to-A substitution in a single codon of SREBP cleavage-activating protein causes sterol resistance in three mutant Chinese hamster ovary cell lines. *Proc Natl Acad Sci USA* 1996; **93**:13709-

- 13714.
- 36 Nohturfft A, Brown MS, Goldstein JL. Sterols regulate processing of carbohydrate chains of wild-type SREBP cleavage-activating protein (SCAP), but not sterol-resistant mutants Y298C or D443N. *Proc Natl Acad Sci USA* 1998; **95**:12848-12853.
- 37 Sever N, Yang T, Brown MS, Goldstein JL, DeBose-Boyd RA. Accelerated degradation of HMG CoA reductase mediated by binding of insig-1 to its sterol-sensing domain. *Mol Cell* 2003; **11**:25-33.
- 38 Sever N, Song BL, Yabe D, *et al*. Insig-dependent ubiquitination and degradation of mammalian 3-hydroxy-3-methylglutaryl-CoA reductase stimulated by sterols and geranylgeraniol. *J Biol Chem* 2003; **278**:52479-52490.
- 39 Lee PC, Sever N, DeBose-Boyd RA. Isolation of sterol-resistant Chinese hamster ovary cells with genetic deficiencies in both Insig-1 and Insig-2. *J Biol Chem* 2005; **280**:25242-25249.
- 40 Song BL, DeBose-Boyd RA. Ubiquitination of 3-hydroxy-3-methylglutaryl-CoA reductase in permeabilized cells mediated by cytosolic E1 and a putative membrane-bound ubiquitin ligase. *J Biol Chem* 2004; **279**:28798-28806.
- 41 Pickart CM. Mechanisms underlying ubiquitination. *Annu Rev Biochem* 2001; **70**:503-533.
- 42 Weissman AM. Themes and variations on ubiquitylation. *Nat Rev Mol Cell Biol* 2001; **2**:169-178.
- 43 Hershko A, Heller H, Elias S, Ciechanover A. Components of ubiquitin-protein ligase system. Resolution, affinity purification, and role in protein breakdown. *J Biol Chem* 1983; **258**:8206-8214.
- 44 Hershko A, Heller H, Eytan E, Reiss Y. The protein substrate binding site of the ubiquitin-protein ligase system. *J Biol Chem* 1986; **261**:11992-11999.
- 45 Hershko A, Ciechanover A, Varshavsky A. Basic Medical Research Award. The ubiquitin system. *Nat Med* 2000; **6**:1073-1081.
- 46 Song BL, Sever N, DeBose-Boyd RA. Gp78, a membrane-anchored ubiquitin ligase, associates with Insig-1 and couples sterol-regulated ubiquitination to degradation of HMG CoA reductase. *Mol Cell* 2005; **19**:829-840.
- 47 Lorick KL, Jensen JP, Fang S, *et al*. RING fingers mediate ubiquitin-conjugating enzyme (E2)-dependent ubiquitination. *Proc Natl Acad Sci USA* 1999; **96**:11364-11369.
- 48 Ponting CP. Proteins of the endoplasmic-reticulum-associated degradation pathway: domain detection and function prediction. *Biochem J* 2000; **351 Pt 2**:527-535.
- 49 Cao J, Wang J, Qi W, *et al*. Ufd1 is a cofactor of gp78 and plays a key role in cholesterol metabolism by regulating the stability of HMG-CoA reductase. *Cell Metab* 2007; **6**:115-128.
- 50 Ye Y, Meyer HH, Rapoport TA. The AAA ATPase Cdc48/p97 and its partners transport proteins from the ER into the cytosol. *Nature* 2001; **414**:652-656.
- 51 Gong Y, Lee JN, Lee PC, *et al*. Sterol-regulated ubiquitination and degradation of Insig-1 creates a convergent mechanism for feedback control of cholesterol synthesis and uptake. *Cell Metab* 2006; **3**:15-24.
- 52 Lee JN, Song B, DeBose-Boyd RA, Ye J. Sterol-regulated degradation of Insig-1 mediated by the membrane-bound ubiquitin ligase gp78. *J Biol Chem* 2006; **281**:39308-39315.
- 53 Schroepfer Jr GJ. Oxysterols: modulators of cholesterol metabolism and other processes. *Physiol Rev* 2000; **80**:361-554.
- 54 Bjorkhem I. Do oxysterols control cholesterol homeostasis? *J Clin Invest* 2002; **110**:725-730.
- 55 Russell DW. Oxysterol biosynthetic enzymes. *Biochim Biophys Acta (BBA) Mol Cell Biol Lipids* 2000; **1529**:126-135.
- 56 Radhakrishnan A, Sun LP, Kwon HJ, Brown MS, Goldstein JL. Direct binding of cholesterol to the purified membrane region of SCAP; mechanism for a sterol-sensing domain. *Mol Cell* 2004; **15**:259-268.
- 57 Adams CM, Goldstein JL, Brown MS. Cholesterol-induced conformational change in SCAP enhanced by Insig proteins and mimicked by cationic amphiphiles. *Proc Natl Acad Sci USA* 2003; **100**:10647-10652.
- 58 Brown AJ, Sun L, Feramisco JD, Brown MS, Goldstein JL. Cholesterol addition to ER membranes alters conformation of SCAP, the SREBP escort protein that regulates cholesterol metabolism. *Mol Cell* 2002; **10**:237-245.
- 59 Radhakrishnan A, Ikeda Y, Kwon HJ, Brown MS, Goldstein JL. From the Cover: Sterol-regulated transport of SREBPs from endoplasmic reticulum to Golgi: Oxysterols block transport by binding to Insig. *PNAS* 2007; **104**:6511-6518.
- 60 Chen HW, Leonard DA, Fischer RT, Trzaskos JM. A mammalian mutant cell lacking detectable lanosterol 14 alpha-methyl demethylase activity. *J Biol Chem* 1988; **263**:1248-1254.
- 61 Leonard DA, Kotarski MA, Tessitore JE, Favata MF, Trzaskos JM. Post-transcriptional regulation of 3-hydroxy-3-methylglutaryl coenzyme A reductase by 3 beta-hydroxy-lanost-8-en-32-al, an intermediate in the conversion of lanosterol to cholesterol. *Arch Biochem Biophys* 1994; **310**:152-157.
- 62 Song BL, Javitt NB, DeBose-Boyd RA. Insig-mediated degradation of HMG CoA reductase stimulated by lanosterol, an intermediate in the synthesis of cholesterol. *Cell Metabolism* 2005; **1**:179-189.
- 63 Gaylor JL. Membrane-bound enzymes of cholesterol synthesis from Lanosterol. *Biochem Biophys Res Commun* 2002; **292**:1139-1146.
- 64 Williams MT, Gaylor JL, Morris HP. Investigation of the rate-determining microsomal reaction of cholesterol biosynthesis from lanosterol in Morris hepatomas and liver. *Cancer Res* 1977; **37**:1377-1383.
- 65 Xu F, Rychnovsky SD, Belani JD, *et al*. Dual roles for cholesterol in mammalian cells. *Proc Natl Acad Sci USA* 2005; **102**:14551-14556.
- 66 Nguyen AD, McDonald JG, Bruick RK, DeBose-Boyd RA. Hypoxia stimulates degradation of 3-hydroxy-3-methylglutaryl-coenzyme A reductase through accumulation of lanosterol and hypoxia-inducible factor-mediated induction of insigs. *J Biol Chem* 2007; **282**:27436-27446.
- 67 Maxwell PH, Wiesener MS, Chang GW, *et al*. The tumour suppressor protein VHL targets hypoxia-inducible factors for oxygen-dependent proteolysis. *Nature* 1999; **399**:271-275.
- 68 Epstein AC, Gleadle JM, McNeill LA, *et al*. *C. elegans* EGL-9 and mammalian homologs define a family of dioxygenases that regulate HIF by prolyl hydroxylation. *Cell* 2001; **107**:43-54.
- 69 Bruick RK, McKnight SL. A conserved family of prolyl-4-hydroxylases that modify HIF. *Science* 2001; **294**:1337-1340.
- 70 Schofield CJ, Ratcliffe PJ. Oxygen sensing by HIF hydroxylases. *Nat Rev Mol Cell Biol* 2004; **5**:343-354.
- 71 Semenza GL. Hydroxylation of HIF-1: oxygen sensing at the molecular level. *Physiology (Bethesda)* 2004; **19**:176-182.

- 72 Jaakkola P, Mole DR, Tian YM, *et al.* Targeting of HIF-alpha to the von Hippel-Lindau ubiquitylation complex by O2-regulated prolyl hydroxylation. *Science* 2001; **292**:468-472.
- 73 Kim H, Lee DK, Choi JW, *et al.* Analysis of the effect of aging on the response to hypoxia by cDNA microarray. *Mech Ageing Dev* 2003; **124**:941-949.
- 74 Mense SM, Sengupta A, Zhou M, *et al.* Gene expression profiling reveals the profound upregulation of hypoxia-responsive genes in primary human astrocytes. *Physiol Genom* 2006; **25**:435-449.
- 75 Sung FL, Hui EP, Tao Q, *et al.* Genome-wide expression analysis using microarray identified complex signaling pathways modulated by hypoxia in nasopharyngeal carcinoma. *Cancer Lett* 2007; **253**:74-88.
- 76 Song BL, DeBose-Boyd RA. Insig-dependent ubiquitination and degradation of 3-hydroxy-3-methylglutaryl coenzyme A reductase stimulated by delta- and gamma-tocotrienols. *J Biol Chem* 2006; **281**:25054-25061.
- 77 Sever N, Lee PCW, Song BL, Rawson RB, DeBose-Boyd RA. Isolation of mutant cells lacking Insig-1 through selection with SR-12813, an agent that stimulates degradation of 3-hydroxy-3-methylglutaryl-coenzyme A reductase. *J Biol Chem* 2004; **279**:43136-43147.
- 78 Seabra MC, Mules EH, Hume AN. Rab GTPases, intracellular traffic and disease. *Trends Mol Med* 2002; **8**:23-30.
- 79 Engelking LJ, Liang G, Hammer RE, *et al.* Schoenheimer effect explained – feedback regulation of cholesterol synthesis in mice mediated by Insig proteins. *J Clin Invest* 2005; **115**:2489-2498.
- 80 Illingworth DR, Tobert JA. HMG-CoA reductase inhibitors. *Adv Protein Chem* 2001; **56**:77-114.
- 81 Kita T, Brown MS, Goldstein JL. Feedback regulation of 3-hydroxy-3-methylglutaryl coenzyme A reductase in livers of mice treated with mevlinolin, a competitive inhibitor of the reductase. *J Clin Invest* 1980; **66**:1094-1100.
- 82 Singer II, Kawka DW, Kazazis DM, *et al.* Hydroxymethylglutaryl-coenzyme A reductase-containing hepatocytes are distributed periportally in normal and mevlinolin-treated rat livers. *Proc Natl Acad Sci USA* 1984; **81**:5556-5560.
- 83 Reihner E, Rudling M, Stahlberg D, *et al.* Influence of pravastatin, a specific inhibitor of HMG-CoA reductase, on hepatic metabolism of cholesterol. *N Engl J Med* 1990; **323**:224-228.