

Journal of Nematology 25(3):315-331. 1993.
© The Society of Nematologists 1993.

Feeding Habits in Soil Nematode Families and Genera—An Outline for Soil Ecologists

G. W. YEATES,¹ T. BONGERS,² R. G. M. DE GOEDE,³ D. W. FRECKMAN,⁴ AND S. S. GEORGIEVA⁵

Abstract: Because research on nematode involvement in trophic interactions, foodweb structure, and biodiversity is constrained by lack of an overview of nematode feeding habits, this outline presents a consensus of current thought on nematode feeding habits. The source of food is fundamental to trophic interactions and provides the basis for our definitions of the essential feeding types: 1) plant feeder, 2) hyphal feeder, 3) bacterial feeder, 4) substrate ingester, 5) predator of animals, 6) unicellular eucaryote feeder, 7) dispersal or infective stage of parasites, and 8) omnivore. Lists of families and genera with their presumed feeding types are given. Major gaps in knowledge of feeding in the smaller tylenchids and many dorylaims are noted.

Key words: bacterial feeding, feeding habit, foodweb, fungal feeding, nematode, omnivore, predator, soil ecology, trophic interaction.

With the increasing interest of soil ecologists in the role of nematodes in ecosystem processes (roles such as nutrient cycling, biological control and economic crop loss), there is an unmet need for a concise summary of current knowledge of nematode feeding habits. The analysis of available information is made increasingly difficult by changes in nematode systematics and recent contributions to nematode ecology. When all species of nematodes can be confidently assigned to feeding groups,

there will follow a better understanding of the role of nematodes in soil and how changes in environmental factors influence the composition of the nematode fauna.

The first comprehensive review of nematode feeding habits was given by Neilsen (77). In an attempt to produce functional groups based on feeding habits, Paramonov (84) applied to nematodes terms such as "pararhizobes" (occur in the rhizosphere and sometimes damage plants) and "dyssaprobates" (feed in decomposing material but may enter healthy tissue). Both Wasilewska (121) and Yeates (127) grouped plant and soil nematodes by feeding habits. The classification of Tylenchida advocated by Siddiqi (99) has a strong "feeding habit" component. Recent ecological studies have revealed that feeding-habit groupings may not be sharply delimited. For example, abundant populations of *Aphelenchoides*, *Tylenchus*, *Tylencholaimus*, and *Ditylenchus* were discovered that could only be classified as "root/fungal feeding nematodes" (105), as well as "predacious" mononchids that multiplied using bacteria as a food source (130). These examples demonstrate the apparently arbitrary nature of traditional nematode feed-

Received for publication 12 January 1993.

Review.

¹ Landcare Research, Private Bag 31902, Lower Hutt, New Zealand.

² Nematology Department, Wageningen Agricultural University, P.O.B. 8123, 6700 ES Wageningen, The Netherlands.

³ Biological Station of Wageningen Agricultural University, Kampsweg 27, 9418 PD Wijkster, The Netherlands.

⁴ Department of Nematology, University of California, Riverside, CA 92521. Present Address: Natural Resource Ecology Laboratory, Colorado State University, Fort Collins, CO 80523.

⁵ Nematology Laboratory, Department of Zoology, Faculty of Biology, University of Sofia, Sofia 1421, Bulgaria.

We are grateful to B. Sohlenius, V. R. Ferris, D. C. Coleman, S. Bostrom, R. K. Niles, K. Prejs, D. Wardle, and P. Arpin for their comments on the manuscript, and to D. J. Carlisle, E. M. Courtright, and T. Gates of D.W.F.'s lab for proofreading and library research. Support from the Foundation for Research, Science and Technology (NZ) to G.W.Y. and from the National Science Foundation (USA) Grant BSR 8818049 to D.W.F. is gratefully acknowledged.

ing groups. Moreover, feeding habits of many nematodes have been inferred rather than confirmed by maintenance over many generations under biologically defined conditions.

Following Petersen and Luxton (86), we use "grazing foodweb" and "detritus foodweb" as terms for communities based on living green plants and dead organic matter, respectively. Their comments on the merging of the two webs are particularly relevant for nematodes, which are so often abundant at interfaces between living and dead material (as in the rhizosphere).

This paper is not a literature review and does not present new results; rather we present a basis for advancing understanding of the role of nematodes in soil ecology. The paper originated in discussions at the Second International Nematology Congress and lists nematode families and genera with our assessment of current understanding of their feeding habits. We hope that this paper will serve as a framework for ecologists to use independently of taxonomic philosophies. Andr assy (2,3), L orenzen (58), Maggenti (62), and Siddiqi (99) all have differing approaches to the general classification of nematodes. The most recent taxonomic overview is contained in the Manual of Agricultural Nematology (76). The generic makeup of the families we use is compatible with these recent works; because they often assign differing taxonomic ranks to these groups, we do not list authorities.

CONFLICTING RESULTS AND OBSERVATIONS

Soil ecologists are primarily concerned with relationships between biological populations and the soil environment, whether it be a high-input agroecosystem, natural ecosystem, or an area managed for sustained production. When these populations are cultured singly or together in the laboratory, many possible interactions and their consequences are reduced (52); thus, results are difficult to extrapolate to field conditions. The following are examples of such difficult extrapolations.

1. Although normally regarded as bacterial feeders, *Chiloplacus* (Acrobelidae) and *Rhabditis* (Rhabditidae) have been cultured on the fungi *Phoma* sp. and *Pythium middletonii*, respectively (39,92). In both cases, the "primary" food source is bacterial; the fact that the apparent secondary food belongs to the same trophic level of the "detritus foodweb" as the primary food shows the value of acknowledging such broad foodwebs. However, our interpretation of this dual feeding habit is i) bacterial contamination of the nematode cultures or ii) direct uptake of nutrients by nematodes, as occurs in axenic culture of *Caenorhabditis elegans* (21,70,132).

2. Hooper and Cowland (45) cultured the foliar nematode *Aphelenchoides ritzeimbosi*, which normally feeds in the "grazing foodweb," on fungi ("detritus foodweb"), reinforcing observations that fungal feeding is the normal situation in Apelenchida. Just as plant root cells are fed on in a variety of ways (see "plant feeding"), so are fungi (1).

3. It is difficult to extrapolate data from closely controlled monoxenic cultures of *Pratylenchus* and *Radopholus* on carrot discs to the heterogenous environment of field populations.

4. Although certain mononchids successfully cultured on bacteria contain living bacteria within the intestines (6), we question whether sufficient aggregations of bacteria exist under field conditions for bacterial ingestion to be of significant nutritional importance to the large mononchids.

5. The intestine of mononchids, rhabditids, dorylaims, etc., often appear pigmented, but such pigmentation has not been observed in Tylenchida or plant-feeding *Trichodorus* and *Longidorus* (76). There has been no attempt to relate this pigmentation to feeding habits.

6. Axenic culture of *Caenorhabditis elegans* and other "bacterial feeding" nematodes (21,70,73,132) highlights problems of interpretation, but there is evidence that development may be slower under axenic conditions (20). Bacterial feeding is re-

garded here as the principal source of nutrition for such nematodes under field conditions. Physical crushing of bacteria has been demonstrated in the pharyngeal bulb of *Acrobeloides nanus*, but bacterial feeding nematodes may defecate living bacteria and there is no general knowledge as to whether some nematodes actually kill, rupture, or lyse bacteria or merely remove adhering organic compounds (126). In several situations, there may be a degree of direct nutrient uptake through the cuticle or epidermis (33,50,79,132).

7. Although identification to family level is usually adequate, identification to species is sometimes necessary to accurately assign nematodes to trophic groups. For example, *Ditylenchus dipsaci* is an economically important plant feeding species, but most *Ditylenchus* spp. are hyphal feeding.

8. The delicate-speared Tylenchidae and Psilenchidae remain problematic. Although Sohlenius et al. (105) regarded several groups of them as "root/fungal feeding nematodes," the literature is conflicting and we have been unable to form a consensus about the classification of these two families. We have generally listed them as feeding on root epidermal cells and root hairs. Specific studies of their feeding in the rhizoplane would provide valuable information.

FEEDING TYPES OF NEMATODES IN PLANT AND SOIL SYSTEMS

Ideally the feeding habits of each nematode species should be determined in each particular ecological setting. Because this task is impractical, we offer the following outline for general use. Regardless of the trophic group, we strongly advocate appending species lists and depositing voucher specimens to make reinterpretation possible.

1. *Plant feeding*: This involves feeding on vascular plants; a tylenchoid stomatostyle or dorylaimoid odontostyle is always present. The posthatching life stages of most species are migratory. In sedentary

species, the feeding site of the female may be undifferentiated, uninucleate, or polynucleate. Males of sedentary species sometimes have a degenerate stylet or reduced oesophagus, but data on the nutrition of such males is inadequate. Plant feeders may be polyphagous or show host specificity. Migratory species may generally be classified as ecto- or endoparasites. Feeding sites may be root-hair, epidermal, cortical, or vascular.

Although apparently not actively feeding, the migratory phases of *Pratylenchus*, infective second-stage *Heterodera*, and resistant stages of *Paratylenchus* are each an essential part of the life-cycle of these obligate plant feeders and are thus included in this category. This group includes algal feeders that have a narrow stylet and do not ingest chloroplasts, in contrast to types that swallow unicellular algae more or less intact.

The group may be subdivided into the following six groups: 1a) sedentary parasites (e.g., females of *Heterodera*, *Globodera*, *Meloidogyne*, *Verutus*, *Sphaeronema*); 1b) migratory endoparasites (e.g., *Pratylenchidae*, some *Anguinidae*); 1c) semi-endoparasites (e.g., *Hoplolaimidae*, *Telotylenchus*); 1d) ectoparasites (e.g., *Dolichodoridae*, *Cephalenchus*, *Criconematidae*, *Hemicyclophoridae*, *Paratylenchidae*, *Trichodoridae*, *Pungentus*, *Longidoridae*); 1e) epidermal cell and root hair feeders (e.g., *Tylenchidae*, *Psilenchidae*, *Atylenchidae*); and 1f) algal, lichen (algal or fungal component), or moss feeders that feed by piercing (e.g., *Tylenchus*, *Laimaphelenchus*, *Anguinidae*).

2. *Hyphal feeding*: This involves penetration of fungal hyphae by a stomatostyle or odontostyle (stylet or spear). In addition to obligate hyphal feeders, this group includes the alternate life cycle of some invertebrate parasites (e.g., *Deladenus*) (75). Yeasts may be included as a food source under this heading, except when ingested whole (see "unicellular eucaryote feeding"). Feeding on hyphae of saprophytic fungi has ecological implications quite different from feeding on mycorrhizal fungi.

There is need for research on such interactions to determine whether the same nematode can feed on both saprophytic and mycorrhizal fungi.

3. *Bacterial feeding*: This category includes species that feed on any procaryote food source, whether through a narrow (*Rhabditis*, *Alaimus*) or broad (*Diplogaster*) mouth; axenic culture indicates that there may be an element of absorption of nutrients (33,50,79,132). Species with a broad mouth may ingest other types of food. The soil stages of certain nematode parasites of vertebrates and invertebrates that feed on bacteria should be included. Although morphological data (75) suggest that infective stages of *Steinernema* may be non-feeding, they may be mass produced on artificial media, providing their bacterial symbionts are present (76). Some Rhabditidae and Diplogasteridae may use a phoretic (transport) host, especially insects.

4. *Substrate ingestion*: This type of feeding occurs in at least diplogasterids and *Daptonema* sp. Substrate ingestion may be incidental to bacterial feeding, predation, and unicellular eucaryote feeding in many groups, because more than a pure food source is ingested. Although soil nematodes may ingest and lyse bacteria, there is no available evidence that digestion of complex organic substrates occurs in the gut of soil nematodes (a fact that may reflect technical difficulty). Consequently, no genera have been given primary classification under this type. The expression "non-selective deposit feeding" used for aquatic nematodes covers a similar situation.

5. *Animal predation*: Some species of nematodes feed on invertebrates such as protozoa, nematodes, rotifers, and enchytraeids, either as "ingesters" (type 5a; e.g., *Diplogaster*, *Mononchus*, *Nyngolaimus*) or as "piercers" (type 5b), sucking body fluids through a narrow stylet (e.g., *Seinura*, *Labronema*, *Laimaphelenchus*). The intestine of piercers never contains distinct prey remnants.

6. *Unicellular eucaryote feeding*: A wide

range of nematodes reportedly feed on diatoms or other algae, but lack of marker structures in the nematode food and the presence of globules, pigmentation, or inclusions in nematode intestines make determination difficult. This feeding type includes ingestion of fungal spores and whole yeast cells.

7. *Dispersal or infective stages of animal parasites*: Other stages of animal parasitic nematodes outside their alternate or definitive hosts may occur in the soil (e.g., *Deladenus*, *Heterorhabditis*) or vertebrates (e.g., *Strongyloides*); we include here entomogenous species. If they feed and contribute to soil processes, these species should be included in other appropriate categories; if they invade a host they leave the soil system (in the narrow sense); if they die in the soil they contribute to nutrient pools. We do not include in this category forms (especially Rhabditidae and Diplogasteridae) that merely use animals as phoretic (transport) hosts.

8. *Omnivorous*: Although some nematodes appear normally to feed on a wide range of foods (particularly combining feeding types 2-6), it is best to restrict use of this term to (a few) dorylaims (31,95); when possible, nematodes should be classified in types 1-7. A similar procedure was used for soil fauna by Petersen et al. (87).

FEEDING TYPES IN NOMINAL ORDERS AND FAMILIES

In the following systematic list of nematode families and feeding type, we give indicative genera for each family; the numbers refer to the eight feeding types. Numbers in parentheses refer to tentative assignments or situations with alternative food sources. For a given family or genus, if two or more types are indicated, they are presented in strictly numerical order. For some groups, detailed information is given (e.g., algal feeding may be a specialization of unicellular eucaryote feeding). Table 1 contains an alphabetical list of nematode genera and their feeding types.

TABLE 1. Feeding types in nematode genera. Numbers refer to the eight feeding types outlined in the text; numbers in parenthesis indicate a tentative assignment. The main food source is given first. For genera maintained for several generations under defined conditions, unqualified reference is made to that publication by number; for genera where the publication number is preceded by "see," a more general (or less specific) support for allocation to the feeding type is indicated; for genera where we have not obtained a direct reference to feeding activity, the family name is given.

| Genus | Feeding type | Family or literature citation |
|-------------------------|-------------------|---|
| <i>Ablechroiuulus</i> | 3 | Rhabditidae |
| <i>Achromadora</i> | (6) | Achromodoridae |
| <i>Acontylus</i> | 1b | (66) |
| <i>Acrobeles</i> | 3 | (111,126) |
| <i>Acrobeloidea</i> | 3 | (71,104,126) |
| <i>Acrobelophis</i> | 3 | Cephalobidae |
| <i>Acrolobus</i> | 3 | Cephalobidae |
| <i>Acromoldavicus</i> | 3 | Cephalobidae |
| <i>Acrostichus</i> | 3 | Diplogasteridae; (88) |
| <i>Actinca</i> | 5, 8 | Actinolaimidae |
| <i>Actinolaimus</i> | 5, 8 | (56); (see 31,100) |
| <i>Aetholaimus</i> | 5 | Nygolaimidae |
| <i>Aglenchus</i> | 1e | Tylenchidae; (122) |
| <i>Alaimus</i> | 3 | (see 77) |
| <i>Allantonema</i> | 7 or 2 | (119) |
| <i>Allodorylaimus</i> | 8 | Dorylaimidae |
| <i>Allotrichodorus</i> | 1d | Trichodoridae; (see Decraemer in 76) |
| <i>Amphidelus</i> | 3 | Alaimidae |
| <i>Amplimerlinius</i> | 1d | (14,36); (see Anderson & Potter in 76) |
| <i>Anaplectus</i> | 3 | (122) |
| <i>Anatonchus</i> | 5a | (18); (see 100) |
| <i>Anguina</i> | 1a or b | (22); (see Krall in 76) |
| <i>Anomyctus</i> | 2? | (see Nickle & Hooper in 76) |
| <i>Antarctylus</i> | 1c | Hoplolaiminae |
| <i>Aorolaimus</i> | 1c | Hoplolaiminae |
| <i>Aphanolaimus</i> | 3 | Halaphanolaimidae |
| <i>Aphasmatylenchus</i> | 1c | (34); (see Fortuner in 76) |
| <i>Aphelenchoides</i> | 2 or 1b, 1e or 1f | (68); (see 122) |
| <i>Aphelenchus</i> | 2 or 1e | (17,23,40,65); (see 122) |
| <i>Aporcelaimellus</i> | 5, 8 | (117,122); (see 31) |
| <i>Aporcelaimium</i> | 8 | Aporcelaimidae |
| <i>Aporcelaimus</i> | 5, 8 | (114,122); (see 31,100) |
| <i>Apratylenchoides</i> | 1b | Pratylenchidae |
| <i>Aprutides</i> | 2 | Aphlenchoididae |
| <i>Aquatides</i> | 5 | (10); (see 100) |
| <i>Atylenchus</i> | 1d | Tylenchidae |
| <i>Aulolaimus</i> | 3 | Cryptonchidae |
| <i>Axonchium</i> | 1, 8? | (see 100) |
| <i>Basiria</i> | 1e | Tylenchidae |
| <i>Bastiana</i> | 3 | Bastianiidae |
| <i>Bathyodontus</i> | 3 | Bathyodontidae; (see 12) |
| <i>Belonolaimus</i> | 1d | (15) |
| <i>Bicirronema</i> | 3 | Chambersiellidae |
| <i>Bitylenchus</i> | 1d | Tylenchorhynchidae; (see Anderson & Potter in 76) |
| <i>Boleodor</i> | 1e or 2(?) | Tylenchidae |
| <i>Brevibucca</i> | 3 | Brevibuccidae |
| <i>Brittonema</i> | 5, 8 | Actinolaimidae |
| <i>Bunonema</i> | 3 | (77,128) |
| <i>Bursaphelenchus</i> | 2 | (32,63); (see Nickle & Hooper in 76) |
| <i>Bursilla</i> | 3 | Rhabditidae |
| <i>Butlerius</i> | 3, 5 | (28,89,101); (see 100) |
| <i>Cacopaurus</i> | 1a | Paratylenchidae; (112); (see Raski in 76) |
| <i>Cactodera</i> | 1a | (19,98); (see Baldwin & Mundo-Ocampo in 76) |
| <i>Caenorhabditis</i> | 3 | (25,78,97); (see 107) |

TABLE 1. *Continued*

| Genus | Feeding type | Family or literature citation |
|-------------------------|---------------|--|
| <i>Caloosia</i> | 1d | Hemicyclophoridae |
| <i>Campydora</i> | 8? | Campydoridae |
| <i>Carcharolaimus</i> | 5 | (28); (see 100) |
| <i>Cephalenchus</i> | 1d | (37,109); (see 122) |
| <i>Cephalobus</i> | 3 | (81,88); (see 72,122) |
| <i>Ceratoplectus</i> | 3 | Plectididae |
| <i>Cervidellus</i> | 3 | (122) |
| <i>Chambersiella</i> | 3 | Chambersiellidae |
| <i>Cheilorhabditis</i> | 3 | Rhabditidae |
| <i>Chiloplacus</i> | 3 | (118,122) |
| <i>Choanolaimus</i> | 5a | Choanolaimidae |
| <i>Chondronema</i> | 7, 2? | (16) |
| <i>Chromadorina</i> | 3, 6? | Chromadoridae; (116) |
| <i>Chromadorita</i> | 6 | (49) |
| <i>Chronogaster</i> | 3 | Leptolaimidae |
| <i>Chrysonemoides</i> | ? | Chrysonematidae |
| <i>Clarkus</i> | 5a | (67,106); (see 101,130) |
| <i>Coarctadera</i> | 3 | Rhabditidae; (see 107) |
| <i>Cobbonchus</i> | 5a | (see 100) |
| <i>Coomansus</i> | 5a | (31) |
| <i>Coslenchus</i> | 1e | (as <i>Aglenchus</i>); (122) |
| <i>Craspedonema</i> | 3 | Butonematidae |
| <i>Criconema</i> | 1d | (see 26,93) |
| <i>Criconemoides</i> | 1d | (see 26,93) |
| <i>Crossonema</i> | 1d | (see 26,93) |
| <i>Cruznama</i> | 3 | (108) |
| <i>Cryphodera</i> | 1a | (see 8,60) |
| <i>Cryptonchus</i> | 3 | Cryptonchidae |
| <i>Curviditis</i> | 3 | Rhabditidae |
| <i>Cuticonema</i> | 3 | Breviibuccidae |
| <i>Cuticularia</i> | 3 | Rhabditidae |
| <i>Cylindrolaimus</i> | 3 | Diplopeltidae |
| <i>Daptonema</i> | 3, 4, 5 or 6 | (13) |
| <i>Deladenus</i> | 2 | (9,129); (see 62) |
| <i>Demaniella</i> | 3 | (88) |
| <i>Deontolaimus</i> | 3 | Leptolaimidae |
| <i>Desmodora</i> | 3, 6 | Desmodoridae; (74) |
| <i>Desmolaimus</i> | 3 | Linhomoeidae |
| <i>Desmoscolex</i> | 3 | Desmoscolecidae |
| <i>Dichromadora</i> | 3, 6? | (13,120) |
| <i>Diphtherophora</i> | 2 | Diphtherophoridae; (see 4) |
| <i>Diplenteron</i> | 3? | (64,125); (see 100) |
| <i>Diplogaster</i> | 3, 5a, 6 or 8 | (see 72,100) |
| <i>Diplogasteritus</i> | 3 | (124) |
| <i>Diplogasteroides</i> | 3 | Diplogasteroididae |
| <i>Diploscapter</i> | 3 | (42,88) |
| <i>Discolaimium</i> | 5? | (see 100) |
| <i>Discolaimus</i> | 5 | (28,56); (see 31,100) |
| <i>Ditylenchus</i> | 2, or 1b | (55,122); (see Sturhan & Brezeski in 76) |
| <i>Dolichodorus</i> | 1d | (82,83,85); (see Smart & Nguyen in 76) |
| <i>Dolichorhabditis</i> | 3 | Rhabditidae |
| <i>Dolichorhynchus</i> | 1d | Dolichodoridae |
| <i>Domorganus</i> | 3 | Diplopeltidae |
| <i>Dorydorella</i> | 8 | Dorylaimidae |
| <i>Dorylaimellus</i> | 1, 2? | Belonidiridae; (see 131) |
| <i>Dorylaimoides</i> | 8? | Leptonchidae |
| <i>Dorylaimus</i> | 8 | (56); (see 77,100) |
| <i>Doryllium</i> | 2? | Leptonchidae |
| <i>Drilocephalobus</i> | 3 | Ostellidae |

TABLE 1. *Continued*

| Genus | Feeding type | Family or literature citation |
|----------------------------|--------------|--|
| <i>Durinema</i> | 5 | (see 77,100) |
| <i>Ecphyadophora</i> | 1d or 2(?) | Ecphyadophoridae |
| <i>Ecumenicus</i> | 8 | Dorylaimidae |
| <i>Elaphonema</i> | 3 | Elaphonematidae |
| <i>Enchodelus</i> | 8 or 6? | Nordiidae |
| <i>Epidorylaimus</i> | 8 | Dorylaimidae |
| <i>Ereptonema</i> | 3 | Ereptonemidae |
| <i>Ethmolaimus</i> | 3, 6? | Ethmolaimidae |
| <i>Eucephalobus</i> | 3 | (122) |
| <i>Eudorylaimus</i> | 5, 8 | (29,44,117); (see 31,100) |
| <i>Eurystomina</i> | (5) | Eurystominidae |
| <i>Eumonhystera</i> | 3, 4 | Monhysteridae |
| <i>Euteratocephalus</i> | 3 | Teratocephalidae |
| <i>Fictor</i> | 3, 5, 6 | (88,103); (see 100) |
| <i>Filenchus</i> | 1e | Tylenchidae |
| <i>Funaria</i> | 2? | Leptonchidae |
| <i>Fungittonchium</i> | 2 | Iontonchiidae (see 99) |
| <i>Geocenamus</i> | 1d | Dolichodoridae |
| <i>Geomonhystera</i> | 3, 4 | Monhysteridae |
| <i>Glauxinema</i> | 6 | Neodiplogasteridae |
| <i>Globodera</i> | 1a | (53); (see Baldwin & Mundo-Ocampo in 76) |
| <i>Goffartia</i> | 3 | Diplogasterididae |
| <i>Goodeyus</i> | 3 | Cylindrocorporidae |
| <i>Gracilacus</i> | 1d | Paratylenchidae; (see Raski in 76) |
| <i>Granonchulus</i> | 5a | Mononchidae; (35) |
| <i>Haliplectus</i> | 3 | Haliplectidae |
| <i>Helicotylenchus</i> | 1c | (51); (see Fortuner in 76) |
| <i>Hemicriconemoides</i> | 1d | Criconematidae; (see 93) |
| <i>Hemicycliophora</i> | 1d | Criconematidae; (see 93) |
| <i>Heterocephalobellus</i> | 3 | Cephalobidae |
| <i>Heterocephalobus</i> | 3 | Cephalobidae |
| <i>Heterodera</i> | 1a | (see Baldwin & Mundo-Ocampo in 76) |
| <i>Heterorhabditis</i> | 7, 3 | (90); (see Wouts in 76) |
| <i>Hexatylus</i> | 2 | (24); (see 72) |
| <i>Hirschmanniella</i> | 1b | (7); (see Loof in 76) |
| <i>Hoplolaimus</i> | 1c | Hoplolaimidae |
| <i>Hoplolytus</i> | 1b | Pratylenchidae |
| <i>Howardula</i> | 7 or 2 | Allantonematidae |
| <i>Huntaphelenchoides</i> | 2 | Aphelenchoididae |
| <i>Iotonchus</i> | 5a | (67,100,101) |
| <i>Ironus</i> | 5a or 6 | (47); (see 100) |
| <i>Isolaimium</i> | (3) | Isolaimidae |
| <i>Kirjanovia</i> | 3 | Cephalobidae |
| <i>Kochinema</i> | 8 | Nordiidae |
| <i>Labronema</i> | 5, 8 | (28,30,88,122); (see 31,100) |
| <i>Laimaphelenchus</i> | 5b, 1f, 2 | (see 100) |
| <i>Laimydorus</i> | 8 | (see 100) |
| <i>Lelenchus</i> | 1e | Tylenchidae |
| <i>Leptolaimus</i> | 3 | (13) |
| <i>Leptonchus</i> | 2 | (see 31) |
| <i>Longidorella</i> | 1d | Nordiidae |
| <i>Longidorus</i> | 1d | Longidoridae |
| <i>Loofia</i> | 1d | Hemicycliophoridae |
| <i>Macrotrophurus</i> | 1d | Dolichodoridae |
| <i>Malenchus</i> | 1e | Tylenchidae |
| <i>Meloidodera</i> | 1a | (see Baldwin & Mundo-Ocampo in 76) |
| <i>Meloidoderita</i> | 1a | (see 76) |
| <i>Meloidogyne</i> | 1a | (see 27,96) |
| <i>Merlinius</i> | 1d | (see Anderson & Potter in 76) |

TABLE 1. *Continued*

| Genus | Feeding type | Family or literature citation |
|---------------------------|--------------|--|
| <i>Mesocriconema</i> | 1d | (see 93) |
| <i>Mesodiplogaster</i> | 3, 5 | (102,103); (see 100) |
| <i>Mesodorylaimus</i> | 8 | (29); (see 31,95,100) |
| <i>Mesorhabditis</i> | 3 | (107,122,126) |
| <i>Metacrobeles</i> | 3 | Cephalobidae |
| <i>Metadiplogaster</i> | 3 | Diplogasteridae |
| <i>Metateratocephalus</i> | 3 | Teratocephalidae |
| <i>Miconchus</i> | 5a | (see 100) |
| <i>Microdorylaimus</i> | 8 | Nordiidae |
| <i>Microlaimus</i> | 3 | Microlaimidae |
| <i>Monhystera</i> | 3, (4) | (122); (see 77,115) |
| <i>Monhystrella</i> | 3, 4 | (13) |
| <i>Monobutlerius</i> | 3 | Diplogasteridae |
| <i>Mononchoides</i> | 3, 5a | (89); (see 100) |
| <i>Mononchulus</i> | 3 | (see 31) |
| <i>Mononchus</i> | 5a | (38,88); (see 31,100) |
| <i>Monotrichodoris</i> | 1d | Trichodoridae; (see Decraemer in 76) |
| <i>Morulaimus</i> | 1d | Belonolaimidae; (see Smart & Nyguen in 76) |
| <i>Mylonchulus</i> | 5a | (48,67); (see 31,100) |
| <i>Myolaimus</i> | 3 | Myolaimidae |
| <i>Nacobbus</i> | 1a | (see Jatala in 76) |
| <i>Nagelus</i> | 1d | Dolichodoridae |
| <i>Namibinema</i> | 3 | Cephalobidae |
| <i>Neoactinolaimus</i> | 5, 8 | (101); (see 100) |
| <i>Neoaplectana</i> | 7, 3 | (see 76) |
| <i>Neodiplogaster</i> | 3, 4 or 5 | Neodiplogasteridae |
| <i>Neopsilenchus</i> | 1e | Tylenchidae |
| <i>Neothada</i> | 1e or 2(?) | Tylenchidae |
| <i>Nothacrobeles</i> | 3 | Cephalobidae |
| <i>Nothotylenchus</i> | 2 | Anguinidae |
| <i>Nullonchus</i> | 5a | (see 100) |
| <i>Nygolaimium</i> | 5 | (see 100) |
| <i>Nygolaimoides</i> | 5 | Nygolaimidae |
| <i>Nygolaimus</i> | 5 | (113,122); (see 31,100) |
| <i>Odontolaimus</i> | 3 or 6? | Odontolaimidae |
| <i>Odontopharynx</i> | 3, 5a | Odontopharyngidae |
| <i>Odontorhabditis</i> | 3 | Odontorhabditidae |
| <i>Ogma</i> | 1d | (see 93) |
| <i>Oionchus</i> | 3 | Mononchulidae |
| <i>Onchulus</i> | 5a or 6 | Onchulidae |
| <i>Opisthodorylaimus</i> | 8 | Dorylaimidae |
| <i>Orrina</i> | 1a or b | (see Krall in 76) |
| <i>Osstella</i> | 3 | Osstellidae |
| <i>Oxydirus</i> | 1, 8? | Belonidiridae |
| <i>Panagrellus</i> | 3 | (21) |
| <i>Panagrobelum</i> | 3 | Panagrolaimidae |
| <i>Panagrobelus</i> | 3 | Panagrolaimidae |
| <i>Panagrocephalus</i> | 3 | Cephalobidae |
| <i>Panagrolaimus</i> | 3 | (88,126) |
| <i>Paracrobeles</i> | 3 | Cephalobidae |
| <i>Paractinolaimus</i> | 5 | (46) (see 100) |
| <i>Paracyatholaimus</i> | 6? | (13) |
| <i>Parahadronchus</i> | 5a | (see 100) |
| <i>Paralongidorus</i> | 1d | (see 11,54) |
| <i>Paramphidelus</i> | 3 | Alaimidae |
| <i>Paraphanolaimus</i> | 3 | Halaphanolaimidae |
| <i>Paraphelenchus</i> | 2 | (110); (see Nickle & Hooper in 76) |
| <i>Paraplectonema</i> | 3 | Leptolaimidae |
| <i>Pararotylenchus</i> | 1c | (see Fortuner in 76) |

TABLE I. *Continued*

| Genus | Feeding type | Family or literature citation |
|----------------------------|--------------|--------------------------------------|
| <i>Paratrichodoros</i> | 1d | Trichodoridae; (see Decraemer in 76) |
| <i>Paratripyla</i> | 5a | Tripylidae |
| <i>Paratrophurus</i> | 1d | Dolichodoridae |
| <i>Paratylenchus</i> | 1d | (94,122); (see Raski in 76) |
| <i>Paravultus</i> | 5? | Nygolaimidae |
| <i>Paraxonchium</i> | 5, 8 | Aporcelaimidae |
| <i>Pareudiplogaster</i> | 6, 3 | (13) |
| <i>Pariogolaimella</i> | 3 | (88) |
| <i>Paurodontus</i> | 2, 7 | (as <i>Neotylenchus</i>); (40) |
| <i>Pellioiditis</i> | 3 | (107) |
| <i>Pelodera</i> | 3 | (88,107,122) |
| <i>Peltamigratus</i> | 1c | (see 76) |
| <i>Phasmarhabditis</i> | 3 | (107) |
| <i>Placodira</i> | 3 | Cephalobidae |
| <i>Plectonchus</i> | 3 | Brevibuccidae |
| <i>Plectus</i> | 3 | (77,88,122) |
| <i>Pleurotylenchus</i> | 1d | Tylodoridae |
| <i>Pratylenchoides</i> | 1b | (see Loof in 76) |
| <i>Pratylenchus</i> | 1b | (see Loof in 76) |
| <i>Prionchulus</i> | 5a | (5,61,69); (see 100) |
| <i>Prismatolaimus</i> | 3? | (see 77) |
| <i>Pristionchus</i> | 3, 5a | Neodiplogasteridae |
| <i>Prochromadora</i> | 3, 6? | Chromadoridae |
| <i>Prodesmodora</i> | 3 | Desmodoridae |
| <i>Prodorylaimium</i> | 8 | Dorylaimidae |
| <i>Prodorylaimus</i> | 8 | Dorylaimidae |
| <i>Proleptonchus</i> | 8? | Leptonchidae |
| <i>Protocylindrocorpus</i> | 3 | Cylindrocorporidae (88) |
| <i>Protorhabditis</i> | 3 | Rhabditidae |
| <i>Pseudacrobeles</i> | 3 | Cephalobidae |
| <i>Pseudhalenchus</i> | 2 | Anguinidae |
| <i>Pseudoaulolaimus</i> | 3 | Cryptonchidae |
| <i>Psilenchus</i> | 1e | Psilenchidae |
| <i>Pterotylenchus</i> | 1a or b | Anguinidae |
| <i>Pterygorhabditis</i> | 3 | Pterygorhabditidae |
| <i>Punctodera</i> | 1a | (see Baldwin & Mundo-Ocampo in 76) |
| <i>Punctodora</i> | 3, 6? | Chromadoridae |
| <i>Pungentus</i> | 1d, 5, 8 | (see 31,100) |
| <i>Quinisulcius</i> | 1d | (see 76) |
| <i>Radopholus</i> | 1b | (see Loof in 76) |
| <i>Rhabditis</i> | 3 | (88,102,107,122) |
| <i>Rhabditoides</i> | 3 | Rhabditidae |
| <i>Rhabditophanes</i> | 3 | Alloionematidae |
| <i>Rhabdolaimus</i> | 3 | (see 77) |
| <i>Rhabdontolaimus</i> | 3 | (88) |
| <i>Rhadinaphelenchus</i> | 1b | (see Nickle & Hooper in 76) |
| <i>Rhodolaimus</i> | 3 | Bunonematidae |
| <i>Rotylenchulus</i> | 1a | (57) |
| <i>Rotylenchus</i> | 1c | Hoplolaiminae; (see Fortuner in 76) |
| <i>Scottinema</i> | 3 | Cephalobidae; (80) |
| <i>Scutellonema</i> | 1c | Hoplolaiminae; (see Fortuner in 76) |
| <i>Scutylenchus</i> | 1d | Dolichodoridae |
| <i>Sectonema</i> | 5, 8 | (see 100) |
| <i>Seimura</i> | 5b | (28,41,43,101,123); (see 100) |
| <i>Seleborca</i> | 3 | Acrobelidae |
| <i>Sphaerolaimus</i> | 5a | Sphaerolaimidae |
| <i>Sphaeronema</i> | 1a | (see Raski in 76) |
| <i>Sphaerularia</i> | 7, 2 | (91) |
| <i>Sporonchulus</i> | 5a | (see 100) |

TABLE 1. *Continued*

| Genus | Feeding type | Family or literature citation |
|--------------------------|--------------|--------------------------------------|
| <i>Stegelleta</i> | 3 | Cephalobidae |
| <i>Stegelletina</i> | 3 | Cephalobidae |
| <i>Steinernema</i> | 7, 3 | (see 76) |
| <i>Stomachoglossa</i> | (5) | (see 100) |
| <i>Subanguina</i> | 1a or b | (see 76) |
| <i>Sulphuretylenchus</i> | 7 or 2 | (see 76) |
| <i>Synonchium</i> | 5 | (see 100) |
| <i>Telotylenchus</i> | 1c | Dolichodoridae |
| <i>Tenunemellus</i> | 1d or 2(?) | Ecphyadorphoridae |
| <i>Teratocephalus</i> | 3 | (see 77) |
| <i>Teratolobus</i> | 3 | Cephalobidae |
| <i>Teratorhabditis</i> | 3 | Rhabditidae |
| <i>Theristus</i> | 3, 4 or 6 | (see 77) |
| <i>Thonus</i> | 5, 8 | (31) |
| <i>Thornenema</i> | 8 | Dorylaimidae |
| <i>Thornia</i> | 8? | (see 100) |
| <i>Tobrilus</i> | 5a or 6 | (see 100) |
| <i>Torumanawa</i> | 8 | Aporcelaimidae |
| <i>Tricephalobus</i> | 3 | Panagrolaimidae |
| <i>Trichodorus</i> | 1d | Trichodoridae; (see Decraemer in 76) |
| <i>Tripilus</i> | 7, 2 | (see 76) |
| <i>Tripyla</i> | 5a | (77, 100) |
| <i>Trischistoma</i> | 5a | Tripylidae; (see 100) |
| <i>Trophonema</i> | 1a | Tylenchulidae; (see Raski in 76) |
| <i>Trophotylenchus</i> | 1a | Tylenchulidae |
| <i>Trophurus</i> | 1d | Dolichodoridae |
| <i>Turbatrix</i> | 3 | (70, 89) |
| <i>Tylencholaimellus</i> | 2 | (122) |
| <i>Tylencholaimus</i> | 2 | (29, 105, 122); (see 31) |
| <i>Tylenchorhynchus</i> | 1d | (see Anderson & Potter in 76) |
| <i>Tylenchulus</i> | 1a | (see Raski in 76) |
| <i>Tylenchus</i> | 1f, 2? | Tylenchidae; (122) |
| <i>Tylocephalus</i> | 3 | (see 77) |
| <i>Tylodorus</i> | 1d | Tylodoridae |
| <i>Tyrolaimophorus</i> | 2 | Diphtherophoridae |
| <i>Tylopharynx</i> | 3 | Tylopharyngidae |
| <i>Verutus</i> | 1a | (see Baldwin & Mundo-Ocampo in 76) |
| <i>Westindicus</i> | (5) | (see 100) |
| <i>Wilsonema</i> | 3 | (77, 122) |
| <i>Xiphinema</i> | 1d | Longidoridae |
| <i>Ypsylonellus</i> | 3 | Acrobelidae |
| <i>Zeldia</i> | 3 | (126) |
| <i>Zygotylenchus</i> | 1b | (see Loof in 76) |

Order Monhysterida

5 predacious

6 algal feeding

Linhomoeidae: *Desmolaimus*

(3) bacterial feeding

*Order Desmoscolecida*Monhysteridae: *Monhystera*, *Geomonhystera*

3 bacterial feeding

4 substrate ingestion

Desmoscolecidae: *Desmoscolex*

(3) bacterial feeding

Sphaerolaimidae: *Sphaerolaimus*

5 predacious

*Order Araeolaimida*Xyalidae: *Theristus*, *Daptonema*

3 bacterial feeding

4 substrate ingestion

All terrestrial forms are apparently type 3.

Diplopeltidae: *Cylindrolaimus*, *Domorganus*

- (3) bacterial feeding
 7 *Domorganus* (sometimes associated with earthworms)
- Halaphanolaimidae: *Aphanolaimus*
 (3) bacterial feeding
- Haliplectidae: *Haliplectus*
 (3) bacterial feeding
- Leptolaimidae: *Leptolaimus*, *Chronogaster*
 3 bacterial feeding
- Plectidae: *Plectus*, *Anaplectus*
 3 bacterial feeding (*Plectus*)
- Rhabdolaimidae: *Rhabdolaimus*
 3 bacterial feeding
- Wilsonematidae: *Wilsonema*, *Tylocephalus*, *Ereptonema*
 3 bacterial feeding
- Order Chromadorida*
- Achromadoridae: *Achromadora*
 6 algal feeding
- Choanolaimidae: *Choanolaimus*, *Synonchium*
 5 predacious
- Cyatholaimidae: *Paracyatholaimus*
 6 algal feeding
- Desmodoridae: *Desmodora*
 (3) bacterial feeding
 6 algal feeding
- Ethmolaimidae: *Ethmolaimus*
 3 bacterial feeding
 6 algal feeding
- Hypodontolaimidae: *Chromadorita*
 6 algal feeding
- Microlaimidae: *Microlaimus*
 3 bacterial feeding
 6 algal feeding
- Order Rhabditida*
- Members of this order that occur in soil are basically type 3 (bacterial feeding).
- Acrobelidae: *Acrobeles*, *Cervidellus*, *Acrobeloides*
 3 bacterial feeding
- Alloionematidae: *Alloionema*, *Rhabditophanes*
 3 bacterial feeding (soil-dwelling stages only)
 7 insect parasites
- Brevibuccidae: *Brevibucca*, *Cuticonema*
 (3) bacterial feeding; insect associates
- Bunonematidae: *Bunonema*, *Rhodolaimus*, *Craspedonema*
 3 bacterial feeding
- Cephalobidae: *Cephalobus*, *Heterocephalobus*, *Eucephalobus*
 3 bacterial feeding
- Chambersiellidae: *Chambersiella*, *Bicirronema*
 3 bacterial feeding
- Cylindrocorporidae: *Protocylindrocorpus*, *Goodeyus*
 (3) bacterial feeding
- Diploscapteridae: *Diploscapter*
 3 bacterial feeding
- Elaphonematidae: *Elaphonema*
 3 bacterial feeding
- Myolaimidae: *Myolaimus*
 3 bacterial feeding
- Odontopharyngidae: *Odontopharynx*
 3 bacterial feeding
 5 predatory
- Odontorhabditidae: *Odontorhabditis*, *Cheilorhabditis*
 3 bacterial feeding
- Osstellidae: *Osstella*
 3 bacterial feeding
- Panagrolaimidae: *Panagrolaimus*, *Panagrelus*, *Turbatrix*
 3 bacterial feeding (many are insect associates)
- Pterygorhabditidae: *Pterygorhabditis*
 3 bacterial feeding
- Rhabditidae: *Rhabditis*, *Mesorhabditis*, *Pelodera*
 3 bacterial feeding (many are associated with earthworms, arthropods, or vertebrates)
- Steinernematidae: *Steinernema* (= *Neoplectana*), *Heterorhabditis*
 7 entomogenous
 3 bacterial feeding in free-living stages
- Teratocephalidae: *Teratocephalus*, *Euteratocephalus*
 3 bacterial feeding
- Order Diplogasterida*
- The basic habit of these is apparently type 3 (bacterial feeding).
- Diplogasteridae: *Diplogaster*, *Butlerius*
 These may use insects for dispersal.

- 3 bacterial feeding
- 5 predacious

Diplogasteroididae: *Diplogasteroides*, *Gof-fartia*

- 3 bacterial feeding

Neodiplogasteridae: *Neodiplogaster*, *Diplen-teron*, *Fictor*, *Pristionchus*

These may use insects for dispersal.

- 3 bacterial feeding
- 4 substrate ingestion
- 5 predacious
- 6 in *Pareudiplogaster*

Tylopharyngidae: *Tylopharynx*

- 3 bacterial feeding

Order Tylenchida

Suborder Tylenchina:

Allantonematidae: *Allantonema*, *Howard-ula*, *Sulphuretylenchus*

- 7 insect parasites
- ? alternate generation possibly feeds in soil

Anguinidae: *Anguina*, *Ditylenchus*, *Nothoty-lenchus*

- 1 parasites of aerial parts of plants
- 2 fungal feeders

Atylenchidae: *Atylenchus*, *Eutylenchus*

- (1) root feeding

Caloosiidae: *Caloosia*

- 1 ectoparasites of roots

Chondronematidae: *Chondronema*

- 7 juveniles parasitic in insects
- ? adults free-living, possible nonfeeding

Criconematidae: *Criconema*, *Ogma*, *Criconemella*, *Hemicriconemoides*

- 1 ectoparasites of roots; males often without stylets

Dolichodoridae: *Dolichodorus*, *Tylenchorhyn-chus*, *Merlinius*, *Belonolaimus*

- 1 migratory ectoparasites of roots

Ecphyadophoridae: *Ecphyadophora*, *Te-nunemellus*

- (1) root feeding

Hemicycliophoridae: *Hemicycliophora*

- 1 ectoparasites of roots; males often without stylet

Heteroderidae: *Heterodera*, *Globodera*, *Cry-phodera*

- 1 obligate endoparasites; female cuticle produces resistant cyst

Hoplolaimidae: *Hoplolaimus*, *Rotylenchus*, *Helicotylenchus*

- 1 ectoparasitic or semi-endoparasitic on roots

Iotonchiidae: *Iotonchium*

- 2 hyphal feeding in some stages
- 7 insect parasites

Meloidogynidae: *Meloidogyne*

- 1 obligate endoparasites; female saccate

Nacobbidae: *Nacobbus*

- 1 gall-forming root endoparasites; sac-cate females

Neotylenchidae: *Fergusobia*, *Deladenus*

- 7 insect parasites with alternate genera-tion being
- 2 hyphal-feeding or
- 1 plant feeding

Paratylenchidae: *Paratylenchus*, *Gracilacus*

- 1 ectoparasites of roots; male often without stylet

Paurodontidae: *Paurodontus*

- 2 hyphal-feeding generation and
- 7 probable insect-parasitic generation

Pratylenchidae: *Pratylenchus*, *Radopholus*

- 1 migratory endoparasites of roots

Psilenchidae: *Psilenchus*

- 1 associates of lower plants (other than fungi); feed on root hairs and epider-mal cells

Rotylenchulidae: *Rotylenchulus*

- 1 ectoparasitic on roots of higher plants; female saccate

Sphaeronematidae: *Sphaeronema*, *Meloido-derita*

- 1 females sedentary ecto- or endopara-sites of roots

Sphaerularidae: *Sphaerularia*, *Tripilus*

- 2 hyphal-feeding generation may occur
- 7 females parasitic in insect haemocoel

Sychnotylenchidae: *Sychnotylenchus*, *Neodi-tylenchus*

- 7 associates of bark beetles

Tylenchidae: *Tylenchus*, *Filenchus*, *Bo-leodorus*

- 1 associates of algae, mosses, lichens, and plant roots

Tylenchulidae: *Tylenchulus*, *Trophonema*

- 1 ectoparasites of roots; female saccate but usually outside root

Tylodoridae: *Tylodorus*, *Cephalenchus*

- 1 root feeding

*Suborder Aphelenchina:*Aphelenchidae: *Aphelenchus*

- 2 hyphal feeding; sometimes feed on fungi in diseased plants

Aphelenchoididae: *Aphelenchoides*,*Bursaphelenchus*, *Anomyctus*,*Rhadinaphelenchus*

- 1 plant feeding in a minority of forms (*Aphelenchoides* may feed on fungi but also on aerial parts of plants including ferns)

- 2 hyphal feeding in terrestrial species; may be insect associated

- 5 predacious (*Laimaphelenchus*)

- 6 algal feeding (*Laimaphelenchus*)

Paraphelenchidae: *Paraphelenchus*

- 2 hyphal feeders

Seinuridae: *Seimura*

- 5 predacious

*Order Enoplida*Alaimidae: *Alaimus*, *Amphidelus*

- 3 bacterial feeding

Cryptonchidae: *Cryptonchus*, *Aulolaimus*

- (3) bacterial feeding (*Aulolaimus*)

Ironidae: *Ironus*

- 5 predacious

Oncholaimidae: *Bastiania*

- (3) bacterial feeding

Onchulidae: *Onchulus*, *Stenonchulus*

- (3) bacterial feeding

- (5) predacious

- (8) omnivorous

Prismatolaimidae: *Prismatolaimus*

- 3 bacterial feeding

Tobrilidae: *Tobrilus*

- (5) predacious

- (6) algal feeding

Tripylidae: *Tripyla*, *Trischistoma*

- (5) predacious

Order Mononchida

Most common forms are type 5 (predacious) but some are type 3 (bacterial feeding).

Anatonchidae: *Anatonchus*

- 5 predacious; prey may be ingested whole

Bathyodontidae: *Bathyodontus*, *Mirolaimus*

- (3) bacterial feeding

Mononchidae: *Mononchus*

- 5 predacious; prey not ingested whole

- (3) may also feed on bacteria

Mononchulidae: *Mononchulus*, *Oionchus*

- (3) bacterial feeding

*Order Dorylaimida*Actinolaimidae: *Actinolaimus*, *Brittonema*, *Actinca*

- 5 predacious

- (8) omnivorous

Aporcelaimidae: *Aporcelaimus*, *Paraxonchium*, *Sectonema*

- 5 predacious

- (8) omnivorous

Belonidiridae: *Axonchium*, *Dorylaimellus*, *Oxydirus*

- (1) plant feeding

Diphtherophoridae: *Diphtherophora*, *Tylo-laimophorus*

- (2) hyphal feeding

Discolaimidae: *Discolaimus*, *Discolaimium*

- (5) predacious

Dorylaimidae: *Dorylaimus*, *Laimydorus*, *Mesodorylaimus*, *Prodorylaimus*, *Thornenema*, *Thornus*

- (8) omnivorous

- 5 predacious (*Labronema*)

Leptonchidae: *Leptonchus*, *Tylencholaimellus*, *Doryllium*

- (2) hyphal feeding

Longidoridae: *Longidorus*, *Xiphinema*

- 1 plant feeding; may transmit plant viruses

Nordiidae: *Eudorylaimus*

- (8) omnivorous (possibly all genera are plant feeding)

- (1) plant feeding

Nygolaimidae: *Nygolaimus*

- 5 predacious, especially on enchytraeids

Trichodoridae: *Trichodorus*, *Paratrichodorus*

- 1 plant feeding; may transmit plant viruses

*Order Isolaimida*Isolaimidae: *Isolaimium*

- (3) bacterial feeding

Order *Stichosomida*Mermithidae: *Mermis*, *Agamomermis*

- 7 parasites of arthropods and other invertebrates; postparasitic stages may occur in soil

LITERATURE CITED

- Anderson, R. V. 1964. Feeding of *Ditylenchus destructor*. *Phytopathology* 4:1121-1126.
- Andrássy, I. 1976. Evolution as a basis for the systematization of nematodes. London: Pitman.
- Andrássy, I. 1984. Klasse Nematoda (Ordnungen Monhysterida, Desmoscolecida, Araeolaimida, Chromadorida, Rhabditida). Stuttgart: Fischer.
- Arpin, P. 1969. Étude préliminaire d'un facteur écologique important pour les Nématodes du sol: L'humidité actuelle du sol. *Revue d'Écologie et du Biologie du Sol* 4:429-435.
- Arpin, P. 1976. Étude et discussion sur un milieu de culture pour Mononchidae (Nematoda). *Revue d'Écologie et du Biologie du Sol* 13:629-634.
- Arpin, P., and G. Kilbertus. 1981. Ultrastructure du contenu digestif et de l'épithélium intestinal chez quelques nématodes prédateurs (Mononchida) et bactériophages. *Revue de Nématologie* 4:131-143.
- Babatola, J. O., and J. Bridge. 1980. Feeding behavior and histopathology of *Hirschmanniella oryzae*, *H. imamuri* and *H. spinicaudata* on rice. *Journal of Nematology* 12:48-53.
- Baldwin, J. G., and M. Mundo-Ocampo. 1991. Heteroderinae, cyst- and non-cyst-forming nematodes. Pp. 275-362 in W. R. Nickle, ed. *Manual of agricultural nematology*. New York: Marcel Dekker.
- Bedding, R. A. 1973. Biology of *Deladenus siricidicola* (Neotylenchidae) an entomophagous-mycetophagous nematode parasitic in siricid woodwasps. *Nematologica* 18:482-493.
- Bilgrami, A. L., I. Ahmad, and M. S. Jairajpuri. 1984. Predatory behaviour of *Aquatides thornei* (Nygolaimina: Nematoda). *Nematologica* 30:457-462.
- Boag, B., I. E. Raschke, and D. J. F. Brown. 1977. Observations on the life cycle and pathogenicity of *Paralongidorus maximus* in a forest nursery in Scotland. *Annals of Applied Biology* 85:389-397.
- Bongers, T. 1988. De nematoden van Nederland. Utrecht: Stichting Uitgeverij Koninklijke Nederlandse Natuurhistorische Vereniging.
- Bouwman, L. A. 1983. Systematics, ecology and feeding biology of estuarine nematodes. BOEDE, Publ en Versl. 3.
- Bridge, J., and N. G. M. Hague. 1974. The feeding behavior of *Tylenchorhynchus* and *Merlinius* species and their effect on growth of perennial ryegrass. *Nematologica* 20:119-130.
- Christie, J. R., A. N. Brooks, and V. G. Perry. 1952. The sting nematode, *Belonolaimus gracilis*, a parasite of major importance on strawberries, celery, and sweet corn in Florida. *Phytopathology* 42:173-176.
- Christie, J. R., and B. G. Chitwood. 1931. *Chondronema passali* (Leidy, 1852) n.g. (Nematoda), with notes on its life history. *Journal of the Washington Academy of Sciences* 21:356-364.
- Christie, J. R., and L. Crossman. 1936. Notes on the strawberry strains of the bud and leaf nematode, *Aphelenchoides fragariae*, I. *Proceedings of the Helminthological Society of Washington* 3:69-72.
- Cobb, N. A. 1916. Notes on new genera and species of nematodes. 4. Subdivisions of *Mononchus*. *Journal of Parasitology* 2:195-196.
- Cordero, D. A., J. G. Baldwin, and M. Mundo-Ocampo. 1991. Fine structure of the posterior cone of females of *Cactodera cacti* Filip'ev & Schuurmans Stekhoven (Nemata: Heteroderinae). *Revue de Nématologie* 14:455-465.
- Croll, N. A., J. M. Smith, and B. M. Zuckerman. 1977. The aging process of the nematode *Caenorhabditis elegans* in bacterial and axenic culture. *Experimental Aging Research* 3:175-189.
- Cryan, W. S., E. Hansen, M. Martin, F. W. Sayre, and E. A. Yarwood. 1963. Axenic cultivation of the dioecious nematode *Panagrellus redivivus*. *Nematologica* 9:313-319.
- Davaine, C. 1857. Recherches sur l'anguille du ble nielle considérée au point de vue de l'histoire naturelle et de l'agriculture. *Comptes Rendus des Séances de la Société de Biologie et de ses Filiales* 3: 201-271.
- Decker, H. 1962. Zur biologie und ökologie von *Aphelenchus avenae* Bastian. *Nematologica* 7:9. (Abstr.).
- Doncaster, C. C., and M. K. Seymour. 1975. Passive ingestion in a plant nematode, *Hexatylus viviparus* (Neotylenchidae: Tylenchida). *Nematologica* 20:297-307.
- Dougherty, E. C., and G. H. Calhoun. 1948. Experiences in culturing *Rhabditis pellio* (Schneider, 1866) Butschli, 1873 and related soil nematodes. *Proceedings of the Helminthological Society of Washington* 15:55-67.
- Dropkin, V. H. 1989. *Introduction to plant nematology*. New York: Wiley.
- Eisenback, J. D., and H. H. Triantaphyllou, Eds. 1991. *Root-knot nematodes: Meloidogyne species and races*. New York: Marcel Dekker.
- Esser, R. P. 1963. Nematode interactions in plates of non-sterile water agar. *Proceedings of the Soil and Crop Science Society of Florida* 23:121-138.
- Ferris, V. R. 1967. Life history studies of certain Dorylaimid nematodes of the Wabash River Basin. *Nematologica* 13:142-143.
- Ferris, V. R. 1968. Biometric analyses in the genus *Labronema* (Nematoda: Dorylaimida) with a description of *L. thornei* n. sp. *Nematologica* 14:276-284.
- Ferris, V. R., and J. M. Ferris. 1989. Why ecologists need systematics: Importance of systematics to ecological research. *Journal of Nematology* 21:308-314.
- Franklin, M. T., and D. J. Hooper. 1962. *Bursaphelenchus fungivorus* n. sp. (Nematoda: Aphelenchoidea) from rotting gardenia buds infected with *Botrytis cinerea* Pers. ex. Fr. *Nematologica* 8:136-142.
- Galtsova, V. V., and O. N. Pavljuk. 1990. The freeliving marine nematodes on the industrial plan-

- tations of pectens. P. 5, Abstracts, 7th International symposium on aquatic nematodes, 8–10 August, 1990, Yerseke, The Netherlands. (Abstr.).
34. Germani, G., and M. Luc. 1982. Études sur la "chlorose voltaïque" des légumineuses due au nematode *Aphasmatylenchus straturatus* Germani. II. Revue de Nématologie 5:195–199.
35. Goodey, T. 1963. Soil and freshwater nematodes, 2nd ed., revised by J. B. Goodey. London: Methuen.
36. Goodey, T. B. 1943. A note on the feeding of the nematode *Anguillulina macura*. Journal of Helminthology 21:17–19.
37. Gowen, S. R. 1970. Observations on the fecundity and longevity of *Tylenchus emarginatus* on sitka spruce seedlings at different temperatures. Nematologica 16:267–272.
38. Grootaert, P., and D. Maertens. 1976. Cultivation and life cycle of *Mononchus aquaticus*. Nematologica 22:173–181.
39. Gupta, M. C., R. S. Singh, and K. Sitaramaiah. 1979. A new endoparasitic fungus on *Xiphinema* and cultivation of *Rhabditis* spp. and *Aphelenchus avenae* on same fungus. Nematologica 25:142.
40. Hechler, H. C. 1962. The description, feeding habits and life history of *Neotylenchus linfordi* n. sp.; a mycophagous nematode. Proceedings of the Helminthological Society of Washington 29:19–27.
41. Hechler, H. C. 1963. Description, developmental biology and feeding habits of *Seinura tenuicaudata* (de Man) J. B. Goodey, 1960 (Nematoda: Aphelenchoididae), a nematode predator. Proceedings of the Helminthological Society of Washington 30:183–195.
42. Hechler, H. C. 1968. Postembryonic development and reproduction in *Diploscapter coronata* (Nematoda: Rhabditidae). Proceedings of the Helminthological Society of Washington 35:24–30.
43. Hechler, H. C., and D. P. Taylor. 1966. The life histories of *Seinura celeris*, *S. oliveirae*, *S. oxura* and *S. steineri* (Nematoda: Aphelenchoididae). Proceedings of the Helminthological Society of Washington 33:71–83.
44. Hollis, J. P. 1957. Cultural studies with *Dorylaimus ettersbergensis*. Phytopathology 47:468–473.
45. Hooper, D. J., and J. A. Cowland. 1986. Fungal hosts for the chrysanthemum nematode *Aphelenchoides ritzemabosi*. Plant Pathology 35:128–129.
46. Hunt, D. J. 1977. Bionomics of *Paractinolaimus vigor* Thorne, 1967 (Dorylaimida: Paractinolaimidae) with a description of *P. dominicus* n. sp. and *Nygolaimum haguei* n. sp. (Dorylaimida: Aporcelaimidae). Nematologica 23:452–462.
47. Hunt, D. J. 1977. Observations on the feeding of *Ironus longicaudatus* (Enoplida: Ironidae). Nematologica 23:478–479.
48. Jairajupuri, M. S., and M. I. Azmi. 1978. Some studies on the predatory behavior of *Mylonchulus dentatus*. Nematologia Mediterranea 6:205–212.
49. Jensen, P. 1981. Phyto-chemical sensitivity and swimming behavior of the free-living marine nematode *Chromadorita tenuis*. Marine Ecology–Progress Series 4:203–206.
50. Jensen, P. 1987. Feeding ecology of free-living aquatic nematodes. Marine Ecology–Progress Series 35:187–196.
51. Jones, R. K. 1978. The feeding behavior of *Helicotylenchus* spp. on wheat roots. Nematologica 24:88–94.
52. Kareiva, P., and R. Sahakian. 1990. Tritrophic effects of a simple architectural mutation in pea plants. Nature 345:433–434.
53. Kühn, J. 1881. Ueber das Vorkommen von Ruebennematoden an den Wurzeln der Halmfruchte. Neue Zeitschrift für Ruebenzucker-Industrie 24:149–153.
54. Lamberti, F., C. E. Taylor, and J. W. Seinhorst, Eds. 1975. Nematode vectors of plant viruses. New York: Plenum Press.
55. Linford, M. B. 1937. Notes on the feeding of *Ditylenchus dipsaci*. (Nematoda: Tylenchidae). Proceedings of the Helminthological Society of Washington 4:46–47.
56. Linford, M. B., and J. M. Oliveira. 1937. The feeding of hollow spear nematodes on other nematodes. Science 85:295–297.
57. Linford, M. B., and J. M. Oliveira. 1940. *Rotylenchulus reniformis* n. gen., n. sp., a nematode parasite of roots. Proceedings of the Helminthological Society of Washington 7:35–42.
58. Lörenzen, S. 1981. Entwurf eines phylogenetischen Systems der freilebenden Nematoden. Veröffentlichungen des Instituts für Meeresforschung in Bremerhaven, Supplement 7:472.
59. Lownsbery, B. F. 1959. Studies of the nematode, *Criconemoides xenoplax*, on peach. Plant Disease Reporter 43:913–917.
60. Luc, M., A. R. Maggenti, and R. Fortuner. 1988. A reappraisal of *Tylenchina* (Nemata). 9. The family Heteropridae Filip'ev & Schuurmans Stekhoven, 1941. Revue de Nématologie 11:159–176.
61. Maertens, D. 1975. Observations on the life cycle of *Prionchulus punctatus* (Cobb, 1917) and some culture conditions. Biologisch Jaarboek Dodona 43:197–218.
62. Maggenti, A. 1981. General nematology. New York: Springer-Verlag.
63. Mamiya, Y. 1972. Pine wood nematode, *Bursaphelenchus lignicolus* Mamiya and Kiyohara, as a causal agent of pine wilting disease. Review of Plant Protection Research 5:46–60.
64. Mankau, R. 1980. Biological control of nematode pests by natural enemies. Annual Review of Phytopathology 18:415–440.
65. Mankau, R., and S. K. Mankau. 1963. The role of mycophagous nematodes in the soil. I. The relationships of *Aphelenchus avenae* to phytopathogenic soil fungi. Pp. 271–280 in J. Doeksen and J. van der Drift, eds. Soil organisms. Amsterdam: North Holland.
66. Meagher, J. W. 1968. *Acontylus vipriensis* n. g., n. sp. (Nematoda: Hoplolaimidae) parasitic on *Eucalyptus* sp. in Australia. Nematologica 14:94–100.
67. Mohandas, C., and N. R. Prabhoo. 1980. The feeding behavior and food preferences of predatory nematodes (Mononchida) from the soils of Kerala (India). Revue d'Écologie et du Biologie du Sol 17: 53–60.

68. Myers, R. F. 1967. Axenic cultivation of *Aphelenchoides sacchari* Hooper. Proceedings of the Helminthological Society of Washington 34:251-255.
69. Nemes, A. J. 1974. Evaluation of the feeding behavior of *Prionchulus punctatus* (Cobb), a nematode predator. Journal of Animal Ecology 43:553-565.
70. Nicholas, W. L. 1956. The axenic culture of *Turbatrix aceti* (the vinegar eelworm). Nematologica 1:337-340.
71. Nicholas, W. L. 1962. A study of a species of *Acrobeloides* (Cephalobidae) in laboratory culture. Nematologica 8:99-109.
72. Nicholas, W. L. 1984. The biology of free-living nematodes, 2nd. ed. Oxford: Clarendon Press.
73. Nicholas, W. L., E. C. Dougherty, and E. L. Hansen. 1959. Axenic cultivation of *Caenorhabditis briggsae* (Nematoda: Rhabditidae) with chemically undefined supplements; comparative studies with other nematodes. Annals of the New York Academy of Sciences 77:218-236.
74. Nicholas, W. L., A. C. Stewart, and T. G. Marples. 1988. Field and laboratory studies of *Desmodora cazca* Gerlach, 1956 (Desmodoridae: Nematoda) from mangrove mud-flats. Nematologica 34:331-349.
75. Nickle, W. R., Ed. 1984. Plant and insect nematodes. New York: Marcel Dekker.
76. Nickle, W. R., Ed. 1991. Manual of agricultural nematology. New York: Marcel Dekker.
77. Nielsen, C. O. 1949. Studies on the soil microfauna. II. The soil inhabiting nematodes. Natura Jutlandica 2:1-131.
78. Nigon, V., and E. C. Dougherty. 1949. Reproductive patterns and attempts in reciprocal crossing of *Rhabditis elegans* Maupas, 1900 and *Rhabditis briggsae* Dougherty & Nigon, 1949. Journal of Experimental Zoology 112:485-503.
79. Nuß, B. 1985. Ultrastrukturuntersuchungen zur Nahrungsabsorption von aquatischen Nematoden. Veröffentlichungen des Instituts Meeresforschung in Bremerhaven 21:1-69.
80. Overhoff, A., D. W. Freckman, and R. A. Virginia. 1993. Life cycle of the microbivorous Antarctic Dry Valley nematode *Scottinema lindsayae* (Timm, 1971). Polar Biology 13:151-156.
81. Palmisano, A. M., and T. Turchetti. 1976. Sviluppo delle Popolazioni di una Specie di *Cephalobus* (Nematoda: Cephalobidae) in Associazione con Batteri Diversi. Redia 59:155-169.
82. Paracer, S. M., M. Waseem, and B. M. Zuckerman. 1967. The biology and pathogenicity of the awl nematode, *Dolichodorus heterocephalus*. Nematologica 13:517-524.
83. Paracer, S. M., and B. M. Zuckerman. 1967. Monoxenic culturing of *Dolichodorus heterocephalus* on corn root callus. Nematologica 13:478-479.
84. Paramonov, A. A. 1962. Plant-parasitic nematodes, I. Origin of nematodes: Ecological and morphological characteristics of plant nematodes: Principles of taxonomy. Moscow: Izdatel'stvo Akedemii Nauk SSR.
85. Perry, V. G. 1953. The awl nematode, *Dolichodorus heterocephalus*, a devastating plant parasite. Proceedings of the Helminthological Society of Washington 20:21-27.
86. Petersen, H., and M. Luxton. 1982. A comparative analysis of soil fauna populations and their roles in decomposition processes. Oikos 39:287-388.
87. Petersen, H., R. V. O'Neil, and R. H. Gardner. 1985. Use of an ecosystem model for testing ecosystem response to inaccuracies of root and microflora productivity estimates. British Ecological Society Special Publication 4:233-242.
88. Pillai, J. K., and D. P. Taylor. 1968. Biology of *Paroigolaimella bernensis* and *Fictor anchicoprophaga* (Diplogasterinae) in laboratory culture. Nematologica 14:159-170.
89. Pillai, J. K., and D. P. Taylor. 1968. *Butlerius micans* n. sp. (Nematoda: Diplogasterinae) from Illinois, with observations on its feeding habits and a key to the species of *Butlerius* Goodey, 1929. Nematologica 14:89-93.
90. Poinar, G. O., Jr. 1976. Description and biology of a new insect parasitic rhabditoid *Heterorhabditis bacteriophora* n. gen., n. sp. (Rhabditida; Heterorhabditidae n. fam.). Nematologica 21:463-470.
91. Poinar, G. O., Jr., and P. A. van der Laan. 1972. Morphology and life history of *Sphaerularia bombi*. Nematologica 18:239-252.
92. Procter, D. L. C. 1986. Fecundity, reproductive effort, age-specific reproductive tactics and intrinsic rate of natural increase of a high Arctic nematode belonging to the genus *Chilopacus*. Holarctic Ecology 9:104-108.
93. Raski, D. J., and M. Luc. 1987. A reappraisal of Tylenchina (Nemata). 10. The superfamily Criconematoidea Taylor, 1936. Revue de Nématologie 10:409-444.
94. Rhoades, H. L., and M. B. Linford. 1961. A study of the parasitic habit of *Paratylenchus projectus* and *P. dianthus*. Proceedings of the Helminthological Society of Washington 28:185-190.
95. Russell, C. C. 1986. The feeding habits of a species of *Mesodorylaimus*. Journal of Nematology 18:641. (Abstr.).
96. Sasser, J. N., and C. C. Carter, Eds. 1985. An advanced treatise on *Meloidogyne*, vol. 1. Biology and control. Raleigh: North Carolina State University Graphics.
97. Schiemer, F. 1982. Food dependence and energetics of free-living nematodes. I. Respiration, growth and reproduction of *Caenorhabditis elegans* (Nematoda) at different levels of food supply. Oecologia 54:108-121.
98. Shmal'co, V. F. 1959. The cactus nematode, *Heterodera cacti* Filipjev et Schuurmans-Stekhoven, 1941. [in Russian]. Trudy Gel'mintologicheskoi Laboratorii Akademiï Nauk SSSR 9:389-390.
99. Siddiqi, M. R. 1986. Tylenchida: Parasites of plants and insects. Slough, UK: Commonwealth Institute of Parasitology.
100. Small, R. W. 1987. A review of the prey of predatory soil nematodes. Pedobiologia 30:179-206.
101. Small, R. W., and P. Grootaert. 1983. Observations on the predation abilities of some soil dwelling predatory nematodes. Nematologica 29:109-118.
102. Sohlenius, B. 1968. Influence of microorganisms and temperature upon some rhabditid nematodes. Pedobiologia 8:137-145.

103. Sohlenius, B. 1968. Studies of the interactions between *Mesodiplogaster* sp. and other rhabditid nematodes and a protozoan. *Pedobiologia* 8:340-344.
104. Sohlenius, B. 1973. Growth and reproduction of a nematode *Acrobeloides* sp. cultivated on agar. *Oikos* 24:64-72.
105. Sohlenius, B., H. Persson, and C. Magnusson. 1977. Distribution of root and soil nematodes in a young Scots pine stand in central Sweden. *Ecological Bulletin (Stockholm)* 25:340-347.
106. Steiner, G., and H. Heinly. 1922. The possibility of control of *Heterodera radicicola* and other plant injurious nematodes by means of predatory nemas, especially *Mononchus papillatus*. *Journal of the Washington Academy of Sciences* 12:367-386.
107. Sudhaus, W. 1976. Vergleichende Untersuchungen zur Phylogenie, Systematik, Ökologie, Biologie und Ethologie der Rhabditidae (Nematoda). *Zoologica, Stuttgart* 43:1-229.
108. Sudhaus, W. 1980. Systematisch-phylogenetische und biologisch-ökologische Untersuchungen an *Rhabditis*-(*Poikilolaimus*-) Arten als Beitrag zur Rassenbildung und Parallelevolution bei Nematoden. *Zoologische Jahrbücher (Systematik)* 107:287-343.
109. Sutherland, J. R., and R. Keable. 1966. The nematode *Tylenchus (Cephalenchus) hexalineatus* (Geraert, 1962) Geraert and Goodey, 1964 found in three Quebec forest nurseries. *Progress Report of Canadian Department of Forestry* 22:2-3.
110. Taylor, D. P., and J. K. Pillai. 1967. *Paraphelenchus acontioides* n. sp. (Nematoda: Paraphelenchidae), a mycophagous nematode from Illinois, with observations on its feeding habits and a key to the species of *Paraphelenchus*. *Proceedings of the Helminthological Society of Washington* 34:51-59.
111. Thomas, P. R. 1965. Biology of *Acrobelus complexus* Thorne, cultivated on agar. *Nematologica* 11:395-408.
112. Thorne, G. 1943. *Cacopaurus pestis*, nov. gen. nov. spec. (Nematoda: Criconematinae), a destructive pest of the walnut, *Juglans regia* Linn. *Proceedings of the Helminthological Society of Washington* 10:78-83.
113. Thorne, G. 1961. *Principles of nematology*. New York: McGraw-Hill.
114. Thorne, G., and H. H. Swanger. 1936. A monograph of the nematode genera *Dorylaimus*, *Aporcelaimus* n. g., *Dorylaimoides* n. g. and *Pungentus*. n. g. *Capita Zoologia* 6:1-223.
115. Tietjen, J. H. 1967. Observations on the ecology of the marine nematode *Monhystera filicaudata* Allgen, 1929. *Transactions of the American Microscopical Society* 86:304-306.
116. Tietjen, J. H., and J. J. Lee. 1977. Life histories of marine nematodes. Influence of temperature and salinity on the reproductive potential of *Chromadorina germanica*. *Mikrofauna Meeresboden* 61:263-270.
117. Tjepkema, J. P., V. R. Ferris, and J. M. Ferris. 1971. Review of the genus *Aporcelaimellus* Heyns, 1965 and six species groups of the genus *Eudorylaimus* Andrassy, 1959 (Nematoda: Dorylaimida). *Purdue University Research Bulletin* 882:1-52.
118. Vanfleteren, J. R. 1978. Axenic culture of free-living, plant-parasitic, and insect-parasitic nematodes. *Annual Review of Phytopathology* 16:131-157.
119. Wachek, F. 1955. Die entoparasitischen Tylenchiden. *Parasitologische Schriftenreihe* 3:1-119.
120. Warwick, R. M., and R. Price. 1979. Ecological and metabolic studies on free-living nematodes from an estuarine mud-flat. *Estuarine Coastal and Marine Science* 9:251-272.
121. Wasilewska, L. 1971. Klasyfikacja troficzna nicieni glebowych i roslinnych. *Wiadomosci Ekologiczne* 17:379-388.
122. Wood, F. H. 1973. Nematode feeding relationships. Feeding relationships of soil-dwelling nematodes. *Soil Biology and Biochemistry* 5:528-537.
123. Wood, F. H. 1974. Biology of *Seimura demani* (Nematoda: Aphelenchoididae). *Nematologica* 20:347-353.
124. Woombs, M., and J. Laybourn-Parry. 1984. Feeding biology of *Diplogasteritus nudicapitatus* and *Rhabditis curvicaudata* (Nematoda) related to food concentration and temperature, in sewage treatment ponds. *Oecologia* 64:163-167.
125. Yeates, G. W. 1969. Predation by *Mononchoidea potohikus* (Nematoda, Diplogasteridae) in laboratory culture. *Nematologica* 15:1-9.
126. Yeates, G. W. 1970. Studies on laboratory cultures of dune sand nematodes. *Journal of Natural History* 4:119-136.
127. Yeates, G. W. 1971. Feeding types and feeding groups in plant and soil nematodes. *Pedobiologia* 11:173-179.
128. Yeates, G. W. 1973. Morphometrics and growth in eight New Zealand soil nematode populations. *New Zealand Journal of Science* 16:711-725.
129. Yeates, G. W. 1974. Studies on a climosequence of soils in tussock-grasslands. 2. Nematodes. *New Zealand Journal of Zoology* 1:171-177.
130. Yeates, G. W. 1987. Significance of developmental stages in the co-existence of three species of Mononchoidea (Nematoda) in a pasture soil. *Biology and Fertility of Soils* 5:225-229.
131. Yeates, G. W., R. N. Watson, and K. W. Steele. 1985. Complementary distribution of *Meloidogyne*, *Heterodera* and *Pratylenchus* (Nematoda: Tylenchida) in roots of white clover. Pp. 71-79, *Proceedings of the 4th Australasian Conference on Grassland Invertebrate Ecology*.
132. Zuckerman, B. M., W. F. Mai, and R. A. Rohde, Eds. 1971. *Plant parasitic nematodes*, vol. 2. Cytogenetics, host-parasite interaction, and physiology. New York: Academic Press.