



Felsic magmatism and uranium deposits

Michel CUNEY

**UNIVERSITE DE LORRAINE – GEORESSOURCES CREGU – CNRS
54 506, Vandoeuvre les NANCY FRANCE**

METAL FRACTIONATION FROM ULTRABASIC ROCKS TO GRANITES

ppm	Ultrabasic	Basic	Intermediate	Granites	X
U	0.021	0.75	2.4	3.3	x 160

Th	0.060	3.5	7.8	17.5
----	-------	-----	-----	------

Earth average Th/U = 4


INCOMPATIBLE BEHAVIOR

WHAT IS AN INCOMPATIBLE ELEMENT ?


OLIVINE >



MANTLE = SILICATES



Mg²⁺ low charge
small ionic radius] ≡ Ni²⁺ = 0.69



Th⁴⁺ high charge
large ionic radius

INCOMPATIBLE with the silicate network

U INCOMPATIBLE BEHAVIOUR


- several major geochemical, geophysical and metallogenic consequences:
- (i) U continuously transferred from the mantle to the Earth crust, & within the continental crust towards its upper part together with Th, K, ...
 - (ii) radiogenic heat production is maximized in the upper crust → radiogenic heat flux production may delineate radioelement enriched crustal blocks,
 - (iii) the most felsic melts tend to be the most enriched in U,
 - (iv) granites & rhyolites = primary U sources for the formation of most U deposits

Despite the strongly incompatible behavior of U, deposits dominantly resulting from magmatic processes are rare.

Average granite (U= 3-4 ppm), U mainly in zircon, apatite, monazite, titanite, ... from which U cannot be leached by most geological fluids.

Some specific granites have higher U contents permitting crystallization of other accessory minerals from which U can be more or less easily leached for the formation of U deposits → “fertile granites” of Moreau [1966]

U continuously transferred from the mantle to the Earth crust



Fondamental fractionation processes in magmas

Partial melting

Fractional crystallisation

Sediment subduction and mantle metasomatism

Mixing with crustal material

Melt/fluid fractionation



Magma aluminous indices to classify magmatic rocks

$Al/(Na+K+2Ca) = A/CNK$ in cations
= ASI Aluminium Saturation Index


$Al/(Na+K)$ ou $(Na+K) Al = AGPAICITY$

why ?

= INDEX OF MAGMA POLYMERISATION

U-rich magma classification using aluminous indices

some specific granites have higher U contents




$Al/(Na+K) = 1$ &
 $Al/(Na+K+2Ca) = 1$
when
Al-Na-K-Ca in feldspars only

$Al/(Na+K+2Ca) > 1$
→ **peraluminous**


$Al/(Na+K+2Ca) < 1$
& $Al/(Na+K) < 1$
→ **peralkaline**

$Al/(Na+K+2Ca) > 1$
& $Al/(Na+K) > 1$
→ **calc-alkaline**


**Why using aluminous indices
for
magma classification ?**



UO₂ SOLUBILITY IN GRANITIC MELTS



MONAZITE SOLUBILITY IN SILICATE MELTS



THREE TYPES OF U – RICH ACIDIC MAGMAS

● PERALKALINE MAGMAS

- $\text{Na} + \text{K} > \text{Al}$
- Riebeckite, Aegyrine, Avfedsonite
- betafite, thorite, complex U,Th,REE,Zr minerals
- **Strongly enriched in U, Th, REE, Zr, ...**

● METALUMINOUS HIGH-K CALC-ALKALINE MAGMAS

- $\text{Al} < \text{Na} + \text{K} + 2\text{Ca}$
- Amphibole, Pyroxene, biotite
- Allanite, U-thorite, titanite, \pm Th-rich uraninite, magnetite
- **Enriched in U, Th, REE**

● PERALUMINOUS FELSIC MAGMAS

- $\text{Al} > \text{Na} + \text{K} + 2\text{Ca}$
- Al-biotite, Muscovite, \pm sill, andalusite, garnet, topaz, tourmaline
- Low-Th uraninite, monazite, ilmenite
- **Enriched in U, poor in Th, REE, Zr ...**

PERALKALINE MAGMAS

$\text{Na} + \text{K} > 1$ + high T \rightarrow highly depolymerized

Very high solubility of accessory minerals

very high U, Th, Zr, REE, Nb, Ta, ... contents
continuously enriched in the residual melts

Volcanic rocks :
U in the glassy matrix

Granites/Syenites : crystalliz. of a complex mineral paragenesis : zircon, U-Th-Zr silicophosphates, Nb-Ta oxydes, ...


Very good U-source

U in refractory sites

Bad U source / high extraction cost

Ex : - Streltsov (Russia)
- McDermitt (USA)

Ex. : - Ilimaussaq, Groenland (syenites)
- Bokan Mountain Alaska (granites)



PERALKALINE MAGMAS

Ilimausacq (Greenland)

peralkaline complexe


U mineralization
in the most fractionated part
where fluid oversaturation occurred

→ simultaneous enrichment in :
U, Th, Zr, REE, Nb, Ta, F ...

U in **steenstrupine** :


Silicophosphate of U, Th, Zr, REE, Nb, Ta

220,000 t U @ 250 ppm




GEOLOGIC MAP OF THE STRELTSOVSKY ORE FIELD

280 000t U in 18 deposits




1. Shironoskoye
2. Streltsovskoye
3. Antei
4. Oktabraskoye
- 5.
6. Martoskoye
7. Malo-Tulukuyev
- 8.
9. Yubilenoye
10. Vesenney
- 11.
12. Pyatletneye
13. KranyKamen
- 14.
15. Zherlovoye
16. Argunskoye
- 17.
18. Dalnee

5 km






U loss evaluation from U-contents of melt inclusions (acidic volcanics compared to altered rhyolites)

STRELTSOV caldera





Conceptual model for U deposits related to peralkaline magmatism







**melt
inclusions**



Melt inclusion geochemistry from sandstone





PERALUMINOUS MAGMAS

Highly polymerized : $A/CNK > 1$ + low T

Accessory minerals low solubility

low Th, Zr, REE,... contents
continuously depleted in the residual melts

Early crystallization of monazite and zircon :
With limited amount of U


U enriched in residual silicate melts (up to some tens of ppm)

**Volcanic rocks: RARE
FERTILE U-SOURCE**

Ex. : - Macusani (Peru)


**Granites: U >> as low-Th URANINITE
▶ FERTILE U-SOURCE**

Ex. : - St Sylvestre (Limousin)
- Erzgebirge (Germany)



GRANITE RELATED DEPOSITS OF THE MID-EUROPEAN VARISCAN BELT

> 300,000 t U @ 0.2 %




- HERCYNIAN GRANITES
- LIMITS PRETRIASSIC BASEMENT
- URANIUM DEPOSITS**
- U SHOWINGS
- U MEDIUM
- U LARGE
- U VERY LARGE

MACUSANI YELLOW CAKE

"KIHITIAN CONCESSION", CHILCUNO-CHICO PROJECT
N-S PROFILE, LOOKING WEST

DRILL HOLES: PT-CH7-TV, PT-CH6-TV, PT-CH5-TV, PT-CH8-TV, PT-CH1-TV, PT-CH10-TV, PT-CH15-TV, PT-CH2-TV, PT-CH18-TV





CUSANI YELLOW CAKE

30,000 t U @ 0.02

HIGH-K CALC-ALKALINE MAGMAS

Intermediate : $A/CNK \sim 1 +$ moderately high T

Accessory minerals intermediate solubility

high U, Th, Zr, REE, Nb, Ta, ...
constant or decreasing in the residual melts

High Ca-
contents

Monazite not stable

REE in Ca-minerals :

- amphibole, allanite, titanite

Volcanic rocks

Variable fertility
According to

glass/accessory mineral ratio

Ex.: Ben Lomond (Australia)

Granites

$Th > U$

Allanite+
U-thorite

BARREN if
not metamict


$U > Th$

\pm uraninite


\pm fertile

Ex.: Hotagen (Sweden)

Th



U



Geologic Map of the Olympic Dam Iron Oxide Cu-Au +U (REE) (IOCG) deposit


2.100 Mt U

2 U-rich sources


high-K calc-alkaline magmas

Roxby Downs Granite		Haematite-matrix-rich granite breccias		Barren quartz - haematite breccias	
Granite and granite breccias		Heterolithic granite and haematite breccias		Volcaniclastics	
Granite-rich breccias		Haematite-rich breccias			

Property Location Map



MICHELIN U DEPOSIT



Associated with **high-K calc-alkaline metavolcanic gneisses** of the Aillik Group
U mineralization associated with hematization + albitization

ANATECTIC PEGMATOIDS

Partial melting of U rich metasediments and/or metavolcanics


Low A/CNK : ~ 1.1 + low T

Accessory minerals intermediate solubility

high U and/or Th and/or Zr and/or REE and/or Nb, Ta, ...
depending of the nature of the source
constant or decreasing in the residual melts


Uraninite \pm Monazite \pm Allanite \pm Uranothorite
 \pm Zircon \pm Nb-Ta minerals \pm ...

Ex.: Rössing (Namibia)



ANATECTIC PEGMATOIDS


Cross-section of the Rossing alaskite body geology & boreholes (drill section zero)





250,000 t U @ 300 ppm


after Berning, 1986

U-Th relationships in the Rossing Alaskites & metasediments



-  Metasediments
-  Alaskites

U - Th FRACTIONATION IN THE 3 TYPES OF U - RICH ACIDIC MAGMAS



Peralkaline

Peraluminous

Metaluminous

CONCLUSIONS

Three types of U-rich felsic magmas identified:

- **Peralkaline (fractionated A1-type)**
 - Weakly fertile granites (U in highly refractory phases)
 - Examples: Ilimausacq (Groenland), Bokan Montain (Alaska)
 - Highly fertile volcanics if high glass/crystal ratio
 - Example: Streltsovka (Russia), Dornot (Mongolia)
- **Metaluminous high-K calc-alkaline (A2-type)**
 - Fertile granites if uraninite-rich or metamict accessories for vein & sedimentary deposits
 - Fertiles volcanics if large glass/crystal ratio
 - Examples : Source for Olympic Dam deposit
- **Peraluminous felsic :** **+ Anatectic pegmatoids**
 - Uraninite bearing granites
 - Very fertile for vein type deposits & sec. dep. in sedimentary basins
 - Example: Variscan belt (Europe), Yenshanian belt (China), ...
 - Peraluminous volcanics :
 - Fertile for vein type deposits, rare
 - Example : Macusani (Peru)