



# Fermatean fuzzy CRITIC-EDAS approach for the selection of sustainable third-party reverse logistics providers using improved generalized score function

Arunodaya Raj Mishra<sup>1</sup> · Pratibha Rani<sup>2</sup> · Kiran Pandey<sup>3</sup>

Received: 7 July 2020 / Accepted: 9 January 2021 / Published online: 9 February 2021  
© The Author(s), under exclusive licence to Springer-Verlag GmbH, DE part of Springer Nature 2021

## Abstract

In today's world, the demand for sustainable third-party reverse logistics providers (S3PRLPs) becomes an increasingly considerable issue for industries seeking improved customer service, cost reduction and sustainability perspectives. However, the assessment and selection of right S3PRLP is a complex uncertain decision-making problem due to involvement of numerous conflicting attributes, imprecise human mind and lack of information. Recently, Fermatean fuzzy set (FFS) has been recognized as one of the suitable tools to tackle the uncertain and inaccurate information. In this paper, we introduce a hybrid methodology based on CRITIC and EDAS methods with Fermatean fuzzy sets (FFSs) to solve the S3PRLP selection problem in which the attributes and decision makers' weights are completely unknown. In this framework, CRITIC approach is applied to calculate the attribute weight and EDAS method is used to evaluate the priority order of S3PRLP options. To do this, a new improved generalized score function (IGSF) is developed with its elegant properties. Also, a formula is discussed to calculate the decision makers' weights based on the developed IGSF. Next, developed framework is applied to assess a case study of S3PRLP selection problem with Fermatean fuzzy information, which elucidates the usefulness and practicality of the proposed method. Finally, comparative study is implemented to show the strength of introduced framework with extant approaches. The outcomes of the work confirm that the introduced approach is more feasible and well-consistent with the other extant approaches.

**Keywords** Fermatean fuzzy sets · Score function · CRITIC · EDAS · Third-party reverse logistics providers

## Abbreviations

3PRLP	Third-party reverse logistics provider	BD	Belongingness degree
AHP	Analytic hierarchy process	BWM	Best–worst method
ANP	Analytic network process	CoCoSo	Combined compromise solution
ARAS	Additive ratio assessment	COPRAS	Complex proportional assessment
AVS	Average solution	CODAS	Combinative distance-based assessment
		CRITIC	Criteria importance through inter-criteria correlation
		DEMATEL	Decision making trial and evaluation laboratory
		DM	Decision maker
		EDAS	Evaluation based on distance from average solution
		ELECTRE	ELimination Et choice translating reality
		EVCSs	Electric vehicle charging stations
		FSs	Fuzzy sets
		FFSs	Fermatean fuzzy sets
		FF-CRITIC-EDAS	Fermatean fuzzy-CRITIC-EDAS
		FF-TOPSIS	Fermatean fuzzy-TOPSIS

✉ Arunodaya Raj Mishra  
arunodaya87@outlook.com

Pratibha Rani  
pratibha138@gmail.com

Kiran Pandey  
kiran.mishra.cse07@iitbhu.ac.in

<sup>1</sup> Department of Mathematics, Government College, Jaitwara, MP, India

<sup>2</sup> Department of Mathematics, NIT, Warangal, Telangana, India

<sup>3</sup> Department of Mathematics, Bioinformatics and Computer Application, MANIT, Bhopal, MP, India

FF-WPM	Fermatean fuzzy-WPM
FFNs	Fermatean fuzzy numbers
HFSs	Hesitant fuzzy sets
HCWD	Health-care waste disposal
IFSs	Intuitionistic fuzzy sets
IGSF	Improved generalized score function
IT2FSs	Interval type-2 fuzzy sets
MCDM	Multi-criteria decision-making
MOORA	Multi-Objective optimization on the basis of ratio analysis
IVPHFSs	Interval-valued pythagorean hesitant fuzzy sets
MOORA-G	Gray Multi-objective optimization by ratio analysis
MULTIMOORA	Multi-objective optimization on the basis of ratio analysis plus full multiplicative form
NDA	Negative distance from average
NBD	Non-belongingness degree
PDA	Positive distance from average
PFSs	Pythagorean fuzzy sets
PULTSs	Probabilistic uncertain linguistic term sets
RL	Reverse logistics
SCM	Supply chain management
SWARA	Step-wise weight assessment ratio analysis
S3PRLP	Sustainable third-party reverse logistics provider
TOPSIS	Technique for order of preference by similarity to ideal solution
VIKOR	VlseKriterijumska Optimizacija I Kaompromisno Resenje
WASPAS	Weighted aggregated sum product assessment
WPM	Weighted product model

## 1 Introduction

During these times, good quality product, satisfaction of customers' requirements and existence in competitive marketplaces are elementary needs for any business. Indeed, these requirements have become business principles, prominent corporations to pursuit for different facets that influence the purchasing options of users. Subsequently, reverse logistics (RL) has become main feature for contributing to the desirable outcomes of several enterprises. RL comprises the actions related with the collection and succeeding retrieval of used products (Fattahi and Govindan 2017). The emerging implication of RL has supervised numerous enterprises to design and reconstruct procedures as a part of their sustainable development initiatives

(Govindan et al. 2015; Banihashemi et al. 2019). Through RL, products are displaced from their final terminus to a new position, where their worth is considered and they are managed to the manufacturing line again or appropriately disposed (Tavana et al. 2016; Kannan et al. 2017). Furthermore, eco-conscious customers incline to give extra on eco-friendly products, increasing the revenue of those businesses that utilize RL to achieve with the needs of such consumers (Mavi et al. 2017; Zarbakhshnia et al. 2018).

RL management is one of the important issues in the SCM, which mostly emphasizes on backward flow of products and raw materials from users to suppliers (Mavi et al. 2017). Growing environmental responsiveness and prospective economic growth have determined ever more corporations to outsource their logistics operations to S3PRLPs (Mavi et al. 2017; Zarbakhshnia et al. 2018; Li et al. 2018). In order to achieve the objectives of cost savings and environmental sustainability, it is significant for the firms to select the best S3PRLPs option. In recent times, the assessment of 3PRLPs selection process has received great attentions from the researchers. Numerous scholarly articles have been presented for selecting the best 3PRLP alternative in the literature, however, more studies are required to manage the prioritizations of different expertise, different environments and knowledge levels on reverse logistics with the consideration of social, environmental and economic dimensions simultaneously. Because of increasing complexity and several constraints, it is not always possible to explore the priorities more proficiently and accurately in the best 3PRLPs selection.

The FSs doctrine (Zadeh 1965) has successfully been employed in diverse 3PRLPs selection problem and proved its powerful ability to tackle with imprecise and uncertain information. As an extension of FSs, the theory of FFSs (Senapati and Yager 2019a) has been proven as one of the powerful platforms to deal with the imprecise and uncertain information. The key characteristic of FFS is the cube addition of BD and NBD is less than or equal to 1. Thus, the FFSs theory is more superior tool than FSs, IFSs and PFSs. Based on its unique advantage, the paper focuses under the environment of FFSs for the assessment of S3PRLP selection. Inspired by the above studies and literature, we introduce Fermatean fuzzy-CRITIC-EDAS framework for assessing the S3PRLPs selection. Thus, this is the first study which proposes a hybrid framework under FFSs. The main contributions of the work are discussed as follows:

1. A New improved generalized score function (IGSF) is proposed for FFNs with their elegant properties.
2. An FF-CRITIC-EDAS framework is introduced to handle the MCDM problem on FFSs.

3. To determine the practicality and effectiveness of the introduced framework, a case study of S3PRLP selection is taken with FFNs.
4. A Comparative discussion is made with the extant models to validate of the developed framework.

The rest of paper is arranged as follows: Sect. 2 depicts a comprehensive review related to present study. Section 3 shows the basic notions on FFSs and proposes a new IGSF with its elegant properties. Section 4 introduces novel Fermatean fuzzy-CRITIC-EDAS approach to elucidate the MCDM problems. Section 5 deliberates a case study of S3PRLP selection and also discusses a comparative discussion with the extant models. Section 6 spectacles the conclusions and scope for future study.

## 2 Related works

In the current section, we present literature survey related to the present study.

### 2.1 3PRLP Selection

Various criteria are involved in the evaluation of 3PRLPs selection process, consequently, this selection process can be observed as a MCDM problem. Existing studies on 3PRLP selection problem confirm the emergent interest of scholars and manufacturers. Over the last few years, copious MCDM models have been established in the setting of 3PRLP assessment problem. Realistic reverse logistics outsourcing assessments are commonly prepared under imprecise and vague environment due to multiple indicators, like as partial ignorance, imprecise estimation, partial or inaccessible decision information (Saen 2010; Azadi and Saen 2011). Consequently, crisp values are usually inappropriate to model such type of practical decision conditions.

FS theory and their extensions have widely been employed to cope with uncertain and vague information occurred in realistic MCDM applications. Senthil et al. (2014) suggested a combined model with AHP and TOPSIS approaches for evaluating an ideal reverse logistics contractor. In a further study by Tajik et al. (2014), a decision-making framework was introduced for choosing most suitable 3PRLP alternative by taking all three aspects of sustainability on FSs. Later, Uygun et al. (2015) planned and selected an outsourcing provider for a telecommunications business by employing DEMATEL and ANP approaches. Tavana et al. (2016) suggested a conceptual analytic network model to thoroughly tackle the complex behavior of interactions among the 3PRLPs assessment factors. Mavi et al. (2017) presented SWARA method for weighting the assessment criteria of 3PRLP in the plastics industry and further,

ranked the sustainable 3PRLP alternatives through MOORA model within FSs context. Tavana et al. (2018) suggested a combined method with the integration of ANP and grey superiority and inferiority methods on IFSs for the assessment of 3PRLPs selection process. Li et al. (2018) used a combined cumulative prospect doctrine with hybrid-information MCDM methodology for the evaluation of 3PRLPs from sustainability perspectives. Zarbakhshnia et al. (2018) weighted the assessment criteria through fuzzy SWARA method and ranked the sustainable 3PRLPs by employing COPRAS method under fuzzy environment. Liu et al. (2019) suggested the BWM to research the selection of 3PRLPs on IVPFSSs. Bai and Sarkis (2019) pioneered multi-stage, multi-method, and multi-metric decision-making tool based on TOPSIS, VIKOR and neighborhood rough set for the evaluation of 3PRLP selection decision. Recently, Zarbakhshnia et al. (2020) established a framework with the AHP and MOORA-G models for the assessment of 3PRLPs under FSs. Zhang and Su (2020) introduced the heterogeneous linguistic model with dominance degree to assess the best S3PRLP for a car manufacture industry.

### 2.2 Fermatean fuzzy sets

The theory of FSs has broadly been received great attention from the researchers for dealing with uncertain and imprecise information. In 1986, Atanassov (1986) presented the idea of IFS, which is termed as belongingness degree (BD), non-belongingness degree (NBD), and holds the condition that the addition of BD and NBD is less than or equal to 1. As an extension of IFSs, Yager (2014) initiated the notion of PFSs. The PFSs are more powerful tool than IFSs for handling the uncertain, vague and imprecise information arisen in the realistic problems. Recently, numerous researchers have explored the different concepts by considering the theoretical and practical aspects of PFSs. For instance, Rani et al. (2019) proposed the VIKOR approach-based on entropy and discrimination measures to handle the renewable energy resources assessment in India. Rani et al. (2020a) developed COPRAS method to select the pharmacological therapies for type-2 diabetes disease. Rani et al. (2020b) assessed and selected the healthcare waste treatment options using SWARA and ARAS methods.

However, in numerous practical concerns, a group of DMs may give the BD to which an alternative satisfies the attribute is 0.8 and the NBD to which an alternative dissatisfies the attribute is 0.7. Here, we observe that  $0.8 + 0.7 > 1$  and  $0.8^2 + 0.7^2 > 1$ , as a result, the IFS and PFS are incapable to tackle this concern. To handle such information, Senapati and Yager (2019a) gave the doctrine of Fermatean fuzzy set (FFS), which is termed as the BD and NBD and the constraint that the cube sum of BD and NBD is less than or equal to 1. Consequently, the FFS can efficiently

solve the aforementioned concern. Also, FFSs can solve the MCDM problems with more effective way and handle the complex uncertain information. For example, Senapati and Yager (2019a) pioneered the doctrine of FFSs and proposed various basic concepts for solving decision-making problem. Senapati and Yager (2019b) discussed numerous aggregation operators with their elegant properties for FFSs and used WPM to solve the MCDM problems (Senapati and Yager 2019c). Garg et al. (2020) studied several Fermatean fuzzy aggregation operators and presented their application in COVID-19 facility selection. Aydemir and Gunduz (2020) proposed Dombi operators and their properties for FFSs. Recently, there is no study related to the S3PRLP assessment on FFSs context.

### 2.3 CRITIC method

In the MCDM process, various criteria weighting methods have been developed in the literature (Suh et al. 2019; Mishra et al. 2020a; Rani et al. 2020c). The criteria weight computation procedures are classified as objective weight and subjective weight (Peng 2019). The CRITIC approach (Diakoulaki et al. 1995) is one of the weighting models to estimate the objective weights of the attributes using the standard deviation and the correlation coefficient to quantify the value of each attribute and computes the attribute weights of MCDM procedure. Recently, several fusion models have been introduced by applying CRITIC with various other MCDM methods (Yalcin and Unlu 2018; Adali and Tus 2019). For example, Ghorabae et al. (2017) discussed a hybrid model with CRITIC and WASPAS methods to solve the 3PRLPs with IT2FSs. Ghorabae et al. (2018) presented a combined MCDM model with the CRITIC, EDAS and SWARA procedures. Peng et al. (2020) gave a CRITIC-CoCoSo method for 5G industry assessment on PFSs. Wei et al. (2020) initiated a hybrid model with GRA and CRITIC methods to assess and choose the suitable site for EVCSs under PULTSs environment. Peng and Huang (2020) discussed a hybrid model with CRITIC and CoCoSo methods for financial risk assessment problem. Liang (2020) gave a hybrid method with CRITIC and EDAS models under IFSS to solve the MCDM procedures.

### 2.4 EDAS method

The objective of MCDM procedure is to select the best option from a set of options under a set of various attributes. Currently, numerous MCDM approaches namely COPRAS (Kumari and Mishra 2020; Mishra et al. 2020a), CODAS (He et al. 2019; Zhou et al. 2020), ELECTRE (Fei et al. 2019; Mishra et al. 2020b), MULTIMOORA (Tian et al. 2017; Wu et al. 2020), TOPSIS (Aydemir and Gunduz 2020; Dammak et al. 2020), TODIM (Zindani et al. 2020;

Mishra et al. 2020c), VIKOR (Rani et al. 2019; Krishankumar et al. 2020), WASPAS (Ghorabae et al. 2017; Mishra and Rani 2018) and others have been discussed to solve the MCDM problems on different uncertain environments. Each MCDM procedure has been developed with different advantages and disadvantages, though, the scholars usually select an approach which is based on the nature and intricacy of the problem.

The EDAS method (Ghorabae et al. 2015) is an original and efficient tool to solve the MCDM problem with conflicting attributes. It utilizes the AVS for prioritizing the options and describes the discrimination between the options and the AVS according to the measures PDA and NDA (Ghorabae et al. 2016). Kahraman et al. (2017) used EDAS method for assessing solid waste disposal location on IFSSs. Gundogdu et al. (2018) extended the EDAS model for assessing and choosing the suitable hospital under HFSSs environment. Mi and Liao (2019) discussed a combined framework with BWM and EDAS method with HFSSs to assess insurance projects. Zhang et al. (2019) extended the EDAS approach to select the best green supplier. Han and Wei (2020) gave multivalued neutrosophic EDAS model for dealing with the MCDM problems. Mishra et al. (2020d) discussed parametric discrimination measure-based EDAS framework for assessing the HCWD method on IFSSs. In this study, we develop a combined framework with CRITIC and EDAS approaches for FFSs.

## 3 A new Fermatean fuzzy score function

This section introduces a new score function for Fermatean Fuzzy Numbers (FFNs) which avoids the shortcoming of existing score function (Senapati & Yager, 2019a,b). Here, the first subsection presents the concept, score and accuracy functions, and operational laws of FFSs. Based on this concept, an improved score function for FFNs is developed in the next subsection.

### 3.1 Prerequisites

**Definition 3.1.** Let  $\Delta$  be a limited universe of discourse. In 2019, Senapati and Yager (2019a) firstly presented the concept of FFS, which is mathematically expressed as.

$$T = \left\{ \langle t_i, (b_T(t_i), n_T(t_i)) \rangle \mid t_i \in \Delta \right\},$$

wherein  $b_T : \Delta \rightarrow [0, 1]$  represent the belongingness degree (BD) of an element  $t_i \in \Delta$  in FFS and  $n_T : \Delta \rightarrow [0, 1]$  signify the non-belongingness degree (NBD) of an element  $t_i \in \Delta$  in FFS. For every  $t_i \in \Delta$ , it satisfies the condition  $0 \leq (b_T(t_i))^3 + (n_T(t_i))^3 \leq 1$ . The indeterminacy degree of FFS is expressed by

$\pi_T(t_i) = \sqrt[3]{1 - b_T^3(t_i) - n_T^3(t_i)}$ ,  $\forall t_i \in \Delta$ . For simplicity, Senapati and Yager (2019a) called  $(b_T(t_i), n_T(t_i))$  as a Fermatean fuzzy number (FFN), given by  $\lambda = (b_\lambda, n_\lambda)$  where  $b_\lambda, n_\lambda \in [0, 1]$ ,  $\pi_\lambda = \sqrt[3]{1 - b_\lambda^3 - n_\lambda^3}$  and  $0 \leq b_\lambda^3 + n_\lambda^3 \leq 1$ .

**Definition 3.2.** Let  $\lambda = (b_\lambda, n_\lambda)$  be a FFN. Then, the score and accuracy functions of  $\lambda$  are defined as follows (Senapati and Yager 2019a, b):

$$\wp_s(\lambda) = (b_\lambda)^3 - (n_\lambda)^3 \text{ and } \wp_a(\lambda) = (b_\lambda)^3 + (n_\lambda)^3, \text{ where } -1 \leq \wp_s(\lambda) \leq 1 \text{ and } 0 \leq \wp_a(\lambda) \leq 1.$$

Thus, to rank two FFNs  $\lambda_1 = (b_{\lambda_1}, n_{\lambda_1})$  and  $\lambda_2 = (b_{\lambda_2}, n_{\lambda_2})$ , we have the following comparative scheme (Senapati and Yager 2019a, b):

- (i) If  $\wp_s(\lambda_1) > \wp_s(\lambda_2)$ , then  $\lambda_1$  is larger than  $\lambda_2$ , denoted by  $\lambda_1 > \lambda_2$ ;
- (ii) If  $\wp_s(\lambda_1) = \wp_s(\lambda_2)$ , then
- (iii) if  $\wp_a(\lambda_1) > \wp_a(\lambda_2)$ , then  $\lambda_1$  is larger than  $\lambda_2$ , denoted by  $\lambda_1 > \lambda_2$ ;
- (iv) if  $\wp_a(\lambda_1) < \wp_a(\lambda_2)$ , then  $\lambda_1$  is smaller than  $\lambda_2$ , denoted by  $\lambda_1 < \lambda_2$ ;
- (v) if  $\wp_a(\lambda_1) = \wp_a(\lambda_2)$ , then  $\lambda_1$  is identical with  $\lambda_2$ , denoted by  $\lambda_1 = \lambda_2$ .

**Definition 3.3.** Assume that  $\lambda = (b_\lambda, n_\lambda)$ ,  $\lambda_1 = (b_{\lambda_1}, n_{\lambda_1})$  and  $\lambda_2 = (b_{\lambda_2}, n_{\lambda_2})$  are three FFNs. Now, the operational laws are summarized as follows:

- 1)  $\lambda^c = (n_\lambda, b_\lambda)$ ;
- 2)  $\lambda_1 \cap \lambda_2 = (\min\{b_{\lambda_1}, b_{\lambda_2}\}, \max\{n_{\lambda_1}, n_{\lambda_2}\})$ ;
- 3)  $\lambda_1 \cup \lambda_2 = (\max\{b_{\lambda_1}, b_{\lambda_2}\}, \min\{n_{\lambda_1}, n_{\lambda_2}\})$ ;
- 4)  $\lambda_1 \oplus \lambda_2 = (\sqrt[3]{b_{\lambda_1}^3 + b_{\lambda_2}^3 - b_{\lambda_1}^3 b_{\lambda_2}^3}, \min\{n_{\lambda_1}, n_{\lambda_2}\})$ ;
- 5)  $\lambda_1 \otimes \lambda_2 = (b_{\lambda_1} b_{\lambda_2}, \sqrt[3]{n_{\lambda_1}^3 + n_{\lambda_2}^3 - n_{\lambda_1}^3 n_{\lambda_2}^3})$ ;
- 6)  $\lambda_1 \ominus \lambda_2 = \begin{cases} \left(\sqrt[3]{\frac{b_{\lambda_1}^3 - b_{\lambda_2}^3}{1 - b_{\lambda_2}^3}}, \frac{n_{\lambda_1}}{n_{\lambda_2}}\right), & b_{\lambda_1} \geq b_{\lambda_2}, n_{\lambda_1} \leq \min\left\{n_{\lambda_2}, \frac{n_{\lambda_2} \pi_{\lambda_1}}{\pi_{\lambda_2}}\right\} \\ (0, 1), & \text{otherwise;} \end{cases}$
- 7)  $\iota \lambda = \left(\sqrt[3]{1 - (1 - b_\lambda^3)^\iota}, (n_\lambda)^\iota\right), \iota > 0$ ;
- 8)  $\lambda^\iota = T\left((b_\lambda)^\iota, \sqrt[3]{1 - (1 - n_\lambda^3)^\iota}\right), \iota > 0$ .

### 3.2 Improved generalized score function (IGSF) for FFN

This section develops an IGSF for any FFN  $\lambda_j = (b_j, n_j)$ , and is presented as

$$\wp_s^*(\lambda_j) = b_j^3 \left[ 1 + (\gamma_1 + \gamma_2) (1 - b_j^3 - n_j^3) \right]. \tag{1}$$

Here,  $\gamma_1, \gamma_2 > 0$  signifies the performance of the proposed IGSF and  $\gamma_1 + \gamma_2 = 1$ . In addition,  $\gamma_1$  and  $\gamma_2$  provide the weighted average of the indeterminacy degree between the BD and NBD.

**Theorem 3.1:** For a FFN  $\lambda_j = (b_j, n_j)$ , the IGSF  $\wp_s^*(\cdot)$  satisfies the following:

- p1.  $\wp_s^*((0, 1)) = 0$  and  $\wp_s^*((1, 0)) = 1$ .
- p2. An IGSF  $\wp_s^*(\lambda_j)$  is increasing monotonically w. r. t.  $b_j$  and is decreasing monotonically w. r. t.  $n_j$ .

**Proof** p1. When  $\lambda_j = (0, 1)$  or  $\lambda_j = (1, 0)$ , then according to Eq. (1), the IGSF  $\wp_s^*(\lambda_j)$  attains the least value ‘0’ or utmost value ‘1’, respectively. It follows that  $0 \leq \wp_s^*(\lambda_j) \leq 1$ .

p2. To prove this part, differentiate Eq. (1) partially w.r.t.  $b_j$ , then we have

$$\frac{\partial \wp_s^*(\lambda_j)}{\partial b_j} = 3 b_j^2 (1 + (\gamma_1 + \gamma_2) (1 - b_j^3 - n_j^3)) - 3 b_j^5 (\gamma_1 + \gamma_2) \geq 0.$$

In the similar way, the first partial derivative of Eq. (1) w.r.t.  $n_j$  is presented as follows:

$$\frac{\partial \wp_s^*(\lambda_j)}{\partial n_j} = -3 b_j^3 n_j^2 (\gamma_1 + \gamma_2) \leq 0.$$

Thus, the theorem is proved.

**Remark 3.1** When we comparing any two FFNs  $\lambda_1 = (0.5, 0.5)$  and  $\lambda_2 = (0.4, 0.4)$ , we can observe that the score function given by Definition 3.2 (Senapati and Yager 2019a, b) is unable to distinguish the difference between the given FFNs because  $\wp_s(\lambda_1) = \wp_s(\lambda_2) = 0$ , whilst the developed IGSF (1) can successfully deal with this example and therefore, we have  $\wp_s^*(\lambda_1) = 0.2188$  and  $\wp_s^*(\lambda_2) = 0.1198$ . Hence,  $\lambda_1 > \lambda_2$ . This verifies the validity of the proposed score function over the existing one.

Here, Fig. 1 presents the value of score function  $\wp_s^*(\lambda_j)$  w. r. t.  $\gamma_1, \gamma_2 > 0, \gamma_1 + \gamma_2 = 1$  and at  $(b_j, n_j) = (0.8, 0.7)$ . The color of each pair  $(\gamma_1, \gamma_2)$  on the simplex demonstrates the score value of the fixed Fermatean fuzzy numbers. As the value of  $\gamma_1$  and  $\gamma_2$  become larger, the value of  $\wp_s^*(\lambda_j)$  becomes better. Similarly, Fig. 2 presents the value of score function  $\wp_s^*(\lambda_j)$  w.r.t.  $b_j$  and  $n_j$  at  $\gamma_1 = \gamma_2 = 0.5$ . The color of each pair  $(b_j, n_j)$  on the simplex presents the deviation on score function of the fixed Fermatean fuzzy numbers. As the value of  $b_j$  become larger, the value of  $\wp_s^*(\lambda_j)$  becomes superior.

Alternatively, the effects of diverse values of  $\gamma_1$  and  $\gamma_2$  on the preference order of the FFNs has been examined and their equivalent score values are summarized in Fig. 1 and Fig. 2. From the Figs. 1 and 2, it has been concluded that with the increase of  $\gamma_1$  from 0 to 1 (or simultaneously decreasing the value of  $\gamma_2$  from 1 to 0) the relative score of the FFNs increases. Also, from Remark 3.1, relative score of the FFNs increases but the overall ranking of the FFNs remain unaltered. Thus, based on these different parameter values, system analysis or decision maker may possess the power to alter the ranks of the FFNs.

### 4 Novel Fermatean fuzzy-based decision making method

Ghorabae et al. (2015) originated the notion of EDAS method as an effective tool for solving MCDM problems with conflicting criteria. In this method, the optimal solution is estimated based on the distance from average

solution. Therefore, there is no need to estimate the ideal and anti-ideal solutions. This is the key benefit of the EDAS approach. As FFS is one of the new ways to deal with the uncertainty of real-life applications. So, in this section, we combine the CRITIC and EDAS approaches with FFSs for solving complicated decision-making problems and named as Fermatean fuzzy CRITIC-EDAS (FF-CRITIC-EDAS) (see Fig. 3).

#### Step 1. Problem description

Consider that  $\{K_1, K_2, \dots, K_p\}$  be a set of options and  $\{N_1, N_2, \dots, N_q\}$  be a criterion set. Let us suppose that a group of decision makers (DMs)  $\{O_1, O_2, \dots, O_r\}$  presents their judgments on each option  $K_i$  concerning a criterion  $N_j$  in terms of FFNs. Assume that  $M_k = (\beta_{ij}^{(k)}) (i = 1, 2, \dots, p, j = 1, 2, \dots, q)$  be the Fermatean fuzzy decision matrix (FF-DM) offered by the  $k^{\text{th}}$  DM, in which  $\beta_{ij}^{(k)}$  signifies the evaluation information of an option  $K_i$  w.r.t. criterion  $N_j$  in the form of FFNs.

#### Step 2. Determination of DMs' weights

Fig. 1 IGSF  $\wp_s^*(\lambda_j)$  w.r.t. parameters  $\gamma_1, \gamma_2$ , at  $(b_j, n_j) = (0.8, 0.7)$ .





Fig. 2 The function  $\wp_s^*(\lambda_j)$  w.r.t. parameters  $b_j, n_j$ , at  $\gamma_1 = \gamma_2 = 0.5$



In order to compute the significance degrees of DMs, first of all, consider the significance degrees of the DMs in term of FFNs. Now, suppose  $W = (b_k, n_k)$  be the importance rating of  $k^{\text{th}}$  DM expressed by an authority in terms of FFN, then the formula for the computation of  $k^{\text{th}}$  DM's weight is presented as follows:

Here,  $\psi_k \geq 0$  and  $\sum_{k=1}^r \psi_k = 1$ .

Step 3. Aggregate the individual opinions of DMs

In the MCDM problem, it is important to merge all the individuals' opinions of DMs into a collective opinion to form the aggregated Fermatean fuzzy decision matrix

$$\psi_k = \frac{b_k^3 [1 + (\gamma_1 + \gamma_2)(1 - b_k^3 - n_k^3)]}{\sum_{k=1}^r (b_k^3 [1 + (\gamma_1 + \gamma_2)(1 - b_k^3 - n_k^3)])}, \quad k = 1, 2, \dots, r, \gamma_1 + \gamma_2 = 1, \gamma_1, \gamma_2 > 0. \tag{2}$$




Fig. 3 Proposed decision making method

(A-FFDM). To facilitate this, FF-weighted averaging operator (FFWAO) is employed on FF-DM and let  $Z = (z_{ij})_{p \times q}$ , ( $i = 1, 2, \dots, p, j = 1, 2, \dots, q$ ) be the A-FFDM, where

$$z_{ij} = \left( \sqrt[3]{1 - \prod_{k=1}^r \left(1 - (b_{ij}^k)^3\right)^{w_k}} \prod_{k=1}^r (n_{ij}^k)^{w_k} \right). \tag{3}$$

Step 4. Evaluate the criteria weight by employing CRITIC technique

Firstly, consider that  $X = (\varpi_1, \varpi_2, \dots, \varpi_q)^T$  be the weight of the criterion set with  $\varpi_j \in [0, 1]$  and  $\sum_{j=1}^q \varpi_j = 1$ .

Now, the computation procedure of CRITIC method is presented in the following steps:

Step 4.1. Evaluate the score matrix  $\Xi = (\kappa_{ij})_{p \times q}$ , wherein

$$\kappa_{ij} = b_{ij}^3 \left[ 1 + (\gamma_1 + \gamma_2) \left( 1 - b_{ij}^3 - n_{ij}^3 \right) \right], \tag{4}$$

Step 4.2. Switch the score matrix  $\Xi$  into a standard Fermatean fuzzy matrix  $\tilde{\Xi} = (\tilde{\kappa}_{ij})_{p \times q}$

$$\tilde{\kappa}_{ij} = \begin{cases} \frac{\kappa_{ij} - \kappa_j^-}{\kappa_j^+ - \kappa_j^-}, & j \in N_b, \\ \frac{\kappa_j^+ - \kappa_{ij}}{\kappa_j^+ - \kappa_j^-}, & j \in N_n, \end{cases} \tag{5}$$

where  $\kappa_j^- = \min_i \kappa_{ij}$  and  $\kappa_j^+ = \max_i \kappa_{ij}$ .

Step 4.3. Estimate the criteria standard deviations with the use of following formula:

$$\sigma_j = \sqrt{\frac{\sum_{i=1}^p (\tilde{\kappa}_{ij} - \bar{\kappa}_j)^2}{p}}, \tag{6}$$

where  $\bar{\kappa}_j = \sum_{i=1}^p \tilde{\kappa}_{ij} / p$ .

Step 4.4. Calculate the correlation between criteria with the use of following formula:

$$r_{jt} = \frac{\sum_{i=1}^p (\tilde{\kappa}_{ij} - \bar{\kappa}_j)(\tilde{\kappa}_{it} - \bar{\kappa}_t)}{\sqrt{\sum_{i=1}^p (\tilde{\kappa}_{ij} - \bar{\kappa}_j)^2 \sum_{i=1}^p (\tilde{\kappa}_{it} - \bar{\kappa}_t)^2}}. \tag{7}$$

Step 4.5. Evaluate the quantity of information of each criterion by using

$$v_j = \sigma_j \sum_{i=1}^q (1 - r_{jt}). \tag{8}$$

The better the  $c_j$  is, the more information an attribute contains, consequently the weight of evaluation attribute is superior than that of other attributes.

Step 4.6. Calculate the weight of each criterion as follows:

$$\varpi_j = \frac{v_j}{\sum_{j=1}^q v_j}. \tag{9}$$

Step 5. Assess the average solution (AVS) associated to the criteria

$$\diamond_i = [\varepsilon_i]_{1 \times q} = \frac{1}{p} \bigoplus_{i=1}^p z_{ij}, \quad i = 1, 2, \dots, p, \quad j = 1, 2, \dots, q. \tag{10}$$

Step 6. Estimate the positive distance and the negative distance from average solution matrix according as the benefit and cost criteria

If  $N_b$  and  $N_n$  are the set of benefit and cost criteria, respectively, then the positive distance from average (PDA) and the negative distance from average (NDA) are computed as below:

$$\Lambda_{PDA} = [h_{ij}]_{p \times q} \text{ and } \Lambda_{NDA} = [\chi_{ij}]_{p \times q}, \tag{11}$$

such that

$$h_{ij} = \begin{cases} \frac{\max(0, z_{ij} \ominus \varepsilon_i)}{\wp_s^*(\varepsilon_i)}, & \text{if } j \in N_b \\ \frac{\max(0, \varepsilon_i \ominus z_{ij})}{\wp_s^*(\varepsilon_i)}, & \text{if } j \in N_n \end{cases} \text{ and} \tag{12}$$

$$\chi_{ij} = \begin{cases} \frac{\max(0, \varepsilon_i \ominus z_{ij})}{\wp_s^*(\varepsilon_i)}, & \text{if } j \in N_b \\ \frac{\max(0, z_{ij} \ominus \varepsilon_i)}{\wp_s^*(\varepsilon_i)}, & \text{if } j \in N_n, \end{cases}$$

wherein  $h_{ij}$  and  $\chi_{ij}$  describe the positive and the negative distances of assessment degrees of  $i^{th}$  option from the AVS on  $j^{th}$  criterion, respectively, and  $\wp_s^*(\varepsilon_i)$  shows the score value of AVS.

Step 7. Compute the weighted sum of PDA ( $Z_i^{(+)}$ ) and weighted sum of NDA ( $Z_i^{(-)}$ ) for all options by using the following formulae:

$$Z_i^{(+)} = \bigoplus_{j=1}^q \varpi_j h_{ij}, \tag{13}$$

$$Z_i^{(-)} = \bigoplus_{j=1}^q \varpi_j \chi_{ij}. \tag{14}$$

Step 8. For all options, determine the normalize values of  $Z_i^{(+)}$  and  $Z_i^{(-)}$ , given as



$$N_i^{(+)} = \frac{Z_i^{(+)}}{\max_i \left[ \wp_s^* \left( Z_i^{(+)} \right) \right]}, \tag{15}$$

$$N_i^{(-)} = 1 - \frac{Z_i^{(-)}}{\max_i \left[ \wp_s^* \left( Z_i^{(-)} \right) \right]}. \tag{16}$$

Here, the terms ‘ $\wp_s^* \left( Z_i^{(+)} \right)$ ’ and ‘ $\wp_s^* \left( Z_i^{(-)} \right)$ ’ denote the score values of weighted sum of PDA and NDA, respectively.

Step 9. Evaluate the appraisal score for all options, given as

$$C_i = \frac{1}{2} \left( N_i^{(+)} \oplus N_i^{(-)} \right), \quad i = 1, 2, \dots, p. \tag{17}$$

Step 10. Based on the score values  $\wp_s^* \left( C_i \right)$  of appraisal scores  $C_i \left( i = 1, 2, \dots, p \right)$ , determine the ranking of the alternatives. As a result, the alternative with the highest appraisal score is the most appropriate choice among the others.

Step 11. End.

### 5 Case study: S3PRLP selection problem

In this section, we present an illustrative example of Indian electronics manufacturing firm to reveal the effectiveness and potentiality of the proposed FF-CRITIC-EDAS methodology. The considered manufacturing firm is situated in Gurugram, India and at present, this firm has five S3PRLP options  $(K_1, K_2, K_3, K_4, K_5)$ , correspondingly. In order to process the present decision-making problem, we form a group of three skilled decision makers  $(O_1, O_2, O_3)$ . The given S3PRLP options are evaluated on the basis of following 13 attributes/criteria: Cost of pollution control  $(N_1)$ , Cost of green product and eco-design  $(N_2)$ , Green warehousing  $(N_3)$ , Green R & D and innovation  $(N_4)$ , Environmental management system  $(N_5)$ , Costs  $(N_6)$ , Flexibility  $(N_7)$ , Quality  $(N_8)$ , Technology capability  $(N_9)$ , Health and safety practices  $(N_{10})$ , Social responsibility  $(N_{11})$ , Education Infrastructure  $(N_{12})$  and Employment Practices  $(N_{13})$ . The preferred criteria in this study are taken from different resources. The descriptions of these attributes are shown in Table 1. In this study, the criteria  $N_1, N_2$  and  $N_6$  are of cost type and remaining others are of benefit type.

The implementation procedures of FF-CRITIC-EDAS method for the evaluation of S3PRLPs are as given below:

Step 1. Let us assume that the importance ratings of DMs’ judgements are given in the form of FFNs which as  $\{(0.80, 0.50, 0.7133), (0.85, 0.45, 0.6655), (0.75, 0.55, 0.7440)\}$ . Now, the FF-DM is constructed in Table 2.

**Steps 2–3:** As the weights of the DMs’ as expressed by the experts are conveyed in the form of FFNs. Next, with the use of Eq. (2), the DMs’ weights are obtained in the following crisp form:  $\{\Psi_1 = 0.3341, \Psi_2 = 0.3807, \Psi_3 = 0.2852\}$ . Further, the evaluation opinions provided by three DMs are aggregated with the use of Eq. (3) and then required A-FFDM are presented in Table 3.

Step 4. In the step, the criteria weights are determined by CRITIC approach, which as

Step 4.1. First, with the use of Eq. (4) and Table 3, the score values of A-FFDM are evaluated.

Step 4.2. Convert the score matrix  $\Xi = (\kappa_{ij})_{p \times q}$  into the standard FF-matrix  $\tilde{\Xi} = (\tilde{\kappa}_{ij})_{p \times q}$  by employing the Eq. (5).

Steps 4.3–4.6. With the use of Eqs. (6)–(8), the standard deviation, correlation coefficient and quantity of information of each factor are computed and presented in Table 4. By employing Eq. (9), the criteria weights  $(\varpi_j)$  are estimated and then shown in Table 4.

Steps 5–9. By employing Eq. (10) and Table 3, the AVS matrix is computed and shown in Table 3. With the use of formulae (11)–(17), the PDA and NDA based on types of criteria, weighted sum of PDA and NDA for all alternatives, the normalize values of weighted sum of PDA and NDA and the appraisal score  $(C_i)$  for all options are estimated and presented in Tables 5, 6, respectively.

Step 10. On the basis of score function  $\wp_s^* \left( C_i \right)$  of appraisal score  $(C_i)$ , the option  $K_3$  is the most optimal choice and the ranking order of the S3PRLP alternatives is  $K_3 > K_4 > K_5 > K_1 > K_2$ .

### 5.1 Comparative analysis

In this section, a comparison with Fermatean Fuzzy-TOPSIS (Senapati & Yager, 2019a) and Fermatean Fuzzy-WPM (Senapati & Yager, 2019b) is presented to verify the robustness of the proposed methodology.

#### 5.1.1 Fermatean fuzzy-TOPSIS (Senapati and Yager 2019a) method

The Fermatean Fuzzy-TOPSIS method involves the following calculation procedures:

Steps 1–4. Same as FF-CRITIC-EDAS framework.

Step 5. In the present case study,  $N_1, N_2$  and  $N_6$  are cost criteria and rest all are benefit criteria, therefore, there is a need to normalize the A-FFDM.

Step 6. Estimate the Fermatean fuzzy positive and negative ideal solutions, presented as  $x^+ = \{(0.578, 0.688), (0.441, 0.722), (0.548, 0.477), (0.701, 0.563), (0.681, 0.556), (0.533, 0.777), (0.707, 0.635), (0.705, 0.556), (0.685, 0.588), (0.685, 0.554), (0.710, 0.604), (0.755, 0.638), (0.724, 0.581)\}$  and  $x^- = \{(0.437, 0.753), (0.616, 0.712), (0.693, 0.544), (0.619,$

**Table 1** Descriptions of the criteria for S3PRLP assessment

Aspects	Criteria	References	Nature
Environmental	Cost of pollution control ( $N_1$ )	Meade and Sarkis (2002), Yayla et al. (2015), Roy et al. (2019), Zarbakhshnia et al. (2018), Zhang and Su (2020)	Non-beneficial
	Cost of green product and eco-design ( $N_2$ )	Saen (2010), Mavi et al. (2017), Tavana et al. (2018), Li et al. (2018),	Non-beneficial
	Green warehousing ( $N_3$ )	Li et al. (2012), Tavana et al. (2016), Zhang and Xu (2020), Zarbakhshnia et al. (2020)	Beneficial
	Green R & D and innovation ( $N_4$ )	Bai and Sarkis (2010), Mavi et al. (2017), Li et al. (2018), Tavana et al. (2018)	Beneficial
	Environmental management system ( $N_5$ )	Amindoust et al. (2012), Kannan et al. (2015), Sen et al. (2017), Roy et al. (2019)	Beneficial
Economic	Costs ( $N_6$ )	Saen (2009), Liou et al. (2011), Azadi et al. (2015), Sen et al. (2017), Zarbakhshnia et al. (2020)	Non-beneficial
	Flexibility ( $N_7$ )	Saen (2010), Liou et al. (2011), Govindan et al. (2012), Zarbakhshnia et al. (2018), Zhang and Xu (2020)	Beneficial
	Quality ( $N_8$ )	Saen (2009, 2010), Mavi et al. (2017), Li et al. (2018), Tavana et al. (2018)	Beneficial
	Technology capability ( $N_9$ )	Kuo et al. (2010), Saen (2010), Tavana et al. (2016), Zarbakhshnia et al. (2018), Zhang and Xu (2020)	Beneficial
Social	Health and safety practices ( $N_{10}$ )	Saen (2010), Govindan et al. (2009, 2012), Diabat et al. (2014), Mavi et al. (2017),	Beneficial
	Social responsibility ( $N_{11}$ )	Kuo et al. (2010), Büyüközkan and Çifçi (2011), Jung et al. (2017), Roy et al. (2019)	Beneficial
	Education infrastructure ( $N_{12}$ )	Saen (2009), Liu and Wang (2009), Kannan et al. (2017), Zhang and Xu (2020), Zarbakhshnia et al. (2020)	Beneficial
	Employment practices ( $N_{13}$ )	Boukherroub et al. (2017), Jung (2017), Tavana et al. (2018), Zarbakhshnia et al. (2018)	Beneficial

0.530), (0.641, 0.547), (0.615, 0.764), (0.671, 0.654), (0.658, 0.538), (0.631, 0.564), (0.664, 0.672), (0.657, 0.610), (0.701, 0.655), (0.625, 0.714)}, respectively.

Now, find out the distances between the alternatives  $K_i$  and the Fermatean fuzzy positive and negative ideal solutions over the attribute  $N_j$ .

Step 7. The relative closeness to the Fermatean fuzzy positive ideal solution is determined by utilizing

$$R(K_i) = \frac{Y_i^-}{Y_i^+ + Y_i^-},$$

where  $Y_i^+ = dis(z_{ij}, x^+) = \sum_{j=1}^n \omega_j \sqrt{\frac{1}{2} \left[ \left( (b_{ij})^3 - (b_j^+)^3 \right)^2 + \left( (n_{ij})^3 - (n_j^+)^3 \right)^2 + \left( (\pi_{ij})^3 - (\pi_j^+)^3 \right)^2 \right]}$  and

$$Y_i^- = dis(z_{ij}, x^-) = \sum_{j=1}^q \omega_j \sqrt{\frac{1}{2} \left[ \left( (b_{ij})^3 - (b_j^-)^3 \right)^2 + \left( (n_{ij})^3 - (n_j^-)^3 \right)^2 + \left( (\pi_{ij})^3 - (\pi_j^-)^3 \right)^2 \right]}.$$

Therefore, the obtained values are  $R(K_1)=0.589$ ,  $R(K_2)=0.343$ ,  $R(K_3)=0.499$ ,  $R(K_4)=0.472$ ,  $R(K_5)=0.480$ .

**Step 8:** The preference order of S3PRLP options are  $K_1 > K_3 > K_5 > K_4 > K_2$ , thus, the option  $K_1$  is the best S3PRLP.

**5.1.2 Fermatean fuzzy-WPM (Senapati and Yager 2019b) method**

Steps 1–5. Same as FF-CRITIC-EDAS approach.

Step 6. Based on Weighted Product Model (WPM), the total relative significance of option  $K_i$  is calculated using  $\delta(K_i) = \otimes_{j=1}^q \omega_j z_{ij}$ ,  $i = 1, 2, \dots, p$ . Thus, we have  $\delta(K_i) = \{(0.654, 0.594), (0.660, 0.630), (0.690, 0.605), (0.677, 0.609), (0.672, 0.598)$ .

Step 7. The score values of relative importance degree of options are calculated as  $\wp_s^*(\delta(K_1)) = 0.422$ ,  $\wp_s^*(\delta(K_2)) = 0.421$ ,  $\wp_s^*(\delta(K_3)) = 0.476$ ,  $\wp_s^*(\delta(K_4)) = 0.454$  and  $\wp_s^*(\delta(K_5)) = 0.450$ . The

**Table 2** Fermatean fuzzy decision matrix for S3PRLPs assessment

	$K_1$	$K_2$	$K_3$	$K_4$	$K_5$
$N_1$	$O_1: (0.50, 0.70)$ $O_2: (0.40, 0.70)$ $O_3: (0.52, 0.75)$	$O_1: (0.40, 0.75)$ $O_2: (0.45, 0.75)$ $O_3: (0.48, 0.76)$	$O_1: (0.60, 0.65)$ $O_2: (0.60, 0.70)$ $O_3: (0.55, 0.72)$	$O_1: (0.50, 0.75)$ $O_2: (0.55, 0.70)$ $O_3: (0.52, 0.76)$	$O_1: (0.55, 0.70)$ $O_2: (0.50, 0.70)$ $O_3: (0.54, 0.75)$
$N_2$	$O_1: (0.48, 0.70)$ $O_2: (0.45, 0.72)$ $O_3: (0.40, 0.75)$	$O_1: (0.60, 0.72)$ $O_2: (0.50, 0.78)$ $O_3: (0.55, 0.75)$	$O_1: (0.65, 0.68)$ $O_2: (0.60, 0.72)$ $O_3: (0.62, 0.74)$	$O_1: (0.55, 0.75)$ $O_2: (0.50, 0.70)$ $O_3: (0.55, 0.75)$	$O_1: (0.60, 0.65)$ $O_2: (0.60, 0.72)$ $O_3: (0.58, 0.70)$
$N_3$	$O_1: (0.60, 0.40)$ $O_2: (0.55, 0.50)$ $O_3: (0.50, 0.55)$	$O_1: (0.70, 0.50)$ $O_2: (0.72, 0.58)$ $O_3: (0.68, 0.55)$	$O_1: (0.74, 0.67)$ $O_2: (0.65, 0.70)$ $O_3: (0.69, 0.58)$	$O_1: (0.70, 0.69)$ $O_2: (0.72, 0.65)$ $O_3: (0.68, 0.60)$	$O_1: (0.70, 0.64)$ $O_2: (0.64, 0.57)$ $O_3: (0.72, 0.65)$
$N_4$	$O_1: (0.60, 0.50)$ $O_2: (0.65, 0.55)$ $O_3: (0.63, 0.54)$	$O_1: (0.67, 0.55)$ $O_2: (0.62, 0.58)$ $O_3: (0.68, 0.52)$	$O_1: (0.72, 0.50)$ $O_2: (0.65, 0.58)$ $O_3: (0.68, 0.55)$	$O_1: (0.64, 0.60)$ $O_2: (0.73, 0.55)$ $O_3: (0.75, 0.54)$	$O_1: (0.64, 0.51)$ $O_2: (0.69, 0.65)$ $O_3: (0.71, 0.54)$
$N_5$	$O_1: (0.70, 0.63)$ $O_2: (0.65, 0.59)$ $O_3: (0.64, 0.52)$	$O_1: (0.60, 0.58)$ $O_2: (0.69, 0.54)$ $O_3: (0.65, 0.52)$	$O_1: (0.73, 0.56)$ $O_2: (0.68, 0.62)$ $O_3: (0.67, 0.60)$	$O_1: (0.70, 0.65)$ $O_2: (0.66, 0.56)$ $O_3: (0.68, 0.60)$	$O_1: (0.66, 0.54)$ $O_2: (0.68, 0.56)$ $O_3: (0.73, 0.57)$
$N_6$	$O_1: (0.58, 0.75)$ $O_2: (0.53, 0.78)$ $O_3: (0.57, 0.79)$	$O_1: (0.50, 0.68)$ $O_2: (0.65, 0.72)$ $O_3: (0.64, 0.75)$	$O_1: (0.51, 0.77)$ $O_2: (0.52, 0.76)$ $O_3: (0.59, 0.81)$	$O_1: (0.65, 0.78)$ $O_2: (0.63, 0.76)$ $O_3: (0.58, 0.75)$	$O_1: (0.58, 0.76)$ $O_2: (0.55, 0.72)$ $O_3: (0.60, 0.74)$
$N_7$	$O_1: (0.68, 0.62)$ $O_2: (0.67, 0.68)$ $O_3: (0.69, 0.66)$	$O_1: (0.74, 0.68)$ $O_2: (0.71, 0.63)$ $O_3: (0.69, 0.62)$	$O_1: (0.74, 0.61)$ $O_2: (0.71, 0.64)$ $O_3: (0.69, 0.66)$	$O_1: (0.67, 0.55)$ $O_2: (0.68, 0.57)$ $O_3: (0.64, 0.61)$	$O_1: (0.64, 0.57)$ $O_2: (0.71, 0.54)$ $O_3: (0.63, 0.56)$
$N_8$	$O_1: (0.69, 0.53)$ $O_2: (0.71, 0.48)$ $O_3: (0.67, 0.56)$	$O_1: (0.69, 0.54)$ $O_2: (0.71, 0.62)$ $O_3: (0.73, 0.50)$	$O_1: (0.69, 0.51)$ $O_2: (0.73, 0.59)$ $O_3: (0.72, 0.57)$	$O_1: (0.67, 0.54)$ $O_2: (0.69, 0.52)$ $O_3: (0.63, 0.56)$	$O_1: (0.67, 0.58)$ $O_2: (0.71, 0.56)$ $O_3: (0.73, 0.54)$
$N_9$	$O_1: (0.68, 0.57)$ $O_2: (0.71, 0.58)$ $O_3: (0.69, 0.62)$	$O_1: (0.67, 0.59)$ $O_2: (0.63, 0.56)$ $O_3: (0.61, 0.54)$	$O_1: (0.68, 0.55)$ $O_2: (0.64, 0.50)$ $O_3: (0.69, 0.54)$	$O_1: (0.69, 0.54)$ $O_2: (0.63, 0.56)$ $O_3: (0.61, 0.52)$	$O_1: (0.65, 0.54)$ $O_2: (0.63, 0.58)$ $O_3: (0.68, 0.59)$
$N_{10}$	$O_1: (0.71, 0.57)$ $O_2: (0.69, 0.53)$ $O_3: (0.68, 0.57)$	$O_1: (0.63, 0.67)$ $O_2: (0.71, 0.69)$ $O_3: (0.67, 0.65)$	$O_1: (0.73, 0.69)$ $O_2: (0.67, 0.64)$ $O_3: (0.68, 0.53)$	$O_1: (0.71, 0.67)$ $O_2: (0.68, 0.63)$ $O_3: (0.64, 0.54)$	$O_1: (0.72, 0.62)$ $O_2: (0.70, 0.63)$ $O_3: (0.68, 0.61)$
$N_{11}$	$O_1: (0.70, 0.63)$ $O_2: (0.72, 0.67)$ $O_3: (0.67, 0.53)$	$O_1: (0.68, 0.61)$ $O_2: (0.73, 0.68)$ $O_3: (0.69, 0.63)$	$O_1: (0.75, 0.63)$ $O_2: (0.69, 0.57)$ $O_3: (0.71, 0.62)$	$O_1: (0.65, 0.58)$ $O_2: (0.71, 0.68)$ $O_3: (0.62, 0.56)$	$O_1: (0.71, 0.66)$ $O_2: (0.68, 0.53)$ $O_3: (0.71, 0.51)$
$N_{12}$	$O_1: (0.71, 0.66)$ $O_2: (0.70, 0.67)$ $O_3: (0.72, 0.63)$	$O_1: (0.78, 0.61)$ $O_2: (0.77, 0.64)$ $O_3: (0.70, 0.69)$	$O_1: (0.69, 0.52)$ $O_2: (0.70, 0.57)$ $O_3: (0.73, 0.68)$	$O_1: (0.74, 0.66)$ $O_2: (0.78, 0.61)$ $O_3: (0.77, 0.65)$	$O_1: (0.72, 0.61)$ $O_2: (0.73, 0.60)$ $O_3: (0.70, 0.62)$
$N_{13}$	$O_1: (0.69, 0.63)$ $O_2: (0.74, 0.67)$ $O_3: (0.68, 0.65)$	$O_1: (0.60, 0.71)$ $O_2: (0.63, 0.72)$ $O_3: (0.67, 0.71)$	$O_1: (0.76, 0.61)$ $O_2: (0.71, 0.63)$ $O_3: (0.66, 0.61)$	$O_1: (0.69, 0.64)$ $O_2: (0.68, 0.63)$ $O_3: (0.67, 0.65)$	$O_1: (0.77, 0.62)$ $O_2: (0.68, 0.55)$ $O_3: (0.74, 0.58)$

**Table 3** A-FFDM and AVS matrix for S3PRLP assessment

	$K_1$	$K_2$	$K_3$	$K_4$	$K_5$	$\diamond_i$
$N_1$	(0.470, 0.714)	(0.437, 0.753)	(0.578, 0.688)	(0.517, 0.733)	(0.523, 0.714)	(0.475, 0.516)
$N_2$	(0.441, 0.722)	(0.546, 0.751)	(0.616, 0.712)	(0.526, 0.731)	(0.586, 0.690)	(0.551, 0.721)
$N_3$	(0.548, 0.477)	(0.693, 0.544)	(0.688, 0.654)	(0.693, 0.648)	(0.679, 0.615)	(0.667, 0.583)
$N_4$	(0.619, 0.530)	(0.648, 0.552)	(0.677, 0.544)	(0.701, 0.563)	(0.672, 0.569)	(0.665, 0.551)
$N_5$	(0.657, 0.582)	(0.641, 0.547)	(0.687, 0.594)	(0.672, 0.600)	(0.681, 0.556)	(0.668, 0.575)
$N_6$	(0.553, 0.773)	(0.597, 0.715)	(0.533, 0.777)	(0.615, 0.764)	(0.568, 0.739)	(0.575, 0.753)
$N_7$	(0.671, 0.654)	(0.707, 0.643)	(0.707, 0.635)	(0.656, 0.574)	(0.656, 0.556)	(0.681, 0.611)
$N_8$	(0.683, 0.518)	(0.701, 0.557)	(0.705, 0.556)	(0.658, 0.538)	(0.695, 0.561)	(0.689, 0.546)
$N_9$	(0.685, 0.588)	(0.631, 0.564)	(0.661, 0.528)	(0.639, 0.542)	(0.644, 0.569)	(0.653, 0.558)
$N_{10}$	(0.685, 0.554)	(0.664, 0.672)	(0.687, 0.622)	(0.672, 0.615)	(0.693, 0.621)	(0.680, 0.616)
$N_{11}$	(0.690, 0.614)	(0.693, 0.642)	(0.710, 0.604)	(0.657, 0.610)	(0.691, 0.564)	(0.689, 0.606)
$N_{12}$	(0.701, 0.655)	(0.747, 0.643)	(0.697, 0.581)	(0.755, 0.638)	(0.709, 0.609)	(0.723, 0.625)
$N_{13}$	(0.697, 0.651)	(0.625, 0.714)	(0.708, 0.618)	(0.672, 0.639)	(0.724, 0.581)	(0.688, 0.639)

**Table 4** The standard FF-matrix  $\tilde{E} = (\tilde{k}_{ij})_{p \times q}$ , SD, quantity of information and weight value for each factor

	$K_1$	$K_2$	$K_3$	$K_4$	$K_5$	$\sigma_j$	$v_j$	$\varpi_j$
$N_1$	0.216	0.000	1.000	0.484	0.551	0.338	2.983	0.0569
$N_2$	1.000	0.501	0.000	0.588	0.168	0.348	5.306	0.1013
$N_3$	1.000	0.000	0.214	0.172	0.212	0.349	3.904	0.0745
$N_4$	0.000	0.334	0.724	1.000	0.601	0.341	5.108	0.0975
$N_5$	0.271	0.000	0.973	0.543	1.000	0.390	3.563	0.0680
$N_6$	0.771	0.060	1.000	0.000	0.488	0.390	3.157	0.0603
$N_7$	0.000	0.945	1.000	0.062	0.132	0.455	5.744	0.1097
$N_8$	0.641	0.905	1.000	0.000	0.749	0.352	3.828	0.0731
$N_9$	1.000	0.000	0.725	0.231	0.248	0.366	3.375	0.0644
$N_{10}$	1.000	0.000	0.728	0.436	0.856	0.355	3.168	0.0605
$N_{11}$	0.592	0.517	1.000	0.000	0.807	0.337	3.074	0.0587
$N_{12}$	0.000	0.831	0.258	1.000	0.354	0.371	6.199	0.1183
$N_{13}$	0.647	0.000	0.796	0.476	1.000	0.339	2.972	0.0567

**Table 5** PDA and NDA matrices for S3PRLP selection

		$K_1$	$K_2$	$K_3$	$K_4$	$K_5$
$N_1$	$\Lambda_{PDA}$	(0.00, 1.00)	(0.00, 1.00)	(0.454, 0.872)	(0.00, 1.00)	(0.396, 0.903)
	$\Lambda_{NDA}$	(0.00, 1.00)	(0.357, 0.616)	(0.00, 1.00)	(0.541, 0.648)	(0.00, 1.00)
$N_2$	$\Lambda_{PDA}$	(0.00, 1.00)	(0.284, 0.837)	(0.00, 1.00)	(0.485, 0.939)	(0.00, 1.00)
	$\Lambda_{NDA}$	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.507, 0.864)
$N_3$	$\Lambda_{PDA}$	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)
	$\Lambda_{NDA}$	(0.00, 1.00)	(0.463, 0.869)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)
$N_4$	$\Lambda_{PDA}$	(0.00, 1.00)	(0.00, 1.00)	(0.359, 0.971)	(0.00, 1.00)	(0.00, 1.00)
	$\Lambda_{NDA}$	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)
$N_5$	$\Lambda_{PDA}$	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.371, 0.931)
	$\Lambda_{NDA}$	(0.359, 0.975)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)
$N_6$	$\Lambda_{PDA}$	(0.475, 0.895)	(0.00, 1.00)	(0.588, 0.859)	(0.00, 1.00)	(0.00, 1.00)
	$\Lambda_{NDA}$	(0.00, 1.00)	(0.443, 0.841)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)
$N_7$	$\Lambda_{PDA}$	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)
	$\Lambda_{NDA}$	(0.354, 0.854)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)
$N_8$	$\Lambda_{PDA}$	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)
	$\Lambda_{NDA}$	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)
$N_9$	$\Lambda_{PDA}$	(0.00, 1.00)	(0.00, 1.00)	(0.319, 0.885)	(0.00, 1.00)	(0.00, 1.00)
	$\Lambda_{NDA}$	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.331, 0.952)
$N_{10}$	$\Lambda_{PDA}$	(0.277, 0.806)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)
	$\Lambda_{NDA}$	(0.00, 1.00)	(0.422, 0.809)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)
$N_{11}$	$\Lambda_{PDA}$	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.213, 0.864)
	$\Lambda_{NDA}$	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)
$N_{12}$	$\Lambda_{PDA}$	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)
	$\Lambda_{NDA}$	(0.476, 0.905)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)
$N_{13}$	$\Lambda_{PDA}$	(0.00, 1.00)	(0.00, 1.00)	(0.437, 0.934)	(0.00, 1.00)	(0.524, 0.839)
	NDA	(0.00, 1.00)	(0.659, 0.722)	(0.00, 1.00)	(0.00, 1.00)	(0.00, 1.00)

preference order of options are as  $K_3 > K_4 > K_5 > K_1 > K_2$ . Hence, the alternative  $K_3$  is the best choice among the given S3PRLPoptions.

Based on Fermatean fuzzy-TOPSIS method, the preference ordering of the S3PRLP alternatives is  $K_1 > K_3 > K_5 > K_4 > K_2$ , and thus, the option  $K_1$  is the

best choice. Similarly, with the use of Fermatean fuzzy-WPM, the final ranking of the S3PRLP alternatives is  $K_3 > K_4 > K_5 > K_1 > K_2$  and thus,  $K_3$  is the most desirable option. Figure 4 presents the graphical representation of ranking of the options by different approaches. Consequently, we can see that the desirable S3PRLP alternative,

**Table 6** Evaluation parameters of FF-CRITIC-EDAS approach for S3PRLP selection

S3PRLP	$Z_i^{(+)}$	$Z_i^{(-)}$	$N_i^{(+)}$	$N_i^{(-)}$	$C_i$	$\phi_s^*(C_i)$	Rankings
$K_1$	(0.201, 0.980)	(0.278, 0.970)	(0.606, 0.543)	(0.277, 0.490)	(0.503, 0.516)	0.221	4
$K_2$	(0.133, 0.982)	(0.339, 0.923)	(0.412, 0.573)	(0.166, 0.825)	(0.336, 0.687)	0.062	5
$K_3$	(0.312, 0.969)	(0.000, 1.000)	(0.851, 0.377)	(1.000, 0.000)	(1.000, 0.000)	1.000	1
$K_4$	(0.230, 0.994)	(0.214, 0.976)	(0.681, 0.821)	(0.422, 0.417)	(0.589, 0.585)	0.326	2
$K_5$	(0.254, 0.971)	(0.254, 0.982)	(0.738, 0.405)	(0.328, 0.325)	(0.622, 0.363)	0.411	3

i.e., ( $K_3$ ) is same by using Fermatean fuzzy-WPM and introduced approach, whereas the ranking result somewhat differ by Fermatean fuzzy-TOPSIS approach and the desirable choice is  $K_1$ . Also, by comparing with Tavana et al. (2018) and Li et al. (2018) approaches, the final ranking order of the S3PRLP alternatives is  $K_3 > K_5 > K_4 > K_2 > K_1$  and  $K_3 > K_4 > K_5 > K_2 > K_1$ , respectively, therefore, the most appropriate choice is  $K_3$  among all other S3PRLPs.

The introduced methodology discussed in this study is found proficient for solving the MCDM problems with conflicting criteria. The main advantages of the proposed method are listed as below:


- To deal with the ambiguity in the MCDM problems, all input variables are taken into account as uncertain issues described by Fermatean fuzzy numbers. The indeterminacy degree is considered necessary independently in the whole method and the options are put in rank utilizing trade-off values of all three parameters, unlike Tavana et al. (2018) wherein the IFSs have been applied, and Li

et al. (2018) wherein the FSs have been used a particular case of the FFSs.

- In the proposed method, the optimal criteria weights are evaluated using the CRITIC approach, which combines the individual contrast intensity and conflict between criteria, thus provides more accurate results, whereas the criteria weights are randomly chosen in Fermatean fuzzy-TOPSIS (Senapati and Yager 2019a) and Fermatean fuzzy-WPM (Senapati and Yager 2019b), and Tavana et al. (2018) applies ANP model to evaluate subjective weights of the criteria and Li et al. (2018) utilizes PCA-AHP model to compute the criteria weights.
- As compared to the Fermatean fuzzy-TOPSIS (Senapati and Yager 2019a) and fuzzy-TOPSIS (Tavana et al. 2018) approaches, in which the “positive ideal solution” and “negative ideal solution” are calculated by the experts as per their own facts, whilst in EDAS method, Fermatean fuzzy weighted averaging operator is employed for the determination of AVS, which is simple and free from the impact of human concerns. Furthermore, the subtraction procedure and IGSF of Fermatean fuzzy numbers is used

**Fig. 4** The significance degrees of alternatives over different methods

Proposed FF-CRITIC-EDAS method      FF-TOPSIS method      FF-WPM method



to estimate the “PDA” and the “NDA” of each alternative. It can fruitfully evade the selecting distance measure which needs to add or reduce the number of objects of FFSs and lead to the distortion of information.

- The developed FF-CRITIC-EDAS framework is not only appropriate for evaluating the MCDM problems under FFSs context, but can also successfully tackle with the MCDM problems under fuzzy sets, intuitionistic fuzzy sets and Pythagorean fuzzy sets contexts. The introduced methodology has the benefits of easy computation process and fast information processing.

Some limitations of the proposed method are as follows:

- In the proposed FF-CRITIC-EDAS method, all criteria are assumed to be independent. In reality, there are inter-relationships among the criteria.
- This method has limitation in order to deal with a large number of criteria.

## 5.2 Discussions

The past period has witnessed ever-increasing issue about the disposal of customer goods, since numerous of these materials comprise both large quantities of waste and considerable amounts of toxic heavy metals. Manufacturers have faced intensifying pressure from both governments and environmentally concentrated committee to ‘reduce’, ‘recycle’, and ‘reuse’ their industrial waste. Presently, RL has been considered as a main concern. The operative management of RL is useful to environmental safety, and it can carry the evident economic benefits to organizations. Many corporations do not hold sufficient resource or capability to accomplish their RL activities, thus they have to select the S3PRLP to those activities.

In this study, a novel FF-CRITIC-EDAS method was used to choose the suitable S3PRLPs for Indian manufacturing company located in Gurugram, India. A case study is taken to display some important insights regarding assessment criteria and prominent S3PRLP options. To do this, proposed IGSF is applied to compute the DMs’ weights, CRITIC model is used to assess the importance value of criteria and proposed EDAS method is implemented to rank the S3PRLP options. The obtained results by proposed method show that the option  $K_3$  is the most appropriate provider for this case. Moreover, the comparative discussion with extant models is also presented to elucidate the rationality of the introduced method. Thus, we found that  $K_3$  is the most suitable choice among a set of S3PRLPs. As a consequence, the proposed model has significant information that can be utilized by administrators in taking strategic or operational decisions in S3PRLPs evaluation.

Without loss of generality, the proposed framework would be correspondingly appropriate to other concerns and different organizations. It can also be applied as a standard procedure for service providers in guiding their modifications to the processes and strategic instructions, so that they can well assistance with client and societal potentials. Simultaneously, administrations and governing bodies can employ the introduced method to study the relationships among economic, environmental, and social concerns, and utilize the outcomes to influence and reassure stronger law and strategy execution on the sustainability.

## 6 Conclusions

The assessment of S3PRLP has become one of important decisions for the enterprises in the modern competitive market. The objective of this study is to introduce a MCDM methodology for assessing and selecting the optimal S3PRLP option on FFSs. To do this, firstly novel IGSF has been proposed to compare the options. Secondly, a hybrid framework based on CRITIC and EDAS methods with FFSs has been developed to solve the MCDM problems, wherein the DMs and attribute weights are completely unknown. In this framework, IGSF-based procedure has been proposed to compute the DMs’ weights and the attribute weights have been calculated by applying CRITIC approach. Further, a case study of S3PRLP selection has been taken to elucidate the practicality and effectiveness of the introduced framework. For this, an evaluation index process for S3PRLPs has been organized, which comprises three prime aspects. These aspects are characterized into five, four and four criteria, respectively, which are broadly deliberated according to the existing literatures. The CRITIC approach determines the weights of the considered criteria, which as Education infrastructure (0.1183), Flexibility (0.1097), Cost of green product and eco-design (0.1013), Green R & D and innovation (0.0975), Green warehousing (0.0745), Quality (0.0731), Environmental management system (0.0680), Technology capability (0.0644), Health and safety practices (0.0605), Costs (0.0603), Social responsibility (0.0587), Cost of pollution control (0.0569) and Employment practices (0.0567). Next, by employing the EDAS method, the priority order of S3PRLPs is obtained as  $K_3 > K_4 > K_5 > K_1 > K_2$ . Next, comparison with extant models has been made to validate the introduced framework. The outcomes verify that the introduced model has good proficiency and strength than the extant models. In addition, the introduced approach not only offers the priority order of the S3PRLP options but also illustrates the attributes performances in the S3PRLP selection.

In future, we will work on diverse MCDM approaches (namely, CoCoSo, WASPAS, MULTIMOORA or DNMA)

to select the optimal S3PRLP under FFSs context. Also, we will implement the introduced framework to the different problems, namely, EVCS site evaluation, HCWD method assessment, green supplier assessment, and other decision-making problems.

## References

- Adali EA, Tus A (2019) Hospital site selection with distance-based multicriteria decision-making methods. *Int J Healthc Manag*. <https://doi.org/10.1080/20479700.2019.1674005>
- Amindoust A, Ahmed S, Saghafinia A, Bahreininejad A (2012) Sustainable supplier selection: a ranking model based on fuzzy inference system. *Appl Soft Comput* 12(6):1668–1677
- Atanassov KT (1986) Intuitionistic fuzzy sets. *Fuzzy Sets Syst* 20:87–96
- Aydemir SB, Gunduz SY (2020) Fermatean fuzzy TOPSIS method with dombi aggregation operators and its application in multi-criteria decision making. *J Intell Fuzzy Syst*. <https://doi.org/10.3233/JIFS-191763>
- Azadi M, Saen RF (2011) A new chance-constrained data envelopment analysis for selecting third-party reverse logistics providers in the existence of dual-role factors. *Expert Syst Appl* 38:12231–12236
- Azadi M, Jafarian M, Saen RF, Mirhedayatian SM (2015) A new fuzzy DEA model for evaluation of efficiency and effectiveness of suppliers in sustainable supply chain management context. *Comput Oper Res* 54:274–285
- Bai C, Sarkis J (2010) Integrating sustainability into supplier selection with grey system and rough set methodologies. *Int J Prod Econ* 124(1):252–264
- Bai C, Sarkis J (2019) Integrating and extending data and decision tools for sustainable third-party reverse logistics provider selection. *Comput Oper Res* 110:188–207
- Banihashemi TA, Fei J, Chen PS-L (2019) Exploring the relationship between reverse logistics and sustainability performance: a literature review. *Mod Supply Chain Res Appl* 1(1):2–27
- Boukherroub K, LeBel, L., Ruiz, A. (2017) A framework for sustainable forest resource allocation: a Canadian case study. *Omega* 66(B):224–235
- Büyükoçkan G, Çifçi G (2011) A novel fuzzy multi-criteria decision framework for sustainable supplier selection with incomplete information. *Comput Ind* 62(2):164–174
- Dammak F, Baccour L, Alimi AM (2020) A new ranking method for TOPSIS and VIKOR under interval valued intuitionistic fuzzy sets and possibility measures. *J Intell Fuzzy Syst* 38(4):4459–4469
- Diabat A, Kannan D, Mathiyazhagan K (2014) Analysis of enablers for implementation of sustainable supply chain management—a textile case. *Journal of Cleaner Production* 83:391–403
- Diakoulaki D, Mavrotas G, Papayannakis L (1995) Determining objective weights in multiple criteria problems: The CRITIC method. *Comput Oper Res* 22:763–770
- Fattahi M, Govindan K (2017) Integrated forward/reverse logistics network design under uncertainty with pricing for collection of used products. *Ann Oper Res* 253:193–225
- Fei L, Xia J, Feng Y, Liu L (2019) An ELECTRE-based multiple criteria decision making method for supplier selection using Dempster-shafer theory. *IEEE Access* 7:84701–84716
- Garg H, Shahzadi G, Akram M (2020) Decision-making analysis based on fermatean fuzzy yager aggregation operators with application in COVID-19 testing facility. *MathProbl Eng* 7279027(27):1–16
- Ghorabae MK, Zavadskas EK, Olfat L, Turskis Z (2015) Multi-criteria inventory classification using a new method of evaluation based on distance from average solution (EDAS). *Informatica* 26(3):435–451
- Ghorabae MK, Zavadskas EK, Amiri M, Turskis Z (2016) Extended EDAS method for fuzzy multi-criteria decision-making: an application to supplier selection. *Int J Comput CommunControl* 11(3):358–371
- Ghorabae MK, Amiri M, Zavadskas EK, Antucheviciene J (2017) Assessment of third-party logistics providers using a CRITIC–WASPAS approach with interval type-2 fuzzy sets. *Transport* 32(1):66–78
- Ghorabae MK, Amiri M, Zavadskas EK, Antucheviciene J (2018) A new hybrid fuzzy MCDM approach for evaluation of construction equipment with sustainability considerations. *Arch Civ Mech Eng* 18(1):32–49
- Govindan K, Palaniappan M, Zhu Q, Kannan D (2012) Analysis of third party reverse logistics provider using interpretive structural modeling. *Int J Prod Econ* 140(1):204–211
- Govindan K, Soleimani H, Kannan D (2015) Reverse logistics and closed-loop supply chain: A comprehensive review to explore the future. *Eur J Oper Res* 240(3):603–626
- Govindan K, Murugesan P, Senthil P, Haq AN (2009) Multicriteria group decision making for the third party reverse logistics service provider in the supply chain model using fuzzy TOPSIS for transportation services. *Int J Serv Technol Manage* 11(2):162–181
- Gundogdu FK, Kahraman C, Civan HN (2018) A novel hesitant fuzzy EDAS method and its application to hospital selection. *J Intell Fuzzy Syst* 35(6):6353–6365
- Han L, Wei C (2020) An extended EDAS method for multicriteria decision-making based on multivalued neutrosophic sets. *Complexity* 7578507:1–9
- He T, Wei G, Wei C, Wang J (2019) CODAS method for Pythagorean 2-tuple linguistic multiple attribute group decision making. *IEEE Access*. <https://doi.org/10.1109/ACCESS.2019.2917588>
- Jung H (2017) Evaluation of third party logistics providers considering social sustainability. *Sustainability* 9(5):777. <https://doi.org/10.3390/su9050777>
- Kahraman C, Ghorabae MK, Zavadskas EK, Onar SC, Yazdani M, Oztaysi B (2017) Intuitionistic fuzzy EDAS method: an application to solid waste disposal site selection. *J Environ Eng Landsc Manag* 25(1):1–12
- Kannan D, Kannan G, Rajendran S (2015) Fuzzy axiomatic design approach based green supplier selection: a case study from Singapore. *J Clean Prod* 96:1–15
- Kannan D, Garg K, Jha PC, Diabat A (2017) Integrating disassembly line balancing in the planning of a reverse logistics network from the perspective of a third party provider. *Ann Oper Res* 253(1):353–376
- Krishankumar R, Premaladha J, Ravichandran KS, Sekar KR, Manikandan R, Gao XZ (2020) A novel extension to VIKOR method under intuitionistic fuzzy context for solving personnel selection problem. *Soft Comput* 24:1063–1081
- Kumari R, Mishra AR (2020) Multi-criteria COPRAS method based on parametric measures for intuitionistic fuzzy sets: application of green supplier selection. *Iran J Sci Technol Trans Electr Eng*. <https://doi.org/10.1007/s40998-020-00312-w>
- Kuo RJ, Wang YC, Tien FC (2010) Integration of artificial neural network and MADA methods for green supplier selection. *J Clean Prod* 18(12):1161–1170
- Li F, Li L, Jin C, Wang R, Wang H, Yang L (2012) A 3PL supplier selection model based on fuzzy sets. *Comput Oper Res* 39(8):1879–1884
- Li Y-L, Ying C-S, Chin K-S, Yang H-T, Xu J (2018) Third party reverse logistics provider selection approach based on

- hybrid-information MCDM and cumulative prospect theory. *J Clean Prod* 195:573–584
- Liang Y (2020) An EDAS method for multiple attribute group decision-making under intuitionistic fuzzy environment and its application for evaluating green building energy-saving design projects. *Symmetry* 12:1–12
- Liou JJH, Wang HS, Hsu CC, Yin SL (2011) A hybrid model for selection of outsourcing provider. *Appl Math Model* 35(10):5121–5133
- Liu HT, Wang WK (2009) An integrated fuzzy approach for provider evaluation and selection in third-party logistics. *Expert Syst Appl* 36(3):4387–4398
- Liu A, Ji X, Lu H, Liu H (2019) The selection of 3PRLs on self-service mobile recycling machine: Interval-valued Pythagorean hesitant fuzzy best-worst multi-criteria group decision-making. *J Clean Prod* 230:734–750
- Mavi RK, Goh M, Zarbakhshnia N (2017) Sustainable third party reverse logistic provider selection with fuzzy SWARA and fuzzy MOORA in plastic industry. *Int J Adv Manuf Technol* 91(5–8):2401–2418
- Meade L, Sarkis J (2002) A conceptual model for selecting and evaluating third-party reverse logistics providers. *Supply Chain Manag* 7(5):283–295
- Mi X, Liao H (2019) An integrated approach to multiple criteria decision making based on the average solution and normalized weights of criteria deduced by the hesitant fuzzy best worst method. *Comput Ind Eng* 133:83–94
- Mishra AR, Rani P (2018) Interval-Valued Intuitionistic Fuzzy WASPAS method: application in reservoir flood control management policy. *Group Decis Negot* 27:1047–1078
- Mishra AR, Mardani A, Rani P, Zavadskas EK (2020a) A novel EDAS approach on intuitionistic fuzzy set for assessment of health-care waste disposal technology using new parametric divergence measures. *J Clean Prod*. <https://doi.org/10.1016/j.jclepro.2020.122807>
- Mishra AR, Rani P, Mardani A, Pardasani KR, Govindan K, Alrasheedi M (2020b) Healthcare evaluation in hazardous waste recycling using novel interval-valued intuitionistic fuzzy information based on complex proportional assessment method. *Comput Ind Eng*. <https://doi.org/10.1016/j.cie.2019.106140>
- Mishra AR, Rani P, Pardasani KR, Mardani A, Stevic Z, Pamucar D (2020c) A novel entropy and divergence measures with multi-criteria service quality assessment using interval-valued intuitionistic fuzzy TODIM method. *Soft Comput*. <https://doi.org/10.1007/s00500-019-04627-7>
- Mishra AR, Singh RK, Motwani D (2020d) Intuitionistic fuzzy divergence measure-based ELECTRE method for performance of cellular mobile telephone service providers. *Neural Comput Appl* 32:3901–3921
- Peng X (2019) New multiparametric similarity measure and distance measure for interval neutrosophic set with IoT industry evaluation. *IEEE Access* 7:28258–28280
- Peng X, Huang H (2020) Fuzzy decision making method based on CoCoSo with CRITIC for financial risk evaluation. *Technol Econ Dev Econy*. <https://doi.org/10.3846/tede.2020.11920>
- Peng X, Zhang X, Luo Z (2020) Pythagorean fuzzy MCDM method based on CoCoSo and CRITIC with score function for 5G industry evaluation. *Artif Intell Rev* 53:3813–3847
- Rani P, Mishra AR, Pardasani KR, Mardani A, Liao HC, Streimikienė D (2019) A novel VIKOR approach based on entropy and divergence measures of Pythagorean fuzzy sets to evaluate renewable energy technologies in India. *J Clean Prod*. <https://doi.org/10.1016/j.jclepro.2019.117936>
- Rani P, Mishra AR, Mardani A (2020a) An extended Pythagorean fuzzy complex proportional assessment approach with new entropy and score function: application in pharmacological therapy selection for type 2 diabetes. *Appl Soft Comput*. <https://doi.org/10.1016/j.asoc.2020.106441>
- Rani P, Mishra AR, Pardasani KR (2020b) A novel WASPAS approach for multi-criteria physician selection problem with intuitionistic fuzzy type-2 sets. *Soft Comput* 24:2355–2367
- Rani P, Mishra AR, Krishankumar R, Ravichandran KS, Gandomi AH (2020c) A new Pythagorean fuzzy based decision framework for assessing healthcare waste treatment. *IEEE Trans Eng Manag*. <https://doi.org/10.1109/TEM.2020.3023707>
- Roy J, Pamucar D, Kar S (2019) Evaluation and selection of third party logistics provider under sustainability perspectives: an interval valued fuzzy-rough approach. *Ann Oper Res*. <https://doi.org/10.1007/s10479-019-03501-x>
- Saen RF (2009) A mathematical model for selecting third-party reverse logistics providers. *Int J Procure Manag* 2(2):180–190
- Saen RF (2010) A new model for selecting third-party reverse logistics providers in the presence of multiple dual-role factors. *Int J Adv Manuf Technol* 46(1):405–410
- Sen DK, Datta S, Mahapatra SS (2017) Decision support framework for selection of 3PL providers: Dominance-based approach in combination with grey set theory. *Int J Inf Technol Decis* 16(01):25–57
- Senapati T, Yager RR (2019a) Fermatean fuzzy sets. *J Ambient Intell Hum Comput*. <https://doi.org/10.1007/s12652-019-01377-0>
- Senapati T, Yager RR (2019b) Some new operations over Fermatean fuzzy numbers and application of Fermatean fuzzy WPM in multiple criteria decision making. *Informatica* 30(2):391–412
- Senapati T, Yager RR (2019c) Fermatean fuzzy weighted averaging/geometric operators and its application in multi-criteria decision-making methods. *Eng Appl Artif Intell* 85:112–121
- Senthil S, Srirangacharyulu B, Ramesh A (2014) A robust hybrid multi-criteria decision making methodology for contractor evaluation and selection in third-party reverse logistics. *Expert Syst Appl* 41(1):50–58
- Suh Y, Park Y, Kang D (2019) Evaluating mobile services using integrated weighting approach and fuzzy VIKOR. *PLoS ONE*. <https://doi.org/10.1371/journal.pone.0217786>
- Tajik G, Azadnia AH, Ma'aram A, Hassan SAHS (2014) Hybrid Fuzzy MCDM Approach for Sustainable Third-Party Reverse Logistics Provider Selection. *Adv Mater Res* 845:521–526
- Tavana M, Zareinejad M, Santos-Arteaga FJ, Kaviani MA (2016) A conceptual analytic network model for evaluating and selecting third-party reverse logistics providers. *Int J AdvManuf Technol* 86(5–8):1705–1721
- Tavana M, Zareinejad M, Santos-Arteaga FJ (2018) An intuitionistic fuzzy-grey superiority and inferiority ranking method for third-party reverse logistics provider selection. *Int J Syst Sci OperLog* 5(2):175–194
- Tian ZP, Wang J, Wang JQ, Zhang HY (2017) An improved MULTIMOORA approach for multi-criteria decision-making based on interdependent inputs of simplified neutrosophic linguistic information. *Neural Comput Appl* 28(1):585–597
- Uygun Ö, Kaçamak H, Kahraman ÜA (2015) An integrated DEMATEL and Fuzzy ANP techniques for evaluation and selection of outsourcing provider for a telecommunication company. *Comput Ind Eng* 86:137–146
- Wei G, Lei F, Lin R, Wang R, Wei Y, Wu J, Wei C (2020) Algorithms for probabilistic uncertain linguistic multiple attribute group decision making based on the GRA and CRITIC method: application to location planning of electric vehicle charging stations. *Econ Res Ekonomska Istraživanja* 33(1):828–846
- Wu S-M, You X-Y, Liu H-C, Wang L-E (2020) Improving quality function deployment analysis with the cloud MULTIMOORA method. *Int Trans Oper Res* 27(3):1600–1621
- Yager RR (2014) Pythagorean membership grades in multicriteria decision making. *IEEE Trans Fuzzy Syst* 22:958–965



- Yalcin N, Unlu U (2018) A multi-criteria performance analysis of initial public offering (IPO) firms using CRITIC and VIKOR methods. *Technol Econ Dev Econ* 24(2):534–560
- Yayla AY, Oztekin A, Gumus AT, Gunasekaran A (2015) A hybrid data analytic methodology for 3PL transportation provider evaluation using fuzzy multi-criteria decision making. *Int J Prod Res* 53(20):6097–6113
- Zadeh LA (1965) Fuzzy sets. *Inf Control* 8(3):338–353
- Zarbakshnia N, Soleimani H, Ghaderi H (2018) Sustainable third-party reverse logistics provider evaluation and selection using fuzzy SWARA and developed fuzzy COPRAS in the presence of risk criteria. *Appl Soft Comput* 65:307–319
- Zarbakshnia N, Wu Y, Govindan K, Soleimani H (2020) A novel hybrid multiple attribute decision-making approach for outsourcing sustainable reverse logistics. *J Clean Prod*. <https://doi.org/10.1016/j.jclepro.2019.118461>
- Zhang X, Su T (2020) The dominance degree-based heterogeneous linguistic decision-making technique for sustainable S3PRLP selection. *Complexity* 2020(6102036):1–18
- Zhang S, Gao H, Wei G, Wei Y, Wei., C. (2019) Evaluation Based on Distance from Average Solution Method for Multiple Criteria Group Decision Making under Picture 2-Tuple Linguistic Environment. *Mathematics* 7:243. <https://doi.org/10.3390/math7030243>
- Zhou J, Li KW, Balezentis T, Streimikienė D (2020) Pythagorean fuzzy combinative distance-based assessment with pure linguistic information and its application to financial strategies of multi-national companies. *Econ Res Ekonomska Istraživanja* 33(1):974–998
- Zindani D, Maity SR, Bhowmik S (2020) Interval-valued intuitionistic fuzzy TODIM method based on Schweizer-Sklar power aggregation operators and their applications to group decision making. *Soft Comput*. <https://doi.org/10.1007/s00500-020-04783-1>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.