

Fermions and supersymmetry in $E_{6(6)}$ exceptional field theory

Edvard T. Musaev^{a,b} and Henning Samtleben^b

^a*Faculty of Mathematics, National Research University Higher School of Economics, 7, st. Vavilova, 117312, Moscow, Russia*

^b*Laboratoire de Physique, Université de Lyon, UMR 5672, CNRS, École Normale Supérieure de Lyon, 46, allée d'Italie, F-69364, Lyon cedex 07, France*

E-mail: emusaev@hse.ru, henning.samtleben@ens-lyon.fr

ABSTRACT: We construct the supersymmetric completion of $E_{6(6)}$ -covariant exceptional field theory. The theory is based on a $(5 + 27)$ -dimensional generalized space-time subject to a covariant section constraint. The fermions are tensors under the local Lorentz group $SO(1, 4) \times USp(8)$ and transform as weighted scalars under the $E_{6(6)}$ (internal) generalized diffeomorphisms. We present the complete Lagrangian and prove its invariance under supersymmetry. Upon explicit solution of the section constraint the theory embeds full $D = 11$ supergravity and IIB supergravity, respectively.

KEYWORDS: Supersymmetry and Duality, Supergravity Models, String Duality, M-Theory

ARXIV EPRINT: [1412.7286](https://arxiv.org/abs/1412.7286)

Contents

1	Introduction	1
2	Gauge structure and connections	3
2.1	Bosonic field content and tensor hierarchy	4
2.2	Fermions and connections	6
2.3	Curvatures	9
3	Supersymmetry transformations and algebra	12
4	Invariant Lagrangian	17
5	Conclusions and discussion	20
A	Details of the supersymmetry calculation	22
A.1	The $\psi\nabla_M\mathcal{D}_\mu\epsilon$ terms	22
A.2	The $\chi\nabla_M\mathcal{D}_\mu\epsilon$ terms	24
A.3	The $\psi\nabla_M\nabla_N\epsilon$ terms	25
A.4	The $\chi\nabla_M\nabla_N\epsilon$ terms	27
B	USp(8) identities	29

1 Introduction

Since the seminal work of Cremmer and Julia [1] it is well known that maximal supergravity compactified on a torus \mathbb{T}^d enjoys a hidden exceptional symmetry $E_{d(d)}$. From the M-theory point of view these U-duality transformations unify the perturbative T-duality, that relates type IIA and type IIB theories, and the S-duality of type IIB string theory. However, such formulation does not provide a natural geometric interpretation of the duality symmetries.

In the series of works [2–5], exceptional field theory, the $E_{d(d)}$ -covariant formulation of the full bosonic sector of maximal supergravity, was constructed for $d = 6, 7, 8$. It brings together the ideas from double field theory [6–10], its extension to exceptional groups [11–14], and extended geometry [15–17] that is an extension of Hitchin’s generalised geometry [18, 19] to the case of exceptional duality groups. These structures are defined on an exceptional space-time parametrized by external and internal coordinates $\{x^\mu, \mathbb{Y}^M\}$, $\mu = 0, \dots, 4$; $M = 1, \dots, 27$, the latter transforming in the fundamental representation of $E_{d(d)}$.¹ This space is dynamically restricted by a covariant differential constraint called

¹In the scheme of [20], generalized space-time is encoded in the infinite-dimensional l_1 representation of E_{11} .

section condition, that allows to systematically drop the extra coordinates and return to the conventional supergravity. The structure of the exceptional field theories resembles the one of the corresponding $(11 - d)$ -dimensional gauged supergravities [21, 22], however with all fields living on the full exceptional space-time. The dynamics of the “internal” sector is formulated in terms of the Lagrangian for a generalised metric, that is constructed from the scalar fields, parametrising the coset space G/K . In this formalism U-duality symmetries are recovered from generalized Lie derivatives in the internal space [15, 23]. Invariance under generalized diffeomorphisms in the external and internal coordinates uniquely fixes all the bosonic couplings of the theory without imposing any supersymmetric structure. Yet, the resulting bosonic system can be supersymmetrized by introducing fermions together with the corresponding connections under the generalized Lorentz group. The supersymmetric version of the full $E_{7(7)}$ EFT has been constructed in [24].

In this work we present the supersymmetric completion of the $E_{6(6)}$ -covariant exceptional field theory that lives on a 5+27-dimensional exceptional space-time. The bosonic theory has been constructed in [2, 3]. Generalized diffeomorphisms in the internal coordinates \mathbb{Y}^M enter the theory as Yang-Mills type gauge symmetries coupled to the Kaluza-Klein vector field \mathcal{A}_μ^M in the fundamental representation of $E_{6(6)}$. Fermions enter the theory as spinors under the generalized $SO(1, 4) \times USp(8)$ Lorentz group. Under generalized diffeomorphisms they transform as weighted scalars. As in $D = 5$ maximal supergravity [21, 25], gravitinos ψ_μ^i and fermions χ^{ijk} transform in the fundamental $\mathbf{8}$ and the antisymmetric traceless $\mathbf{42}$ representation of $USp(8)$, respectively. However, unlike in the five-dimensional truncation, they live on the full exceptional space-time modulo the covariant section condition, which effectively reduces the number of physical coordinates down to ten or eleven. Accordingly, the coupling of fermions requires a set of spin connections

$$\begin{array}{c|cc}
 & \mathcal{D}_\mu & \mathcal{D}_M \\
 \hline
 SO(1, 4) & \omega_\mu^{ab} & \omega_M^{ab} \\
 USp(8) & \mathcal{Q}_\mu^{ij} & \mathcal{Q}_M^{ij}
 \end{array} \quad , \tag{1.1}$$

in the external and internal directions, and for the two factors of the Lorentz group, respectively. These connections are defined in terms of the bosonic frame fields, the fünfbein e_μ^a , and the $E_{6(6)}$ -valued 27-bein \mathcal{V}_M^{ij} . The $SO(1, 4)$ connection ω_μ^{ab} is defined by the usual vanishing torsion condition

$$\mathcal{D}_{[\mu} e_{\nu]}^a = 0 \quad \iff \quad \Gamma_{[\mu\nu]}^\rho = 0 \quad , \tag{1.2}$$

however modified by the fact, that the derivative is covariantized also w.r.t. internal generalized diffeomorphisms under which the fünfbein e_μ^a transforms as a weighted scalar. For the internal sector on the other hand, vanishing torsion translates into the projection condition [15]

$$\Gamma_{MN}^K \Big|_{\mathbf{351}} = 0 \quad , \tag{1.3}$$

for the generalized Christoffel connection, decomposed into irreducible $E_{6(6)}$ representations. The off-diagonal blocks in (1.1) finally are determined by demanding that the

algebra-valued currents

$$\mathcal{J}_M^{ab} \equiv e^{a\mu} \mathcal{D}[\omega]_M e_\mu^b, \quad \mathcal{J}_{\mu kl}{}^{ij} \equiv \mathcal{V}_{kl}{}^M \mathcal{D}[\mathcal{A}, \mathcal{Q}]_\mu \mathcal{V}_M{}^{ij}, \quad (1.4)$$

of the frame fields live in the complement of the maximal compact subalgebra within $\text{GL}(5) \times \text{E}_{6(6)}$:

$$\mathcal{J}_M^{ab} \Big|_{\text{so}(1,4)} = 0, \quad \mathcal{J}_{\mu kl}{}^{ij} \Big|_{\text{usp}(8)} = 0. \quad (1.5)$$

Based on these connections we construct the supersymmetry transformation laws and the full supersymmetric Lagrangian in $\text{E}_{6(6)}$ -covariant form. Upon explicit solutions of the section condition, the Lagrangian reduces to full $D = 11$ supergravity and the IIB theory, respectively.

The paper is organized as follows. In section 2 we briefly review the structure of the bosonic $\text{E}_{6(6)}$ exceptional field theory. We give explicit expressions for the $\text{SO}(1,4)$ and $\text{USp}(8)$ connections (1.1) and the associated curvatures which are the building blocks for the bosonic field equations. In section 3 we present the supersymmetry transformations for all the fields of the theory in a U-duality covariant form based on the connections (1.1). The supersymmetry algebra closes with the following schematic form

$$\begin{aligned} [\delta(\epsilon_1), \delta(\epsilon_2)] = & \xi^\mu \mathcal{D}_\mu + \delta_{\text{so}(1,4)}(\Omega^{ab}) + \delta_{\text{usp}(8)}(\Lambda^{ij}) + \delta_{\text{susy}}(\epsilon_3) \\ & + \delta_{\text{gauge}}(\Lambda^M) + \delta_{\text{gauge}}(\Xi_\mu M) + \delta_{\text{gauge}}(\Xi_{\mu\nu} \alpha) + \delta_{\text{gauge}}(\Xi_{\mu\nu} M), \end{aligned} \quad (1.6)$$

into the local bosonic symmetries of the theory, with the explicit transformation parameters listed in (3.4) below. The geometry of the extended space deforms the supersymmetry algebra in a non-trivial way, although its structural form remains the same as that of the maximal gauged $D = 5$ supergravity [21, 25]. The full U-duality covariant supersymmetric Lagrangian is then given in section 4. In particular, we observe that all Pauli couplings of the fermions to the field strength $\mathcal{F}_{\mu\nu}{}^M$ can be absorbed by a shift of the internal spin connection according to

$$\omega_M^\pm{}^{ab} \equiv \omega_M{}^{ab} \pm \frac{1}{2} \mathcal{M}_{MN} \mathcal{F}_{\mu\nu}{}^N e^{\mu a} e^{\nu b}. \quad (1.7)$$

We sketch the relevant steps in the proof of supersymmetry invariance while the full calculational details are collected in appendix A. The results are discussed in section 5.

2 Gauge structure and connections

We start by giving a brief review of the bosonic field content and gauge symmetry of the $\text{E}_{6(6)}$ -covariant exceptional field theory, constructed in [2, 3] (to which we refer for details). Next we set up the $\text{USp}(8) \times \text{E}_{6(6)}$ -covariant geometrical formalism and in particular define the $\text{SO}(1,4)$ and $\text{USp}(8)$ spin connections required for the coupling of fermions. We then work out their various curvatures which are the building blocks for the bosonic field equations.

2.1 Bosonic field content and tensor hierarchy

The bosonic field content of $E_{6(6)}$ exceptional field theory is given by

$$\{e_\mu^a, \mathcal{V}_M^{ij}, \mathcal{A}_\mu^M, \mathcal{B}_{\mu\nu M}\}, \quad (2.1)$$

with indices $\mu, \nu = 0, \dots, 4$, and $M = 1, \dots, 27$, labelling external and internal coordinates, respectively, while indices $a = 0, \dots, 4$, and $i, j = 1, \dots, 8$, label fundamental indices of the $SO(1,4)$ and $USp(8)$ Lorentz group, respectively. The fünfbein e_μ^a defines the five-dimensional ‘external’ metric as $g_{\mu\nu} \equiv e_\mu^a e_\nu^b \eta_{ab}$ with the flat Minkowski metric η_{ab} . Similarly, the pseudo-real 27-bein \mathcal{V}_M^{ij} defines an ‘internal’ metric as

$$\mathcal{M}_{MN} = \mathcal{V}_M^{ij} \mathcal{V}_N^{ij}, \quad (2.2)$$

where $\mathcal{V}_M^{ij} \equiv (\mathcal{V}_M^{ij})^*$. The 27-bein \mathcal{V}_M^{ij} can be viewed as an $E_{6(6)}/USp(8)$ coset representative with the properties

$$\mathcal{V}_M^{ij} = \mathcal{V}_M^{[ij]}, \quad \mathcal{V}_M^{ij} \Omega_{ij} = 0, \quad \mathcal{V}_M^{ij} \equiv (\mathcal{V}_M^{ij})^* = \mathcal{V}_M^{kl} \Omega_{ki} \Omega_{lj}, \quad (2.3)$$

where $\Omega_{ij} = \Omega_{[ij]}$ denotes the symplectic invariant tensor. Thus \mathcal{M}_{MN} in (2.2) is real and symmetric. We further define the inverse 27-bein as

$$\mathcal{V}_M^{ij} \mathcal{V}_{ij}^N = \delta_M^N, \quad \mathcal{V}_M^{kl} \mathcal{V}_{ij}^M = \delta_{ij}^{kl} - \frac{1}{8} \Omega_{ij} \Omega^{kl}, \quad (2.4)$$

where we use conventions $\delta_{kl}^{ij} = \frac{1}{2}(\delta_k^i \delta_l^j - \delta_l^i \delta_k^j)$ and $\Omega_{ik} \Omega^{jk} = \delta_i^j$. The fact that the 27-bein is an $E_{6(6)}$ group-valued matrix is most efficiently encoded in the structure of its infinitesimal variation,

$$\delta \mathcal{V}_M^{ij} = -2 \delta q_k^{[i} \mathcal{V}_M^{j]k} + \delta p^{ijkl} \mathcal{V}_M^{kl}, \quad (2.5)$$

with δq_i^j and p^{ijkl} spanning the **36** and **42** of $USp(8)$, respectively, i.e.

$$\delta q_i^j = -\delta q_l^k \Omega_{ik} \Omega^{jl}, \quad \delta p^{ijkl} = \delta p^{[ijkl]}, \quad (2.6)$$

and corresponding to the compact and non-compact generators of $\mathfrak{e}_{6(6)}$, respectively. Double brackets $[[ijkl]]$ here and in the following indicate projection onto the totally antisymmetric and Ω -traceless part, i.e. $\delta p^{ijkl} \Omega_{kl} = 0$.

All fields (2.1) formally depend on the five external coordinates x^μ , and 27 internal coordinates \mathbb{Y}^M , with the latter transforming in the fundamental representation of $E_{6(6)}$. The \mathbb{Y}^M -dependence is strongly restricted by the $E_{6(6)}$ covariant section condition [11, 23, 26]

$$d^{KMN} \partial_M \partial_N A = 0, \quad d^{KMN} \partial_M A \partial_N B = 0, \quad (2.7)$$

for any fields or gauge parameters A, B . Here, d^{KMN} is the totally symmetric cubic invariant of $E_{6(6)}$. These constraints admit (at least) two inequivalent solutions, in which the fields depend on a subset of six or five of the internal coordinates. The resulting theories

are the full $D = 11$ supergravity and the type IIB theory, respectively. For later use we note that the cubic $E_{6(6)}$ invariant d^{MNK} is related to the symplectic tensor Ω_{ij} via

$$\begin{aligned} d^{MNP} &= \frac{2}{\sqrt{5}} \mathcal{V}_{ij}{}^M \mathcal{V}_{kl}{}^N \mathcal{V}_{mn}{}^P \Omega^{jk} \Omega^{lm} \Omega^{ni}, \\ d_{MNP} &= \frac{2}{\sqrt{5}} \mathcal{V}_M{}^{ij} \mathcal{V}_N{}^{kl} \mathcal{V}_P{}^{mn} \Omega_{jk} \Omega_{lm} \Omega_{ni}, \end{aligned} \quad (2.8)$$

as a consequence of the group property of $\mathcal{V}_M{}^{ij}$. We use normalization such that $d_{MKL} d^{NKL} = \delta_M{}^N$. With (2.8), the section constraint (2.7) can be rewritten as

$$\Omega_{[i} \mathcal{V}_{j]k}{}^M \mathcal{V}^{klN} \partial_M A \partial_N B = \frac{1}{8} \Omega_{ij} \mathcal{M}^{MN} \partial_M A \partial_N B, \quad \text{etc.}, \quad (2.9)$$

which is a form that we will often use in the following.

The exceptional field theory is invariant under generalized diffeomorphisms in the internal coordinates which act according to [15]

$$(\mathcal{L}_\Lambda V)^M = \Lambda^N \partial_N V^M - 6 \mathbb{P}^M{}_N{}^K{}_L \partial_K \Lambda^L V^N + \lambda_V \partial_P \Lambda^P V^M, \quad (2.10)$$

on a vector V^M of weight λ_V . Here, $\mathbb{P}^M{}_N{}^K{}_L = (t_\alpha)_N{}^M (t^\alpha)_L{}^K$ denotes the projector onto the adjoint representation of $E_{6(6)}$, $(t_\alpha)_N{}^M$ denoting the representation matrix in the fundamental representation. The diffeomorphism parameter Λ^M in (2.10) may depend on internal and external coordinates. As a result, all external derivatives are covariantized according to

$$D_\mu = \partial_\mu - \mathcal{L}_{\mathcal{A}_\mu}, \quad (2.11)$$

with the vector field $\mathcal{A}_\mu{}^M$ from (2.1). Accordingly, non-abelian field strengths for vector and two-form fields are defined as

$$\begin{aligned} \mathcal{F}_{\mu\nu}{}^M &= 2 \partial_{[\mu} \mathcal{A}_{\nu]}{}^M - 2 \mathcal{A}_{[\mu}{}^K \partial_K \mathcal{A}_{\nu]}{}^M + 10 d^{MKR} d_{NLR} \mathcal{A}_{[\mu}{}^N \partial_K \mathcal{A}_{\nu]}{}^L \\ &\quad + 10 d^{MNK} \partial_K \mathcal{B}_{\mu\nu N}, \\ \mathcal{H}_{\mu\nu\rho M} &= 3 D_{[\mu} \mathcal{B}_{\nu\rho] M} - 3 d_{MKL} \mathcal{A}_{[\mu}{}^K \partial_\nu \mathcal{A}_{\rho]}{}^L + 2 d_{MKL} \mathcal{A}_{[\mu}{}^K \mathcal{A}_{\nu}{}^P \partial_P \mathcal{A}_{\rho]}{}^L \\ &\quad - 10 d_{MKL} d^{LPR} d_{RNQ} \mathcal{A}_{[\mu}{}^K \mathcal{A}_{\nu}{}^N \partial_P \mathcal{A}_{\rho]}{}^Q + \dots, \end{aligned} \quad (2.12)$$

with the dots indicating terms that vanish under projection with $d^{KMN} \partial_N$. Here, vector fields and two-forms carry weight $\lambda_{\mathcal{A}} = \frac{1}{3}$, $\lambda_{\mathcal{B}} = \frac{2}{3}$, respectively, and the same weight is carried by their respective gauge parameters. The field strengths (2.12) transform covariantly under the non-abelian gauge transformations

$$\begin{aligned} \delta \mathcal{A}_\mu{}^M &= D_\mu \Lambda^M - 10 d^{MNK} \partial_K \Xi_{\mu N}, \\ \delta \mathcal{B}_{\mu\nu M} &= 2 D_{[\mu} \Xi_{\nu] M} + d_{MKL} (\Lambda^K \mathcal{F}_{\mu\nu}{}^L - \mathcal{A}_{[\mu}{}^K \delta \mathcal{A}_{\nu]}{}^L) + \mathcal{O}_{\mu\nu M}, \end{aligned} \quad (2.13)$$

with $d^{KMN} \partial_M \mathcal{O}_{\nu\nu N} = 0$. The parameter $\mathcal{O}_{\mu\nu M}$ can be viewed as the tensor gauge parameter of the three-forms of the theory which we have not explicitly introduced, since they

do not enter the Lagrangian. More precisely, this parameter may be decomposed according to the field content of three-forms, as

$$\mathcal{O}_{\mu\nu M} = (t^\alpha)_M{}^N \partial_N \Xi_{\mu\nu\alpha} + \Xi_{\mu\nu M}, \quad (2.14)$$

with gauge parameter $\Xi_{\mu\nu\alpha}$ in the adjoint representation, and a constrained gauge parameter $\Xi_{\mu\nu M}$ satisfying the same section condition (2.7) as the internal derivatives, i.e. $d^{KMN} \Xi_M \partial_N = 0$, etc.. This is analogous to the structure of two-forms in $E_{7(7)}$ EFT and vector fields in $E_{8(8)}$ EFT, respectively, cf. [4, 5]. The two forms $\mathcal{B}_{\mu\nu M}$ enter the Lagrangian only under projection $d^{KMN} \partial_M \mathcal{B}_{\nu N}$, such that their shift symmetry $\delta_{\mathcal{O}}$ constitutes a trivial symmetry of the action.

Under generalized diffeomorphisms, the field strengths $\mathcal{F}_{\mu\nu}{}^M$ and $\mathcal{H}_{\mu\nu\rho M}$ transform according to (2.10) as contravariant and covariant vector of weight $\lambda_{\mathcal{F}} = \frac{1}{3}$ and $\lambda_{\mathcal{H}} = \frac{2}{3}$, respectively. In contrast, both are inert under tensor gauge transformations $\Xi_{\mu M}$. The remaining bosonic fields in (2.1) transform as scalars under generalized diffeomorphisms with vanishing weight for $\mathcal{V}_M{}^{ij}$ and weight $\frac{1}{3}$ for the fünfbein $e_\mu{}^a$.

Furthermore, the non-abelian field strengths (2.12) satisfy the Bianchi identities

$$\begin{aligned} 3 D_{[\mu} \mathcal{F}_{\nu\rho]}{}^M &= 10 d^{MKN} \partial_K \mathcal{H}_{\mu\nu\rho N}, \\ 4 D_{[\mu} \mathcal{H}_{\nu\rho\sigma]}{}_M &= -3 d_{MKL} \mathcal{F}_{[\mu\nu}{}^K \mathcal{F}_{\rho\sigma]}{}^L + \dots \end{aligned} \quad (2.15)$$

In addition to the generalized internal diffeomorphisms and tensor gauge transformations (2.10), (2.13), the theory is invariant under external diffeomorphisms in the coordinates x^μ , under which the fields transform as

$$\begin{aligned} \delta e_\mu{}^a &= \xi^\nu D_\nu e_\mu{}^a + D_\mu \xi^\nu e_\nu{}^a, \\ \delta \mathcal{M}_{MN} &= \xi^\mu D_\mu \mathcal{M}_{MN}, \\ \delta \mathcal{A}_\mu{}^M &= \xi^\nu \mathcal{F}_{\nu\mu}{}^M + \mathcal{M}^{MN} g_{\mu\nu} \partial_N \xi^\nu, \\ \delta \mathcal{B}_{\mu\nu M} &= \frac{1}{2\sqrt{10}} \xi^\rho e \varepsilon_{\mu\nu\rho\sigma\tau} \mathcal{F}^{\sigma\tau N} \mathcal{M}_{MN} - d_{MKL} \mathcal{A}_{[\mu}{}^K \delta \mathcal{A}_{\nu]}{}^L, \end{aligned} \quad (2.16)$$

according to a modified version of the standard five-dimensional diffeomorphisms, with parameter ξ^μ which also is a function of x^μ and Y^M .

2.2 Fermions and connections

The fermionic fields of the theory comprise 8 gravitino fields ψ_μ^i and 42 spin- $\frac{1}{2}$ fermions $\chi^{ijk} = \chi^{\llbracket ijk \rrbracket}$. With respect to generalized internal diffeomorphisms (2.10) the fermionic fields transform as weighted scalars of weight $\lambda_\psi = \frac{1}{6}$, $\lambda_\chi = -\frac{1}{6}$. With respect to the (external and internal) Lorentz group, the fermions are $\text{SO}(1,4)$ spinors and transform in the corresponding representations of $\text{USp}(8)$. Like the bosonic fields, also the fermions depend on all coordinates x^μ , Y^M , modulo the section condition (2.7). We use the conventions of [21] from five-dimensional gauged supergravity.²

²The only exception is our convention for the Levi-Civita density where we follow [3], with the two conventions related by $\varepsilon_{\mu\nu\rho\sigma\tau}^{\llbracket 1312.0614 \rrbracket} = -i \varepsilon_{\mu\nu\rho\sigma\tau}^{\llbracket \text{hep-th}/0412173 \rrbracket}$. Accordingly, γ -matrices satisfy $\gamma^{abcde} = i \varepsilon^{abcde}$.

In particular, we use symplectic Majorana spinors subject to the reality constraint

$$C^{-1}\bar{\psi}_i^T = \Omega_{ij}\psi^j, \quad \psi^{iT}C = \Omega^{ij}\bar{\psi}_j, \quad C^{-1}\bar{\chi}_{ijk}^T = \Omega_{il}\Omega_{jm}\Omega_{kn}\chi^{lmn}, \quad (2.17)$$

where the charge conjugation matrix C is defined by the following relations

$$C\gamma_a C^{-1} = \gamma_a^T, \quad C^T = -C, \quad C^\dagger = C^{-1}. \quad (2.18)$$

This implies the following relation for fermionic bilinears with spinor fields ψ^i and φ^i

$$\bar{\psi}_i\Gamma\varphi^j = -\Omega_{ik}\Omega^{jl}\bar{\varphi}_l(C^{-1}\Gamma^T C)\psi^k, \quad (2.19)$$

for any expression of gamma matrices Γ .

According to the structure of the internal and external Lorentz group there are four different blocks of the spin connection

$$\begin{bmatrix} \omega_\mu & \mathcal{Q}_\mu \\ \omega_M & \mathcal{Q}_M \end{bmatrix}, \quad (2.20)$$

that ensure $\text{SO}(1,4)$ and $\text{USp}(8)$ covariance of external and internal derivatives, respectively. Let us discuss them one by one. The external $\text{SO}(1,4)$ connection ω_μ^{ab} is defined by the vanishing torsion condition

$$\mathcal{D}_{[\mu}e_{\nu]}^a \equiv D_{[\mu}e_{\nu]}^a + \omega_{[\mu}^{ab}e_{\nu]b} \stackrel{!}{=} 0, \quad (2.21)$$

as in standard Riemannian geometry albeit with derivatives D_μ covariantized according to (2.11). Furthermore, the external Christoffel connection Γ_μ can be defined by imposing the vielbein postulate for the fünfbein $\mathcal{D}_\mu e_\nu^a - \Gamma_{\mu\nu}^\lambda e_\lambda^a = 0$. The internal spin connection on the other hand is defined via

$$e^{\mu[a}\mathcal{D}_M e_\mu^{b]} \stackrel{!}{=} 0 \iff \omega_M^{ab} = e^{\mu[a}\partial_M e_\mu^{b]}. \quad (2.22)$$

Its presence guarantees that internal spinor derivatives transform as $\text{SO}(1,4)$ spinors. As a general notation in the following we will use \mathcal{D} to indicate (internal or external) derivatives including all spin connections while D_μ will only refer to the covariantization (2.11). Moreover, in the following it will be useful to define the modified internal spin connections

$$\omega_M^\pm{}^{ab} \equiv \omega_M^{ab} \pm \frac{1}{2}\mathcal{M}_{MN}\mathcal{F}_{\mu\nu}{}^N e^{\mu a}e^{\nu b}, \quad (2.23)$$

shifted by the non-abelian field strength (2.12). We will denote the corresponding covariant derivatives by \mathcal{D}_M^\pm .

Similar relations define the $\text{USp}(8)$ connections. The external connection $\mathcal{Q}_\mu{}^{ij}$ is defined in analogy to $D = 5$ gauged supergravity [21] by imposing that the covariant derivative of the 27-bein takes the form

$$\mathcal{D}_\mu\mathcal{V}_M{}^{ij} \equiv D_\mu\mathcal{V}_M{}^{ij} + 2\mathcal{Q}_{\mu k}{}^{[i}\mathcal{V}_M{}^{j]k} \stackrel{!}{=} \mathcal{P}_\mu{}^{ijkl}\mathcal{V}_M{}^{kl}, \quad (2.24)$$

with an $E_{6(6)}/\text{USp}(8)$ coset current $\mathcal{P}_\mu^{ijkl} = \mathcal{P}_\mu^{\llbracket ijkl \rrbracket}$. After proper contractions of indices it is straightforward to find the explicit expressions

$$\mathcal{Q}_\mu{}^i{}_j = \frac{1}{3} \mathcal{V}_{ik}{}^M D_\mu \mathcal{V}_M{}^{jk}, \quad \mathcal{P}_\mu^{ijkl} = D_\mu \mathcal{V}_M^{\llbracket ij} \mathcal{V}^{kl \rrbracket M}. \quad (2.25)$$

Note the use of the covariant derivative $D_\mu = \partial_\mu - \mathcal{L}_{\mathcal{A}_\mu}$ to preserve invariance under generalized diffeomorphisms. These equations imply the following Maurer-Cartan integrability conditions

$$\begin{aligned} \mathcal{Q}_{\mu\nu}{}^i{}_j &\equiv 2 \partial_{[\mu} \mathcal{Q}_{\nu]}{}^i{}_j + 2 \mathcal{Q}_{[\mu i}{}^k \mathcal{Q}_{\nu]}{}^j{}_k = -\frac{2}{3} \mathcal{P}_{\llbracket \mu i k l m \rrbracket} \mathcal{P}_{\nu]}{}^{j k l m} - \frac{1}{3} \mathcal{V}_{ki}{}^M \mathcal{L}_{\mathcal{F}_{\mu\nu}} \mathcal{V}_M{}^{kj}, \\ \mathcal{D}_{[\mu} \mathcal{P}_{\nu]}{}^{ijkl} &= -\frac{1}{2} \mathcal{L}_{\mathcal{F}_{\mu\nu}} \mathcal{V}_M^{\llbracket ij} \mathcal{V}^{kl \rrbracket M}, \end{aligned} \quad (2.26)$$

with the field strength $\mathcal{F}_{\mu\nu}{}^M$ from (2.12). It is straightforward to check that the $\mathcal{B}_{\mu\nu}{}^M$ contribution in the action $\mathcal{L}_{\mathcal{F}_{\mu\nu}} \mathcal{V}_M{}^{ij}$ drops out due to the section condition (2.7).

Finally, the internal $\text{USp}(8)$ connection \mathcal{Q}_M is defined by an analogue of the vanishing torsion condition (2.21) for the internal vielbein [15, 16]. To this end, it is convenient to define the full internal covariant derivative on an $E_{6(6)} \times \text{USp}(8)$ tensor $X_N{}^i$ of weight λ_X as

$$\nabla_M X_N{}^i \equiv \partial_M X_N{}^i - \mathcal{Q}_{Mj}{}^i X_N{}^j - \Gamma_{MN}{}^K X_K{}^i - \frac{3}{4} \lambda_X \Gamma_{KM}{}^K X_N{}^i, \quad (2.27)$$

with the algebra valued Christoffel connection $\Gamma_{MN}{}^K \equiv \Gamma_M^\alpha (t_\alpha)_N{}^K$. Such defined covariant derivative transforms as a generalized tensor of the weight $\lambda = \lambda_X - \frac{1}{3}$ under generalised diffeomorphisms. Vanishing torsion corresponds to imposing the relation

$$\mathcal{T}_{NK}{}^M \equiv \Gamma_{NK}{}^M - 6 \mathbb{P}^M{}_{K^P} \Gamma_{PN}{}^L + \frac{3}{2} \mathbb{P}^M{}_{K^Q} \Gamma_{PQ}{}^P \stackrel{!}{=} 0, \quad (2.28)$$

which transforms as a tensor under generalized diffeomorphisms (2.10). The vanishing torsion condition can equivalently be rewritten as [15, 17, 27]

$$(\mathbb{P}_{351})_M{}^{\alpha N} \Gamma_N{}^\beta = 0, \quad (2.29)$$

with the explicit form of the projector \mathbb{P}_{351} onto the **351** representation of $E_{6(6)}$ given by

$$(\mathbb{P}_{351})_M{}^{\alpha N}{}_\beta = -\frac{6}{5} (t^\alpha)_P{}^N (t_\beta)_M{}^P + \frac{3}{10} (t^\alpha)_M{}^P (t_\beta)_P{}^N + \frac{1}{5} \delta_M^N \delta_\beta^\alpha. \quad (2.30)$$

A particular consequence of (2.29) is

$$d^{MNK} \Gamma_{NK}{}^L = -\frac{1}{2} d^{MKL} \Gamma_{NK}{}^N. \quad (2.31)$$

The vanishing torsion conditions (2.21) can be explicitly solved upon imposing the generalized vielbein postulate for the 27-bein

$$\nabla_M \mathcal{V}_N{}^{ij} \equiv \partial_M \mathcal{V}_N{}^{ij} + 2 \mathcal{Q}_{Mk}{}^{\llbracket i} \mathcal{V}_N{}^{j \rrbracket k} - \Gamma_{MN}{}^K \mathcal{V}_K{}^{ij} \stackrel{!}{=} 0, \quad (2.32)$$

which allows to express the Christoffel connection in terms of the 27-bein and the internal USp(8) connection. In turn, the vanishing torsion conditions (2.21) translate into the conditions

$$\begin{aligned} \mathcal{D}_N \mathcal{V}_K^{[[ij] \mathcal{V}^{kl}] K} &= 6 \mathcal{D}_K \mathcal{V}_N^{[[ij] \mathcal{V}^{kl}] K} - \frac{3}{2} \mathcal{V}_N^{[[ij] \mathcal{V}^{kl}] M} \Gamma_{KM}^K, \\ \mathcal{V}_{ik}^K \mathcal{D}_N \mathcal{V}_K^{jk} &= 3 \left(\mathcal{V}_{ik}^M \mathcal{D}_M \mathcal{V}_N^{jk} - \mathcal{V}^{jkM} \mathcal{D}_M \mathcal{V}_{Nik} \right) \\ &\quad - \frac{3}{4} \Gamma_{KM}^K \left(\mathcal{V}_{ik}^M \mathcal{V}_N^{jk} - \mathcal{V}^{jkM} \mathcal{V}_{Nik} \right), \end{aligned} \quad (2.33)$$

for the USp(8) connection $\mathcal{Q}_M i^j$. These equations determine (part of) the USp(8) connection $\mathcal{Q}_M i^j$ in terms of the standard decomposition of the Cartan form $\mathcal{V}^{-1} \partial_M \mathcal{V}$ along the compact and non-compact parts of the $E_{6(6)}$ Lie algebra

$$q_M i^j \equiv \frac{1}{3} \mathcal{V}_{ik}^N \partial_M \mathcal{V}_N^{jk}, \quad p_M^{ijkl} \equiv \partial_M \mathcal{V}_N^{[ij] \mathcal{V}^{kl] N}. \quad (2.34)$$

Explicitly, parametrizing the connection as

$$\mathcal{Q}_M j^i = q_M j^i + \mathcal{V}_M^{kl} \Omega^{im} q_{kl, jm}, \quad (2.35)$$

with $q_{kl, ij} = q_{[[kl], (ij)}$, it is straightforward to verify that equations (2.33) are verified provided that³

$$\begin{aligned} q_{kl, mn} &= -p_{Mklp(m} \mathcal{V}_{n)q}^M \Omega^{pq} - \frac{1}{4} \mathcal{V}^{pqM} (p_{Mpqk(m} \Omega_{n)l} - p_{Mpq l(m} \Omega_{n)k}) \\ &\quad + \frac{1}{4} \Gamma_{KM}^K (\mathcal{V}_{k(m}^M \Omega_{n)l} - \mathcal{V}_{l(m}^M \Omega_{n)k}) + u_{kl, mn}. \end{aligned} \quad (2.36)$$

Here, $u_{kl, mn}$ denotes the undetermined part of the connection, satisfying

$$u_{kl, jm} = u_{[[kl], (jm)}, \quad u_{[kl, m]n} = 0, \quad u_{kl, jm} \Omega^{lj} = 0, \quad (2.37)$$

i.e. transforming in the **594** of USp(8), and dropping out from equations (2.33). Vanishing torsion thus determines the USp(8) connection (and thereby the Christoffel connection) up to a block transforming in the **594** of USp(8) [15, 16, 27]. The undetermined part of this connection drops out of all field equations and supersymmetry variations. Finally, one may fix the trace part in the Christoffel connection by demanding

$$\nabla_M e \stackrel{!}{=} 0 \quad \Longrightarrow \quad \Gamma_{KM}^K = \frac{4}{5} e^{-1} \partial_M e. \quad (2.38)$$

2.3 Curvatures

Let us recollect the notation for the various covariant derivatives introduced in the previous sections for the external and internal coordinates

$$\begin{aligned} D_\mu &= D[A_\nu]_\mu, \\ \mathcal{D}_\mu &= \mathcal{D}[A_\nu, \omega_\nu, \mathcal{Q}_\nu]_\mu, & \mathcal{D}_M &= \mathcal{D}[\omega_N, \mathcal{Q}_N]_M, \\ \nabla_\mu &= \nabla[A_\nu, \omega_\nu, \mathcal{Q}_\nu, \Gamma_\nu]_\mu, & \nabla_M &= \nabla[\omega_N, \mathcal{Q}_N, \Gamma_N]_\mu, \end{aligned} \quad (2.39)$$

³An explicit form of $\mathcal{Q}_M i^j$ in terms of the GL(6) components of \mathcal{V}_M^{ij} has been given in [15].

with vector field \mathcal{A}_μ^M gauging generalized diffeomorphisms as (2.11) and the composite connections ω , \mathcal{Q} , defined by (2.21), (2.22), (2.25), (2.35), in terms of the fünfbein e_μ^a and the 27-bein \mathcal{V}_M^{ij} . In addition, we recall the modified covariant derivatives \mathcal{D}_M^\pm and ∇_M^\pm , defined with the shifted internal spin connection ω_M^\pm from (2.23), that will come to play their role below.

The external curvature can be evaluated in the standard way by the commutator of covariant derivatives on an $\text{SO}(1,4) \times \text{USp}(8)$ spinor ϵ^i of weight λ_ϵ

$$[\mathcal{D}_\mu, \mathcal{D}_\nu] \epsilon^i = \frac{1}{4} \mathcal{R}_{\mu\nu}{}^{ab} \gamma_{ab} \epsilon^i - \mathcal{Q}_{\mu\nu j}{}^i \epsilon^j - \mathcal{F}_{\mu\nu}{}^M \partial_M \epsilon^i - \lambda_\epsilon \partial_M \mathcal{F}_{\mu\nu}{}^M \epsilon^i, \quad (2.40)$$

in terms of the Riemann curvature, $\text{USp}(8)$ curvature $\mathcal{Q}_{\mu\nu i}{}^j$, and the non-abelian field strength $\mathcal{F}_{\mu\nu}{}^M$ from (2.12). As it stands however, none of the terms on the r.h.s. is simultaneously covariant under generalized diffeomorphisms and local $\text{SO}(1,4) \times \text{USp}(8)$ transformations. In particular, the naive Riemann curvature defined as the curvature of the external spin connection

$$\mathcal{R}_{\mu\nu}{}^{ab} = 2 D_{[\mu} \omega_{\nu]}{}^{ab} + 2 \omega_{[\mu}{}^{ac} \omega_{\nu]c}{}^b, \quad (2.41)$$

transforms as $\delta_\lambda \mathcal{R}_{\mu\nu}{}^{ab} = \mathcal{F}_{\mu\nu}{}^M \partial_M \lambda^{ab}$ under $\text{SO}(1,4)$ Lorentz transformations. Using (2.26) and (2.32), the terms on the r.h.s. of (2.40) can be rearranged into the manifestly covariant expressions

$$[\mathcal{D}_\mu, \mathcal{D}_\nu] \epsilon^i = \frac{1}{4} \widehat{\mathcal{R}}_{\mu\nu}{}^{ab} \gamma_{ab} \epsilon^i + \frac{2}{3} \mathcal{P}_{[\mu j k l m} \mathcal{P}_{\nu]}{}^{i k l m} \epsilon^j + \nabla_M \mathcal{F}^N \left(\mathcal{V}_N{}^{jk} \mathcal{V}_{ik}{}^M - \mathcal{V}_N{}^{ik} \mathcal{V}^{jkM} \right) \epsilon^j - \mathcal{F}_{\mu\nu}{}^M \nabla_M \epsilon^i - \lambda_\epsilon \nabla_M \mathcal{F}_{\mu\nu}{}^M \epsilon^i, \quad (2.42)$$

with the full covariant internal derivatives ∇_M from (2.27) and the ‘improved’ Riemann tensor defined by [3, 28]

$$\widehat{\mathcal{R}}_{\mu\nu}{}^{ab} \equiv \mathcal{R}_{\mu\nu}{}^{ab} + \mathcal{F}_{\mu\nu}{}^M \omega_M{}^{ab}, \quad (2.43)$$

transforming covariantly under local Lorentz transformations. For later use, we note that this tensor and the associated Ricci tensor $\widehat{\mathcal{R}}_{\mu\nu} \equiv \mathcal{R}_{\mu\rho}{}^{ab} e_a{}^\rho e_{\nu b}$, satisfy the modified Bianchi identities

$$\begin{aligned} \widehat{\mathcal{R}}_{[\mu\nu]} &= -\frac{1}{2} g_{\rho[\mu} \nabla_M \mathcal{F}_{\nu]}{}^{\rho M}, \\ \widehat{\mathcal{R}}_{[\mu\nu\rho]}{}^a &= -\mathcal{F}_{[\mu\nu}{}^K \nabla_K e_{\rho]}{}^a - \frac{1}{3} e_{[\mu}{}^a \nabla_K \mathcal{F}_{\nu\rho]}{}^K. \end{aligned} \quad (2.44)$$

In contrast, the symmetric part of the Ricci tensor $\widehat{\mathcal{R}}_{(\mu\nu)}$ will appear in the Einstein field equations in the standard way. Similar to (2.42), the Maurer-Cartan integrability relations for the coset currents (2.26) can be rewritten in the manifestly covariant form

$$\mathcal{D}_{[\mu} \mathcal{P}_{\nu]}{}^{ijkl} = -3 \mathcal{V}_N{}^{[ij} \mathcal{V}{}^{kl]M} \nabla_M \mathcal{F}_{\mu\nu}{}^N. \quad (2.45)$$

Let us now discuss the mixed components of the curvature, i.e. the tensors obtained by commuting internal with external covariant derivatives. We only consider combinations of

commutators in which the undetermined part of the USp(8) spin connection (2.35), (2.36) drops out. This is the case for

$$\mathcal{V}_{ij}{}^M [\mathcal{D}_M^-, \mathcal{D}_\mu] \epsilon^j = \frac{1}{2} \mathcal{V}^{jkM} \mathcal{D}_M \mathcal{P}_\mu{}^{ijkn} \epsilon^n + \frac{1}{4} \mathcal{R}_{M\mu}{}^{-ab} \gamma_{ab} \epsilon^j . \quad (2.46)$$

Indeed, the undetermined connection on the l.h.s. appears as $\Omega^{jm} u_{ij,km} = 0$. On the r.h.s., the two terms describe the mixed USp(8) and SO(1,4) curvature, respectively, with the second term defined by the tensor

$$\mathcal{R}_{M\mu}{}^{-ab} \equiv \partial_M \omega_\mu{}^{ab} - \mathcal{D}_\mu \omega_M{}^{-ab} . \quad (2.47)$$

Evaluating this curvature gives rise to its Bianchi identity

$$\mathcal{R}_{M[\nu\rho\sigma]}^- = \frac{1}{2} \mathcal{D}_{[\nu} (\mathcal{F}_{\rho\sigma]}{}^N \mathcal{M}_{NM}) , \quad (2.48)$$

and the mixed Ricci tensor

$$\mathcal{R}_{M\nu}{}^{-\mu\nu} = -\frac{1}{2} \hat{\mathcal{J}}^\mu{}_M + \frac{1}{2} e_a{}^\mu e_b{}^\nu \mathcal{D}_\nu (\mathcal{M}_{MN} \mathcal{F}{}^{abN}) , \quad (2.49)$$

with the current $\hat{\mathcal{J}}^\mu{}_M$ defined by

$$\hat{\mathcal{J}}^\mu{}_M \equiv -2e_a{}^\mu e_b{}^\nu \left(\partial_M \omega_\nu{}^{ab} - \mathcal{D}_\nu (e^{\rho[a} \partial_M e_\rho{}^{b]}) \right) , \quad (2.50)$$

that will feature in the vector field equations. Similar to (2.46), we can evaluate the following combination of commutators

$$\begin{aligned} \mathcal{V}^{[ijM} [\mathcal{D}_M^-, \mathcal{D}_\mu] \epsilon^{k]} &= -2 \mathcal{V}_{mn}{}^M \Omega_{pr} \Omega^{m[i} \mathcal{D}_M \mathcal{P}_\mu{}^{jk]np} \epsilon^r - \frac{1}{2} \mathcal{V}_{mn}{}^M \mathcal{D}_M \mathcal{P}_\mu{}^{mn[ij} \epsilon^{k]} \\ &\quad + \frac{1}{4} \mathcal{R}_{M\mu}{}^{-ab} \mathcal{V}^{[ijM} \gamma_{ab} \epsilon^{k]} , \end{aligned} \quad (2.51)$$

in terms of the coset current $\mathcal{P}_\mu{}^{ijkl}$ and the mixed curvature (2.47). Again, the undetermined part of the USp(8) spin connection drops out on the l.h.s..

Let us finally discuss the internal components of the curvature. These are obtained from commutators of internal derivatives in combinations such that the undetermined part of the spin connections drops out. The relevant combinations are given by [15–17]

$$\begin{aligned} \mathcal{V}^{ikM} \mathcal{V}_{kj}{}^N [\nabla_M, \nabla_N] \epsilon^j &+ \left(4 \mathcal{V}^{ikM} \mathcal{V}_{kj}{}^N + \frac{1}{2} \mathcal{M}^{MN} \delta_j^i \right) \nabla_{(M} \nabla_{N)} \epsilon^j \\ &= \frac{1}{4} \mathcal{V}^{ikM} \mathcal{V}_{kj}{}^N \mathcal{R}_{MN}{}^{ab} \gamma_{ab} \epsilon^j - \frac{1}{16} \mathcal{R} \epsilon^i , \end{aligned} \quad (2.52)$$

$$\begin{aligned} \mathcal{V}^{[ijN} \mathcal{V}^{k]lM} \Omega_{ln} [\nabla_M, \nabla_N] \epsilon^n &+ 2 \Omega_{lm} \mathcal{V}^{l[iM} \mathcal{V}^{j]mN} \nabla_{(M} \nabla_{N)} \epsilon^{k]} \\ &= \frac{1}{4} \mathcal{V}^{[ijN} \mathcal{V}^{k]lM} \Omega_{ln} \mathcal{R}_{MN}{}^{ab} \gamma_{ab} \epsilon^n - \frac{1}{4} \mathcal{R}^{ijkl} \Omega_{ln} \epsilon^n . \end{aligned} \quad (2.53)$$

The combinations on the l.h.s. are such that the undetermined part $u_{kl,mn}$ of the USp(8) spin connection is projected out while the leading two-derivative terms vanish due to the

section condition (2.9). The first terms on the r.h.s. refer to the curvature of the internal spin connection (2.22) which takes the form [24]

$$\mathcal{R}_{MN}{}^{ab} = -\frac{1}{2} e^{\mu[a} e^{b]\nu} g^{\sigma\tau} \nabla_M g_{\mu\sigma} \nabla_N g_{\nu\tau} . \quad (2.54)$$

The generalized scalar curvatures \mathcal{R} and \mathcal{R}^{ijkl} in (2.53) can be evaluated using the explicit expressions for the USp(8) spin connection (2.35), (2.36), leading to

$$\begin{aligned} \mathcal{R} &= -2 \mathcal{V}_{ij}{}^M \mathcal{V}_{kl}{}^N \left(\partial_M p_N{}^{ijkl} + 4 q_M m^i p_N{}^{jklm} \right) + \frac{1}{6} \mathcal{M}^{MN} p_M{}^{ijkl} p_N{}^{ijkl} \\ &\quad + 2 \mathcal{V}_{ij}{}^M \mathcal{V}{}^{klN} p_M{}^{ijmn} p_N{}^{klmn} - \frac{16}{5} \mathcal{V}_{ij}{}^M \mathcal{V}_{kl}{}^N e^{-1} \partial_M e p_N{}^{ijkl} \\ &\quad + \frac{8}{5} \mathcal{M}^{MN} e^{-1} \partial_M \partial_N e - \frac{4}{5} \mathcal{M}^{MN} e^{-2} \partial_M e \partial_N e , \\ \mathcal{R}^{ijkl} &= \frac{1}{3} \mathcal{M}^{MN} e^{-1} \left(\partial_M (e p_N{}^{ijkl}) + 4 e q_M m^i p_N{}^{jklm} \right) \\ &\quad - 4 \mathcal{V}_{mn}{}^M e^{-1} \left(\partial_M (e p_N{}^{mn[ij]}) + 2 e p_N{}^{mnp[i} q_M p^j + 2 e q_M p^m p_N{}^{np[ij]} \right) \mathcal{V}{}^{kl]N} \\ &\quad + 2 \mathcal{V}{}^{[ijM} \mathcal{V}{}^{kl]N} e^{-1} \partial_M \partial_N e - \frac{8}{5} \mathcal{V}{}^{[ijM} \mathcal{V}{}^{kl]N} e^{-2} \partial_M e \partial_N e \\ &\quad - \frac{2}{3} \mathcal{V}_{mn}{}^M \mathcal{V}_{pq}{}^N p_M{}^{ijkl} p_N{}^{mnpq} + \frac{32}{3} \mathcal{V}_{mn}{}^{[M} \mathcal{V}_{pq}{}^{N]} p_M{}^{mnp[i} p_N{}^{jkl]q} \\ &\quad + 4 \mathcal{V}_{mn}{}^M \mathcal{V}_{pq}{}^N p_M{}^{mn[ij} p_N{}^{kl]pq} + \frac{1}{3} \mathcal{V}{}^{[ijM} \mathcal{V}{}^{kl]N} p_M{}^{mnpq} p_N{}^{mnpq} , \end{aligned} \quad (2.55)$$

in terms of the 27-bein and its derivatives. Their explicit calculation requires a number of non-trivial USp(8) identities, some of which are collected in appendix B. Together with (2.54), these curvatures appear in the Einstein and the scalar field equations, respectively. For the following, it is also useful to note the relation between the curvature components \mathcal{R} and \mathcal{R}^{ijkl} : under a non-compact $\mathfrak{e}_{6(6)}$ transformation of the form

$$\delta \mathcal{V}_M{}_{ij} = -\Sigma_{ijkl} \mathcal{V}_M{}^{kl} , \quad (2.56)$$

the scalar curvature \mathcal{R} transforms as

$$\delta \mathcal{R} = \mathcal{R}^{ijkl} \Sigma_{ijkl} + \nabla_M \mathcal{J}_\Sigma^M , \quad (2.57)$$

into the \mathcal{R}^{ijkl} curvature, up to a boundary current of weight $\lambda_{\mathcal{J}_\Sigma} = -\frac{1}{3}$. Moreover, the dependence of \mathcal{R} on the external metric is such that

$$\delta(e\mathcal{R}) = (\delta e) \mathcal{R} + \text{total derivatives} . \quad (2.58)$$

3 Supersymmetry transformations and algebra

As the main result of this section we present the supersymmetry transformation rules for all the fields of the $E_{6(6)}$ exceptional field theory and verify that their algebra consistently

closes into generalized diffeomorphisms and gauge transformations. The full set of supersymmetry transformations is given by

$$\begin{aligned}
\delta_\epsilon \psi_\mu^i &= \mathcal{D}_\mu \epsilon^i - i\sqrt{2} \mathcal{V}^{ijM} \left(\nabla_M^- (\gamma_\mu \epsilon^k) - \frac{1}{3} \gamma_\mu \nabla_M^- \epsilon^k \right) \Omega_{jk} , \\
\delta_\epsilon \chi^{ijk} &= \frac{i}{2} \mathcal{P}_\mu^{ijkl} \Omega_{lm} \gamma^\mu \epsilon^m + \frac{3}{\sqrt{2}} \mathcal{V}^{[ijM} \nabla_M^- \epsilon^{k]} , \\
\delta_\epsilon e_\mu^a &= \frac{1}{2} \bar{\epsilon}_i \gamma^a \psi_\mu^i , \quad \delta_\epsilon \mathcal{V}_M^{ij} = 4i \Omega^{im} \Omega^{jn} \mathcal{V}_M^{kl} \Omega_{p[k} \bar{\chi}_{lmn]} \epsilon^p , \\
\delta_\epsilon \mathcal{A}_\mu^M &= \sqrt{2} \left(i \Omega^{ik} \bar{\epsilon}_k \psi_\mu^j + \bar{\epsilon}_k \gamma_\mu \chi^{ijk} \right) \mathcal{V}_{ij}^M , \\
\delta_\epsilon \mathcal{B}_{\mu\nu M} &= -\frac{1}{\sqrt{5}} \mathcal{V}_M^{ij} \left(2 \bar{\psi}_{i[\mu} \gamma_{\nu]} \epsilon^k \Omega_{jk} + i \bar{\chi}_{ijk} \gamma_{\mu\nu} \epsilon^k \right) - d_{MNP} \mathcal{A}_{[\mu}^N \delta_\epsilon \mathcal{A}_{\nu]}^P ,
\end{aligned} \tag{3.1}$$

in terms of the covariant derivatives defined above. Spinor conventions were summarized in section 2.2. Upon dropping all internal derivatives $\partial_M \rightarrow 0$, these transformation rules precisely reproduce those of $D = 5$ maximal supergravity [21, 25].⁴ It is interesting to note that just as for the supersymmetric $E_{7(7)}$ theory [24], all appearance of the gauge field strength $\mathcal{F}_{\mu\nu}^M$ in the transformation rules can be absorbed into the homogeneous shift (2.23) of the internal spin connection. In the next section, we will see that the supersymmetric Lagrangian in contrast carries the opposite derivative ∇_M^+ as well.

The internal derivatives ∇_M appear in the supersymmetry transformations only in particular combinations such that the undetermined part of the $USp(8)$ connection (2.35) drops out [15, 16]. With the explicit parametrization of $\mathcal{Q}_M i^j$ from (2.36) we may explicitly evaluate these derivatives in terms of the Cartan form (2.34) of the 27-bein as

$$\begin{aligned}
\mathcal{V}_{ij}^M \nabla_M \epsilon^j &= \mathcal{V}_{ij}^M \left(\partial_M \epsilon^j - q_M k^j \epsilon^k \right) - \frac{1}{2} \mathcal{V}^{jkM} p_{Mijkn} \epsilon^n + \frac{1}{4} (2 - 3\lambda_\epsilon) \mathcal{V}_{ij}^M \Gamma_{KM}^K \epsilon^j , \\
\mathcal{V}^{[ijM} \nabla_M \epsilon^{k]} &= \mathcal{V}^{[ijM} \left(\partial_M \epsilon^{k]} - q_M l^{k]} \epsilon^l \right) + 2 \mathcal{V}_{mn}^M \Omega_{pr} \Omega^m [i p_M^{jk]} n^p \epsilon^r \\
&\quad + \frac{1}{2} \mathcal{V}_{mn}^M p_M^{mn [ij} \epsilon^{k]} + \frac{1}{8} (1 - 6\lambda_\epsilon) \Gamma_{KM}^K \mathcal{V}^{[ijM} \epsilon^{k]} ,
\end{aligned} \tag{3.2}$$

where we have suppressed all ω_M contributions (which enter canonically), and used (B.4) to simplify the expression in the second line.

The algebra of the supersymmetry transformations closes on the $(1+4)$ -dimensional general coordinate transformations (2.16), generalized internal diffeomorphisms (2.10), covariant gauge transformations of the p -form fields (2.13), local $SO(1,4)$ and $USp(8)$ rotations, and an additional supersymmetry transformation, higher order in the fermions. The structural form of the supersymmetry algebra is the same as for the five-dimensional theory [21]

$$\begin{aligned}
[\delta(\epsilon_1), \delta(\epsilon_2)] &= \xi^\mu \mathcal{D}_\mu + \delta_{so(1,4)}(\Omega^{ab}) + \delta_{usp(8)}(\Lambda^{ij}) + \delta_{susy}(\epsilon_3) \\
&\quad + \delta_{\text{gauge}}(\Lambda^M) + \delta_{\text{gauge}}(\Xi_\mu M) + \delta_{\text{gauge}}(\Xi_{\mu\nu} \alpha) + \delta_{\text{gauge}}(\Xi_{\mu\nu} M) .
\end{aligned} \tag{3.3}$$

⁴To be precise, we note the rescaling of gauge and tensor fields $\mathcal{A}_\mu^M [1312.0614] = \frac{1}{\sqrt{2}} A_\mu^M [hep-th/0412173]$, $\mathcal{B}_{\mu\nu M} [1312.0614] = -\frac{1}{4} B_{\mu\nu M} [hep-th/0412173]$ together with rescaling of the associated symmetry parameters, in order to translate the notation from [21] into [3]. In this paper, we will stick to the conventions of [3] for the normalization of the gauge fields.

The transformation parameters on the r.h.s. can be explicitly given as combinations of the spinors $\epsilon_{1,2}$, their covariant derivatives, and the external and internal vielbeins e_μ^a , \mathcal{V}_M^{ij} , as

$$\begin{aligned}
 \xi^\mu &= \frac{1}{2} \bar{\epsilon}_{2i} \gamma^\mu \epsilon_1^i, \\
 \Omega^{ab} &= -\frac{\sqrt{2}i}{3} \left(\bar{\epsilon}_{1i} \gamma^{ab} \nabla_M^- \epsilon_2^k - \nabla_M^- \bar{\epsilon}_{1i} \gamma^{ab} \epsilon_2^k \right) \mathcal{V}^{ijM} \Omega_{jk} - \Lambda^M \omega_M^{-ab}, \\
 \Lambda^M &= -\sqrt{2}i \mathcal{V}^{ijM} \Omega_{jk} \bar{\epsilon}_{2i} \epsilon_1^k, \\
 \Xi_{\mu M} &= \frac{1}{\sqrt{5}} \mathcal{V}_M^{kl} \Omega_{lm} (\bar{\epsilon}_{2k} \gamma_\mu \epsilon_1^m), \\
 \Xi_{\mu\nu\alpha} &= \frac{3i}{\sqrt{10}} (t_\alpha)^M{}_N \mathcal{V}_M{}_{li} \mathcal{V}^{kiN} (\bar{\epsilon}_{2k} \gamma_{\mu\nu} \epsilon_1^l), \\
 \Xi_{\mu\nu M} &= -\frac{i}{\sqrt{10}} \left(\bar{\epsilon}_{2k} \gamma_{\mu\nu} \partial_M \epsilon_1^k - \partial_M \bar{\epsilon}_{2k} \gamma_{\mu\nu} \epsilon_1^k - (\bar{\epsilon}_{2k} \epsilon_1^k) e^a{}_{[\mu} \partial_M e_{\nu]a} \right. \\
 &\quad \left. - \frac{2}{3} \mathcal{V}^{kiN} \partial_M \mathcal{V}_{Nli} (\bar{\epsilon}_{2k} \gamma_{\mu\nu} \epsilon_1^l) \right).
 \end{aligned} \tag{3.4}$$

In the rest of this section we provide the explicit calculations that show closure of the supersymmetry algebra (3.3), (3.4), thereby confirming the supersymmetry transformation laws (3.1). Let us start with closure on the external vielbein e_μ^a

$$\begin{aligned}
 [\delta_{\epsilon_1}, \delta_{\epsilon_2}] e_\mu^a &= \frac{1}{2} \bar{\epsilon}_{2i} \gamma^a \mathcal{D}_\mu \epsilon_1^i - \frac{i}{\sqrt{2}} \bar{\epsilon}_{2i} \mathcal{V}_M^{ij} \gamma^a \left(\nabla_M^- (\gamma_\mu \epsilon_2^k) - \frac{1}{3} \gamma_\mu \nabla_M^- \epsilon_1^k \right) \Omega_{jk} - (1 \leftrightarrow 2) \\
 &= \frac{1}{2} \mathcal{D}_\mu (\bar{\epsilon}_{2i} \gamma^\nu \epsilon_1^i e_\nu^a) - \sqrt{2}i \left(\bar{\epsilon}_{2i} \epsilon_1^k \mathcal{V}^{ijM} \Omega_{jk} \right) \nabla_M^- e_\mu^a \\
 &\quad - \frac{\sqrt{2}i}{3} \left(\bar{\epsilon}_{2i} \gamma^{ab} \nabla_M^- \epsilon_1^k - \nabla_M^- \bar{\epsilon}_{2i} \gamma^{ab} \epsilon_1^k \right) \mathcal{V}^{ijM} \Omega_{jk} e_{\mu b} \\
 &\quad - \frac{\sqrt{2}i}{3} \nabla_M^- \left(\bar{\epsilon}_{2i} \epsilon_1^k \right) \mathcal{V}^{ijM} \Omega_{jk} e_\mu^a.
 \end{aligned} \tag{3.5}$$

Taking into account that the term $\bar{\epsilon}_{2i} \epsilon_1^k$ has all spinor indices contracted, the generalized vielbein postulate (2.32), and the vanishing torsion (2.21), we may rewrite the above expression as follows

$$[\delta_{\epsilon_1}, \delta_{\epsilon_2}] e_\mu^a = e_\nu^a \mathcal{D}_\mu \xi^\nu + \xi^\nu \mathcal{D}_\nu e_\mu^a + \Lambda^N \partial_N e_\mu^a + \frac{1}{3} \partial_N \Lambda^N e_\mu^a + \Omega^{ab} e_{\mu b}, \tag{3.6}$$

reproducing the correct transformation under external and internal diffeomorphisms. In particular, we obtain the correct value $\lambda = 1/3$ for the weight of the fünfbein.

Next we check closure of the supersymmetry on the generalized vielbein \mathcal{V}_M^{ij} . We directly project the variation onto its coset valued part, since any remaining part can be absorbed into a local USp(8) transformation. The result is

$$\begin{aligned}
 \mathcal{V}^{[klM} [\delta_{\epsilon_1}, \delta_{\epsilon_2}] \mathcal{V}_M^{ij}] &= 2 \mathcal{P}_\mu{}^m [^{ijk} \Omega^l]{}^p \Omega_{mn} (\bar{\epsilon}_{2p} \gamma^\mu \epsilon_1^n) + 6\sqrt{2}i \mathcal{V}^{[klM} \Omega^j]{}^p \bar{\epsilon}_{2p} \nabla_M^- \epsilon_1^i - (1 \leftrightarrow 2) \\
 &= \xi^\mu \mathcal{P}_\mu{}^{klj} + 6 \mathcal{V}^{[klM} \nabla_M (\mathcal{V}_N^{ij}] \Lambda^N),
 \end{aligned} \tag{3.7}$$

where we used the identity $\mathcal{P}_\mu^{[ijkl]\Omega^{mn}} = 0$ and the vielbein postulate. The first term in the expression above gives just a (covariantized) diffeomorphism along ξ^μ , while the second can be rewritten using the generalised vanishing torsion condition (2.33) which gives

$$\begin{aligned}
 & \mathcal{V}^{[klM} [\delta_{\epsilon_1}, \delta_{\epsilon_2}] \mathcal{V}_M^{ij}] \\
 &= \xi^\mu \mathcal{P}_\mu^{kl ij} + 6 \mathcal{V}^{[klM} \mathcal{D}_M (\mathcal{V}_N^{ij}) \Lambda^N + 6 \mathcal{V}^{[klM} \mathcal{V}_N^{ij]} \left(\partial_M \Lambda^N - \frac{1}{4} \Lambda^N \Gamma_{KM}^K \right) \\
 &= \xi^\mu \mathcal{P}_\mu^{kl ij} + (\mathcal{V}^{[klM} \mathcal{D}_K \mathcal{V}_M^{ij}]) \Lambda^N + 6 \mathcal{V}^{[klM} \mathcal{V}_N^{ij]} \partial_M \Lambda^N \\
 &= \xi^\mu \mathcal{P}_\mu^{kl ij} + \mathcal{V}^{[klM} \delta_\Lambda \mathcal{V}_M^{ij]} .
 \end{aligned} \tag{3.8}$$

The weight term that comes from the derivative of $\mathcal{V}_N^{ij} \Lambda^N$ is cancelled by the same contribution from the vanishing torsion condition. Again, we find the correct transformation with the same gauge parameters as in (3.6).

Now we turn to the gauge field sector and investigate closure of the supersymmetry algebra on the vector field \mathcal{A}_μ^M . A direct calculation gives

$$\begin{aligned}
 [\delta_{\epsilon_1}, \delta_{\epsilon_2}] \mathcal{A}_\mu^M &= \sqrt{2} i \Omega^{ik} (\bar{\epsilon}_{2k} \mathcal{D}_\mu \epsilon_1^j) \mathcal{V}_{ij}^M + \sqrt{2} i \bar{\epsilon}_{2k} \gamma_\mu \mathcal{P}_\nu^{ijkl} \gamma^\nu \Omega_{lm} \epsilon_1^m \mathcal{V}_{ij}^M \\
 &\quad + 3 \bar{\epsilon}_{2k} \gamma_\mu \left(\Omega^{m[i} \mathcal{V}^{jk]N} - \frac{1}{3} \Omega^{[ij} \mathcal{V}^{k]mN} \right) \Omega_{mr} \nabla_N^- \epsilon_1^r \mathcal{V}_{ij}^M \\
 &\quad + 2 \Omega^{ik} \bar{\epsilon}_{2k} \mathcal{V}^{jrN} \left(\nabla_N^- (\gamma_\mu \epsilon_1^s) - \frac{1}{3} \gamma_\mu \nabla_N^- \epsilon_1^s \right) \Omega_{rs} \mathcal{V}_{ij}^M - (1 \leftrightarrow 2) \\
 &= \mathcal{D}_\mu \Lambda^M + \frac{1}{2} \partial_N (\bar{\epsilon}_{2k} \gamma_\mu \epsilon_1^k) \mathcal{M}^{MN} - (\bar{\epsilon}_{2k} \gamma^\nu \epsilon_1^k) \mathcal{M}^{MN} e_{\nu a} \nabla_N^- e_\mu^a \\
 &\quad + 2 \left(\mathcal{V}^{Mik} \mathcal{V}_{ij}^N + \mathcal{V}^{Nik} \mathcal{V}_{ij}^M - \frac{1}{4} \delta_j^k \mathcal{M}^{MN} \right) \nabla_N^- (\bar{\epsilon}_{2k} \gamma_\mu \epsilon_1^j) \\
 &= \mathcal{D}_\mu \Lambda^M + g_{\mu\nu} \partial_N \xi^\nu \mathcal{M}^{MN} - 2 \xi^\nu \mathcal{M}^{MN} e_{a[\nu} \nabla_N^- e_{\mu]}^a \\
 &\quad - 2 \nabla_N^- \left[\left(\mathcal{V}^{ikM} \mathcal{V}_{ij}^N + \mathcal{V}^{ikN} \mathcal{V}_{ij}^M - \frac{1}{4} \delta_j^k \mathcal{M}^{MN} \right) (\bar{\epsilon}_{2k} \gamma_\mu \epsilon_1^j) \right] .
 \end{aligned} \tag{3.9}$$

Finally using the relations (2.8) and (2.22), the above expression can be written in the following form

$$\begin{aligned}
 [\delta_{\epsilon_1}, \delta_{\epsilon_2}] \mathcal{A}_\mu^M &= g_{\mu\nu} \partial_N \xi^\nu \mathcal{M}^{MN} - 2 \xi^\nu \mathcal{M}^{MN} e_{a[\nu} \nabla_N^- e_{\mu]}^a \\
 &\quad + \mathcal{D}_\mu \Lambda^M - 10 d^{MNK} \partial_N \Xi_{\mu K} \\
 &= \xi^\nu \mathcal{F}_{\nu\mu}^M + g_{\mu\nu} \partial_N \xi^\nu \mathcal{M}^{MN} + \mathcal{D}_\mu \Lambda^M - 10 d^{MNK} \partial_N \Xi_{\mu K},
 \end{aligned} \tag{3.10}$$

with the parameter $\Xi_{\mu M}$ from (3.4), thus precisely reproducing the $E_{6(6)}$ covariant gauge transformation (2.13) of the gauge field coming from tensor hierarchy. The first two terms in the expression correspond to the transformation (2.16) of the gauge field under external diffeomorphisms.

Finally, we investigate transformations of the two-form field $\mathcal{B}_{\mu\nu M}$ that give

$$\begin{aligned}
 [\delta_{\epsilon_1}, \delta_{\epsilon_2}] \mathcal{B}_{\mu\nu M} &= -\frac{1}{\sqrt{5}} \left[2 \mathcal{V}_M^{ij} \bar{\epsilon}_{2i} \gamma_{[\mu} \mathcal{D}_{\nu]} \epsilon_1^k \Omega_{jk} + \frac{1}{2} \mathcal{V}_M^{lm} \bar{\epsilon}_{2n} \gamma_{\mu\nu} \gamma^\rho \mathcal{P}_\rho^{lmnp} \Omega_{pq} \epsilon_1^q \right] \\
 &\quad - \frac{4i}{\sqrt{10}} \mathcal{V}_M^{ij} \mathcal{V}_{jp}{}^N \left(\bar{\epsilon}_{2i} \gamma_{[\mu} \nabla_N^- (\gamma_{\nu]} \epsilon_1^p) - \frac{1}{3} \bar{\epsilon}_{2i} \gamma_{\mu\nu} \nabla_N^- \epsilon_1^p \right) \\
 &\quad - \frac{3i}{\sqrt{10}} \mathcal{V}_M^{lm} \bar{\epsilon}_{2n} \gamma_{\mu\nu} \left(\Omega^{k[l} \mathcal{V}^{mn]N} - \frac{1}{3} \Omega^{[lm} \mathcal{V}^{n]kN} \right) \Omega_{kp} \nabla_N^- \epsilon_1^p - (1 \leftrightarrow 2) \\
 &= \frac{1}{\sqrt{5}} \mathcal{D}_{[\mu} \left(\mathcal{V}_M^{ij} \bar{\epsilon}_{2i} \gamma_{\nu]} \epsilon_1^k \Omega_{jk} \right) - \frac{4i}{\sqrt{10}} \mathcal{V}_{jl}{}^N \mathcal{V}^{kjM} \left(\bar{\epsilon}_{2k} \gamma_a \gamma_b \epsilon_1^l \right) e_{[\mu}{}^a \nabla_N^- e_{\nu]}{}^b \\
 &\quad - \frac{2i}{\sqrt{10}} \left(\mathcal{V}_M{}^{ni} \mathcal{V}^{kiN} - \mathcal{V}_{ni}{}^N \mathcal{V}_M{}^{ki} \right) \left(\bar{\epsilon}_{2k} \gamma_{\mu\nu} \nabla_N^- \epsilon_1^n \right) + \frac{i}{\sqrt{10}} \left(\bar{\epsilon}_{2k} \gamma_{\mu\nu} \nabla_M^- \epsilon_1^k \right) \\
 &\quad - (1 \leftrightarrow 2) \\
 &= 2 \mathcal{D}_{[\mu} \Xi_{\nu]M} - \frac{2i}{\sqrt{10}} \left(\mathcal{V}_M{}^{ni} \mathcal{V}^{kiN} - \mathcal{V}_{ni}{}^N \mathcal{V}_M{}^{ki} \right) \nabla_N^- \left(\bar{\epsilon}_{2k} \gamma_{\mu\nu} \epsilon_1^n \right) \\
 &\quad - \frac{4i}{\sqrt{10}} \left(\mathcal{V}_M{}^{ni} \mathcal{V}^{kiN} - \mathcal{V}_{ni}{}^N \mathcal{V}_M{}^{ki} \right) \left(\bar{\epsilon}_{2k} \epsilon_1^n \right) e_{[\mu}{}^a \nabla_N^- e_{\nu]a} \\
 &\quad + \frac{i}{\sqrt{10}} \left(\bar{\epsilon}_{2k} \gamma_{\mu\nu} \nabla_M^- \epsilon_1^k - \nabla_M^- \bar{\epsilon}_{2k} \gamma_{\mu\nu} \epsilon_1^k \right). \tag{3.11}
 \end{aligned}$$

Here, we have systematically ignored the contribution from the last term $d_{MNP} \mathcal{A}_{[\mu}{}^N \delta \mathcal{A}_{\nu]}{}^P$ in the supersymmetry variation, which will simply reproduce the corresponding terms in the action of generalized diffeomorphisms and gauge transformations, due to the fact that the algebra closes on the vector field $\mathcal{A}_\mu{}^M$. To simplify the second term in (3.11) it is helpful to consider the following identity

$$\begin{aligned}
 &\left(\mathcal{V}_M{}^{ni} \mathcal{V}^{kiN} - \mathcal{V}_{ni}{}^N \mathcal{V}_M{}^{ki} \right) \left(\bar{\epsilon}_{2k} \gamma_{\mu\nu} \epsilon_1^n \right) \\
 &= \left(\mathcal{V}_M{}^{ni} \mathcal{V}^{kiN} \Omega_{mk} + \mathcal{V}_M{}^{mi} \mathcal{V}^{kiN} \Omega_{nk} \right) \Omega^{mp} \left(\bar{\epsilon}_{2p} \gamma_{\mu\nu} \epsilon_1^n \right) \\
 &= \frac{3}{2} \mathbb{P}_M{}^N \mathcal{Q}{}^P \mathcal{V}_{Pni} \mathcal{V}^{kiQ} \left(\bar{\epsilon}_{2k} \gamma_{\mu\nu} \epsilon_1^n \right), \tag{3.12}
 \end{aligned}$$

where in the second line we notice that the expression in brackets is symmetric in (mn) and hence is an element of the $\mathfrak{usp}(8)$ part of $\mathfrak{e}_{6(6)}$. The traceless antisymmetrisation on the r.h.s. can be replaced by the usual antisymmetrization giving the same result. Using this relation and the vanishing torsion condition we may express the corresponding term as follows

$$\begin{aligned}
 &\nabla_N \left[\left(\mathcal{V}_M{}^{ni} \mathcal{V}^{kiN} - \mathcal{V}_{ni}{}^N \mathcal{V}_M{}^{ki} \right) \left(\bar{\epsilon}_{2k} \gamma_{\mu\nu} \epsilon_1^n \right) \right] \\
 &= \partial_N \left[\left(\mathcal{V}_M{}^{ni} \mathcal{V}^{kiN} - \mathcal{V}_{ni}{}^N \mathcal{V}_M{}^{ki} \right) \left(\bar{\epsilon}_{2k} \gamma_{\mu\nu} \epsilon_1^n \right) \right] - \frac{1}{3} \mathcal{V}_{Nni} \mathcal{D}_M \mathcal{V}^{kiN} \left(\bar{\epsilon}_{2k} \gamma_{\mu\nu} \epsilon_1^n \right) \\
 &= \frac{\sqrt{10}i}{2} (t_\alpha)_M{}^N \partial_N \Xi_{\mu\nu}{}^\alpha - \frac{1}{3} \mathcal{V}_{Nni} \mathcal{D}_M \mathcal{V}^{kiN} \left(\bar{\epsilon}_{2k} \gamma_{\mu\nu} \epsilon_1^n \right), \tag{3.13}
 \end{aligned}$$

with $\Xi_{\mu\nu}{}^\alpha$ from (3.4). Next, using the identity (2.8), the second line of the last equation

in (3.11) can be rewritten in the following suggestive form

$$\begin{aligned}
 & -\frac{4i}{\sqrt{10}} \left(\mathcal{V}_{Mni} \mathcal{V}^{kiN} + \mathcal{V}_{ni}{}^N \mathcal{V}_M{}^{ki} \right) (\bar{\epsilon}_{2k} \epsilon_1^n) e_{[\mu}{}^a \nabla_N^- e_{\nu]a} \\
 & = d_{MKN} \Lambda^N \mathcal{F}_{\mu\nu}{}^K - \frac{i}{\sqrt{10}} (\bar{\epsilon}_{2k} \epsilon_1^k) e_{[\mu}{}^a \nabla_M^- e_{\nu]a}.
 \end{aligned} \tag{3.14}$$

Finally, we focus on the last term in the last equation of (3.11) and notice that its USp(8) connection part cancels that of the last term in (3.13). Hence, we may take into account only the spin connection $\omega_M{}^{\rho\sigma}$ that includes the SO(1,4) connection and the field strength $\mathcal{F}^{\rho\sigma M}$. After some gamma-matrices algebra we obtain the following expression

$$\begin{aligned}
 \bar{\epsilon}_{2k} \gamma_{\mu\nu} \nabla_M^- \epsilon_1^k - \nabla_M^- \bar{\epsilon}_{2k} \gamma_{\mu\nu} \epsilon_1^k & = -\frac{1}{4} \mathcal{M}_{MN} (\bar{\epsilon}_{2k} \gamma_{\mu\nu\rho\sigma} \epsilon_1^k) \mathcal{F}^{\rho\sigma N} + (\bar{\epsilon}_{2k} \epsilon_1^k) e_{[\mu}{}^a \nabla_M^- e_{\nu]a} \\
 & \quad + \bar{\epsilon}_{2k} \gamma_{\mu\nu} \partial_M \epsilon_1^k - \partial_M \bar{\epsilon}_{2k} \gamma_{\mu\nu} \epsilon_1^k - (\bar{\epsilon}_{2k} \epsilon_1^k) e_{[\mu}{}^a \partial_M e_{\nu]a} \\
 & = -\frac{i}{2} \xi^\lambda e \varepsilon_{\lambda\mu\nu\rho\sigma} \mathcal{M}_{MN} \mathcal{F}^{\rho\sigma N} + (\bar{\epsilon}_{2k} \epsilon_1^k) e_{[\mu}{}^a \nabla_M^- e_{\nu]a} \\
 & \quad + \bar{\epsilon}_{2k} \gamma_{\mu\nu} \partial_M \epsilon_1^k - \partial_M \bar{\epsilon}_{2k} \gamma_{\mu\nu} \epsilon_1^k - (\bar{\epsilon}_{2k} \epsilon_1^k) e_{[\mu}{}^a \partial_M e_{\nu]a}.
 \end{aligned} \tag{3.15}$$

The first term here represents the diffeomorphism transformation (2.16) of the field $\mathcal{B}_{\mu\nu}$, the second term precisely cancels the last term in (3.14). The rest can be packaged into a tensor $\mathcal{O}_{M\mu\nu}$ constrained by

$$d^{MNK} \partial_N \mathcal{O}_{K\mu\nu} = 0, \tag{3.16}$$

as a consequence of the section condition. Collecting everything together, the commutator of supersymmetry transformations (3.11) of the two-form field takes the following form

$$\begin{aligned}
 [\delta_{\epsilon_1}, \delta_{\epsilon_2}] \mathcal{B}_{\mu\nu M} & = 2 \mathcal{D}_{[\mu} \bar{\Xi}_{\nu]M} + \frac{1}{2\sqrt{10}} \xi^\lambda e \varepsilon_{\lambda\mu\nu\rho\sigma} \mathcal{M}_{MN} \mathcal{F}^{\rho\sigma N} + (t_\alpha)_{M^N} \partial_N \bar{\Xi}_{\mu\nu}{}^\alpha \\
 & \quad + d_{MKN} \Lambda^N \mathcal{F}^K{}_{\mu\nu} + \mathcal{O}_{M\mu\nu} - d_{MKL} \mathcal{A}_{[\mu}{}^K [\delta_{\epsilon_1}, \delta_{\epsilon_2}] \mathcal{A}_{\nu]}{}^L,
 \end{aligned} \tag{3.17}$$

up to terms of higher order in the fermions. This confirms the supersymmetry algebra (3.3), (3.4).

4 Invariant Lagrangian

We now have all the ingredients to present the full supersymmetric Lagrangian for $E_{6(6)}$ exceptional field theory. Its bosonic part has been constructed in [2, 3], here we give the supersymmetric extension based on the fermionic structures introduced in the previous sections. The final result reads

$$\begin{aligned}
 e^{-1} \mathcal{L} & = \widehat{\mathcal{R}} - \frac{1}{4} \mathcal{M}_{MN} \mathcal{F}_{\mu\nu}{}^M \mathcal{F}^{\mu\nu N} - \frac{1}{6} \mathcal{P}_\mu{}^{ijkl} \mathcal{P}^\mu{}_{ijkl} + \frac{\sqrt{10}}{8} e^{-1} \mathcal{L}_{\text{top}} - V(\mathcal{M}, g) \\
 & \quad - \bar{\psi}_{\mu i} \gamma^{\mu\nu\rho} \mathcal{D}_\nu \psi_\rho^i + 2\sqrt{2} i \mathcal{V}_{ij}{}^M \Omega^{ik} \bar{\psi}_{\mu k} \gamma^{[\mu} \nabla_M^+ (\gamma^{\nu]} \psi_\nu^j) \\
 & \quad - \frac{4}{3} \bar{\chi}_{ijk} \gamma^\mu \mathcal{D}_\mu \chi^{ijk} + 8\sqrt{2} i \mathcal{V}_{mn}{}^M \Omega^{np} \bar{\chi}_{pkl} \nabla_M^+ \chi^{mkl} \\
 & \quad + \frac{4i}{3} \mathcal{P}_\mu{}^{ijkl} \bar{\chi}_{ijk} \gamma^\nu \gamma^\mu \psi_\nu{}^m \Omega_{lm} + 4\sqrt{2} \mathcal{V}^{ijM} \bar{\chi}_{ijk} \gamma^\mu \nabla_M^- \psi_\mu{}^k,
 \end{aligned} \tag{4.1}$$

up to quartic fermion terms. The latter are expected to coincide with the quartic terms of the $D = 5$ theory [25]. Let us explain the various terms of (4.1). The first line describes the bosonic couplings, with the modified Ricci scalar $\widehat{\mathcal{R}}$ obtained from contracting (2.43), Yang-Mills term for the field strength (2.12) and the scalar kinetic term

$$-\frac{1}{6} \mathcal{P}_\mu^{ijkl} \mathcal{P}^\mu_{ijkl} = \frac{1}{24} \mathcal{D}_\mu \mathcal{M}_{MN} \mathcal{D}^\mu \mathcal{M}^{MN}. \quad (4.2)$$

We note, that variation of the Einstein-Hilbert term and the scalar kinetic term w.r.t. the vector fields is given by

$$\delta \left(e \widehat{\mathcal{R}} - \frac{1}{6} e \mathcal{P}_\mu^{ijkl} \mathcal{P}^\mu_{ijkl} \right) = e \left(\widehat{\mathcal{J}}^\mu_M + \mathcal{J}^\mu_M \right) \delta \mathcal{A}_\mu^M, \quad (4.3)$$

with the current $\widehat{\mathcal{J}}^\mu_M$ from (2.50) and the scalar current given by

$$\mathcal{J}^\mu_M = -2 \mathcal{V}_{Mij} \mathcal{V}_{kl}^N \nabla_N \left(g^{\mu\nu} \mathcal{P}_\nu^{ijkl} \right). \quad (4.4)$$

The topological term in (4.1) is most compactly defined by its variation

$$\begin{aligned} \delta \mathcal{L}_{\text{top}} = & \varepsilon^{\mu\nu\rho\sigma\tau} \left(d_{MKN} \mathcal{F}_{\mu\nu}^M \mathcal{F}_{\rho\sigma}^N \delta \mathcal{A}_\tau^K \right. \\ & \left. + \frac{20}{3} d^{MKN} \partial_N \mathcal{H}_{\mu\nu\rho M} \left(\delta \mathcal{B}_{\sigma\tau K} + d_{K PQ} \mathcal{A}_\sigma^P \delta \mathcal{A}_\tau^Q \right) \right), \end{aligned} \quad (4.5)$$

equivalently, the associated action can be expressed as the boundary contribution of a manifestly covariant integral over six external dimensions. The scalar potential V has been given in [2, 3] in the explicit form

$$\begin{aligned} V(\mathcal{M}, g) = & -\frac{1}{24} \mathcal{M}^{MN} \partial_M \mathcal{M}^{KL} \partial_N \mathcal{M}_{KL} + \frac{1}{2} \mathcal{M}^{MN} \partial_M \mathcal{M}^{KL} \partial_L \mathcal{M}_{NK} \\ & - \frac{1}{2} g^{-1} \partial_M g \partial_N \mathcal{M}^{MN} - \frac{1}{4} \mathcal{M}^{MN} g^{-1} \partial_M g g^{-1} \partial_N g - \frac{1}{4} \mathcal{M}^{MN} \partial_M g^{\mu\nu} \partial_N g_{\mu\nu}, \end{aligned} \quad (4.6)$$

and can be rewritten in the following manifestly covariant form

$$V(\mathcal{M}, g) = \mathcal{R} - \frac{1}{4} \mathcal{M}^{MN} \nabla_M g_{\mu\nu} \nabla_N g^{\mu\nu} + \nabla_M I^M, \quad (4.7)$$

with the curvature scalar \mathcal{R} from (2.55), up to boundary contributions I^M and terms that vanish due to the section condition. The explicit calculation confirming (4.7) requires a number of non-trivial $\text{USp}(8)$ identities, some of which are collected in appendix B.

The kinetic fermion terms in (4.1) are such that upon dropping all internal derivatives, the Lagrangian $\mathcal{L}_0 \equiv \mathcal{L}|_{\partial_M \rightarrow 0}$ reduces to the five-dimensional theory [21, 25]. The fermion terms carrying internal derivatives ∇_M are then obtained by imposing invariance of the Lagrangian under the supersymmetry transformations (3.1).⁵ In the limit $\partial_M \rightarrow 0$, these terms reduce to the Pauli couplings of fermions to the field strength via (2.23) and again reproduce the couplings from the $D = 5$ theory. It is interesting to observe that in the

⁵See also [16] for these couplings in a $\text{Cliff}(10, 1; \mathbb{R})$ formulation.

full theory, and unlike for the supersymmetry transformations (3.1), these $\mathcal{F}_{\mu\nu}^M$ couplings cannot entirely be absorbed into a homogeneous shift of the internal spin connection (2.23), but require both ∇_M^+ and ∇_M^- derivatives, however in a very systematic pattern.

By construction, the full Lagrangian (4.1) is manifestly invariant under generalized internal diffeomorphisms. To show that it is invariant under supersymmetry, one has to go through rather tedious calculations, that we sketch in the remainder of this section. For the full detailed calculations the reader is referred to appendix A. The proof of supersymmetry of the Lagrangian is most conveniently organized order by order in the internal derivatives ∇_M .⁶ Internal derivatives enter in \mathcal{L} in two different ways: first they render the Lagrangian \mathcal{L}_0 covariant under generalized diffeomorphisms by virtue of (2.11) and (2.12), second they give rise to explicit couplings such as the bilinear fermion terms and the scalar potential V . I.e. the Lagrangian schematically organizes as

$$\mathcal{L} = \mathcal{L}_0^{\text{cov}} + \mathcal{L}_1[\bar{\psi}\nabla_M\psi] + \mathcal{L}_2[\nabla_M\mathcal{M}\nabla_N\mathcal{M}]. \quad (4.8)$$

Similarly, the supersymmetry transformations (3.1) organize as

$$\delta = \delta_0^{\text{cov}} + \delta_1[\nabla_M\epsilon], \quad (4.9)$$

where δ_0 describe the supersymmetry transformation laws of the five-dimensional theory. In lowest order in ∇_M , supersymmetry of the Lagrangian amounts to the corresponding property of the five-dimensional theory [21, 25]. In first and second order in ∇_M , the contributions from $\delta_0^{\text{cov}}\mathcal{L}_1$, $\delta_1\mathcal{L}_0^{\text{cov}}$, and $\delta_1\mathcal{L}_1$ can be organized according to their fermion structure

$$\psi\mathcal{D}_\mu\nabla_M\epsilon, \quad \chi\mathcal{D}_\mu\nabla_M\epsilon, \quad \psi\nabla_M\nabla_N\epsilon, \quad \chi\nabla_M\nabla_N\epsilon, \quad (4.10)$$

and we discuss the four classes of terms separately in appendices A.1–A.4. The latter terms combine with the second order contributions from $\delta_0^{\text{cov}}\mathcal{L}_2$ arising from variation of the scalar potential (4.7). These are obtained by using the properties of the scalar curvature (2.57), (2.58) as

$$\begin{aligned} \delta_\epsilon(eV) = & \frac{1}{2}e\left(g^{\mu\nu}\mathcal{R} - \frac{1}{4}g^{\mu\nu}\mathcal{M}^{MN}\nabla_Mg^{\mu\nu}\nabla_Ng_{\mu\nu} + \nabla_N(\mathcal{M}^{MN}\nabla_Mg^{\mu\nu})\right)\delta_\epsilon g_{\mu\nu} \\ & + e\Sigma_{ijkl}^\epsilon\left(\mathcal{R}^{ijkl} - \frac{1}{2}\mathcal{V}^{ijM}\mathcal{V}^{klN}\nabla_Mg_{\mu\nu}\nabla_Ng^{\mu\nu}\right), \end{aligned} \quad (4.11)$$

up to total derivatives, and with $\Sigma_{ijkl}^\epsilon \equiv -4i\Omega_{m[i}\bar{\chi}_{jkl]}\epsilon^m$ describing the supersymmetry variation of the scalar fields (3.1).

In addition, we have further contributions from $\delta_0^{\text{cov}}\mathcal{L}_0^{\text{cov}}$ due to the fact that covariant derivatives \mathcal{D}_μ no longer commute. Such contributions arise from variation of the fermionic kinetic term with (2.45)

$$-\frac{4i}{3}\bar{\chi}_{ijk}\gamma^{\mu\nu}\epsilon^m\mathcal{D}_\mu\mathcal{P}_\nu{}^{ijkl}\Omega_{lm} = 4i\bar{\chi}_{ijk}\gamma^{\mu\nu}\epsilon^m\Omega_{lm}\mathcal{V}_N{}^{[ij}\mathcal{V}^{kl]M}\nabla_M\mathcal{F}_{\mu\nu}{}^N, \quad (4.12)$$

⁶This is very much in parallel with the analogous calculation in gauged supergravity [21] order by order in the coupling constant.

but also from variation of the Rarita-Schwinger term upon using the commutator (2.42)

$$\begin{aligned}
& \bar{\psi}_{\mu i} \gamma^{\mu\nu\rho} [\mathcal{D}_\nu, \mathcal{D}_\rho] \epsilon^i \\
&= \bar{\psi}_{\mu i} \gamma_\nu \epsilon^i \left(\widehat{\mathcal{R}}^{\nu\mu} + \frac{1}{2} g^{\mu\nu} \widehat{\mathcal{R}} \right) + \frac{2}{3} \mathcal{P}_\nu{}^{iklm} \mathcal{P}_{\rho jklm} \bar{\psi}_{\mu i} \gamma^{\mu\nu\rho} \epsilon^j \\
&\quad - \mathcal{F}_{\nu\rho}{}^M \bar{\psi}_{\mu i} \gamma^{\mu\nu\rho} \nabla_M \epsilon^i + \nabla_M \mathcal{F}_{\nu\rho}{}^N \left(\mathcal{V}_N{}^{jk} \mathcal{V}_{ik}{}^M - \mathcal{V}_N{}_{ik} \mathcal{V}^{jkM} \right) \bar{\psi}_{\mu i} \gamma^{\mu\nu\rho} \epsilon^j \\
&\quad - \frac{1}{2} \bar{\psi}_{\mu i} \gamma^{\mu\rho}{}_\nu \epsilon^i \nabla_M \mathcal{F}_\rho{}^{\nu M} + \frac{1}{4} \bar{\psi}_{\mu i} \gamma^{\nu\rho\sigma} \epsilon^i \mathcal{F}_{\nu\rho}{}^K g^{\mu\tau} \nabla_K g_{\sigma\tau} .
\end{aligned} \tag{4.13}$$

Here the first two terms cancel as in the $D = 5$ theory (where it is important though that $\widehat{\mathcal{R}}^{\mu\nu}$ arises with indices contracted in the proper order since $\widehat{\mathcal{R}}^{[\mu\nu]} \neq 0$), while all remaining terms cancel against terms of the form (4.10) as discussed in appendix A.3, A.4.

Finally, there are the contributions that arise from variation of the vector gauge field in the minimal couplings of (2.11), and from variation of the two-form gauge field in the vector kinetic term and the topological term. The first appear proportional to the currents from (4.3)

$$\delta_A \left(e \widehat{\mathcal{R}} - \frac{1}{6} e \mathcal{P}_\mu{}^{ijkl} \mathcal{P}^\mu{}_{ijkl} \right) = e \left(\widehat{\mathcal{J}}^\mu{}_M + \mathcal{J}^\mu{}_M \right) \delta \mathcal{A}_\mu{}^M , \tag{4.14}$$

and the latter are proportional to the first order duality equation between vectors and tensors

$$\delta_B \mathcal{L} = 5 d^{MKN} \nabla_N \left(e \mathcal{M}_{MN} \mathcal{F}^{\mu\nu N} + \frac{\sqrt{10}}{6} \varepsilon^{\mu\nu\rho\sigma\tau} \mathcal{H}_{\rho\sigma\tau M} \right) \left(\delta_\epsilon \mathcal{B}_{\mu\nu K} + d_{KPQ} \mathcal{A}_\mu{}^P \delta_\epsilon \mathcal{A}_\nu{}^Q \right) . \tag{4.15}$$

All these terms cancel against terms of the form (4.10) as discussed in appendix A.1–A.4.

As a final result, we find that the Lagrangian (4.1) is supersymmetric under the transformations (3.1) up to terms of higher order in the fermions. Remarkably, and unlike in the reduced theory, invariance of the Lagrangian under generalized diffeomorphisms (2.10), (2.16) already fixes all the bosonic couplings without reference to supersymmetry. The present construction gives the fermionic completion which turns the bosonic Lagrangian of [2, 3] into a supersymmetric system.

5 Conclusions and discussion

In this paper we have constructed the supersymmetric completion of $E_{6(6)}$ -covariant exceptional field theory, with the final result given by the Lagrangian (4.1) and the supersymmetry transformation laws (3.1). The section condition (2.7) effectively constrains the geometry of the extended space. It admits at least two independent maximal solutions which restrict the number of internal coordinates to six and five, respectively [2, 3].⁷ They are identified upon splitting the **27** representation of $E_{6(6)}$ under the action of the subgroup $GL(6)$ and $GL(5) \times SL(2)$, respectively. Upon imposing the former solution,

⁷The same is true for the $E_{7,8}$ cases and the higher dimensional $SO(5,5)$ and $SL(5)$ EFT's [4, 5, 29, 30].

the Lagrangian (4.1) reproduces the full Lagrangian of $D = 11$ supergravity, as explicitly demonstrated for its bosonic part in [3]. With the latter solution, the Lagrangian (4.1) describes the full supersymmetric IIB theory. It may at first appear surprising that one and the same set of fermions and couplings encodes both type IIA and type IIB, despite the crucial difference of their fermion chiralities. This is due to the fact that the $E_{6(6)}$ -covariant formulation (4.1) does not preserve the original $D = 10$ Lorentz invariance. As a consequence, its fermions can consistently encode the fermions of the type IIA and type IIB theory in the same way that both type IIA and type IIB give rise to the same supersymmetric theory in $D = 5$ upon dimensional reduction.

Upon the most straightforward solution of the section constraint, that is $\partial_M = 0$, the Lagrangian (4.1) directly reduces to the maximal $D = 5$ supergravity of [25]. In the context of generalized Scherk-Schwarz reductions, it has been proposed to relax the section condition (2.7) from a differential constraint into the known algebraic constraints on the embedding tensor, that naturally appears as a generalized torsion [17, 31–33]. Although, the generalized torsion formally reproduces all the gaugings, it remains an open question, to which extent they can be embedded into higher-dimensional supergravity via the corresponding EFT. The work [34], where the structure of the space of T-duality orbits was analysed, suggests that in principle one should be able to catch non-geometric compactifications by generalized Scherk-Schwarz reductions of EFT. On the other hand, a generalized Scherk-Schwarz ansatz that is consistent with the section condition (2.7), describes a consistent truncation of the exceptional field theory (4.1) and by virtue of the section condition translates into a consistent truncation of the conventional higher-dimensional supergravities. For the $SO(p, q)$ gauged supergravities, this ansatz has been constructed in [35]. It yields their higher-dimensional embedding as sphere and hyperboloid compactifications of the higher-dimensional supergravities [35],⁸ and naturally extends to the full Lagrangian (4.1).

In discussion of geometry of the extended space let us mention the works [37–39] where the geometrical meaning of the T-duality group $O(d, d)$ has been investigated. It was conjectured that the d -dimensional torus is just one of possible solutions of the field equations of double field theory, precisely the one that preserves the whole $O(d, d)$ group. Following this direction one may try to construct other solutions of DFT or EFT that preserve less duality symmetries and compare these with the known examples. Recently, in [40] it was shown that the brane solutions of $D = 4$ supergravity can be uplifted to a single solution of $E_{7(7)}$ exceptional field theory, that solves the twisted self-duality constraint. A possible direction of further research would be the investigation of similar uplifts in the presented $E_{6(6)}$ theory adding, possibly, winding coordinates, that should lead to non-geometric branes. Following the lines of [24] and the result of this paper one may explicitly investigate the supersymmetry properties of the obtained solutions in the EFT sense. In this context, we also mention the recent [41] for the embedding of supersymmetric flux backgrounds in exceptional geometry.

⁸See also [36] for the explicit uplift of several vacua of these theories.

Acknowledgments

On its last stage the work of ETM was partially supported by the German Science Foundation (DFG) under the Collaborative Research Center (SFB) 676 Particles, Strings and the Early Universe. ETM would like to thank Theoretical Department of CERN, Theory Department of DESY and II. Institut für Theoretische Physik, and personally Jan Louis, for warm hospitality during completion of part of this work. We thank Arnaud Baguet, Olaf Hohm, Hermann Nicolai, and Oscar Varela for inspiring discussions.

A Details of the supersymmetry calculation

In this section we provide most of the technical details of the rather lengthy calculations required to verify supersymmetry invariance of the Lagrangian (4.1) under the transformations (3.1). We discuss the various cancellations according to the different types of terms (4.10) that arise in the variation of the Lagrangian.

A.1 The $\psi \nabla_M \mathcal{D}_\nu \epsilon$ terms

The relevant contributions of this type from variation of the Rarita-Schwinger term are

$$\begin{aligned}
 & \delta_\epsilon (-e \bar{\psi}_{\mu i} \gamma^{\mu\nu\rho} \mathcal{D}_\nu \psi_\rho^i) \\
 & \longrightarrow -\bar{\psi}_{\mu i} \mathcal{D}_\nu (e \gamma^{\mu\nu\rho}) \delta_\epsilon \psi_\rho^i - 2e \bar{\psi}_{\mu i} \gamma^{\mu\nu\rho} \mathcal{D}_\nu \delta_\epsilon \psi_\rho^i \\
 & \longrightarrow -2\sqrt{2}ie \mathcal{D}_\nu \bar{\psi}_{\mu i} \gamma^{\mu\nu\rho} \left(\frac{2}{3} \gamma_\rho \nabla_M^- \epsilon^k \mathcal{V}^{ijM} \Omega_{jk} + \nabla_M^- \gamma_\rho \epsilon^k \mathcal{V}^{ijM} \Omega_{jk} \right) \\
 & = -4\sqrt{2}ie \mathcal{D}_\nu \bar{\psi}_{\mu i} \gamma^{\mu\nu} \nabla_M^- \epsilon^k \mathcal{V}^{ijM} \Omega_{jk} - 2\sqrt{2}ie \mathcal{D}_\nu \bar{\psi}_{\mu i} \gamma^{\mu\nu\rho} \nabla_M^- \gamma_\rho \epsilon^k \mathcal{V}^{ijM} \Omega_{jk},
 \end{aligned} \tag{A.1}$$

where the term $\mathcal{D}_\nu (e \gamma^{\mu\nu\rho})$ vanishes due to the vanishing torsion condition. The other contributions of this type come from the following variations

$$\begin{aligned}
 & \delta_\epsilon \left(4\sqrt{2} \mathcal{V}^{ijM} \bar{\chi}_{ijk} \gamma^\mu \nabla_M^- \psi_\mu^k \right) \longrightarrow 2\sqrt{2}i \mathcal{D}_\nu \mathcal{V}_{kl}^M \Omega^{lm} \nabla_M^- \bar{\psi}_{\mu m} \gamma^\mu \gamma^\nu \epsilon^k, \\
 & \delta_\epsilon \left(2\sqrt{2}ie \mathcal{V}_{ij}^M \Omega^{ik} \bar{\psi}_{\mu k} \gamma^{[\mu} \nabla_M^+ (\gamma^{\nu]} \psi_\nu^j) \right) \longrightarrow -4\sqrt{2}ie \Omega^{ik} \mathcal{V}_{ij}^M \nabla_M^+ (\bar{\psi}_{\mu k} \gamma^{[\mu} \gamma^{\nu]}) \delta_\epsilon \psi_\nu^j \\
 & \quad - 2\sqrt{2}i \nabla_M^+ e \mathcal{V}_{ij}^M \Omega^{ik} \bar{\psi}_{\mu k} \gamma^{\mu\nu} \delta_\epsilon \psi_\nu^j, \\
 & \delta_\epsilon \left(\frac{4i}{3} \mathcal{P}_\mu^{ijkl} \bar{\chi}_{ijk} \gamma^\nu \gamma^\mu \psi_\nu^m \Omega_{lm} \right) \longrightarrow 2\sqrt{2}i \mathcal{D}_\mu \mathcal{V}_{kl}^M \Omega^{lm} \bar{\psi}_{\nu m} \gamma^\mu \gamma^\nu \nabla_M^- \epsilon^k \\
 & \quad - \frac{16}{3} \mathcal{P}_\mu^{ijkl} \bar{\chi}_{ijk} \gamma^\mu \nabla_M^- \epsilon^r \mathcal{V}_{lr}^M \\
 & \quad - \frac{4\sqrt{2}}{3} \mathcal{P}_\mu^{ijkl} \bar{\chi}_{ijk} (\gamma^\mu \gamma^\nu \nabla_M^- \gamma_\nu) \epsilon^r \mathcal{V}_{lr}^M.
 \end{aligned} \tag{A.2}$$

Let us first separately verify cancellation of the $\mathcal{F}_{\mu\nu}{}^M$ terms against the variation of the vector kinetic term and of the topological term. From the above expressions we have

$$\begin{aligned}
 & \Omega^{lm} \left(\mathcal{V}_{kl}{}^M \mathcal{D}_\mu \bar{\psi}_{\nu m} (\gamma^{\mu\nu} \gamma^{\rho\sigma} - 2\gamma^{\mu\nu\rho} \gamma^\sigma) \epsilon^k \mathcal{F}_{\rho\sigma M} + \mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} \gamma^{[\mu} \gamma^{\rho\sigma} \gamma^{\nu]} \mathcal{D}_\mu \epsilon^k \mathcal{F}_{M\rho\sigma} \right. \\
 & \quad \left. + \frac{1}{2} \mathcal{D}_\mu \mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} (\gamma^{\rho\sigma} \gamma^\nu \gamma^\mu - \gamma^\mu \gamma^\nu \gamma^{\rho\sigma}) \epsilon^k \mathcal{F}_{\rho\sigma M} \right) \\
 & = \left(\mathcal{V}_{kl}{}^M \mathcal{D}_\mu \bar{\psi}_{\nu m} (\gamma^{\mu\nu\rho\sigma} - 2g^{\mu\sigma} g^{\nu\rho}) \epsilon^k \mathcal{F}_{\rho\sigma M} - \mathcal{D}_\mu \mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} (\gamma^{\mu\nu\rho\sigma} + 2g^{\mu\sigma} g^{\nu\rho}) \epsilon^k \mathcal{F}_{\rho\sigma M} \right. \\
 & \quad \left. + \mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} (\gamma^{\mu\nu\rho\sigma} - 2g^{\mu\sigma} g^{\nu\rho}) \mathcal{D}_\mu \epsilon^k \mathcal{F}_{\rho\sigma M} \right) \Omega^{lm} \\
 & = \Omega^{lm} \mathcal{D}_\mu (e \mathcal{V}_{Mkl} \bar{\psi}_{\nu m} \gamma^{\mu\nu\rho\sigma} \epsilon^k) \mathcal{F}_{\rho\sigma}{}^M + 2\Omega^{lm} \mathcal{D}_\mu \left(\mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} \epsilon^k \right) \mathcal{F}^{\mu\nu}{}_M, \tag{A.3}
 \end{aligned}$$

where we have defined $\mathcal{F}_{\mu\nu}{}^M \equiv \mathcal{F}_{\mu\nu}{}^N \mathcal{M}_{MN}$. The last term above is already present in the $D = 5$ reduced theory and cancels the $\bar{\epsilon} \psi$ part of the lowest order variation of the vector kinetic term. The first term can be rewritten upon partial integration and use of the Bianchi identities (2.15)

$$\frac{5\sqrt{2}}{3} \Omega^{lm} \varepsilon^{\mu\nu\rho\sigma\tau} \bar{\psi}_{\nu m} \gamma_\tau \epsilon^k \mathcal{V}_{Mkl} d^{MNK} \partial_N \mathcal{H}_{\mu\rho\sigma K}, \tag{A.4}$$

which precisely cancels the corresponding part in the variation (4.15) of the topological term.

To check the remaining terms one first notes the following relations

$$e\gamma^{\mu\nu\rho} \nabla_M \gamma_\rho = \nabla_M (e\gamma^{\mu\nu}), \quad \nabla_M (\gamma^{\mu\nu}) = 2(\nabla_M \gamma^{[\mu} \gamma^{\nu]}), \tag{A.5}$$

which can be used to bring the remainder into the following form

$$\begin{aligned}
 & 2e\mathcal{V}_{kl}{}^M \Omega^{lm} \mathcal{D}_\mu \bar{\psi}_{\nu m} \gamma^{\mu\nu} \nabla_M \epsilon^k + \mathcal{V}_{kl}{}^M \Omega^{lm} \mathcal{D}_\mu \bar{\psi}_{\nu m} \nabla_M (e\gamma^{\mu\nu}) \epsilon^k \\
 & - e\mathcal{D}_\mu \mathcal{V}_{kl}{}^M \Omega^{lm} \left(\nabla_M \bar{\psi}_{\nu m} \gamma^{\mu\nu} \epsilon^k - \bar{\psi}_{\nu m} \gamma^{\mu\nu} \nabla_M \epsilon^k \right) + eg^{\mu\nu} \mathcal{D}_\mu \mathcal{V}_{kl}{}^M \Omega^{lm} \nabla_M (\bar{\psi}_{\nu m} \epsilon^k) \\
 & - 2e\Omega^{lm} \mathcal{V}_{kl}{}^M \nabla_M \bar{\psi}_{\nu m} \gamma^{\mu\nu} \mathcal{D}_\mu \epsilon^k - \mathcal{V}_{kl}{}^M \Omega^{lm} \bar{\psi}_{\nu m} \nabla_M (e\gamma^{\mu\nu}) \mathcal{D}_\mu \epsilon^k.
 \end{aligned} \tag{A.6}$$

Now integrating by parts of \mathcal{D}_μ in the first term and of ∇_M in the fourth and the seventh term we get

$$\begin{aligned}
 & \Omega^{lm} \left(-2e\mathcal{D}_\mu \mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} \gamma^{\mu\nu} \nabla_M \epsilon^k - 2e\mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} \gamma^{\mu\nu} \mathcal{D}_\mu \nabla_M \epsilon^k + \mathcal{V}_{kl}{}^M \mathcal{D}_\mu \bar{\psi}_{\nu m} \nabla_M (e\gamma^{\mu\nu}) \epsilon^k \right. \\
 & \quad + e\nabla_M \mathcal{D}_\mu \mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} \gamma^{\mu\nu} \epsilon^k + \mathcal{D}_\mu \mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} \nabla_M (e\gamma^{\mu\nu}) \epsilon^k + 2e\mathcal{D}_\mu \mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} \gamma^{\mu\nu} \nabla_M \epsilon^k \\
 & \quad + 2\mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} \nabla_M (e\gamma^{\mu\nu}) \mathcal{D}_\mu \epsilon^k + 2e\mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} \gamma^{\mu\nu} \nabla_M \mathcal{D}_\mu \epsilon^k - \mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} \nabla_M (e\gamma^{\mu\nu}) \mathcal{D}_\mu \epsilon^k \\
 & \quad \left. + eg^{\mu\nu} \mathcal{D}_\mu \mathcal{V}_{kl}{}^M \nabla_M (\bar{\psi}_{\nu m} \epsilon^k) \right). \tag{A.7}
 \end{aligned}$$

Here it is straightforward to construct a commutator from the second terms in the first and the third lines, while the terms with $\nabla_M \epsilon^k$ cancel. What is left can be collected into the following expression

$$\begin{aligned}
 & \Omega^{lm} \left(2e\mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} \gamma^{\mu\nu} [\nabla_M, \mathcal{D}_\mu] \epsilon^k + e\nabla_M \mathcal{D}_\mu \mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} \gamma^{\mu\nu} \epsilon^k + eg^{\mu\nu} \mathcal{D}_\mu \mathcal{V}_{kl}{}^M \nabla_M (\bar{\psi}_{\nu m} \epsilon^k) \right. \\
 & \quad \left. + \mathcal{V}_{kl}{}^M \mathcal{D}_\mu \bar{\psi}_{\nu m} \nabla_M (e\gamma^{\mu\nu}) \epsilon^k + \mathcal{D}_\mu \mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} \nabla_M (e\gamma^{\mu\nu}) \epsilon^k + \mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} \nabla_M (e\gamma^{\mu\nu}) \mathcal{D}_\mu \epsilon^k \right).
 \end{aligned}$$

Integrating \mathcal{D}_μ and ∇_M by parts in the second line this simplifies into

$$\Omega^{lm} \left(e\mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} \gamma^{\mu\nu} [\nabla_M, \mathcal{D}_\mu] \epsilon^k - e\mathcal{V}_{kl}{}^M [\nabla_M, \mathcal{D}_\mu] \bar{\psi}_{\nu m} \gamma^{\mu\nu} \epsilon^k + eg^{\mu\nu} \mathcal{D}_\mu \mathcal{V}_{kl}{}^M \nabla_M (\bar{\psi}_{\nu m} \epsilon^k) \right). \quad (\text{A.8})$$

Upon using the expression (2.46) for the commutator of covariant derivatives together with (2.49) and restoring the prefactor $2\sqrt{2}i$, these terms reduce to

$$2\sqrt{2}i\Omega^{lm} \left(-\frac{1}{2} e\mathcal{V}_{kl}{}^M \bar{\psi}_{\nu m} \epsilon^k \hat{J}^\nu{}_M + eg^{\mu\nu} \mathcal{D}_\mu \mathcal{V}_{kl}{}^M \nabla_M (\bar{\psi}_{\nu m} \epsilon^k) \right). \quad (\text{A.9})$$

Upon partial integration in the second term, these remaining contributions precisely cancel the corresponding terms in (4.14). In what follows we drop the e -factor for simpler presentation as the corresponding terms cancel out in the very similar way as above.

A.2 The $\chi \nabla_M \mathcal{D}_\mu \epsilon$ terms

There are four fermionic terms from the Lagrangian which contribute such terms

$$\begin{aligned} (1) &= -\frac{4}{3} \bar{\chi}_{ijk} \gamma^\mu \mathcal{D}_\mu \chi^{ijk}, & (2) &= 8\sqrt{2} i \mathcal{V}_{mn}{}^M \Omega^{np} \bar{\chi}_{pkl} \nabla_M^+ \chi^{mkl}, \\ (3) &= \frac{4i}{3} \mathcal{P}_\mu{}^{ijkl} \bar{\chi}_{ijk} \gamma^\nu \gamma^\mu \psi_\nu{}^m \Omega_{lm}, & (4) &= 4\sqrt{2} \mathcal{V}_{kl}{}^M \Omega^{ki} \Omega^{lj} \bar{\chi}_{ijk} \gamma^\mu \nabla_M^- \psi_\mu{}^k. \end{aligned} \quad (\text{A.10})$$

The relevant terms in supersymmetry variations of these expressions have the following form

$$\begin{aligned} \delta_\epsilon^1(1) &= -4\sqrt{2} \bar{\chi}_{ijk} \gamma^\mu \mathcal{D}_\mu (\mathcal{V}^{ij}{}^M \nabla_M^- \epsilon^k) \\ &= -4\sqrt{2} \bar{\chi}_{ijk} \gamma^\mu \mathcal{V}^{ij}{}^M \mathcal{D}_\mu \nabla_M^- \epsilon^k + 4\sqrt{2} \bar{\chi}_{ijk} \gamma^\mu \nabla_M^- \epsilon^k \mathcal{P}_\mu{}^{ijmn} \mathcal{V}_{mn}{}^M \\ \delta_\epsilon^0(2) &= -8\sqrt{2} \mathcal{V}_{mn}{}^M \Omega^{np} \bar{\chi}_{pkl} \nabla_M^+ (\mathcal{P}_\mu{}^{mklq} \gamma^\mu \Omega_{qr} \epsilon^r) \\ &= -4\sqrt{2} \left(2\mathcal{V}_{mn}{}^M \Omega^{np} \bar{\chi}_{pkl} \gamma^\mu \Omega_{qr} \epsilon^r \nabla_M \mathcal{P}_\mu{}^{mklq} + 2\mathcal{V}_{mn}{}^M \Omega^{np} \bar{\chi}_{pkl} \mathcal{P}_\mu{}^{mklq} \gamma^\mu \Omega_{qr} \nabla_M^+ \epsilon^r \right. \\ &\quad \left. + \mathcal{V}_{mn}{}^M \Omega^{np} \bar{\chi}_{pkl} \gamma_\nu \Omega_{qr} \epsilon^r \mathcal{P}_\mu{}^{mklq} \nabla_M g^{\mu\nu} - \mathcal{V}_{Mmn} \Omega^{np} \bar{\chi}_{pkl} \mathcal{P}_\mu{}^{mklq} \gamma_\nu \Omega_{qr} \epsilon^r \mathcal{F}^{\mu\nu M} \right), \\ \delta_\epsilon^1(3) &= \frac{4\sqrt{2}}{3} \mathcal{P}_\mu{}^{ijkl} \bar{\chi}_{ijk} \gamma^\nu \gamma^\mu \mathcal{V}_{rl}{}^N \left(\nabla_N^- (\gamma_\nu \epsilon^r) - \frac{1}{3} \gamma_\nu \nabla_N^- \epsilon^r \right) \\ &= -\frac{4\sqrt{2}}{3} \mathcal{P}_\mu{}^{ijkl} \bar{\chi}_{ijk} \gamma_\nu \epsilon^r \nabla_N g^{\mu\nu} \mathcal{V}_{rl}{}^N - \frac{8\sqrt{2}}{3} \mathcal{P}_\mu{}^{mijk} \mathcal{V}_{mr}{}^N \bar{\chi}_{ijk} \gamma^\mu \nabla_N^- \epsilon^r \\ &\quad - \frac{2\sqrt{2}}{3} \mathcal{P}_\mu{}^{ijkl} \bar{\chi}_{ijk} \mathcal{V}_{Mrl} \mathcal{F}_{\nu\rho}{}^M \gamma^{\mu\nu\rho} \epsilon^r, \\ \delta_\epsilon^0(4) &= 4\sqrt{2} \mathcal{V}^{ij}{}^M \bar{\chi}_{ijk} \gamma^\mu \nabla_M^- \mathcal{D}_\mu \epsilon^k. \end{aligned} \quad (\text{A.11})$$

The variations of (1) and (4) give rise to a commutator of type (2.51)

$$\begin{aligned} &4\sqrt{2} \mathcal{V}^{ij}{}^M \bar{\chi}_{ijk} \gamma^\mu [\nabla_M^-, \mathcal{D}_\mu] \epsilon^k \\ &= 8\sqrt{2} \mathcal{V}_{nm}{}^M \Omega^{mi} \nabla_M \mathcal{P}_\mu{}^{jkn} \bar{\chi}_{ijk} \gamma^\mu \Omega_{pr} \epsilon^r + 2\mathcal{V}_{mn}{}^M \nabla_M \mathcal{P}^{\mu mnij} \mathcal{V}_{Nij} \delta_\epsilon^{(\bar{\chi})} \mathcal{A}_\mu{}^N \\ &\quad + 2\sqrt{2} \mathcal{V}_{mn}{}^M \mathcal{P}_\mu{}^{mnij} \nabla_M g^{\mu\nu} \bar{\chi}_{ijk} \gamma_\nu \epsilon^k + \sqrt{2} \mathcal{V}^{ij}{}^M \mathcal{R}_{M\mu}{}^{ab} \bar{\chi}_{ijk} \gamma^\mu \gamma_{ab} \epsilon^k, \end{aligned} \quad (\text{A.12})$$

of which the first term cancels the corresponding term in $\delta_\epsilon^0(2)$, and the second term cancels with the \mathcal{J}_M contribution in (4.14). The $\nabla_M g^{\mu\nu}$ can be seen to cancel against the

contributions from $\delta_\epsilon^0(2)$ and $\delta_\epsilon^1(3)$ by virtue of the $\text{USp}(8)$ identity (B.4). By the same identity, the three $\nabla_M \epsilon$ terms in (A.11) would cancel if they came with the same spin connection ω_M^- , i.e. they induce an extra term in the field strength $\mathcal{F}_{\mu\nu}^M$.

Collecting all resulting terms, we arrive at

$$\begin{aligned}
 (\text{A.11}) &= \\
 &= \sqrt{2} \mathcal{V}^{ijM} \mathcal{R}_{M\mu}^-{}^{ab} \bar{\chi}_{ijk} \gamma^\mu \gamma_{ab} \epsilon^k - 2\sqrt{2} \mathcal{V}_{Mmn} \Omega^{np} \bar{\chi}_{pkl} \mathcal{P}_\mu{}^{mklq} \gamma^\mu \Omega_{qr} \gamma^{\nu\rho} \epsilon^r \mathcal{F}_{\nu\rho}^M \\
 &\quad + 4\sqrt{2} \mathcal{V}_{Mmn} \Omega^{np} \bar{\chi}_{pkl} \mathcal{P}_\mu{}^{mklq} \gamma_\nu \Omega_{qr} \epsilon^r \mathcal{F}^{\mu\nu M} - \frac{2\sqrt{2}}{3} \mathcal{P}_\mu{}^{ijkl} \bar{\chi}_{ijk} \mathcal{V}_{Mrl} \mathcal{F}_{\nu\rho}^M \gamma^{\mu\nu\rho} \epsilon^r,
 \end{aligned} \tag{A.13}$$

with the second term coming from converting ∇_M^+ into ∇_M^- . Now the curvature term can be expanded with (2.48), (2.49) as

$$\begin{aligned}
 &\sqrt{2} \mathcal{V}^{ijM} \mathcal{R}_{M\mu}^-{}^{ab} \bar{\chi}_{ijk} \gamma^\mu \gamma_{ab} \epsilon^k \\
 &= \sqrt{2} \mathcal{V}^{ijM} \mathcal{R}_{M\mu\nu\rho}^- \bar{\chi}_{ijk} \gamma^{\mu\nu\rho} \epsilon^k + 2\sqrt{2} \mathcal{V}^{ijM} \mathcal{R}_{M\mu}^-{}^{\mu\nu} \bar{\chi}_{ijk} \gamma_\nu \epsilon^k \\
 &= \frac{1}{\sqrt{2}} \mathcal{D}_{[\mu} (\mathcal{F}_{\nu\rho]}^N \mathcal{M}_{NM}) \mathcal{V}^{ijM} \bar{\chi}_{ijk} \gamma^{\mu\nu\rho} \epsilon^k \\
 &\quad + \sqrt{2} \mathcal{V}^{ijM} \hat{J}^\mu{}_M \bar{\chi}_{ijk} \gamma_\mu \epsilon^k - \sqrt{2} \mathcal{V}^{ijM} e_a{}^\nu e_b{}^\mu \mathcal{D}_\mu (\mathcal{M}_{MN} \mathcal{F}^{abN}) \bar{\chi}_{ijk} \gamma_\nu \epsilon^k.
 \end{aligned} \tag{A.14}$$

The last two terms cancel against the vector field variation from the Einstein-Hilbert term (4.14) and from the vector kinetic term. The first term gives

$$\begin{aligned}
 &\rightarrow \frac{1}{\sqrt{2}} \mathcal{V}_M{}^{ij} \mathcal{D}_{[\mu} \mathcal{F}_{\nu\rho]}^M \bar{\chi}_{ijk} \gamma^{\mu\nu\rho} \epsilon^k + \sqrt{2} \mathcal{P}_\mu{}^{ijmn} \mathcal{V}_{Mmn} \mathcal{F}_{\nu\rho}^M \bar{\chi}_{ijk} \gamma^{\mu\nu\rho} \epsilon^k \\
 &= \frac{1}{\sqrt{2}} \mathcal{V}_M{}^{ij} \mathcal{D}_{[\mu} \mathcal{F}_{\nu\rho]}^M \bar{\chi}_{ijk} \gamma^{\mu\nu\rho} \epsilon^k \\
 &\quad + \frac{2}{3} \sqrt{2} \mathcal{V}_{Mmn} \mathcal{P}_\mu{}^{mijk} \mathcal{F}_{\nu\rho}^M \bar{\chi}_{ijk} \gamma^{\mu\nu\rho} \epsilon^n - 2\sqrt{2} \mathcal{V}_{Mmn} \mathcal{F}_{\nu\rho}^M \Omega_{pr} \bar{\chi}_{ijk} \gamma^{\mu\nu\rho} \epsilon^r \Omega^{mi} \mathcal{P}_\mu{}^{jkn\rho}
 \end{aligned} \tag{A.15}$$

where we have once more used the algebraic identity (B.4). The last two terms precisely cancel the $\mathcal{F}_{\mu\nu}^M$ terms from (A.13). We remain with the first term of (A.15) which can be rewritten with the Bianchi identity (2.15) and cancels against the corresponding $\mathcal{H}_{\mu\nu\rho M}$ term from variation of the topological term in (4.15).

A.3 The $\psi \nabla_M \nabla_N \epsilon$ terms

These terms arise from the $\nabla_M \epsilon$ variation of the following two terms from the Lagrangian (4.1)

$$(1) = 2\sqrt{2} i \mathcal{V}_{ij}{}^M \Omega^{ik} \bar{\psi}_{\mu k} \gamma^{[\mu} \nabla_M^+ (\gamma^{\nu]} \psi_{\nu}{}^j), \quad (2) = 4\sqrt{2} \mathcal{V}_{ij}{}^M \bar{\psi}_{\mu k} \nabla_M^- (\gamma^\mu \chi^{ijk}). \tag{A.16}$$

Explicitly, with (3.1) this gives

$$\begin{aligned}
 \delta_\epsilon^1(1) &= 8 \mathcal{V}_{nj}{}^M \mathcal{V}^{jkN} \bar{\psi}_{\mu k} \gamma^{[\mu} \nabla_N^+ \left(\gamma^{\nu]} \left(\nabla_M^- (\gamma_\nu \epsilon^n) - \frac{1}{3} \gamma_\nu \nabla_M^- \epsilon^n \right) \right), \\
 \delta_\epsilon^1(2) &= -12 \left(\mathcal{V}^{[ijN} \Omega^{k]m} - \frac{1}{3} \mathcal{V}^{m[iN} \Omega^{jk]} \right) \Omega_{mr} \mathcal{V}_{ij}{}^M \bar{\psi}_{\mu k} \nabla_M^- (\gamma^\mu \nabla_N^- \epsilon^r) \\
 &= 4 \left(\mathcal{M}^{MN} \delta_n^k + 2 \mathcal{V}_{nj}{}^M \mathcal{V}^{jkN} + \frac{2}{3} \mathcal{V}_{ij}{}^M \mathcal{V}^{jmN} \Omega^{ik} \Omega_{mn} \right) \bar{\psi}_{\mu k} \nabla_M^- (\gamma^\mu \nabla_N^- \epsilon^n) \\
 &= 4 \left(\mathcal{M}^{MN} \delta_n^k + 2 \mathcal{V}_{nj}{}^M \mathcal{V}^{jkN} + \frac{2}{3} \mathcal{V}_{nj}{}^N \mathcal{V}^{jkM} \right) \bar{\psi}_{\mu k} \nabla_M^- (\gamma^\mu \nabla_N^- \epsilon^n).
 \end{aligned} \tag{A.17}$$

Let us now consider the terms containing ∇_M and the gauge field flux $\mathcal{F}_{\mu\nu}{}^M$ separately. For the derivative terms and ignoring all derivatives on the external metric we have

$$\begin{aligned}
 \delta_\epsilon(1) + \delta_\epsilon(2) &\rightarrow \left(\frac{64}{3} + \frac{8}{3} \right) \mathcal{V}_{nj}{}^M \mathcal{V}^{Njk} \bar{\psi}_{\mu k} \gamma^\mu \nabla_N \nabla_M \epsilon^n \\
 &\quad + 8 \mathcal{V}_{nj}{}^M \mathcal{V}^{jkN} \bar{\psi}_{\mu k} \gamma^\mu \nabla_M \nabla_N \epsilon^n + 4 \mathcal{M}^{MN} \bar{\psi}_{\mu k} \gamma^\mu \nabla_M \nabla_N \epsilon^k \\
 &= \left(-\frac{32}{3} - \frac{4}{3} + 4 \right) \mathcal{V}_{nj}{}^M \mathcal{V}^{jkN} \bar{\psi}_{\mu k} \gamma^\mu [\nabla_M, \nabla_N] \epsilon^n \\
 &\quad \left(+\frac{64}{3} + \frac{8}{3} + 8 \right) \mathcal{V}_{nj}{}^M \mathcal{V}^{jkN} \bar{\psi}_{\mu k} \gamma^\mu \nabla_{(M} \nabla_{N)} \epsilon^n + 4 \mathcal{M}^{MN} \bar{\psi}_{\mu k} \gamma^\mu \nabla_M \nabla_N \epsilon^k \\
 &= -8 \mathcal{V}_{nj}{}^M \mathcal{V}^{jkN} \bar{\psi}_{\mu k} \gamma^\mu [\nabla_M, \nabla_N] \epsilon^n \\
 &\quad + 32 \mathcal{V}_{nj}{}^M \mathcal{V}^{jkN} \bar{\psi}_{\mu k} \gamma^\mu \nabla_{(M} \nabla_{N)} \epsilon^n + 4 \mathcal{M}^{MN} \bar{\psi}_{\mu k} \gamma^\mu \nabla_M \nabla_N \epsilon^k \\
 &= -\frac{1}{2} \mathcal{R} \bar{\psi}_{\mu k} \gamma^\mu \epsilon^k,
 \end{aligned} \tag{A.18}$$

upon using (2.52).

That cancels the corresponding variations of the scalar potential. Now for the \mathcal{FF} terms altogether we obtain

$$\begin{aligned}
 \delta_\epsilon(1) + \delta_\epsilon(2) &\rightarrow \frac{1}{16} \left(\mathcal{M}^{MN} \delta_n^k + 8 \mathcal{V}_{nj}{}^N \mathcal{V}^{jkM} \right) \bar{\psi}_{\mu k} \gamma^{\mu\kappa\lambda\rho\sigma} \epsilon^n \mathcal{F}_{\kappa\lambda M} \mathcal{F}_{\rho\sigma N} \\
 &\quad - \frac{1}{8} \mathcal{M}^{MN} \bar{\psi}_{\mu k} \gamma^\mu \epsilon^k \mathcal{F}_{\rho\nu M} \mathcal{F}^{\rho\nu N} - \frac{1}{2} \mathcal{M}^{MN} \bar{\psi}_{\mu k} \gamma^\rho \epsilon^k \mathcal{F}^{\mu\nu M} \mathcal{F}_{\nu\rho N} \\
 &= -\frac{i}{4} \sqrt{5} \epsilon^{\mu\kappa\lambda\rho\sigma} d^{MNK} \mathcal{V}_{ij}{}^K \Omega^{ir} \bar{\epsilon}_r \psi_\mu{}^j \mathcal{F}_{\kappa\lambda}{}^M \mathcal{F}_{\rho\sigma}{}^N \\
 &\quad + \frac{1}{4} (e^{-1} \delta_\epsilon e) \mathcal{M}^{MN} \mathcal{F}_{\rho\nu M} \mathcal{F}^{\rho\nu N} + \frac{1}{4} \delta_\epsilon (\gamma^{\mu\nu} g^{\rho\sigma}) \mathcal{M}^{MN} \mathcal{F}_{\mu\rho M} \mathcal{F}_{\nu\sigma N},
 \end{aligned} \tag{A.19}$$

that precisely cancels the variation from the kinetic and topological vector term. Let us

turn to the terms of the form $\nabla\mathcal{F}$ that give

$$\begin{aligned}
 \delta_\epsilon(1) + \delta_\epsilon(2) &\rightarrow \mathcal{V}_{nj}{}^M \mathcal{V}^{jkN} \bar{\psi}_{\mu k} \left(\gamma^{\rho\sigma} \gamma^\mu \epsilon^n + \frac{1}{3} \gamma^\mu \gamma^{\rho\sigma} \epsilon^n \right) \nabla_N \mathcal{F}_{\rho\sigma M} \\
 &\quad - \frac{1}{2} \left(\mathcal{M}^{MN} \delta_n^k + 2\mathcal{V}_{nj}{}^N \mathcal{V}^{jkM} + \frac{2}{3} \mathcal{V}_{nj}{}^M \mathcal{V}^{jkN} \right) \bar{\psi}_{\mu k} \gamma^\mu \gamma^{\rho\sigma} \epsilon^n \nabla_N \mathcal{F}_{\rho\sigma M} \\
 &= -\frac{1}{2} \mathcal{M}^{MN} \bar{\psi}_{\mu k} \gamma^{\mu\rho\sigma} \epsilon^k \nabla_M \mathcal{F}_{\rho\sigma N} - 2\mathcal{V}_{nj}{}^M \mathcal{V}^{jkN} \bar{\psi}_{\mu k} \gamma^{\mu\rho\sigma} \epsilon^n \nabla_{[M} \mathcal{F}_{\rho\sigma N]} \\
 &\quad - 4\mathcal{V}_{nj}{}^M \mathcal{V}^{jkN} \bar{\psi}_{\mu k} \gamma_\nu \epsilon^n \nabla_{(M} \mathcal{F}^{\mu\nu}{}_{N)} - \frac{1}{2} \mathcal{M}^{MN} \bar{\psi}_{\mu k} \gamma_\nu \epsilon^k \nabla_M \mathcal{F}^{\mu\nu}{}_N \\
 &= -2\mathcal{V}_{nj}{}^M \mathcal{V}^{jkN} \bar{\psi}_{\mu k} \gamma^{\mu\rho\sigma} \epsilon^n \nabla_{[M} \mathcal{F}_{\rho\sigma N]} - \frac{1}{2} \bar{\psi}_{\mu k} \gamma^{\mu\rho\sigma} \epsilon^k \nabla_M \mathcal{F}_{\rho\sigma}{}^M \\
 &\quad - 5d^{MNK} \delta_\epsilon^{(\bar{\psi}\epsilon)} \mathcal{B}_{\mu\nu K} \nabla_{(M} \mathcal{F}^{\mu\nu}{}_{N)}. \tag{A.20}
 \end{aligned}$$

The first line here precisely cancels against the corresponding terms in (4.13) from variation of the Rarita-Schwinger term. The second line cancels against the corresponding contribution in (4.15). Finally, for the terms of type $\mathcal{F}\nabla\epsilon$, we obtain

$$\begin{aligned}
 \delta_\epsilon(1) &\rightarrow \mathcal{V}_{nj}{}^M \mathcal{V}^{jkN} \bar{\psi}_{\mu k} \gamma^{[\mu} \gamma^{\rho\sigma} \left(\gamma^{\nu]} \left(\frac{2}{3} \gamma_\nu \nabla_M \epsilon^n \right) \right) \mathcal{F}_{\rho\sigma N} \\
 &\quad - \mathcal{V}_{nj}{}^N \mathcal{V}^{jkM} \bar{\psi}_{\mu k} \gamma^{[\mu} \left(\gamma^{\nu]} \left(\gamma^{\rho\sigma} \gamma_\nu - \frac{1}{3} \gamma_\nu \gamma^{\rho\sigma} \right) \right) \nabla_M \epsilon^n \mathcal{F}_{\rho\sigma N} \\
 &= \frac{4}{3} \mathcal{V}_{nj}{}^M \mathcal{V}^{jkN} \bar{\psi}_{\mu k} \gamma^{\mu\rho\sigma} \nabla_M \epsilon^n \mathcal{F}_{\rho\sigma N} + \frac{4}{3} \mathcal{V}_{nj}{}^M \mathcal{V}^{jkN} \bar{\psi}_{\mu k} \gamma^\sigma \nabla_M \epsilon^n \mathcal{F}^\mu{}_{\sigma N} \\
 &\quad + \frac{4}{3} \mathcal{V}_{nj}{}^N \mathcal{V}^{jkM} \bar{\psi}_{\mu k} \gamma^{\mu\rho\sigma} \nabla_M \epsilon^n \mathcal{F}_{\rho\sigma N} - \frac{4}{3} \mathcal{V}_{nj}{}^N \mathcal{V}^{jkM} \bar{\psi}_{\mu k} \gamma^\sigma \nabla_M \epsilon^n \mathcal{F}^\mu{}_{\sigma N}, \tag{A.21} \\
 \delta_\epsilon(2) &\rightarrow -\frac{1}{2} \left(\mathcal{M}^{MN} \delta_n^k + 2\mathcal{V}_{nj}{}^N \mathcal{V}^{jkM} + \frac{2}{3} \mathcal{V}_{nj}{}^M \mathcal{V}^{Njk} \right) \bar{\psi}_{\mu k} \gamma^{\rho\sigma} \gamma^\mu \nabla_M \epsilon^n \mathcal{F}_{\rho\sigma N} \\
 &\quad - \frac{1}{2} \left(\mathcal{M}^{MN} \delta_n^k + 2\mathcal{V}_{nj}{}^M \mathcal{V}^{jkN} + \frac{2}{3} \mathcal{V}_{nj}{}^N \mathcal{V}^{jkM} \right) \bar{\psi}_{\mu k} \gamma^\mu \gamma^{\rho\sigma} \nabla_M \epsilon^n \mathcal{F}_{\rho\sigma N}.
 \end{aligned}$$

Together these contribution simplify to the following nice expression

$$\delta_\epsilon(1) + \delta_\epsilon(2) \rightarrow -\bar{\psi}_{\mu k} \gamma^{\mu\rho\sigma} \mathcal{F}_{\rho\sigma}{}^M \nabla_M \epsilon^k, \tag{A.22}$$

which precisely cancels the corresponding contribution from (4.13).

A.4 The $\chi\nabla_M\nabla_N\epsilon$ terms

As the final check we collect the $\chi\nabla_M\nabla_N\epsilon$ terms which originate from the $\nabla_M\epsilon$ variation of the following two terms

$$(1) = 8\sqrt{2}i \mathcal{V}_{mn}{}^M \Omega^{np} \bar{\chi}_{pkl} \nabla_M^+ \chi^{mkl}, \quad (2) = 4\sqrt{2} \mathcal{V}_{ij}{}^M \bar{\psi}_{\mu k} \nabla_M^- \left(\gamma^\mu \chi^{ijk} \right), \tag{A.23}$$

of the Lagrangian (4.1). Their supersymmetry variation gives

$$\begin{aligned}
 \delta_\epsilon(1) &= -48 i \mathcal{V}_{mn}{}^M \Omega^{np} \left(\mathcal{V}^{N[lm\Omega^k]j} - \frac{1}{3} \mathcal{V}^{Nj[l\Omega^{mk}]} \right) \Omega_{jr} \bar{\chi}_{pkl} \nabla_M^+ \nabla_N^- \epsilon^r \\
 &= 16 i \left(2 \mathcal{V}^{kjN} \Omega_{jn} \mathcal{V}^{nlM} \delta_r^p + \mathcal{V}^{klN} \mathcal{V}^{pjM} \Omega_{jr} + \frac{2}{3} \mathcal{V}^{klM} \mathcal{V}^{pjN} \Omega_{jr} \right) \bar{\chi}_{pkl} \nabla_M^+ \nabla_N^- \epsilon^r, \\
 \delta_\epsilon(2) &= -8 i \mathcal{V}^{klM} \mathcal{V}^{pjN} \Omega_{jr} \bar{\chi}_{klp} \gamma^\mu \nabla_M^- \left(\frac{2}{3} \gamma_\mu \nabla_N^- \epsilon^r + \gamma_a \epsilon^r \nabla_N^- e_\mu{}^a \right). \tag{A.24}
 \end{aligned}$$

Again for simplicity we start from analysis for the terms that do not contain the field strength

$$\begin{aligned}
 \delta_\epsilon(1) + \delta_\epsilon(2) &\rightarrow 32 i \mathcal{V}^{kjM} \Omega_{jn} \mathcal{V}^{nlN} \bar{\chi}_{pkl} \nabla_{(M} \nabla_{N)} \epsilon^p + 16 i \mathcal{V}^{klN} \mathcal{V}^{pjM} \Omega_{jr} \bar{\chi}_{pkl} [\nabla_M, \nabla_N] \epsilon^r \\
 &\quad - 8 i \mathcal{V}^{klM} \mathcal{V}^{pjN} \Omega_{jr} \bar{\chi}_{klp} \gamma^\mu \gamma_a \epsilon^r \nabla_M \nabla_N e_\mu{}^a \\
 &= 16 i \left(\mathcal{V}^{klN} \mathcal{V}^{pjM} \Omega_{jr} \bar{\chi}_{pkl} [\nabla_M, \nabla_N] \epsilon^r + 2 \mathcal{V}^{kjM} \Omega_{jn} \mathcal{V}^{nlN} \bar{\chi}_{pkl} \nabla_{(M} \nabla_{N)} \epsilon^p \right) \\
 &\stackrel{(2.51)}{=} -4 i \mathcal{R}^{ijkl} \Omega_{ri} \bar{\chi}_{jkl} \epsilon^r + 2 i \mathcal{V}^{klM} \mathcal{V}^{pjN} \Omega_{jr} \bar{\chi}_{klp} \epsilon^r \nabla_M g_{\mu\nu} \nabla_N g^{\mu\nu}, \tag{A.25}
 \end{aligned}$$

with the curvature \mathcal{R}^{ijkl} in the **42** representation. These terms, after using the section constraint for the second one, precisely cancel the variation of the scalar potential (4.11). Collecting now the $\nabla\mathcal{F}$ terms, we get

$$\begin{aligned}
 &\delta_\epsilon(1) + \delta_\epsilon(2) \\
 &\rightarrow -4 i \mathcal{V}^{klM} \mathcal{V}_K{}^{pj} \Omega_{r[k} \bar{\chi}_{l]pj} \gamma^{\mu\nu} \epsilon^r \nabla_M \mathcal{F}_{\mu\nu}{}^K - 4 i \mathcal{V}^{klM} \mathcal{V}_K{}^{pj} \Omega_{r[k} \bar{\chi}_{l]pj} \gamma^\mu \lambda \epsilon^r \mathcal{F}^{\lambda\rho K} \nabla_M g_{\mu\rho} \\
 &\quad - 4 i \mathcal{V}^{kjM} \Omega_{jn} \mathcal{V}_K{}^{nl} \bar{\chi}_{pkl} \gamma^{ab} \epsilon^p \nabla_M \mathcal{F}_{ab}{}^K \\
 &= -4 i \mathcal{V}^{klM} \mathcal{V}_K{}^{pj} \Omega_{r[k} \bar{\chi}_{l]pj} \gamma^{\mu\nu} \epsilon^r \nabla_M \mathcal{F}_{\mu\nu}{}^K - 4 i \mathcal{V}^{klM} \mathcal{V}_K{}^{pj} \Omega_{r[k} \bar{\chi}_{l]pj} \gamma^\mu \lambda \epsilon^r \mathcal{F}^{\lambda\rho K} \nabla_M g_{\mu\rho} \\
 &\quad - 2 i \mathcal{V}^{kjM} \Omega_{jn} \mathcal{V}_K{}^{nl} \bar{\chi}_{pkl} \gamma_{\mu\nu} \epsilon^p \nabla_M \mathcal{F}^{\mu\nu K} \\
 &= -4 i \mathcal{V}^{klM} \mathcal{V}_K{}^{pj} \Omega_{r[k} \bar{\chi}_{l]pj} \gamma^{\mu\nu} \epsilon^r \nabla_M \mathcal{F}_{\mu\nu}{}^K - 4 i \mathcal{V}^{klM} \mathcal{V}_K{}^{pj} \Omega_{r[k} \bar{\chi}_{l]pj} \gamma^\mu \lambda \epsilon^r \mathcal{F}^{\lambda\rho K} \nabla_M g_{\mu\rho} \\
 &\quad - 5 d^{KMN} \delta_\epsilon B_{\mu\nu N} \nabla_M (\mathcal{M}_{KL} \mathcal{F}^{\mu\nu L}). \tag{A.26}
 \end{aligned}$$

The last term precisely cancels the corresponding variation of the vector kinetic term, the second term cancels against (A.28) below, the first one upon using the identity (B.5) cancels against the contribution from (4.12). Collecting the $\mathcal{F}\mathcal{F}$ terms we obtain (again with $\mathcal{F}_M \equiv \mathcal{M}_{MN} \mathcal{F}^N$)

$$\begin{aligned}
 &-(\delta_\epsilon(1) + \delta_\epsilon(2)) \tag{A.27} \\
 &\rightarrow \frac{i}{4} \left(2 \mathcal{V}^{kjN} \Omega_{jn} \mathcal{V}^{nlM} \delta_r^p + \mathcal{V}^{klN} \mathcal{V}^{pjM} \Omega_{jr} + \frac{2}{3} \mathcal{V}^{klM} \mathcal{V}^{pjN} \Omega_{jr} \right) \bar{\chi}_{pkl} \gamma^{\mu\nu} \gamma^{\rho\sigma} \epsilon^r \mathcal{F}_{\mu\nu M} \mathcal{F}_{\rho\sigma N} \\
 &\quad \frac{i}{8} \mathcal{V}^{klM} \mathcal{V}^{pjN} \Omega_{jr} \bar{\chi}_{klp} \left(\gamma^\tau \gamma^{\mu\nu} \gamma^{\rho\sigma} \gamma_\tau - \frac{1}{3} \gamma^\tau \gamma^{\mu\nu} \gamma_\tau \gamma^{\rho\sigma} \right) \epsilon^r \\
 &= \frac{i}{4} \left(2 \mathcal{V}^{kjN} \Omega_{jn} \mathcal{V}^{nlM} \delta_r^p + \mathcal{V}^{klN} \mathcal{V}^{pjM} \Omega_{jr} \right) \bar{\chi}_{pkl} (\gamma^{\mu\nu\rho\sigma} + 4 \gamma^{\mu\sigma} g^{\nu\rho} - 2 g^{\mu\rho} g^{\nu\sigma}) \epsilon^r \mathcal{F}_{\mu\nu M} \mathcal{F}_{\rho\sigma N} \\
 &\quad \frac{i}{8} \mathcal{V}^{klM} \mathcal{V}^{pjN} \Omega_{jr} \bar{\chi}_{klp} (-2 \gamma^{\mu\nu\rho\sigma} + 8 \gamma^{\mu\sigma} g^{\nu\rho} - 12 g^{\mu\rho} g^{\nu\sigma}) \epsilon^r \mathcal{F}_{\mu\nu M} \mathcal{F}_{\rho\sigma N} \\
 &= \frac{i}{2} \mathcal{V}^{kjN} \Omega_{jn} \mathcal{V}^{nlM} \delta_r^p \bar{\chi}_{pkl} \gamma^{\mu\nu\rho\sigma} \epsilon^r \mathcal{F}_{\mu\nu}{}^M \mathcal{F}_{\rho\sigma}{}^N - 2 i \mathcal{V}_M{}^{kl} \mathcal{V}_N{}^{pj} \Omega_{r[[k} \bar{\chi}_{l]pj]} \epsilon^r \mathcal{F}_{\mu\nu}{}^M \mathcal{F}^{\mu\nu N}.
 \end{aligned}$$

These cancel against the corresponding variation of the kinetic and the topological vector terms. Finally, for the $\mathcal{F}\nabla$ terms, we write

$$\begin{aligned}
 & \delta_\epsilon(1) + \delta_\epsilon(2) \\
 \rightarrow & \frac{2}{3}i\mathcal{V}^{klN}\mathcal{V}_M{}^{pj}\Omega_{jr}\mathcal{F}_{\mu\nu}{}^M\bar{\chi}_{pkl}\gamma^{\mu\nu}\nabla_N\epsilon^r - \frac{2}{3}i\mathcal{V}_M{}^{kl}\mathcal{V}^{pjN}\Omega_{jr}\mathcal{F}_{\mu\nu}{}^M\bar{\chi}_{pkl}\gamma^{\mu\nu}\nabla_N\epsilon^r \\
 & + i\mathcal{V}^{klM}\mathcal{V}^{pjN}\Omega_{jr}\bar{\chi}_{klp}\gamma^{\nu\rho}\nabla_N\epsilon^r\mathcal{F}_{\nu\rho M} - \frac{i}{3}\mathcal{V}^{klM}\mathcal{V}^{pjN}\Omega_{jr}\bar{\chi}_{klp}\gamma^{\nu\rho}\nabla_N\epsilon^r\mathcal{F}_{\nu\rho M} \\
 & + i\mathcal{V}^{klN}\mathcal{V}^{pjM}\Omega_{jr}\bar{\chi}_{klp}\gamma^{\nu\rho}\nabla_N\epsilon^r\mathcal{F}_{\nu\rho M} - \frac{5i}{3}\mathcal{V}^{klN}\mathcal{V}^{pjM}\Omega_{jr}\bar{\chi}_{klp}\gamma^{\nu\rho}\nabla_N\epsilon^r\mathcal{F}_{\nu\rho M} \quad (\text{A.28}) \\
 & + i\mathcal{V}^{klM}\mathcal{V}^{pjN}\Omega_{jr}\bar{\chi}_{klp}\gamma^\mu\gamma^{\nu\rho}\gamma_a\epsilon^r\mathcal{F}_{\nu\rho M}\nabla_N e_\mu{}^a \\
 & - \frac{i}{3}\mathcal{V}^{klN}\mathcal{V}^{pjM}\Omega_{jr}\bar{\chi}_{klp}\gamma^\mu\gamma_a\gamma^{\nu\rho}\epsilon^r\mathcal{F}_{\nu\rho M}\nabla_N e_\mu{}^a \\
 = & 4i\mathcal{V}^{klM}\mathcal{V}^{pjN}\Omega_{jr}\bar{\chi}_{klp}\gamma^{\mu\nu}\epsilon^r\mathcal{F}_{\nu\rho(M}g^{\rho\lambda}\nabla_N)g_{\mu\lambda},
 \end{aligned}$$

that precisely cancels the second term above in (A.26).

B USp(8) identities

In this section some useful algebraic relations, that follow from the structure of USp(8) representations. Their derivation was facilitated in part by using the computer algebra system Cadabra [42, 43]. Some of the more complicated algebraic relations were obtained using an explicitly chosen USp(8) representation.

We first recall the notation of double brackets

$$P[[ijkl]] = P^{[ijkl]} - (\Omega\text{-traces}), \quad \text{etc.}, \quad (\text{B.1})$$

in order to define the irreducible USp(8) representations. E.g. the tensor $P^{ijkl} = P[[ijkl]]$ defines the irreducible **42** representation of USp(8) and can explicitly be constructed by making use of the corresponding projector

$$\begin{aligned}
 P[[ijkl]] &= \mathbb{P}_{\mathbf{42}}^{ijkl}{}_{mnpq} P^{mnpq}, \\
 \mathbb{P}_{\mathbf{42}}^{ijkl}{}_{mnpq} &\equiv \delta^{ijkl}{}_{mnpq} - \frac{3}{2}\Omega^{[ij}\delta^{kl]}{}_{[mn}\Omega_{pq]} + \frac{1}{8}\Omega^{[ij}\Omega^{kl]}\Omega_{[mn}\Omega_{pq]}. \quad (\text{B.2})
 \end{aligned}$$

Several of the USp(8) identities are not straightforward to derive but most conveniently derived by identifying the underlying representation structure. A simple example of such an identity is

$$P^{[ijkl}\Omega^{mn]} = 0, \quad (\text{B.3})$$

for $P^{ijkl} = P[[ijkl]]$ in the **42** of USp(8). The identity (B.3) follows straightforwardly from the fact that there is no **42** representation in the six-fold antisymmetric tensor product.

In the same manner, one may derive the identity

$$0 = \frac{3}{4}\mathcal{V}_{mn}{}^M\epsilon^{[i}P^{jk]mn} - \frac{1}{2}\mathcal{V}_{mn}{}^M P^{mijk}\epsilon^n + \frac{3}{2}\mathcal{V}_{mn}{}^M\Omega_{pr}\epsilon^r\Omega^m{}^{[i}P^{jk]np}, \quad (\text{B.4})$$

for $P^{ijkl} = P^{\llbracket ijkl \rrbracket}$, whose existence follows from the fact that there is no **42** representation in the tensor product $\mathbf{27} \times \mathbf{42}$, and as a consequence there are only two singlets in $\mathbf{8} \otimes \mathbf{27} \otimes \mathbf{48} \otimes \mathbf{42}$. The coefficients in (B.4) can then be fixed by employing an explicit realization of these objects, or by using the explicit form (B.2) of the projector.

Similarly, one shows the identity

$$\begin{aligned} \mathcal{V}_K^{\llbracket ij \rrbracket} \mathcal{V}^{kl} M \Omega_{ri} \bar{\chi}_{jkl} &= \mathcal{V}^{mn} M \mathcal{V}_K^{pq} \left(\Omega_{r[m} \bar{\chi}_{npq]} - \frac{3}{2} \Omega_{ri} \bar{\chi}_{jkl} \Omega^{[ij} \delta^{kl]}_{[mn} \Omega_{pq]} \right) \\ &= \mathcal{V}^{mn} M \mathcal{V}_K^{pq} \Omega_{r[m} \bar{\chi}_{npq]} + \frac{1}{2} \mathcal{V}^{mn} M \Omega_{nq} \mathcal{V}_K^{qp} \bar{\chi}_{rmp} . \end{aligned} \quad (\text{B.5})$$

In the main text, the calculation of the scalar potential (4.7) and its properties like (2.57) require further USp(8) identities. E.g. one derives

$$\begin{aligned} \mathcal{V}_{ij}^M \mathcal{V}_{kl}^N \partial_{(MPN)}^{jklm} \epsilon_m - \Omega_{im} \mathcal{V}_{pj}^M \mathcal{V}_{kl}^N \partial_{(MPN)}^{jklm} \Omega^{pq} \epsilon_q \\ = -\frac{1}{4} \mathcal{V}_{kl}^M \mathcal{V}_{mn}^N \partial_{(MPN)}^{klmn} \epsilon_i + \Omega_{ir} \mathcal{V}_{km}^M \Omega^{mn} \mathcal{V}_{nl}^N \partial_{(MPN)}^{rjkl} \epsilon_j , \end{aligned} \quad (\text{B.6})$$

that follows from the fact that in the above $\mathcal{V}_{ij}^M \mathcal{V}_{kl}^N$ appears only projected onto the **42** due to the section condition (2.9), and furthermore there is only **1** and no **36** in $(\mathbf{42} \otimes \mathbf{42})_{\text{sym}}$.

Another set of relations is required for the evaluation of the commutator (2.53) that contains

$$\begin{aligned} \rightarrow \mathcal{V}_{mn}^M \mathcal{V}^{kl} N \Omega_{pq} \epsilon^p \chi_{klr} P_{MN}^{mnqr} - \mathcal{V}^{ij} M \mathcal{V}^{kl} N \Omega^{mn} \epsilon^p \chi_{ikm} P_{MN}^{jlnp} \\ - \frac{1}{2} \mathcal{V}_{mn}^M \mathcal{V}^{kl} N \Omega_{kp} \epsilon^q \chi_{lqr} P_{MN}^{mnpr} - \frac{1}{16} \mathcal{M}^{MN} \Omega_{ij} \epsilon^i \chi_{klm} P_{MN}^{jklm} \\ + \frac{1}{4} \mathcal{V}_{mn}^M \mathcal{V}^{kl} N \Omega_{kp} \epsilon^p \chi_{lqr} P_{MN}^{mnqr} + \frac{3}{2} \mathcal{V}_{mn}^M \mathcal{V}^{kl} N \Omega_{kp} \epsilon^m \chi_{lqr} P_{MN}^{npqr} , \end{aligned} \quad (\text{B.7})$$

where we denote $p_{MN}^{ijkl} \equiv \partial_{(MPN)}^{ijkl}$. Next, one notes a non-trivial USp(8) identity

$$\begin{aligned} 0 &= \frac{1}{2} \mathcal{V}_{mn}^M \mathcal{V}^{kl} N \Omega_{pq} \epsilon^p \chi_{klr} P_{MN}^{mnqr} - \mathcal{V}^{ij} M \mathcal{V}^{kl} N \Omega^{mn} \epsilon^p \chi_{ikm} P_{MN}^{jlnp} \\ &\quad - \mathcal{V}_{mn}^M \mathcal{V}^{kl} N \Omega_{kp} \epsilon^q \chi_{lqr} P_{MN}^{mnpr} - \frac{1}{12} \mathcal{M}^{MN} \Omega_{ij} \epsilon^i \chi_{klm} P_{MN}^{jklm} \\ &\quad - \frac{1}{2} \mathcal{V}_{mn}^M \mathcal{V}^{kl} N \Omega_{kp} \epsilon^p \chi_{lqr} P_{MN}^{mnqr} + \mathcal{V}_{mn}^M \mathcal{V}^{kl} N \Omega_{kp} \epsilon^m \chi_{lqr} P_{MN}^{npqr} . \end{aligned} \quad (\text{B.8})$$

Finally, one employs the relations

$$\begin{aligned} \mathcal{V}_{mn}^M \mathcal{V}^{kl} N \Omega_{kp} \epsilon^q \chi_{lqr} P_{MN}^{mnpr} &= -\frac{1}{3} \mathcal{V}^{i[j} M \mathcal{V}^{l]k} N \Omega_{ik} \Omega_{lq} \epsilon^q \Omega^{rs} \chi_{rnm} P_{MN}^{mn}{}_{js} \\ &\quad - \mathcal{V}^{i[j} M \mathcal{V}^{l]k} N \Omega_{ik} \epsilon^r \chi_{lmn} P_{MN}^{mn}{}_{jr} , \end{aligned} \quad (\text{B.9})$$

and

$$\begin{aligned} \mathcal{V}_{mn}^M \mathcal{V}^{kl} N \Omega_{pq} \epsilon^p \chi_{klr} P_{MN}^{mnqr} &- 2 \mathcal{V}^{ij} M \mathcal{V}^{kl} N \Omega^{mn} \epsilon^p \chi_{ikm} P_{MN}^{jlnp} \\ &- \frac{1}{12} \mathcal{M}^{MN} \Omega_{ij} \epsilon^i \chi_{klm} P_{MN}^{jklm} \\ &= 2 \mathcal{V}^{i[j} M \mathcal{V}^{l]k} N \Omega_{ik} \epsilon^r \chi_{lmn} P_{MN}^{mn}{}_{jr} , \end{aligned}$$

of which both r.h.s. vanish due to the section constraint. Together, we conclude that the expression (B.7) contains

$$\longrightarrow \mathcal{V}_{mn}{}^M \mathcal{V}^{klN} \Omega_{p[q} \epsilon^p \chi_{ktr]} p_{MN}{}^{mnr} - \frac{1}{12} \mathcal{M}^{MN} \Omega_{ij} \epsilon^i \chi_{klm} p_{MN}{}^{jklm}, \quad (\text{B.10})$$

which is precisely the contribution from $-\frac{1}{4} \mathcal{R}^{ijkl}$ from (2.55).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] E. Cremmer and B. Julia, *The $N = 8$ Supergravity Theory. 1. The Lagrangian*, *Phys. Lett.* **B 80** (1978) 48 [[INSPIRE](#)].
- [2] O. Hohm and H. Samtleben, *Exceptional Form of $D = 11$ Supergravity*, *Phys. Rev. Lett.* **111** (2013) 231601 [[arXiv:1308.1673](#)] [[INSPIRE](#)].
- [3] O. Hohm and H. Samtleben, *Exceptional Field Theory I: $E_{6(6)}$ covariant Form of M -theory and Type IIB*, *Phys. Rev.* **D 89** (2014) 066016 [[arXiv:1312.0614](#)] [[INSPIRE](#)].
- [4] O. Hohm and H. Samtleben, *Exceptional field theory. II. $E_{7(7)}$* , *Phys. Rev.* **D 89** (2014) 066017 [[arXiv:1312.4542](#)] [[INSPIRE](#)].
- [5] O. Hohm and H. Samtleben, *Exceptional field theory. III. $E_{8(8)}$* , *Phys. Rev.* **D 90** (2014) 066002 [[arXiv:1406.3348](#)] [[INSPIRE](#)].
- [6] W. Siegel, *Superspace duality in low-energy superstrings*, *Phys. Rev.* **D 48** (1993) 2826 [[hep-th/9305073](#)] [[INSPIRE](#)].
- [7] C. Hull and B. Zwiebach, *Double field theory*, *JHEP* **09** (2009) 099 [[arXiv:0904.4664](#)] [[INSPIRE](#)].
- [8] C. Hull and B. Zwiebach, *The gauge algebra of double field theory and Courant brackets*, *JHEP* **09** (2009) 090 [[arXiv:0908.1792](#)] [[INSPIRE](#)].
- [9] O. Hohm, C. Hull and B. Zwiebach, *Background independent action for double field theory*, *JHEP* **07** (2010) 016 [[arXiv:1003.5027](#)] [[INSPIRE](#)].
- [10] O. Hohm, C. Hull and B. Zwiebach, *Generalized metric formulation of double field theory*, *JHEP* **08** (2010) 008 [[arXiv:1006.4823](#)] [[INSPIRE](#)].
- [11] D.S. Berman and M.J. Perry, *Generalized Geometry and M -theory*, *JHEP* **06** (2011) 074 [[arXiv:1008.1763](#)] [[INSPIRE](#)].
- [12] D.S. Berman, H. Godazgar and M.J. Perry, *$SO(5, 5)$ duality in M -theory and generalized geometry*, *Phys. Lett.* **B 700** (2011) 65 [[arXiv:1103.5733](#)] [[INSPIRE](#)].
- [13] D.S. Berman, H. Godazgar, M. Godazgar and M.J. Perry, *The local symmetries of M -theory and their formulation in generalised geometry*, *JHEP* **01** (2012) 012 [[arXiv:1110.3930](#)] [[INSPIRE](#)].
- [14] D.S. Berman, E.T. Musaev and M.J. Perry, *Boundary Terms in Generalized Geometry and doubled field theory*, *Phys. Lett.* **B 706** (2011) 228 [[arXiv:1110.3097](#)] [[INSPIRE](#)].

- [15] A. Coimbra, C. Strickland-Constable and D. Waldram, $E_{d(d)} \times \mathbb{R}^+$ generalised geometry, connections and M-theory, *JHEP* **02** (2014) 054 [[arXiv:1112.3989](#)] [[INSPIRE](#)].
- [16] A. Coimbra, C. Strickland-Constable and D. Waldram, Supergravity as Generalised Geometry II: $E_{d(d)} \times \mathbb{R}^+$ and M-theory, *JHEP* **03** (2014) 019 [[arXiv:1212.1586](#)] [[INSPIRE](#)].
- [17] G. Aldazabal, M. Graña, D. Marqués and J.A. Rosabal, Extended geometry and gauged maximal supergravity, *JHEP* **06** (2013) 046 [[arXiv:1302.5419](#)] [[INSPIRE](#)].
- [18] N. Hitchin, Generalized Calabi-Yau manifolds, *Quart. J. Math. Oxford Ser.* **54** (2003) 281 [[math/0209099](#)] [[INSPIRE](#)].
- [19] M. Gualtieri, Generalized complex geometry, [math/0401221](#) [[INSPIRE](#)].
- [20] P.C. West, E_{11} , $SL(32)$ and central charges, *Phys. Lett. B* **575** (2003) 333 [[hep-th/0307098](#)] [[INSPIRE](#)].
- [21] B. de Wit, H. Samtleben and M. Trigiante, The maximal $D = 5$ supergravities, *Nucl. Phys. B* **716** (2005) 215 [[hep-th/0412173](#)] [[INSPIRE](#)].
- [22] B. de Wit, H. Samtleben and M. Trigiante, The maximal $D = 4$ supergravities, *JHEP* **06** (2007) 049 [[arXiv:0705.2101](#)] [[INSPIRE](#)].
- [23] D.S. Berman, M. Cederwall, A. Kleinschmidt and D.C. Thompson, The gauge structure of generalised diffeomorphisms, *JHEP* **01** (2013) 064 [[arXiv:1208.5884](#)] [[INSPIRE](#)].
- [24] H. Godazgar, M. Godazgar, O. Hohm, H. Nicolai and H. Samtleben, Supersymmetric $E_{7(7)}$ Exceptional Field Theory, *JHEP* **09** (2014) 044 [[arXiv:1406.3235](#)] [[INSPIRE](#)].
- [25] E. Cremmer, Supergravities in 5 dimensions, in *Supergravities in diverse dimensions. Vol. 1*, A. Salam and E. Sezgin eds., World Scientific, Singapore (1980), pp. 422.
- [26] D.S. Berman, H. Godazgar, M.J. Perry and P. West, Duality Invariant Actions and Generalised Geometry, *JHEP* **02** (2012) 108 [[arXiv:1111.0459](#)] [[INSPIRE](#)].
- [27] M. Cederwall, J. Edlund and A. Karlsson, Exceptional geometry and tensor fields, *JHEP* **07** (2013) 028 [[arXiv:1302.6736](#)] [[INSPIRE](#)].
- [28] O. Hohm and H. Samtleben, U-duality covariant gravity, *JHEP* **09** (2013) 080 [[arXiv:1307.0509](#)] [[INSPIRE](#)].
- [29] C.D.A. Blair, E. Malek and J.-H. Park, M-theory and Type IIB from a Duality Manifest Action, *JHEP* **01** (2014) 172 [[arXiv:1311.5109](#)] [[INSPIRE](#)].
- [30] A. Abzalov, I. Bakhmatov and E. Musaev, Exceptional geometry for supergravity: groups $SO(5, 5)$ and $SL(5)$, to appear.
- [31] M. Graña and D. Marques, Gauged Double Field Theory, *JHEP* **04** (2012) 020 [[arXiv:1201.2924](#)] [[INSPIRE](#)].
- [32] D.S. Berman, E.T. Musaev, D.C. Thompson and D.C. Thompson, Duality Invariant M-theory: Gauged supergravities and Scherk-Schwarz reductions, *JHEP* **10** (2012) 174 [[arXiv:1208.0020](#)] [[INSPIRE](#)].
- [33] E.T. Musaev, Gauged supergravities in 5 and 6 dimensions from generalised Scherk-Schwarz reductions, *JHEP* **05** (2013) 161 [[arXiv:1301.0467](#)] [[INSPIRE](#)].
- [34] G. Dibitetto, J.J. Fernandez-Melgarejo, D. Marques and D. Roest, Duality orbits of non-geometric fluxes, *Fortsch. Phys.* **60** (2012) 1123 [[arXiv:1203.6562](#)] [[INSPIRE](#)].

- [35] O. Hohm and H. Samtleben, *Consistent Kaluza-Klein Truncations via Exceptional Field Theory*, *JHEP* **01** (2015) 131 [[arXiv:1410.8145](#)] [[INSPIRE](#)].
- [36] W.H. Baron and G. Dall'Agata, *Uplifting non-compact gauged supergravities*, *JHEP* **02** (2015) 003 [[arXiv:1410.8823](#)] [[INSPIRE](#)].
- [37] D.S. Berman, M. Cederwall and M.J. Perry, *Global aspects of double geometry*, *JHEP* **09** (2014) 066 [[arXiv:1401.1311](#)] [[INSPIRE](#)].
- [38] M. Cederwall, *The geometry behind double geometry*, *JHEP* **09** (2014) 070 [[arXiv:1402.2513](#)] [[INSPIRE](#)].
- [39] M. Cederwall, *T-duality and non-geometric solutions from double geometry*, *Fortsch. Phys.* **62** (2014) 942 [[arXiv:1409.4463](#)] [[INSPIRE](#)].
- [40] D.S. Berman and F.J. Rudolph, *Strings, Branes and the Self-dual Solutions of Exceptional Field Theory*, [arXiv:1412.2768](#) [[INSPIRE](#)].
- [41] A. Coimbra, C. Strickland-Constable and D. Waldram, *Supersymmetric Backgrounds and Generalised Special Holonomy*, [arXiv:1411.5721](#) [[INSPIRE](#)].
- [42] K. Peeters, *A Field-theory motivated approach to symbolic computer algebra*, *Comput. Phys. Commun.* **176** (2007) 550 [[cs/0608005](#)] [[INSPIRE](#)].
- [43] K. Peeters, *Introducing Cadabra: A Symbolic computer algebra system for field theory problems*, [hep-th/0701238](#) [[INSPIRE](#)].