Fifteen years of ocean observations with the global Argo array

Riser Stephen C. ¹, Freeland Howard J. ^{2,*}, Roemmich Dean ³, Wijffels Susan ⁴, Troisi Ariel ⁵, Belbeoch Mathieu ⁶, Gilbert Denis ⁷, Xu Jianping ⁸, Pouliquen Sylvie ⁹, Ann Thresher ⁴, Le Traon Pierre-Yves ^{10, 11}, Maze Guillaume ⁹, Klein Birgit ¹², Ravichandran M. ¹³, Grant Fiona ¹⁴, Poulain Pierre-Marie ¹⁵, Suga Toshio ^{16, 17}, Lim Byunghwan ¹⁸, Sterl Andreas ¹⁹, Sutton Philip ²⁰, Mork Kjell-Arne ²¹, Joaquin Velez-Belch Pedro ²², Ansorge Isabelle ²³, King Brian ²⁴, Turton Jon ²⁵, Baringer Molly ²⁶, Jayne Steven R. ²⁷

¹ Univ Washington, Dept Oceanog, Seattle, WA 98195 USA.

² Inst Ocean Sci, Fisheries & Oceans Canada, North Saanich, BC V8L 4B2, Canada.

³ Scripps Inst Oceanog, 9500 Gilman Dr,0230, La Jolla, CA 92093 USA.

⁴ CSIRO, Ctr Australian Weather & Climate Res, Hobart, Tas 7004, Australia.

⁵ Serv Hidrog Naval, A Montes Oca 2124, Buenos Aires C1270 ABV, Buenos Aires, DF, Argentina.

⁶ Joint Commiss Oceanog & Marine Meteorol Operat JC, BP 70, F-29280 Plouzane, France.

⁷ Inst Maurice Lamontagne, Fisheries & Oceans Canada, Mont Joli, PQ G5H 3Z4, Canada.

⁸ SOA, Inst Oceanog 2, 36 Baochubei Rd, Hangzhou 310012, Zhejiang, Peoples R China.

⁹ IFREMER, BP 70, Plouzane, France.

¹⁰ IFREMER, 8-10 Rue Hermes Parc Technol Canal, F-31520 Ramonville St Agne, France.

¹¹ Mercator Ocean, 8-10 Rue Hermes Parc Technol Canal, F-31520 Ramonville St Agne, France.

¹² Bundesamt Seeschifffahrt & Hydrog, Bernhard Nocht Str 78, D-20359 Hamburg, Germany.

¹³ Indian Natl Ctr Ocean Informat Serv, Hyderabad 500090, Andhra Pradesh, India.

¹⁴ Inst Marine, Int Programmes, Wilton Pk House, Wilton Pl, Dublin 2, Ireland.

¹⁵ OGS, 42-C Sgonico, I-20359 Trieste, Italy.

¹⁶ JAMSTEC, Aoba Ku, Aramaki Aza Aoba 6-3, Sendai, Miyagi 9808578, Japan.

¹⁷ Tohoku Univ, Aoba Ku, Aramaki Aza Aoba 6-3, Sendai, Miyagi 9808578, Japan.

¹⁸ Natl Inst Meteorol Sci KMA, 33 Seohobuk Ro, Seogwipo Si 63568, Jeju Do, South Korea.

¹⁹ KNMI, POB 201, NL-3730 AE De Bilt, Netherlands.

²⁰ Natl Inst Water & Atmospher Reseach, 301 Evans Bay Parade, Greta Point, Wellington 6021, New Zealand. ²¹ Marine Res Inst, POB 1870 Nordes, N-5817 Bergen, Norway.

²² Inst Espanol Oceanog, Via Espaldon, Darsena Pesquera, Parcela 8, Santa Cruz de Tenerife 38180, Spain.

Univ Cape Town, Inst Marine Res, Dept Oceanog, ZA-7701 Rondebosch, South Africa.

²⁴ Natl Oceanog Ctr, Southampton S014 3ZH, Hants, England.

²⁵ Met Off, FitzRoy Rd, Exeter EX1 3PB, Devon, England.

²⁶ NOAA, Atlantic Oceanog & Meteorol Lab, 4301 Rickenbacker Causeway, Miami, FL 33149 USA.

²⁷ Woods Hole Oceanog Inst, Woods Hole, MA 02543 USA.

* Corresponding author : Howard Freland, email address : Howard.Freeland@dfo-mpo.gc.ca

Abstract :

More than 90% of the heat energy accumulation in the climate system between 1971 and the present has been in the ocean. Thus, the ocean plays a crucial role in determining the climate of the planet. Observing the oceans is problematic even under the most favourable of conditions. Historically, shipboard ocean sampling has left vast expanses, particularly in the Southern Ocean, unobserved for long periods of time. Within the past 15 years, with the advent of the global Argo array of pro ling oats, it has become possible to sample the upper 2,000 m of the ocean globally and uniformly in space and time. The primary goal of Argo is to create a systematic global network of pro ling oats that can be integrated with other elements of the Global Ocean Observing System. The network provides freely available temperature and salinity data from the upper 2,000 m of the ocean with global coverage. The data are available within 24 hours of collection for use in a broad range of applications that focus on examining climate-relevant variability on seasonal to decadal timescales, multidecadal climate change, improved initialization of coupled ocean–atmosphere climate models and constraining ocean analysis and forecasting systems.

32 Introduction

In the late 1990s it was recognized that little progress would be made monitoring the changing climate of the Earth without systematic global observations. A small group of oceanographers proposed that some readilyavailable technology could be adapted and, with sufficient international cooperation, be used to create a global array of floats that could supply real-time global views of the ocean. Thus was born the Argo program. This article reviews the progress made in the first 15 years of Argo and attempts to provide an outline of how the program is likely to change over the next decade.

39

40 History and present status

The Argo program¹ is a major component of the Global Ocean Observing System² and strives to monitor the 41 42 evolving temperature and salinity fields of the upper ocean The profiling floats used in Argo are 2-meter long, 43 freely drifting robotic devices that adjust their depth in the ocean by changing their buoyancy. In the context of 44 Argo, the majority of these instruments are programmed to drift at a nominal depth of 1000 meters (known as 45 the parking depth). A typical duty cycle of an Argo float is shown in Figure 1. Floats are launched at the sea 46 surface and dive to a so-called parking depth of 1000 meters. After a ~9-day interval at the parking depth, the 47 floats descend to 2000 meters, then rise over a period of roughly 6 hours to the sea surface, sampling ocean 48 properties such as temperature and salinity during their ascent. At the surface the data are transmitted to land 49 stations via satellite; the floats then descend back to the parking depth to begin another cycle. A typical float will 50 repeat this 10-day cycle in excess of 200 times over the course of five years or more. Over 10,000 floats have 51 been deployed by the nations participating in the program since the beginning of Argo, with almost 3900 floats presently operating over the world-ocean. A defining aspect of Argo is that all data are reported in near real-52 time to meteorological forecasting centers and to the two Argo Global Data Assembly Centers (GDACs, 53 54 localized in the USA and France), from which the accumulated data are made freely available without 55 limitation.

Argo float deployments began in 1999. The original program plan called for an array with global coverage of 56 57 about 3000 floats by the year 2007, a goal that was achieved in November of that year. The resulting improvement in sampling the ocean achieved by Argo compared with the previous century of measurements is 58 59 dramatic (Fig. 2). Historically, oceanographic sampling has had a bias towards areas that are more easily 60 sampled, so the Northern Hemisphere and coastal regions are over-represented. Though Argo floats cannot 61 sample in shallow coastal waters, the program has helped eliminate these spatial biases. Argo has also 62 eliminated a major seasonal bias in sampling, particularly in the polar oceans; presently Argo delivers more 63 winter profiles in the Southern Ocean in one year than the total of all winter data collected during the 100 years

prior to Argo, and the majority of the temperature and salinity observations from the Southern Ocean (defined
here as the ocean south of 30°S) in the global database now originate from Argo.

In Chapter 3 of its Fifth Assessment Report (hereinafter AR5)⁵, the Intergovernmental Panel on Climate Change 66 67 (IPCC) Working Group 1 included an Appendix on the Availability of Observations for Assessment of Change 68 in the Oceans showing the distribution of temperature and salinity profile data from the 1950-1955 to the 2005-69 2010 pentads; these results indicate the general lack of observations prior to 2000, as well the Northern 70 Hemispheric bias within the pre-Argo observing system. Although not shown in AR5, there is also a known 71 seasonal bias, with the local winter season being generally under-represented in the pre-Argo observing system. 72 It is apparent from this report that the recent expansion of the ocean observing system clearly sets AR5 apart from its predecessor⁶, with most of this change ascribed to the existence of the Argo array. Some issues remain 73 74 that are being addressed. Since Argo floats must reach the sea surface to transmit their data, Argo could not 75 originally target ice-covered parts of the oceans; now floats are successfully operating in the Antarctic seasonal ice zone^{7,8}. Within the temperate oceans coverage is not homogeneous, but there are efforts to increase coverage 76 77 in under-sampled regions such as near western boundary currents.

78 While the volume of Argo data easily overwhelms traditional sources, the quantity is of value only if the quality 79 of the data is high. The original Argo target called for temperature and salinity accuracies of 0.005°C and 0.01 80 salinity units, with a pressure accuracy of 2.5 decibars (equivalent to a depth error of about 2.5 meters). 81 Experience has shown that about 80% of the raw profile data transmitted from the floats meet these standards, 82 with little or no correction required. The other 20% of the data are corrected using delayed-mode quality control procedures developed over the past decade^{9,10,11}, with nearly all of these profiles eventually meeting the 83 accuracy goals. Agencies developing operational forecast systems require data in a timely fashion. Currently, 84 about 90% of Argo profiles are distributed electronically within 24 hours of acquisition. 85

86

87 Recent Results and Findings

88 Perhaps the single most powerful metric of the value of Argo is the widespread use of the data produced by the 89 program: since the beginning of Argo in the late 1990s, more than 2100 papers in the refereed science literature 90 have used Argo observations, attesting to the array's value in expanding our understanding of the oceans and 91 climate. The key ocean state variables, temperature and salinity varying with pressure and velocity were 92 measured (or for velocity, mostly inferred) in the world ocean for at least a century prior to the initial 93 deployment of the Argo array. During the 1980s and 1990s the World Ocean Circulation Experiment (WOCE) 94 produced about 8000 high quality ship-based profiles along a number of tracks across the ocean that provide a 95 high quality baseline estimate of the state of the ocean at that time. However, a great deal of the ocean was left unobserved in WOCE due to the limited number of lines that could be occupied, particularly in the Southern
Ocean. In this context, the high sampling density and repeat frequency of the Argo observations, in conjunction
with their overall quality, have led to important new insights into the nature of the general circulation of the
ocean.

100 While compilations of thousands of ship-based observations of ocean temperature and salinity have been averaged and made available as digital atlases since at least the 1980s¹², they suffer from the sampling biases 101 previously noted. With data collected by Argo now becoming the overwhelming majority of observations in the 102 103 global database, there is an unprecedented opportunity to map the detailed structure of the global ocean temperature and salinity fields, at both surface and subsurface levels, and both globally and locally¹³. These 104 estimates allow the development of climate indicators such as the recent changes in ocean heat content and 105 thermosteric sea level. The global velocity dataset produced by Argo floats¹⁴ is beginning to be used in 106 systematic (and previously impossible) studies^{15,16} to quantify directly the subsurface interior flow fields. The 107 Argo data have also been used to improve our view of the complex structure of oceanic variability at spatial 108 109 scales smaller than the climate scale. A number of studies have examined properties of mesoscale eddies using Argo profiles^{17,18}, and it has been possible to describe the spatial variability in the internal gravity wave field¹⁹, 110 even though such applications were not envisioned when Argo began. 111

112 Argo complements the other elements of the global ocean observing system, in particular satellite altimetry. 113 The combination of in situ Argo data with sea surface height anomalies derived from satellites allows the construction of time series of the dynamical state of the ocean circulation (such as the North Atlantic meridional 114 overturning circulation²⁰) and high-resolution three-dimensional temperature fields²¹. Techniques have been 115 presented^{22,23} whereby geostrophic computations relative to 1000 decibars can be merged with a combination of 116 117 satellite altimetry observations and trajectory data to yield estimates of the absolute circulation fields, in wellsampled areas. Argo data are now systematically used together with altimeter data for ocean analysis and 118 119 forecasting. Most climate models assimilate the subsurface temperature observations from Argo, leading to improved forecasts of intraseasonal waves in the atmosphere, monsoon activity, and ocean-atmosphere 120 interactions such as the El Nino Southern Oscillation (ENSO)²²⁴. 121

Given that they provide a comprehensive baseline of today's ocean state, the Argo observations have been especially useful in examining ocean changes on time scales of decades and longer. A stunning example²⁵ has used contemporary Argo observations in conjunction with data from the HMS *Challenger* expedition, carried out in the second half of the 19th Century. The study reveals a warming of the ocean over the past 135 years of nearly 0.6°C near the sea surface, tapering to near zero at depths near 1000 m (Fig. 3). Over the upper 900 m of much of the ocean there is an average increase in temperature of over 0.3°C over the 135-year study interval. This work, further supported by analogous results in climate models²⁶, underscores the changing nature of ocean properties and the need to sustain global observing systems over long periods and, further, that recent changes inocean temperature likely predate the sparse global-scale ocean datasets of the past half century.

A more detailed comparison of Argo profiles available in the time frame 2004-2008 with data in the ship-based data archive (containing data collected through 2001²⁷) shows global-scale changes in both temperature and salinity²⁸ in the upper kilometer of the world ocean in recent decades. The comparison suggests that most regions of the world ocean are warmer in the near surface-layer than in previous decades, by over 1°C in some places. A few areas, such as the eastern Pacific from Chile to Alaska, have cooled by as much as 1°C, yet overall the upper ocean globally has warmed by nearly 0.2°C since the mid-20th century. A recent update of this work²⁹ shows the warming continues to be observed through at least 2013.

Prior to the beginning of Argo, ocean salinity was much more poorly sampled than temperature. The highest 138 139 quality salinity measurements that existed prior to Argo were made during WOCE in the 1980s and 1990s, with 140 periodic reoccupation of some of these sections continuing. With the advent of Argo, it is possible to examine the global variability of salinity above 2000 m with some confidence for the first time. As with temperature, a 141 comparison of Argo results with earlier data³⁰ clearly shows decadal-scale changes in ocean salinity. A related 142 study that compared the Argo results with the data in the older archives³¹ identified a pattern of change in upper 143 ocean salinity in each of the major ocean basins. This change appears consistent with a general warming of the 144 145 surface layers, causing a poleward migration of wintertime density outcrops. Because there are pole-to-equator 146 temperature and salinity gradients at the sea surface, the surface waters at any latitude are subducted below those 147 at lower latitudes as they circulate towards the equator, with a poleward migration of density yielding a generally lower salinity at higher latitudes. At mid-latitudes, surface warming is accompanied by increased 148 149 evaporation from the sea surface, thus increasing the salinity of the near-surface waters. When compared, surface and basin-scale contrasts in salinity are found consistently to be strengthening 29,30,31 . This amplification, 150 151 with fresh areas getting fresher and salty areas becoming more saline, is consistent with an overall net increase 152 of water vapor transport through the atmosphere, and is also seen to occur in climate models that very closely follow Clausius-Clapeyron thermodynamics³². 153

154 On more regional scales, Argo data have been used to examine the variable nature of deep ocean ventilation, which is important in sequestering heat and gases into the vast subsurface layers of the ocean. This process 155 occurs at a few high latitude sites in the North Atlantic and the Antarctic, and the resulting circulation induced 156 by this sinking has long been thought to have an important influence on century-scale climate variability³³. The 157 Labrador/Irminger Sea region in the high-latitude N. Atlantic has been extensively examined by Argo in the past 158 159 decade, with the observations (combined with ship-based datasets and atmospheric heat flux estimates) showing the high sensitivity of the strength of deep convection to year-to-year variations in the wintertime atmospheric 160 conditions³⁴. That work has recently been brought up to date³⁵ and the result is a spectacular demonstration (Fig. 161

4) of the power of Argo to show variability of deep convection in the Labrador Sea and in 2014 the deepestconvection observed since 2008.

164 More detailed, regional views of the effects of changing near-surface temperature and salinity, from the central 165 North Pacific, using Argo profiles together with older, ship-based data in the region, have shown a general 166 decrease in upper ocean salinity in the tropics and subtropics, caused by a freshening of the source waters for the upper ocean in the region, essentially water in the surface mixed layer at higher latitudes in winter^{36,37}. This 167 168 freshening can be attributed to a combination of factors, including increased local precipitation or the poleward migration of the source region due to regional warming. The data availability in this region (including Argo and 169 170 other datasets) is rich enough to allow an actual heat budget to be constructed, suggesting that precipitation and 171 wintertime entrainment of subsurface water into the mixed layer are the most important factors in determining 172 the surface density over time, and hence also the surface salinity. It appears likely that these factors will change 173 in response to a changing climate state. Elsewhere in the North Pacific, temperature and salinity properties along a track originating off Vancouver Island have been regularly surveyed by research vessels since the 1950s, 174 175 and in the past decade these surveys have been augmented by Argo observations. The combination of the older data and Argo³⁸ also show that subpolar surface waters are warming and freshening, resulting in a lower surface 176 density (and higher near-surface stratification), thus limiting the direct influence of the atmosphere on the 177 178 subsurface ocean.

179 As noted, a number of observation-based studies have concluded that the upper layers of the global ocean have warmed in recent decades, or even over a longer term. Yet it is the change in heat storage by the ocean 180 181 (essentially the vertical integral of temperature change) that is likely to be directly related to the planetary radiation imbalance^{39,40}. It appears that the observed increase in ocean heat content in the upper 700 m over the 182 183 past 40 years, inferred from Argo and hydrographic observations, is the dominant term in the global inventory of heating changes, with over 90% of the excess heat in the climate system being stored in the oceans⁴¹. Without 184 Argo, it is unlikely that such a conclusion could have been drawn. The data have also allowed temporal spatial 185 variations in ocean heat content to be discerned⁴², suggesting that most of the increase in heat content in the past 186 decade has occurred in the Southern Ocean (which was poorly sampled prior to Argo); it has also been noted⁴³ 187 188 that ENSO variability in the tropical Pacific has for now somewhat obscured the global increase in sea surface temperature. Some of these ideas have been tested using Argo data alone (Fig. 5^{29}), by examining the change in 189 heat content of the upper 2000 m of the ocean in three latitudinal bands from 2006 to the present time; during 190 the Argo era it can be seen that the ocean is warming, mostly south of 20°S. In Figure 5 the Argo-only plots 191 192 (shown as an inset) are only for the years 2006-2014, inclusive, the period when global coverage from Argo exists. So these short plots are overlaid on a plot of heat content estimates 0-2000 m for the period 1955-2010⁴⁴. 193 194 The Argo estimates show a very similar trend. This is a crucial result in making an assessment of the ocean's role in climate change, one that would have likely been impossible prior to Argo. Indeed, a recent detailed and systematic analysis of the change in ocean heat content and our ability to observe it⁴⁵ concludes that estimates of the upper ocean heat content based on data collected prior to Argo systematically underestimate the amount of heat content change, mostly due to under-sampling of the Southern Hemisphere ocean; with the advent of a significant amount of Argo data becoming available around 2004, the estimates are greatly improved and show much improved agreement with climate models.

201

202 Future prospects

203 When Argo began in the late 1990s, it was by no means clear that the project would be successful in deploying 204 and sustaining an array of 3000 floats over the global ocean, as the required technology was in its infancy and 205 the degree of international cooperation required was unprecedented in the oceanographic community. Now, in 206 the second decade of the 21st century, the float technology is well proven, and over 30 countries are contributing 207 resources to Argo, making it sensible to contemplate expansions of its mission. The Argo Steering Team (hereinafter AST) has provided a roadmap⁴⁶ for how the project might evolve and expand in the next decade, 208 209 and some of this proposed development is now underway via test deployments or regional pilot arrays. One project is to support an increase in the spatial sampling resolution in special parts of the world ocean where the 210 211 ocean is especially turbulent (challenging the array's signal resolving power) and the interaction of the ocean 212 and the atmosphere and the resulting climate impacts are especially strong. Improved technology also allows us 213 to expand Argo into previously unsampled regions such as marginal seas and the seasonal ice-zone, implying 214 that the array is more truly global in its coverage than its original design.

215 Intense western boundary currents such as the Gulf Stream and Kuroshio are some of the most striking features 216 of the ocean circulation, serving to transport heat poleward as they carry warm tropical and subtropical waters to 217 higher latitudes. A sizeable portion of this heat is exchanged with the atmosphere along the path of the flow, especially after these currents separate from the boundary and enter the interior of the ocean, altering the tracks 218 of storms and perhaps ameliorating continental climates⁴⁷. When Argo data are used to map monthly and 219 220 seasonal evolution of the large-scale ocean structures in these regions, the errors are relatively high compared to 221 other regions due to the presence of the vigorous ocean turbulence associated with these boundary current 222 regimes and the lack of spatial resolution of the Argo array. This limits the ability of Argo to achieve one of its 223 main goals, which is to observe the slow evolution of the large-scale ocean structure in these important regions. 224 The AST has suggested that extra float resources should be committed to the study of these regions, and some of 225 the nations that contribute to the array have responded to this call. In the western North Pacific, for example, 226 the boundary domain adjacent to China, South Korea, and Japan is already sampled at a spatial density nearly 227 three times the standard Argo protocol. In the western North Atlantic, between Cape Hatteras and the midAtlantic Ridge, the sampling is similarly dense. The ability of the program to enhance sampling in all western
boundary current regions is unclear, but these pilot activities will help demonstrate the gains made and thus
guide the future sampling recommendations.

231 A similar increase in sampling density is called for in near-Equatorial regions, where strong air-sea coupling can 232 produce global-scale atmospheric anomalies. The tropical Pacific, for example, is the gestation site of the ENSO phenomenon, an irregular seesaw of oceanic warming and cooling that can, via strong coupled 233 234 amplification, drive global climate variability and cause economic impacts at sites far removed from the low-235 latitude Pacific. The Indian Ocean Dipole (IOD) is an analogous phenomenon that impacts Indian Ocean rim nations⁴⁶. The surface temperature and salinity associated with these phenomena can be monitored via satellites, 236 237 but Argo is now the main source of information on their deeper, subsurface expressions. Argo data are used in 238 model forecasts for ENSO and the IOD, and, along with the existing tropical moored arrays, are central to 239 providing and improving predictions of the onset and strength of these scientifically interesting yet socially 240 disruptive events.

Additionally, the AST has encouraged the deployment of floats in marginal seas, environments that are 241 242 regionally important to natural resources and trade. During the early stages of Argo it was decided to avoid deployments in such areas, due to the likely premature loss of floats and troublesome political issues. More than 243 244 a decade later some of these issues remain, yet the scientific success of Argo has encouraged a number of groups 245 to begin Argo-like programs in several of the marginal seas of the world. Within the Euro-Argo European 246 Research Infrastructure Consortium that is coordinating European contributions to Argo, the MedArgo initiative 247 has maintained an array of more than 50 floats in these regions since 2008 and has plans for expansion. This has 248 helped to clarify the details of water mass formation in the Mediterranean and to improve predictions of the 249 basin-scale circulation by assimilating profile data into numerical models of the circulation. Argo-type efforts in 250 the marginal seas of East Asia have yielded new insights into the interplay of the local ocean and the regional 251 climate in these regions and the nature and causes of long-term changes in their properties, and data resulting 252 from deployments now underway in the Gulf of Mexico will likely eventually be useful in improved hurricane 253 prediction.

Argo was originally planned to cover the open ocean equatorward of 60°, thus purposefully avoiding regions seasonally covered in sea ice. A float trapped under sea ice cannot transmit its data, and, for floats that are somehow able to get to the surface in the presence of ice, the danger of being crushed between floes is sizable. In the past decade algorithms have been developed⁴⁸ that greatly increase a float's chances of surviving the winter in the ice zone by inferring the presence of ice from the near-surface temperature structure. Simply stated, if the float determines that ice is present through an on-board analysis of the stratification, then it avoids the surface (thus avoiding the ice), stores its profile and descends for another cycle. In the spring, when the ice retreats and the float can reach the surface, all of its winter data can be transmitted, though without tracking information. This concept has worked well^{7,8} with floats in the seasonal ice zone surviving at about the same rate as floats at lower latitudes. Based on this success, the large fraction of the area of the global ocean that exists at high southern latitudes, and the importance of ice-covered regions around the Antarctic in setting the conditions for meridional overturning, the AST has suggested that deployments in such regions should be systematically increased. Several float deployment groups are now staging pilot deployments in these regions.

267 Beyond deployments in new regions and advances in float software, technological developments have provided 268 new capabilities to profiling floats that allow a larger range of scientific questions to be examined. As 269 originally designed, Argo was intended to be a program to examine the temperature, salinity, and heat content of 270 the world ocean and the climate implications of the variability in these quantities. In the past decade new 271 sensors have come into existence that also allow biogeochemical variables such as dissolved oxygen, nitrate, 272 chlorophyll, and pH to be measured from Argo-type floats. These sensors are small in both physical size and power consumption and barely affect the duration of the basic float mission, and thus floats equipped with such 273 274 sensors allow the prospect of examining the impact of physical aspects of the ocean circulation on key 275 biogeochemical processes sensitive to the climate state, such as the biogeochemical cycle of carbon, ocean deoxygenation, and ocean acidification⁵⁰. These floats will also contribute to the improvement of the capabilities 276 of biogeochemical models⁵¹ and to the extension of surface biogeochemical properties monitored by ocean color 277 satellites to deeper levels of the water column⁵². The AST has endorsed the general idea of taking advantage of 278 279 these new developments, while at the same time moving cautiously as the Argo data system gains experience in 280 ingesting and disseminating these new data types. While it is expected that the number of biogeochemical Argo floats deployed will increase dramatically in the coming years, pilot projects in a few key oceanic areas where 281 282 carbon uptake is crucial will be carried out before envisaging the implementation of a biogeochemical float 283 program at the global scale. Several European efforts in this regard are underway in the North Atlantic. 284 Similarly, the mating of the under-ice capabilities and biogeochemical floats has led to plans by the Southern Ocean Carbon and Climate Observations and Models (SOCCOM) group to examine the carbon cycle around the 285 Southern Ocean, a region responsible for roughly 40% of the global oceanic carbon uptake.^{53,54} 286

In recent years a series of papers^{55,56,57,58} have shown that the deep ocean well below 2000 m contributes a significant fraction of the total water column increases in heat content and thermosteric sea level rise, especially in (but not confined to) the high southern latitudes. This warming, in many cases present at all depths below 3000 m, has been deduced from an analysis of sparse but repeated high-quality ship-based observations of temperature and salinity conducted since the 1980s. While the capability of making such measurements from ships has existed for several decades, the cost of the vessels carrying out such work is high, often as much as US\$35K per day. Argo floats, on the other hand, are a seemingly more economical way to observe the ocean, 294 although the present generation of Argo floats generally samples no deeper than 2000 m. The economy may or 295 may not be illusory depending on the strategy used to deploy floats. If floats are deployed on a ship of 296 opportunity there may be no incremental launch costs, but strategic launches do often require a dedicated vessel. 297 In order to begin to explore the abyssal ocean and to refine present estimates of the warming of the deep sea, the 298 AST, has since 2012, supported efforts to develop floats capable of profiling deeper than 2000 m (to as deep as 299 6000 m) and to begin deploying these floats increasingly in 2015 and beyond. This is by far the most ambitious, 300 and technically challenging, development in Argo since the initial float deployments took place in the late 1990s. At the present time several prototypes have been tested that use new technologies, such as carbon-epoxy 301 302 filament wound cylindrical hulls for a 4000 m prototypes being developed in Europe and Japan, or hulls made 303 from glass spheres, for two 6000 m versions being developed in the US. An additional issue is that the 304 temperature, salinity, and pressure sensors on these floats must be more accurate than for standard Argo floats, 305 since the variability in the abyssal ocean is likely to be considerably smaller than in the upper 2000 m of the 306 water column. Prototypes for the 4000 m versions have been successfully deployed in the N. Atlantic and N. 307 Pacific. A dedicated test cruise for deployment of 6000 m prototype floats and for calibration and validation of 308 new ultra-accurate temperature/salinity/pressure sensors took place in the western S. Pacific in mid-2014 and 309 successfully showed the promise of both the new deep floats and sensors. While the deep floats will surely be 310 more costly than present Argo floats, it is hoped that by 2020 systematic sampling of the abyssal ocean will be 311 implemented, with as much as 30% of the Argo array comprised of these deep floats. High-quality sampling of 312 the deep sea from dedicated hydrographic vessels will need to continue far into the future, however, even after 313 the full complement of deep floats is in place; the shipboard surveys together with Argo will provide a more 314 complete global sampling of the abyssal ocean, and the ship-based data will be necessary to monitor the quality 315 of the Argo data in the abyss.

It is possible that a decade or two from now the science community will barely recognize the deployment strategies or the instruments being deployed by Argo or its successor programs at that time. For the next few years, the AST has suggested continuing with its long-term plan while also moving in new directions that are part evolution (deploying floats in new and special regions, and the addition of new sensors) and part revolution (designing and testing floats capable of operation in the abyss). The goals of all of this work are to sustain the present systematic observations of the global ocean and to further improve our assessment of the ocean's role in climate.

The development of Deep Argo appears to be an especially crucial step in the evolution of the program. The present array has done a credible job of providing estimates of the heat content change in the upper 2000 m of the world ocean over the past few decades, with the measured increases being significant but somewhat smaller than the predictions of model-based global heat budgets. Other studies^{57,58} point to the even deeper waters of the world ocean, beneath the present limits of Argo sampling, as a possible repository of the increased heating

necessary to close the global heat budget. Assessing this question, and making progress in other new directions

such as observing the oceanic carbon cycle from floats, helping to make improved forecasts of ENSO and IOD,

- and probing the depths of marginal seas, will likely challenge the abilities of future leaders of Argo to push the
- 331 limits of evolution and revolution of the program in order to further increase our understanding of the circulation
- 332 of the ocean and its role in climate.
- 333 In conclusion, it is important to remember that the prize we are aiming for is an understanding of how the ocean
- is changing both naturally and as a result of human activities. That said, Argo is indeed a major accomplishment
- but it does not stand alone. Recently five programs have been brought together in the JCOMMOPS (Joint
- 336 Commission on Oceanography and Marine Meteorology Operations) office. This includes moored buoy
- activities such as the equatorial arrays and the individual moorings sponsored by the OceanSites program, also
- the XBT surveys and the repeat ship-board hydrographic surveys sponsored by the GO-SHIP program. Many of
- these supply data in real-time in parallel with the Argo program. With all of these programs now operating
- under a single umbrella the supply of data from the oceans is undergoing a profound transformation.
- 341

Author Contributions: The paper was written by SCR (60%) and HJF(40%), with editorial comments and
 suggestions provided by the remaining co-authors whohave all served as members of the Argo Steering Team
 and so have contributed significantly to the Argo program.

345

346 **Author Affiliations:**

- 347 Dept. of Oceanography, University of Washington, Seattle, WA 98195, U.S.A.
- 348 Steve Riser349
- Fisheries and Oceans Canada, Institute of Ocean Sciences, North Saanich, BC V8L 4B2 CANADA
 Howard Freeland
- 352
- Scripps Institution of Oceanography, 9500 Gilman Dr.,#0230, La Jolla, CA 92093-0230, USA
 Dean Roemmich
- 355
- Centre for Australian Weather and Climate Research, CSIRO, Hobart, Tasmania, 7004, Australia
 Susan Wijffels and Ann Thresher
- Servicio de Hidrografia Naval, A. Montes de Oca 2124, Buenos Aires, ARGENTINA
 Ariel Troisi
- 361

- 362 JCOMMOPS, BP70, Plouzane, 29280 FRANCE
- 363 Mathieu Belbeoch
- 364
- 365 Denis Gilbert
- 366 Fisheries and Oceans Canada, Institut Maurice-Lamontagne, Mont-Joli, Quebec, G5H 3Z4 Canada
- 367

368 369 370	The Second Institute of Oceanography, SOA, No.36 Baochubei Road, Hangzhou, Zhejiang, CHINA 310012 Jianping Xu
371 372	IFREMER & MERCATOR OCEAN, 8-10 rue Hermes Parc Technologique du Canal, Ramonville St. Agne, FRANCE 31520
373 374	Pierre-Yves Le Traon
375 376 377	IFREMER, BP70, Plouzané, 29280 FRANCE Guillaume Maze and Sylvie Pouliquen
378 379 380	Bundesamt fuer Seeschifffahrt und Hydrographie, bernhard-Nocht-str., 78, Hamburg, Germany 20359 Birgit Klein
381 382 383	Indian National Centre for Ocean Information Services, Hyderabad, Andhra Pradesh, 500090, India M. Ravichandran
384 385 386	International Programmes, Marine Institute, Wilton Park House, Wilton Place, Dublin 2, Ireland Fiona Grant
387 388 389	OGS, Borgo Grotta Gigante, 42/c, Sgonico, Trieste, ITALY 34010 Pierre-Marie Poulain
390 391 392	JAMSTEC and Tohoku University, Aramaki-Aza-Aoba 6-3, Aoba-Ku, Sendai, Miyagi, 980-8578 Japan Toshio Suga
393 394	National Institute of Meteorological Research/KMA, 45 Gisangcheong-gil, Dongjak-gu, Seoul, KOREA, 156-720
395 396	Byunghwan Lim
397 398 399	KNMI, P.O. Box 201, 3730 AE de Bilt, Netherlands Andreas Sterl
400 401 402	National Institute of Water and Atmospheric Reseach, 301 Evans Bay Parade, Greta Point, Wellington, NZ, 6021 Philip Sutton
403 404 405 406	Institute of Marine Research, P.O. Box 1870 Nordnes, 5817 Bergen, Norway Kjell-Arne Mork
400 407 408	Instituto Español de Oceanografía, Vía Espaldón, Dársena Pesquera, Parcela 8, 38180 Santa Cruz de Tenerife, España
409 410	Pedro Joaquín Vélez-Belchí
411 412 413	University of Cape Town, 7701 Rondebosch, South Africa Isabelle Ansorge
414 415 416	National Oceanography Centre, Southampton, Empress Dock, Southampton, Hampshire, S014 3ZH UK Brian King
417 418	Met Office, FitzRoy Rd, Exeter, Devon EX1 3PB UK Jon Turton

- 419
 420 AOML/NOAA, 4301 Rickenbacker Causeway, Miami, FL 33149 USA
 421 Molly Baringer
 422
 423 Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA
 424 Steve Jayne
 425
 426
 427 Author Information: All data used in this paper are freely available
- 427 Author Information: All data used in this paper are freely available from the Argo Global Data Assembly 428 Center at *http://www.coriolis.eu.org/Data-Products*. The authors of this work have no competing financial 429 interests related to the material published here. Correspondence and requests for materials should be addressed 430 to HJF at *howard.freeland@dfo-mpo.gc.ca*. Reprints and permissions information is available at 431 *http://www.nature.com/reprints*.

- Acknowledgments: Argo data were collected and made freely available by the International Argo Program and
 the national programs that contribute to it. (http://www.argo.ucsd.edu, http://argo.jcommops.org). The Argo
 Program is part of the Global Ocean Observing System.
- 436
- 437
- 438
- 439

References

- 442 1. Argo Science Team. On the design and Implementation of Argo An initial Plan for a global array of
- 443 *Profiling Floats.* International CLIVAR project Office ICPO Report No.21. GODAE Report No 5. Published by
- the GODAE International Project office, c/o Bureau of Meteorology, Melbourne, Australia, 32pp (1998).
- 445 2. Intergovernmental Oceanographic Commission (IOC) Toward a global ocean observing system: the
 446 approach to GOOS. IOC publication IOC-XVII/8, UNESCO, Paris (1993).
- 447 3. Boyer, T. et al. World Ocean Database 2009. S. Levitus, ed. *NOAA Atlas NESDIS* 66, U.S. Gov.
- 448 Printing Office, Wash., D.C., 216 pp (2009).
- 449 4. <u>http://dx.doi.org/10.12770/ca035889-880d-463e-a523-10aabc3d6be3</u>
- 450 5. Intergovernmental Panel on Climate Change (IPCC). *Climate Change 2014: Synthesis Report*. IPCC
- 451 Geneva, pp 151 (2014).
- 452 6. Intergovernmental Panel on Climate Change (IPCC). *Climate Change 2007: Synthesis Report*. IPCC
 453 Geneva, 104 pp (2007).
- 454 7. Wong, A. & Riser, S. Profiling float observations of the upper ocean under sea ice off the Wilkes Land coast
 455 of Antarctica. *Journal of Physical Oceanography*, 41, 1102-1115 (2011).
- 456 8. Wong, A. & Riser, S. Modified shelf water on the continental slope north of Mac Robertson Land, East
 457 Antarctica. *Geophysical Research Letters*, 40, 6186-6190 (2013).
- Wong, A., Johnson G., & Owens, W.B. Delayed-mode calibration of autonomous CTD profiling float
 salinity data by theta-S climatology. *Journal of Atmospheric and Oceanic Technology*, 20, 308-318 (2003).
- 460 10. Owens, W.B. & Wong, A. An improved calibration method for the drift of the conductivity sensor on
 461 autonomous CTD profiling floats by theta-S climatology, *Deep-Sea Research Part I: Oceanographic Research*462 *Papers*, 56, 450-457 (2009).
- 463 11. Gaillard, F. et al. Quality Control of Large Argo Datasets. *Journal of Atmospheric and Oceanic Technology*,
 464 26, 337-351(2009).
- 465 12. Levitus, S. Climatological Atlas of the World Ocean, NOAA Professional Paper 13, U.S. Gov. Printing
 466 Office, Rockville, M.D., 190 pp (1982).
- von Schuckmann, K., Gaillard, F., & Le Traon, P-Y. Global hydrographic variability patterns during 2003-*Journal of Geophysical Research-Oceans*, **114**, C09007, doi:10.1029/2008JC005237 (2009).

- 469 14. Ollitrault, M. & Rannou, J-P. ANDRO: An Argo-based deep displacement dataset. *Journal of Atmospheric*470 *and Oceanic Technology*, **30**, 759-788 (2013).
- 471 15. Ollitrault, M. & De Verdiere, C. The ocean general circulation near 1000 m depth. *Journal of Physical*472 *Oceanography*, 44, 384-409 (2014).
- 473 16. Gray, A. & Riser, S. A global analysis of Sverdrup balance using absolute geostrophic velocities from
 474 Argo. *Journal of Physical Oceanography*, 44, 1213–1229 (2014).
- 475 17. Castelao, R. Mesoscale eddies in the South Atlantic Bight and the Gulf Stream recirculation region: vertical
 476 structure. *Journal of Geophysical Research-Oceans*, doi: 10.1002/2014JC009796 (2014).
- 477 18. Zhang, Z., Wang, W., & Qiu, B. Oceanic mass transport by mesoscale eddies. *Science*, 345, 322-324
 478 (2014).
- 479 19. Hennon, T., Riser, S., & Alford, M. Observations of internal gravity waves from Argo floats. *Journal of*480 *Physical Oceanography*, 44, 2370-2386 (2014).
- 481 20. Mercier, H. et al. Variability of the meridional overturning circulation at the Greenland--Portugal OVIDE
 482 section from 1993 to 2010. *Progress in Oceanography*, **132**, 250-261 (2015).
- 483 21. Guinehut, S., Dhomps A., Larnicol, G., & Le Traon, P-Y. High resolution 3-D temperature and salinity
 484 fields derived from in situ and satellite observations. *Ocean Science*, 68, 845-857 (2012).
- 485 22. Willis, J. K., and L.-L. Fu (2008), Combining altimeter and subsurface float data to estimate the 486 timeaveraged circulation in the upper ocean, J. Geophys. Res., 113, C12017, doi:10.1029/2007JC004690.
- 487 23. Willis, (2010). GEOPHYSICAL RESEARCH LETTERS, VOL. 37, L06602, doi:10.1029/2010GL042372, 2010
- 488 24. Chang, Y., Zhang, S., Rosati, A., Delworth, T., and Stern, W. An assessment of oceanic variability for
- 1960-2010 from the GFDL ensemble coupled data assimilation. *Climate Dynamics*, **40**, 775-803 (2013).
- 25. Roemmich, D, Gould, W.J., & Gilson, J. 135 years of global ocean warming between the Challenger
 expedition and the Argo Programme. *Nature Climate Change*, 2, 425-428 (2012).
- 492 26. Hobbs, W. & Willis, J. Detection of an observed 135 year ocean temperature change from limited data.
 493 *Geophysical Research Letters*, 40, 2252-2258 (2013).
- 494 27. Levitus, S., et al. Anthropogenic warming of earth's climate system. *Science*, **292**, 267-270 (2001).
- 495 28. Roemmich, D, & Gilson, J. The 2004-2008 mean and annual cycle of temperature, salinity, and steric
- 496 height in the global ocean from the Argo Program. *Progress in Oceanography*, **82**, 81-100 (2009).

- 497 29. Roemmich, D. et al. Unabated Planetary Warming and its Anatomy since 2006. *Nature Climate Change*, 5,
 498 240-245 (2015).
- 30. Durack, P. & Wijffels, S. Fifty-year trends in global ocean salinities and their relationship to broad-scale
 warming. *Journal of Climate*, 23, 4342-4362 (2010).
- 31. Hosoda, S., Suga, T., Shikama, N., and Mizuno, K. Global surface layer salinity change detected by Argo
 and its implication for hydrological cycle intensification. *Journal of Oceanography*, 65, 579-586 (2009).
- 503 32. Durack, P., Wijffels, S. & Matear, R. Ocean salinities reveal strong global water cycle intensification during
 504 1950-2000. *Science*, 336, 455-458 (2012).
- 505 33. Manabe, S. & Stouffer, R. Multiple-century response of a coupled ocean-atmosphere model to an increase
- of atmospheric carbon dioxide. *Journal of Climate*, 7, 5-23 (1994).
- 34. Yashayaev, I. & and Loder, J. Enhanced production of Labrador Sea Water in 2008. *Geophysical Research Letters*, 36, doi:10.1029/2008GL036162 (2009).
- 509 35. Kieke, D., Yashayaev, I. Studies of Labrador Sea Water formation and variability in the subpolar North
- 510 Atlantic in the light of international partnership and collaboration. Prog. Oceanogr. (2015),
- 511 doi:10.1016/j.pocean.2014.12.010
- 36. Ren, L. & Riser, S. Seasonal salt budget in the northeast Pacific Ocean. *Journal of Geophysical Research*, 114, doi:10.1029/2009JC005307 (2009).
- S14 37. Ren, L. & Riser, S. Observations of decadal-scale salinity changes in the thermocline of the North Pacific
 S15 Ocean. *Deep-Sea Research II*, 57, 1161-1170 (2010).
- 516 38. Freeland, H. Evidence of change in the winter mixed layer in the Northeast Pacific Ocean: a problem
 517 revisited. *Atmosphere-Ocean*, 51, 126-133 (2013).
- 518 39. Hansen, J., Lacis, A., & Rind, D. Climate trends due to increasing greenhouse gases. In: Proceedings of the
- *Third Symposium on Coastal and Ocean Management*, ASCE/San Diego, California, June 1-4, 1983, 2796-2810
 (1984).
- 40. Palmer, M., McNeall, D., & Dunstone, N. Importance of the deep ocean for estimating decadal changes in
 Earth's radiation balance. *Geophysical Research Letters*, 38, doi:10.1029/2011GL047835 (2011).
- 41. Rhein, M. et al. Observations: Ocean. In: *Climate Change 2013: The Physical Science Basis. Contribution*
- 524 of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.
- 525 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA (2013).

- 526 42. Sutton, P. & Roemmich, D. Decadal steric and sea surface height changes in the Southern Hemisphere.
 527 *Geophyical Research Letters*, **38**, doi: 10.1029/2011GL046802 (2011).
- 43. Kosaka, Y. & and Xie, S-P. Recent global-warming hiatus tied to equatorial Pacific surface cooling. *Nature*, 501,
 403–407 (2013).
- 45. Durack, P., Gleckler, P., Landerer, F., & Taylor, K. Quantifying underestimates of long-term upper ocean
 warming. *Nature Climate Change*, doi:10.1038/NCLIMATE2389 (2014).
- 46. Freeland, H. et al. Argo-a decade of progress. In: *Proceedings of OceanObs'09: Sustained Ocean Observations and Information for Society* (V2), Venice, Italy, 21-25 September 2009: ESA Publication WPP306 10.5270/OceanObs09.cwp.32 (2010).
- 47. Kwon, Y-O. et al. Role of Gulf Stream, Kuroshio-Oyashio, and their extensions in large-scale atmosphereocean interaction : a review. *Journal of Climate*, **23**, 3249-3281 (2010).
- 537 48. Saji, N., Goswami, B., Vinayachandran, P., & Yamagata, T. A dipole mode in the tropical Indian Ocean.
 538 *Nature*, 401, 360–363 (1999).
- 49. Klatt, O., Boebel, O., & Fahrbach, E. A profiling float's sense of ice. *Journal of Atmospheric and Oceanic Technology*, 24, 1301-1308 (2007).
- 541 50. Johnson, K. et al. Observing biogeochemical cycles at global scales with profiling floats and gliders:
 542 prospects for a global array. *Oceanography*, 22, 216-225 (2009).
- 543 51. Brasseur, P. et al. Integrating biogeochemistry and ecology into ocean data assimilation systems.
 544 *Oceanography*, 22, 206-215 (2009).
- 545 52. International Ocean Colour Coordinating Group (IOCCG). Bio-Optical Sensors on Argo Floats. Reports of
- 546 *the International Ocean-Colour Coordinating Group*, **11**, H. Claustre, ed. IOCCG, Dartmouth, Canada (2011).
- 547 53. Frolicher, T. et al. Dominance of the Southern Ocean in anthropogenic carbon and heat uptake in CMIP5
 548 models. *Journal of Climate*, 28 862-886 (2015).
- 549 54. Morrison, A., Frolicher, T. and Sarmiento, J. Upwelling in the Southern Ocean. Physics Today, 68, 27-29550 (2015).
- 55. Fukasawa, M., et al. Bottom water warming in the North Pacific Ocean. *Nature*, **427**, 825-827 (2004).
- 552 56. Johnson, G., Purkey, S., & Bullister, J. Warming and freshening in the abyssal southeastern Indian Ocean.
- 553 *Journal of Climate*, **21**, 5351-5363 (2008).

554	57. Purkey, S. & Johnson, G. Warming of global abyssal and deep Southern Ocean waters between the 1990s
555	and 2000s: contributions to global heat and sea level rise budgets. <i>Journal of Climate</i> , 23 , 6336-6353 (2010).
556	58. Purkey, S., & Johnson, G. Antarctic Bottom Water warming and freshening: contributions to sea level rise,
557	ocean freshwater budgets, and global heat gain. Journal of Climate, 26, 6105-6122 (2013).
558	
559	
560	
561	
562	
563	
564	
565	
566	
567	
568	
569	
570	
571	
572	
573	
574	
575	
576	
577	
578	
579	

List of Figures



581

Figure 1. In the cycle typical of an Argo float, a float starts at the surface and dives to a depth of 1000 meters (the parking depth) where it rests for 9 to 10 days. After 9 days at rest it dives to a depth of 2000 m, turns on its sampling equipment and measures ocean properties as it rises to the surface where it rests for sufficient time to transmit the data collected to Argos or Iridium satellite systems. It then returns to the parking depth to start another cycle, the typical duration of a complete cycle is 10 days.

587

- 588
- 589

590



Figure 2. The sampling density of profiles reported by Argo floats⁴ (upper panel; ~1.5 million collected between 1999 to Oct. 2015), and (lower panel) by the most complete assembly of all previous historical efforts (0.5 million largely shipboard observations collected over the past 100 years from the World Ocean Data Base 2009³). This sampling density is computed as the total number of samples in each 1° latitude × 1° longitude square and is color coded according to the legend on each panel. The analysis only includes profiles that sample both temperature and salinity to a depth

592



Figure 3: Contours of the mean temperature over the period 2004-2010 in the global ocean²⁶ at (*a*) the sea surface, (*b*) a depth of 366 m, and (*c*) a depth of 914 m, using Argo data; the dots denote the Argo minus *Challenger* temperature difference, according to the color scale. Reprinted from (23).



Figure 4. The seasonal to interannual evolution of potential temperature (a) and salinity (b) in the upper 2000 m
of the Labrador Sea during the period 2002-2014, compiled from Argo observations and recomputed from data
forming the original figure published by Kieke and Yashayaev³³.



Figure 5: Estimates of the global heat content anomaly²⁷, for latitude ranges 60° S– 20° S (black), 20° S– 20° N (red), and 20° N– 60° N (blue), computed as 12-month running mean values and shown as an inset. Smoothed trends are shown as the dashed lines on the figure in the same colors; the colored numbers in parentheses refer to the estimate of the trend of the same color in units of 10^{22} Joules yr⁻¹. In the main diagram the mean values are changed so that they overlay the 0-2000m heat content computed by Levitus et al.⁴⁴