

Aplikace matematiky

Jaroslav Haslinger

Finite element analysis for unilateral problems with obstacles on the boundary

Aplikace matematiky, Vol. 22 (1977), No. 3, 180–188

Persistent URL: <http://dml.cz/dmlcz/103691>

Terms of use:

© Institute of Mathematics AS CR, 1977

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

FINITE ELEMENT ANALYSIS FOR UNILATERAL PROBLEMS
WITH OBSTACLES ON THE BOUNDARY

JAROSLAV HASLINGER

(Received June 24, 1976)

INTRODUCTION

In the present paper the finite element analysis for unilateral problems with obstacles on the boundary Γ of a polygonal domain $Q \subset R^2$ is given. Using the technique of [9], [7], the rate of convergence is proved provided the exact solution is smooth enough. We obtain the same results as in [2], [3], where the technique of [5], [6] was used for investigating the primary and the dual problems. In the present paper the case of nonhomogeneous obstacles as well as the case of two obstacles, "lower and upper", is studied. As the regularity hypotheses are not fulfilled in general, we prove — for one class of problems — the convergence of finite element approximations to the exact solution without any regularity assumptions.

1. SETTING OF THE PROBLEM

Let $Q \subset R^m$ be a bounded domain. $H^k(Q)$ ($k \geq 0$ integer) will denote the space of all functions, the derivatives of which up to the order k (in the sense of distributions) are square integrable in Q . The norm of $u \in H^k(Q)$ (defined by the usual manner) will be denoted by $\|u\|_k$. For simplicity we write $H^0(Q) = L^2(Q)$ and the scalar product of $u, v \in L^2(Q)$ will be denoted by (u, v) . A repeated Latin index implies always summation over the range $1, \dots, n$.

Let $Q \subset R^2$ be a bounded domain with a Lipschitz boundary $\Gamma = \Gamma_1 \cup \Gamma_2 \cup \mathcal{R}$, where Γ_1, Γ_2 are open in Γ , $\Gamma_2 \neq \emptyset$, $\Gamma_1 \cap \Gamma_2 = \emptyset$, $\text{mes}_1 \mathcal{R} = 0$ (one-dimensional Lebesgue measure) and $\bar{\Gamma}_1 \cap \bar{\Gamma}_2 = \{A, B\}$ if Γ_1 is non-empty.¹⁾

¹⁾ After slight modifications of proofs of this paper, one can easily extend its results to the case of multiply connected domains.

Let us set

$$V = \{v \in H^1(Q) : v = 0 \text{ on } \Gamma_1\},$$

$$K = \{v \in V : v \geq \psi \text{ on } \Gamma_2\},$$

where ψ is a given function defined on Γ_2 such that $\psi(A) = \psi(B) = 0$.

Let us define

$$J(v) = \int_Q \left(a_{ij} \frac{\partial v}{\partial x_j} \frac{\partial v}{\partial x_i} + a_0 v^2 \right) dx - 2 \int_Q f v dx,$$

where

$$(1.1) \quad f \in L^2(Q),$$

$$(2.1) \quad a_{ij}, a_0 \in L^\infty(Q), \quad i, j = 1, 2,$$

$$(3.1) \quad a_{ij}(x) = a_{ji}(x) \text{ a.e. in } Q, \quad i, j = 1, 2,$$

$$(4.1) \quad a_0 \geq 0 \text{ and either } \Gamma_1 \neq \emptyset \text{ or } \exists c = \text{const.} > 0,$$

$$a_0(x) \geq c \text{ a.e. in } Q,$$

$$(5.1) \quad \exists \alpha = \text{const.} > 0 : \forall \zeta \in R^2$$

$$a_{ij} \zeta_i \zeta_j \geq \alpha \|\zeta\|^2$$

a.e. in Q .

We shall consider the following problem (P):

$$(P) \quad \text{find } u \in K : J(u) = \min_{v \in K} J(v).$$

Theorem 1.1. *If (1.1)–(5.1) hold, then there exists a unique solution of (P), which is characterized by*

$$(6.1) \quad u \in K : a(u, v - u) \geq (f, v - u) \quad \forall v \in K.$$

where

$$a(u, v) = \int_Q \left(a_{ij} \frac{\partial u}{\partial x_j} \frac{\partial v}{\partial x_i} + a_0 uv \right) dx.$$

2. APPROXIMATION OF (P)

Let $Q \subset R^2$ be a polygonal domain, $\bar{\Gamma}_2 = \bigcup_{j=1}^m A_j A_{j+1}$, $A_1 = A$, $A_{m+1} = B$ (if $\Gamma_1 \neq \emptyset$). Let $\{\mathcal{T}_h\}$, $h \in (0, 1)$ be a system of regular triangulations of \bar{Q} , satisfying

the usual requirements concerning the mutual position of triangles T_i and such that A, B are vertices of \mathcal{T}_h for every $h \in (0, 1)$. We define

$$V_h = \{v \in C(\bar{Q}) \cap V : v|_{T_i} \text{ is linear in } T_i, \forall T_i \in \mathcal{T}_h\},$$

$$K_h = \{v \in V_h : v(a_i) \geq \psi(a_i), \text{ where } a_i \text{ are vertices of } \mathcal{T}_h \text{ on } \Gamma_2, i = 1, \dots, n\}.$$

In general, $K_h \not\subset K$.

We define the problem (P_h) (an approximation of (P)) in the following manner:

$$(P_h) \text{ find } u_h \in K_h : J(u_h) = \min_{v \in K_h} J(v).$$

Theorem 1.2. *There exists a unique solution of (P_h) , which is characterized by*

$$(1.2) \quad u_h \in K_h : a(u_h, v_h - u_h) \geq (f, v_h - u_h) \quad \forall v_h \in K_h.$$

For the proof see [1]. Our aim is to estimate the rate of convergence of $\|u - u_h\|_1$.

Lemma 1.2. *It holds*

$$(2.2) \quad \|u - u_h\|_1^2 \leq c\{(f, u - v_h) + (f, u_h - v) + a(u_h - u, v_h - u) + \\ + a(u, v - u_h) + a(u, v_h - u)\} \quad \forall v \in K, v_h \in K_h,$$

where c is an absolute constant.

Proof. Let $v \in K, v_h \in K_h$. Then

$$\alpha \|u - u_h\|_1^2 \leq a(u - u_h, u - u_h) = a(u, u) + a(u_h, u_h) - a(u, u_h) - \\ - a(u_h, u) \leq a(u, v) + (f, u - v) + a(u_h, v_h) + (f, u_h - v_h) - \\ - a(u, u_h) - a(u_h, u) = (f, u - v_h) + (f, u_h - v) + a(u, v - u_h) + \\ + a(u_h - u, v_h - u) + a(u, v_h - u).$$

Consequence. *If $K_h \subset K, h \in (0, 1)$, then substituting $v = u_h$ in (2.2) we obtain for all $v_h \in K_h$:*

$$(3.2) \quad \|u - u_h\|_1^2 \leq c\{(f, u - v_h) + a(u_h - u, v_h - u) + a(u, v_h - u)\}.$$

Theorem 2.2. *Let $\psi = 0$ on Γ_2 , let the solution u satisfy $u \in K \cap H^2(Q)$ and $u|_{\Gamma_2} \in H^2(A_j A_{j+1}) \quad \forall j = 1, \dots, m$. Then*

$$\|u - u_h\|_1 = O(h).$$

Proof. The Green formula yields

$$a(u, v_h - u) = \int_Q \left(- \frac{\partial}{\partial x_i} \left(a_{ij} \frac{\partial u}{\partial x_j} \right) (v_h - u) + a_0 u (v_h - u) \right) dx + \int_{\Gamma_2} \frac{\partial u}{\partial n_A} (v_h - u) ds,$$

where $\partial u / \partial n_A = a_{ij} (\partial u / \partial x_j) n_i$, n_i being the components of the unit outward normal to Γ . It is easy to verify that

$$(4.2) \quad - \frac{\partial}{\partial x_i} \left(a_{ij} \frac{\partial u}{\partial x_j} \right) + a_0 u = f \quad \text{a.e. in } Q.$$

This together with (3.2) implies

$$(5.2) \quad \begin{aligned} \|u - u_h\|_1^2 &\leq c \left\{ a(u_h - u, v_h - u) + \int_{\Gamma_2} \frac{\partial u}{\partial n_A} (v_h - u) ds \right\} \leq \\ &\leq c \{ \|u_h - u\|_1 \|v_h - u\|_1 + \|v_h - u\|_{L_2(\Gamma_2)} \} \leq \\ &\leq c\varepsilon \|u - u_h\|_1^2 + \frac{c}{\varepsilon} \|u - v_h\|_1^2 + c \|v_h - u\|_{L_2(\Gamma_2)}, \end{aligned}$$

where $\varepsilon > 0$ is arbitrary.

Let us set $v_h = R_h u$, where $R_h u \in V_h$ is defined by the relation

$$R_h u = \Pi_{T_i} u \quad \text{on } T_i, \quad \forall T_i \in \mathcal{T}_h.$$

$\Pi_{T_i} u$ denotes the linear Lagrange interpolate of u on T_i . As $v_h \in K_h$, the assertion of the theorem follows from the well-known properties of Lagrange interpolation, (5.2) and the regularity assumptions.

Let us consider the general case $K_h \not\subset K$.

Theorem 3.2. Let $\psi \in H^2(A_j A_{j+1}) \cap H^1(\Gamma_2)$, $u \in K \cap H^2(Q)$, $u|_{\Gamma_2} \in H^2(A_j A_{j+1})$, $j = 1, \dots, m$. Then

$$\|u - u_h\|_1 = O(h).$$

Proof follows from the Green formula, (2.2) and (4.2):

$$(6.2) \quad \begin{aligned} \|u - u_h\|_1^2 &\leq c \left\{ a(u_h - u, v_h - u) + \int_{\Gamma_2} \frac{\partial u}{\partial n_A} (v_h - u) ds + \right. \\ &\quad \left. + \int_{\Gamma_2} \frac{\partial u}{\partial n_A} (v_h - u) ds \right\} \quad \forall v \in K, \quad \forall v_h \in K_h. \end{aligned}$$

The first and the third member on the right hand side of (6.2) can be estimated in the same manner as in Theorem 2.2 (we substitute $v_h = R_h u$). Let us consider the second member. We define

$$\begin{aligned}\bar{v} &= \sup(u_h, \psi) \quad \text{on } \Gamma_2, \\ \bar{v} &= 0 \quad \text{on } \Gamma - \Gamma_2.\end{aligned}$$

Then $\bar{v} \in H^1(\Gamma)$, $\bar{v} \geq \psi$ on Γ_2 and there exists a function $v \in H^1(Q)$ such that $v = \bar{v}$ on Γ . Hence $v \in K$ and

$$u_h - \bar{v} = \begin{cases} 0 & \text{if } u_h \geq \psi \\ u_h - \psi & \text{if } u_h < \psi. \end{cases}$$

As $u_h(a_i) \geq \psi(a_i)$, $i = 1, \dots, n$, it is $u_h \geq r_h \psi$ on Γ_2 , where $r_h \psi|_{a_i a_{i+1}}$ is the linear Lagrange interpolate of ψ on $a_i a_{i+1}$. Now

$$\int_{\Gamma_2} (\bar{v} - u_h)^2 ds = \int_{\Gamma_2^+} (\psi - u_h)^2 ds,$$

where

$$\Gamma_2^+ = \{x \in \Gamma_2 : u_h(x) < \psi(x)\}.$$

Since $0 < (\psi - u_h)(x) \leq (\psi - r_h \psi)(x)$ on Γ_2^+ , we have

$$\int_{\Gamma_2} (\bar{v} - u_h)^2 ds = \int_{\Gamma_2^+} (\psi - u_h)^2 ds \leq \int_{\Gamma_2^+} (\psi - r_h \psi)^2 ds = O(h^4).$$

Remark. Supposing $u \in H^2(Q)$, we can prove that $\|u - u_h\|_1 = O(h^{3/4})$. See also [8].

Now let us consider the case of "two obstacles" on Γ_2 . Let

$$K = \{v \in V : \varphi \leq v \leq \psi \quad \text{on } \Gamma_2\},$$

where φ, ψ are functions defined on Γ_2 , $\varphi(x) \leq \psi(x)$ for each $x \in \Gamma_2$, $\varphi(A) = \varphi(B) = \psi(A) = \psi(B) = 0$ if $\Gamma_1 \neq \emptyset$. We define the problem (\tilde{P}):

$$(\tilde{P}) \quad \text{find } u \in K : J(u) = \min_{v \in K} J(v).$$

Let $K_h = \{v \in C(\bar{Q}) \cap V, v|_{T_i} \text{ is linear on } T_i \in \mathcal{T}_h, \varphi(a_i) \leq v(a_i) \leq \psi(a_i), i = 1, \dots, n\}$.

Let us recall that a_1, \dots, a_n are the vertices of \mathcal{T}_h on Γ_2 .

We define the problem (\tilde{P}_h):

$$(\tilde{P}_h) \quad \text{find } u_h \in K_h : J(u_h) = \min_{v \in K_h} J(v).$$

Theorem 4.2. Let the solution u satisfy $u \in K \cap H^2(Q)$, $u|_{\Gamma_2} \in H^2(A_j A_{j+1})$; let $\varphi, \psi \in H^2(A_j A_{j+1}) \cap H^1(\Gamma_2)$, $j = 1, \dots, m$. Then

$$\|u - u_h\|_1 = O(h).$$

Proof. We use (6.2). It is sufficient to estimate

$$\int_{\Gamma_2} \frac{\partial u}{\partial n_A} (v - u_h) \, ds.$$

Let us set

$$\bar{v} = \begin{cases} \max(\min(u_h, \psi), \varphi) & \text{on } \Gamma_2 \\ 0 & \text{on } \Gamma - \Gamma_2. \end{cases}$$

It is easy to see that $\varphi \leq \bar{v} \leq \psi$ on Γ_2 , $\bar{v} \in H^1(\Gamma)$ so that there exists a function $v \in H^1(Q)$: $v = \bar{v}$ on Γ . Hence $v \in K$ and

$$u_h - \bar{v} = \begin{cases} u_h - \psi & \text{if } u_h > \psi \\ 0 & \text{if } \varphi \leq u_h \leq \psi \\ u_h - \varphi & \text{if } u_h < \varphi. \end{cases}$$

We can write

$$\int_{\Gamma_2} (u_h - \bar{v})^2 \, ds = \int_{\Gamma_2^+} (u_h - \psi)^2 \, ds + \int_{\Gamma_2^-} (u_h - \varphi)^2 \, ds,$$

where $\Gamma_2^+ = \{x \in \Gamma_2 : u_h(x) > \psi(x)\}$, $\Gamma_2^- = \{x \in \Gamma_2 : u_h(x) < \varphi(x)\}$. The inequality $u_h(a_i) \leq \psi(a_i)$ implies $u_h \leq r_h \psi$ on Γ_2 and similarly $u_h \geq r_h \varphi$ on Γ_2 .

Hence

$$0 < u_h(x) - \psi(x) \leq r_h \psi(x) - \psi(x) \quad \text{on } \Gamma_2^+,$$

$$0 < \varphi(x) - u_h(x) \leq \varphi(x) - r_h \varphi(x) \quad \text{on } \Gamma_2^-$$

and

$$\int_{\Gamma_2^+} (u_h - \psi)^2 \, ds \leq \int_{\Gamma_2} (r_h \psi - \psi)^2 \, ds = O(h^4),$$

$$\int_{\Gamma_2^-} (u_h - \varphi)^2 \, ds \leq \int_{\Gamma_2} (\varphi - r_h \varphi)^2 \, ds = O(h^4).$$

APPENDIX

In the above analysis we needed very strong regularity assumptions concerning the solution u in order to be able to prove the rate of convergence. Unfortunately there are no reasons to expect such a great smoothness in general case. In this Appendix we prove the convergence of u_h to u for a particular problem without any regularity assumptions. We shall consider the problem (P) with

$$K = \{v \in H^1(Q) : v \geq \psi \text{ on } \Gamma\},$$

where (A 1) ψ is the trace of a function $\Psi \in H^{1+\varepsilon}(Q)$ ($\varepsilon > 0$). First we prove two auxiliary lemmas.

Lemma A1. Let $\tilde{K} = \{v \in H^1(Q) : v \geq 0 \text{ on } \Gamma\}$. Then \tilde{K} is the closure in $H^1(\bar{Q})$ of the set

$$\mathcal{E}_+(\bar{Q}) = \{v \in \mathcal{E}(\bar{Q}) : v \geq 0 \text{ on } \Gamma\}^1)$$

Proof. Let $u \in \tilde{K}$ be arbitrary. Then $u|_{\Gamma} \in H^{1/2}(\Gamma)$ (for the definition see [4]), $u|_{\Gamma} \geq 0$ and there exists a function $U \in H^1(Q)$ such that $U = u$ on Γ , $U \geq 0$ in Q . For the construction of U see [4], p. 100. We can write

$$u = U + Z,$$

where $Z \in H_0^1(Q)$. The density of $D(Q)$ in $H_0^1(Q)^2$ implies that there exist $Z_h \in D(Q)$ such that $Z_h \rightarrow Z$ in $H^1(Q)$. The regularization U_h of U are also non-negative in Q and if U is suitably extended on a domain G , $\bar{Q} \subset G$ then $U_h \rightarrow U$ in $H^1(Q)$. Setting $u_h = U_h + Z_h \in \mathcal{E}_+(\bar{Q})$ we have $u_h \rightarrow u$ in $H^1(Q)$.

Lemma A2. Let φ be a continuous function defined on $\langle a, b \rangle$ ($-\infty < a < b < \infty$), $D_n : a = x_0^n < x_1^n < \dots < x_n^n = b$ a division of $\langle a, b \rangle$, $v(D_n) = \max_{i=1, \dots, n} |x_i^n - x_{i-1}^n| \rightarrow 0$ for $n \rightarrow \infty$. Let $\{\psi_n\}_{n=1}^{\infty}$ be a sequence of piecewise linear functions with nodes at x_n^i such that $\psi_n(x_i^n) \geq \varphi(x_n^i) \forall i = 0, \dots, n; n = 1, 2, \dots$. Let $\psi_n \rightarrow \psi$ a.e. in $\langle a, b \rangle$. Then $\psi \geq \varphi$ a.e. in $\langle a, b \rangle$.

Proof. Let $M \subseteq \langle a, b \rangle$, $\mu(M) = b - a$ (one-dimensional Lebesgue measure) and such that $\forall x \in M : \psi_n(x) \rightarrow \psi(x)$. We shall show that for all $x \in M$, $\psi(x) \geq \varphi(x)$. Let there exist $x \in M$ such that $\psi(x) < \varphi(x)$. Since φ is continuous there exists $\delta > 0$ such that

$$(A2) \quad \varphi(x) > \psi(\bar{x}) + 1/2(\varphi(\bar{x}) - \psi(\bar{x})) \quad \text{for } \forall x \in (\bar{x} - \delta, \bar{x} + \delta), \quad \delta > 0.$$

On the other hand, there exists n_0 such that

$$(A3) \quad \begin{aligned} \psi_n(\bar{x}) &< \psi(\bar{x}) + 1/2(\varphi(\bar{x}) - \psi(\bar{x})), \\ v(D_n) &< \delta \end{aligned}$$

for $\forall n \geq n_0$. We distinguish two cases:

¹⁾ $\mathcal{E}(\bar{Q}) = \bigcap_{k=1}^{\infty} C^k(\bar{Q})$ ($k \geq 0$ integer), where $C^k(\bar{Q})$ denotes the space of continuous functions, the derivatives of which up to the order k are also continuous and continuously extendible onto \bar{Q} .

²⁾ $H_0^1(Q)$ is the subspace of $H^1(Q)$ of functions, the traces of which are equal to zero on Γ . $D(Q) \subset \mathcal{E}(\bar{Q})$ is the space of functions with compact support in Q .

1. Let there exist $n \geq n_0$ such that $\bar{x} \in (x_{i_0}^n, x_{i_0+1}^n)$ for some $i_0 \in \{0, 1, \dots, n\}$. The restriction of ψ_n on $x_{i_0}^n x_{i_0+1}^n$ is linear and $\psi_n(x_{i_0}^n) \geq \varphi(x_{i_0}^n)$, $\psi_n(\bar{x}) < \varphi(\bar{x})$ and (A2), (A3) yield $\varphi(x_{i_0}^n) > \psi_n(\bar{x})$. Hence $\psi_n(x_{i_0+1}^n) < \psi_n(\bar{x}) < \psi(\bar{x}) + 1/2(\varphi(\bar{x}) - \psi(\bar{x})) < \varphi(x)$ $\forall x \in (\bar{x} - \delta, \bar{x} + \delta)$. In particular, for $x = x_{i_0+1}^n$ we obtain

$$\psi_n(x_{i_0+1}^n) < \varphi(x_{i_0+1}^n),$$

i.e. a contradiction with the assumptions of Lemma A2.

2. Let $\bar{x} = x_{i_0}^n$ for all $n \geq n_0$. Then $\psi_n(\bar{x}) \geq \varphi(\bar{x}) \Rightarrow \psi(\bar{x}) \geq \varphi(\bar{x})$ which is a contradiction.

Now we can prove the main result of this Appendix.

Theorem A1. *Let (A1) be satisfied. Then $\|u - u_h\|_1 \rightarrow 0$ for $h \rightarrow 0+$.*

Proof. according to [1], p. 142, it is sufficient to prove

- (i) $\forall v \in K \exists v_h \in K_h : \|v - v_h\|_1 \rightarrow 0$ for $h \rightarrow 0+$,
- (ii) $v_h \in K_h, v_h \rightarrow v$ (weakly) in $H^1(Q) \Rightarrow v \in K$.

(ad i) Let $v \in K$ be arbitrary. Then $v - \Psi \in H^1(Q)$ and $v - \Psi = v - \psi \geq 0$ on Γ . Lemma A1 implies that there exist $\varphi_H \in \mathcal{E}(\bar{Q})$: $\varphi_H \rightarrow v - \Psi$ in $H^1(Q)$. As $H^{1+\epsilon}(Q) \supset C(\bar{Q})$, we can construct $R_h \Psi$ in the same manner as in Theorem 2.2 and

$$\|\Psi - R_h \Psi\|_1 \rightarrow 0, \quad h \rightarrow 0+.$$

The same we do for φ_H :

$$\|\varphi_H - R_h \varphi_H\|_1 \rightarrow 0, \quad h \rightarrow 0+.$$

Setting $v_h = R_h \Psi + R_h \varphi_H$ we have $v_h \in K_h$ for each $h > 0$ and

$$\|v - v_h\|_1 \leq \|(v - \Psi) - \varphi_H + \varphi_H - R_h \varphi_H\|_1 + \|\Psi - R_h \Psi\|_1 \rightarrow 0$$

for $h, H \rightarrow 0+$.

(ad ii) Let $v_h \rightarrow v$ in $H^1(Q)$, $v_h \in K_h$. Then $v_h \rightarrow v$ in $L^2(\Gamma)^1$ and we can extract a subsequence $\{v_{h_n}\}$ such that $v_{h_n} \rightarrow v$ a.e. in Γ . Using Lemma A2 we prove $v \geq \psi$ a.e. in Γ , i.e. $v \in K$.

References

- [1] Céa J.: Optimisation, théorie et algorithmes. Dunod, Paris 1971.
- [2] Hlaváček, I.: Dual finite element analysis for unilateral boundary value problems. To appear in Apl. Mat.
- [3] Hlaváček I.: Dual finite element analysis for elliptic problems with obstacles on the boundary, I. To appear in Apl. Mat.
- [4] Nečas, J.: Les méthodes directes en théorie des équations elliptiques. Academie, Prague 1967.

¹⁾ By virtue of the complete continuity of the mapping $\gamma : H^1(Q) \rightarrow L^2(\Gamma)$, see [4] p. 107.

- [5] *Mosco U., Strang G.*: One sided approximations and variational inequalities. Bull. Am. Math. Soc. 80 (1974), 308—312.
- [6] *Strang, G.*: One-sided approximations and plate bending. Computing methods in applied sciences and engineering-Part I. Versailles 1973.
- [7] *Raoult-Puech*: Approximation des inequations variationnelles, Seminaire Ciarlet-Glowinski-Raviart 1974.
- [8] *Scarpini F. - Vivaldi M.*: Error estimates for the approximations of some unilateral problems. To appear in R.A.I.R.O.
- [9] *Falk R. S.*: Error estimates for approximation of a class of a variational inequalities. Math. of Comp. 28 (1974), 963—971.

Souhrn

ANALÝZA JEDNOSTRANNÝCH ÚLOH S PŘEKÁŽKAMI NA HRANICI METODOU KONEČNÝCH PRVKŮ

JAROSLAV HASLINGER

Práce se zabývá aplikací metody konečných prvků pro řešení (i) jednostranných úloh s obecně nehomogenní překážkou na hranici Γ (ii) dvoustranných úloh, kdy je na Γ zadána „horní a dolní“ překážka. Jsou-li řešení uvedených problémů dostatečně hladká, potom je dokázáno, že konvergence přibližných řešení k přesnému je řádu $O(h)$. Přitom používáme po částech lineárních konečných prvků. Protože oblast, na níž problémy řešíme je polygonální, není obecně zaručena tak vysoká hladkost řešení. Proto v závěru práce je proveden důkaz konvergence (bez odhadu její rychlosti) pro problémy typu (i), aniž bychom předpokládali dodatečnou hladkost.

Author's address: Dr. Jaroslav Haslinger, katedra matematické fyziky MFF UK, Malostranské nám. 25, 118 00 Praha 1.