



Finite generation of symmetric ideals

Matthias Aschenbrenner and Christopher J. Hillar

In memoriam Karin Gatermann (1965–2005).

ABSTRACT. Let A be a commutative Noetherian ring, and let $R = A[X]$ be the polynomial ring in an infinite collection X of indeterminates over A . Let \mathfrak{S}_X be the symmetric group of X . The group \mathfrak{S}_X acts on R in a natural way, and this in turn gives R the structure of a left module over the group ring $R[\mathfrak{S}_X]$. We prove that all ideals of R invariant under the action of \mathfrak{S}_X are finitely generated as $R[\mathfrak{S}_X]$ -modules. The proof involves introducing a certain partial order on monomials and showing that it is a well-quasi-ordering. We also consider the concept of an invariant chain of ideals for finite-dimensional polynomial rings and relate it to the finite generation result mentioned above. Finally, a motivating question from chemistry is presented, with the above framework providing a suitable context in which to study it.

RÉSUMÉ. Soit A un anneau Noetherien commutatif, et $R = A[X]$ l'anneau des polynômes en une infinité d'indéterminées X sur A . Soit \mathfrak{S}_X le groupe symétrique de X . Le groupe \mathfrak{S}_X agit sur R de manière naturelle, ce qui donne à R la structure d'un module gauche sur l'anneau $R[\mathfrak{S}_X]$. Nous prouvons que tous les idéaux de R invariants sous l'action de \mathfrak{S}_X sont finitement engendrés comme $R[\mathfrak{S}_X]$ -modules. La démonstration utilise le fait qu'un certain ordre partiel sur les monômes est un quasi-ordre. Nous utilisons aussi le concept de chaîne invariante des idéaux pour les anneaux de polynômes de dimension finie, que nous relierons au résultat de génération finie mentionné plus haut. Finalement, nous présentons une motivation pour notre travail issue de la chimie.

1. Introduction

A pervasive theme in invariant theory is that of finite generation. A fundamental example is a theorem of Hilbert stating that the invariant subrings of finite-dimensional polynomial algebras over finite groups are finitely generated [5, Corollary 1.5]. In this article, we study invariant ideals of infinite-dimensional polynomial rings. Of course, when the number of indeterminates is finite, Hilbert's basis theorem tells us that any ideal (invariant or not) is finitely generated.

Our setup is as follows. Let X be an infinite collection of indeterminates, and let \mathfrak{S}_X be the group of permutations of X . Fix a commutative Noetherian ring A and let $R = A[X]$ be the polynomial ring in the indeterminates X . The group \mathfrak{S}_X acts naturally on R : if $\sigma \in \mathfrak{S}_X$ and $f \in A[x_1, \dots, x_n]$ where $x_i \in X$, then

$$\sigma f(x_1, x_2, \dots, x_n) = f(\sigma x_1, \sigma x_2, \dots, \sigma x_n) \in R.$$

This in turn gives R the structure of a left module over the (non-commutative) group ring $R[\mathfrak{S}_X]$. An ideal $I \subseteq R$ is called *invariant under \mathfrak{S}_X* (or simply *invariant*) if

$$\mathfrak{S}_X I := \{\sigma f : \sigma \in \mathfrak{S}_X, f \in I\} \subseteq I.$$

Notice that invariant ideals are simply the $R[\mathfrak{S}_X]$ -submodules of R . We may now state our main result.

1991 *Mathematics Subject Classification.* 13E05, 13E15, 20B30, 06A07.

Key words and phrases. Invariant ideal, well-quasi-ordering, symmetric group.

The first author is partially supported by the National Science Foundation Grant DMS 03-03618. The work of the second author is supported under a National Science Foundation Graduate Research Fellowship.

THEOREM 1.1. *Every ideal of $R = A[X]$ invariant under \mathfrak{S}_X is finitely generated as an $R[\mathfrak{S}_X]$ -module. (Stated more succinctly, R is a Noetherian $R[\mathfrak{S}_X]$ -module.)*

For the purposes of this work, we will use the following notation. Let B be a ring and let G be a subset of a B -module M . Then $\langle f : f \in G \rangle_B$ will denote the B -submodule of M generated by elements of G .

EXAMPLE 1.2. Suppose that $X = \{x_1, x_2, \dots\}$. The invariant ideal $I = \langle x_1, x_2, \dots \rangle_R$ is clearly not finitely generated over R , however, it does have the compact representation $I = \langle x_1 \rangle_{R[\mathfrak{S}_X]}$.

The outline of this paper is as follows. In Section 2, we define a partial order on monomials and show that it can be used to obtain a well-quasi-ordering of the monomials in R . Section 3 then goes on to detail our proof of Theorem 1.1, using the main result of Section 2 in a fundamental way. In the penultimate section, we discuss a relationship between invariant ideals of R and chains of increasing ideals in finite-dimensional polynomial rings. The notions introduced there provide a suitable framework for studying a problem arising from chemistry, the subject of the final section of this article.

2. The Symmetric Cancellation Ordering

We begin this section by briefly recalling some basic order-theoretic notions. We also discuss some fundamental results due to Higman and Nash-Williams and some of their consequences. We define the ordering mentioned in the section heading, and give a sufficient condition for it to be a well-quasi-ordering; this is needed in the proof of Theorem 1.1.

2.1. Preliminaries. A *quasi-ordering* on a set S is a binary relation \leq on S which is reflexive and transitive. A *quasi-ordered set* is a pair (S, \leq) consisting of a set S and a quasi-ordering \leq on S . When there is no confusion, we will omit \leq from the notation, and simply call S a quasi-ordered set. If in addition the relation \leq is *anti-symmetric* ($s \leq t \wedge t \leq s \Rightarrow s = t$, for all $s, t \in S$), then \leq is called an *ordering* (sometimes also called a *partial ordering*) on the set S . The *trivial* ordering on S is given by $s \leq t \iff s = t$ for all $s, t \in S$. A quasi-ordering \leq on a set S induces an ordering on the set $S/\sim = \{s/\sim : s \in S\}$ of equivalence classes of the equivalence relation $s \sim t \iff s \leq t \wedge t \leq s$ on S . If s and t are elements of a quasi-ordered set, we write as usual $s \leq t$ also as $t \geq s$, and we write $s < t$ if $s \leq t$ and $t \not\leq s$.

A map $\varphi: S \rightarrow T$ between quasi-ordered sets S and T is called *increasing* if $s \leq t \Rightarrow \varphi(s) \leq \varphi(t)$ for all $s, t \in S$, and *strictly increasing* if $s < t \Rightarrow \varphi(s) < \varphi(t)$ for all $s, t \in S$. We also say that $\varphi: S \rightarrow T$ is a *quasi-embedding* if $\varphi(s) \leq \varphi(t) \Rightarrow s \leq t$ for all $s, t \in S$.

An *antichain* of S is a subset $A \subseteq S$ such that $s \not\leq t$ and $t \not\leq s$ for all $s \not\sim t$ in A . A *final segment* of a quasi-ordered set (S, \leq) is a subset $F \subseteq S$ which is closed upwards: $s \leq t \wedge s \in F \Rightarrow t \in F$, for all $s, t \in S$. We can view the set $\mathcal{F}(S)$ of final segments of S as an ordered set, with the ordering given by reverse inclusion. Given a subset M of S , the set $\{t \in S : \exists s \in M \text{ with } s \leq t\}$ is a final segment of S , the final segment *generated by M* . An *initial segment* of S is a subset of S whose complement is a final segment. An initial segment I of S is *proper* if $I \neq S$. For $a \in S$ we denote by $S^{\leq a}$ the initial segment consisting of all $s \in S$ with $s \leq a$.

A quasi-ordered set S is said to be *well-founded* if there is no infinite strictly decreasing sequence $s_1 > s_2 > \dots$ in S , and *well-quasi-ordered* if in addition every antichain of S is finite. The following characterization of well-quasi-orderings is classical (see, for example, [8]). An infinite sequence s_1, s_2, \dots in S is called *good* if $s_i \leq s_j$ for some indices $i < j$, and *bad* otherwise.

PROPOSITION 2.1. *The following are equivalent, for a quasi-ordered set S :*

- (1) S is well-quasi-ordered.
- (2) Every infinite sequence in S is good.
- (3) Every infinite sequence in S contains an infinite increasing subsequence.
- (4) Any final segment of S is finitely generated.
- (5) $(\mathcal{F}(S), \supseteq)$ is well-founded (i.e., the ascending chain condition holds for final segments of S). \square

Let (S, \leq_S) and (T, \leq_T) be quasi-ordered sets. If there exists an increasing surjection $S \rightarrow T$ and S is well-quasi-ordered, then T is well-quasi-ordered, and if there exists a quasi-embedding $S \rightarrow T$ and T is well-quasi-ordered, then so is S . Moreover, the cartesian product $S \times T$ can be turned into a quasi-ordered set by using the cartesian product of \leq_S and \leq_T :

$$(s, t) \leq (s', t') \iff s \leq_S s' \wedge t \leq_T t', \quad \text{for } s, s' \in S, t, t' \in T.$$

Using Proposition 2.1 we see that the cartesian product of two well-quasi-ordered sets is again well-quasi-ordered.

Of course, a total ordering \leq is well-quasi-ordered if and only if it is well-founded; in this case \leq is called a *well-ordering*. Every well-ordered set is isomorphic to a unique ordinal number, called its *order type*. The order type of $\mathbb{N} = \{0, 1, 2, \dots\}$ with its usual ordering is ω .

2.2. A lemma of Higman. Given a set X , we let X^* denote the set of all finite sequences of elements of X (including the empty sequence). We may think of the elements of X^* as *non-commutative words* $x_1 \cdots x_m$ with letters x_1, \dots, x_m coming from the alphabet X . With the concatenation of such words as operation, X^* is the free monoid generated by X . A quasi-ordering \leq on X yields a quasi-ordering \leq_H (the *Higman quasi-ordering*) on X^* as follows:

$$x_1 \cdots x_m \leq_H y_1 \cdots y_n \quad :\iff \quad \left\{ \begin{array}{l} \text{there exists a strictly increasing function} \\ \varphi: \{1, \dots, m\} \rightarrow \{1, \dots, n\} \text{ such that} \\ x_i \leq y_{\varphi(i)} \text{ for all } 1 \leq i \leq m. \end{array} \right.$$

If \leq is an ordering on X , then \leq_H is an ordering on X^* . The following fact was shown by Higman [6] (with an ingenious proof due to Nash-Williams [12]):

LEMMA 2.2. *If \leq is a well-quasi-ordering on X , then \leq_H is a well-quasi-ordering on X^* .* □

It follows that if \leq is a well-quasi-ordering on X , then the quasi-ordering \leq^* on X^* defined by

$$x_1 \cdots x_m \leq^* y_1 \cdots y_n \quad :\iff \quad \left\{ \begin{array}{l} \text{there exists an injective function} \\ \varphi: \{1, \dots, m\} \rightarrow \{1, \dots, n\} \text{ such} \\ \text{that } x_i \leq y_{\varphi(i)} \text{ for all } 1 \leq i \leq m \end{array} \right.$$

is also a well-quasi-ordering. (Since \leq^* extends \leq_H .)

We also let X^\diamond be the set of *commutative words* in the alphabet X , that is, the free commutative monoid generated by X (with identity element denoted by 1). We sometimes also refer to the elements of X^\diamond as *monomials* (in the set of indeterminates X). We have a natural surjective monoid homomorphism $\pi: X^* \rightarrow X^\diamond$ given by simply “making the indeterminates commute” (i.e., interpreting a non-commutative word from X^* as a commutative word in X^\diamond). Unlike \leq_H , the quasi-ordering \leq^* is compatible with π in the sense that $v \leq^* w \Rightarrow v' \leq^* w'$ for all $v, v', w, w' \in X^*$ with $\pi(v) = \pi(v')$ and $\pi(w) = \pi(w')$. Hence $\pi(v) \leq^\diamond \pi(w) :\iff v \leq^* w$ defines a quasi-ordering \leq^\diamond on $X^\diamond = \pi(X^*)$ making π an increasing map. The quasi-ordering \leq^\diamond extends the divisibility relation in the monoid X^\diamond :

$$v|w \quad :\iff \quad uv = w \text{ for some } u \in X^\diamond.$$

If we take for \leq the trivial ordering on X , then \leq^\diamond corresponds exactly to divisibility in X^\diamond , and this ordering is a well-quasi-ordering if and only if X is finite. In general we have, as an immediate consequence of Higman’s lemma (since π is a surjection):

COROLLARY 2.3. *If \leq is a well-quasi-ordering on the set X , then \leq^\diamond is a well-quasi-ordering on X^\diamond .* □

2.3. A theorem of Nash-Williams. Given a totally ordered set S and a quasi-ordered set X , we denote by $\text{Fin}(S, X)$ the set of all functions $f: I \rightarrow X$, where I is a proper initial segment of S , whose range $f(I)$ is *finite*. We define a quasi-ordering \leq_H on $\text{Fin}(S, X)$ as follows: for $f: I \rightarrow X$ and $g: J \rightarrow X$ from $\text{Fin}(S, X)$ put

$$f \leq_H g \quad :\iff \quad \left\{ \begin{array}{l} \text{there exists a strictly increasing function } \varphi: I \rightarrow J \\ \text{such that } f(i) \leq g(\varphi(i)) \text{ for all } i \in I. \end{array} \right.$$

We may think of an element of $\text{Fin}(S, X)$ as a sequence of elements of X indexed by indices in some proper initial segment of S . So for $S = \mathbb{N}$ with its usual ordering, we can identify elements of $\text{Fin}(\mathbb{N}, X)$ with words in X^* , and then \leq_H for $\text{Fin}(\mathbb{N}, X)$ agrees with \leq_H on X^* as defined above. We will have occasion to use a far-reaching generalization of Lemma 2.2:

THEOREM 2.4. *If X is well-quasi-ordered and S is well-ordered, then $\text{Fin}(S, X)$ is well-quasi-ordered.* □

This theorem was proved by Nash-Williams [13]; special cases were shown earlier in [4, 11, 14].

2.4. Term orderings. A *term ordering* of X^\diamond is a well-ordering \leq of X^\diamond such that

- (1) $1 \leq x$ for all $x \in X$, and
- (2) $v \leq w \Rightarrow xv \leq xw$ for all $v, w \in X^\diamond$ and $x \in X$.

Every ordering \leq of X^\diamond satisfying (1) and (2) extends the ordering \leq^\diamond obtained from the restriction of \leq to X . In particular, \leq extends the divisibility ordering on X^\diamond . By the corollary above, a total ordering \leq of X^\diamond which satisfies (1) and (2) is a term ordering if and only if its restriction to X is a well-ordering.

EXAMPLE 2.5. Let \leq be a total ordering of X . We define the induced *lexicographic ordering* \leq_{lex} of monomials as follows: given $v, w \in X^\diamond$ we can write $v = x_1^{a_1} \cdots x_n^{a_n}$ and $w = x_1^{b_1} \cdots x_n^{b_n}$ with $x_1 < \cdots < x_n$ in X and all $a_i, b_i \in \mathbb{N}$; then

$$v \leq_{\text{lex}} w \quad :\Leftrightarrow \quad (a_n, \dots, a_1) \leq (b_n, \dots, b_1) \text{ lexicographically (from the left).}$$

The ordering \leq_{lex} is total and satisfies (1), (2); hence if the ordering \leq of X is a well-ordering, then \leq_{lex} is a term ordering of X^\diamond .

REMARK 2.6. Let \leq be a total ordering of X . For $w \in X^\diamond$, $w \neq 1$, we let

$$|w| := \max \{x \in X : x|w\} \quad (\text{with respect to } \leq).$$

We also put $|1| := -\infty$ where we set $-\infty < x$ for all $x \in X$. One of the perks of using the lexicographic ordering as a term ordering on X^\diamond is that if v and w are monomials with $v \leq_{\text{lex}} w$, then $|v| \leq |w|$. Below, we often use this observation.

The previous example shows that for every set X there exists a term ordering of X^\diamond , since every set can be well-ordered by the Axiom of Choice. In fact, every set X can be equipped with a well-ordering every proper initial segment of which has strictly smaller cardinality than X ; in other words, the order type of this ordering (a certain ordinal number) is a cardinal number. We shall call such an ordering of X a *cardinal well-ordering* of X .

LEMMA 2.7. *Let X be a set equipped with a cardinal well-ordering, and let I be a proper initial segment of X . Then every injective function $I \rightarrow X$ can be extended to a permutation of X .*

PROOF. Since this is clear if X is finite, suppose that X is infinite. Let $\varphi: I \rightarrow X$ be injective. Since I has cardinality $|I| < |X|$ and X is infinite, we have $|X| = \max\{|X \setminus I|, |I|\} = |X \setminus I|$. Similarly, since $|\varphi(I)| = |I| < |X|$, we also have $|X \setminus \varphi(I)| = |X|$. Hence there exists a bijection $\psi: X \setminus I \rightarrow X \setminus \varphi(I)$. Combining φ and ψ yields a permutation of X as desired. \square

2.5. A new ordering of monomials. Let G be a permutation group on a set X , that is, a group G together with a faithful action $(\sigma, x) \mapsto \sigma x: G \times X \rightarrow X$ of G on X . The action of G on X extends in a natural way to a faithful action of G on X^\diamond : $\sigma w = \sigma x_1 \cdots \sigma x_n$ for $\sigma \in G$, $w = x_1 \cdots x_n \in X^\diamond$. Given a term ordering \leq of X^\diamond , we define a new relation on X^\diamond as follows:

DEFINITION 2.8. (The symmetric cancellation ordering corresponding to G and \leq .)

$$v \preceq w \quad :\Leftrightarrow \quad \begin{cases} v \leq w \text{ and there exist } \sigma \in G \text{ and a monomial} \\ u \in X^\diamond \text{ such that } w = u\sigma v \text{ and for all } v' \leq v, \\ \text{we have } u\sigma v' \leq w. \end{cases}$$

REMARK 2.9. Every term ordering \leq is *linear*: $v \leq w \Leftrightarrow uv \leq uw$ for all monomials u, v, w . Hence the condition above may be rewritten as: $v \preceq w$ and there exists $\sigma \in G$ such that $\sigma v|w$ and $\sigma v' \leq \sigma v$ for all $v' \leq v$. (We say that “ σ witnesses $v \preceq w$.”)

EXAMPLE 2.10. Let $X = \{x_1, x_2, \dots\}$ be a countably infinite set of indeterminates, ordered such that $x_1 < x_2 < \dots$, and let $\leq = \leq_{\text{lex}}$ be the corresponding lexicographic ordering of X^\diamond . Let also G be the group of permutations of $\{1, 2, 3, \dots\}$, acting on X via $\sigma x_i = x_{\sigma(i)}$. As an example of the relation \preceq , consider the following chain:

$$x_1^2 \preceq x_1 x_2^2 \preceq x_1^3 x_2 x_3^2.$$

To verify the first inequality, notice that $x_1 x_2^2 = x_1 \sigma(x_1^2)$, in which σ is the transposition (12). If $v' = x_1^{a_1} \cdots x_n^{a_n} \leq x_1^2$ with $a_1, \dots, a_n \in \mathbb{N}$, $a_n > 0$, then it follows that $n = 1$ and $a_1 \leq 2$. In particular, $x_1 \sigma v' = x_1 x_2^{a_1} \leq x_1 x_2^2$. For the second relationship, we have that $x_1^3 x_2 x_3^2 = x_1^3 \tau(x_1 x_2^2)$, in which τ is the cycle (123). Additionally, if $v' = x_1^{a_1} \cdots x_n^{a_n} \leq x_1 x_2^2$ with $a_1, \dots, a_n \in \mathbb{N}$, $a_n > 0$, then $n \leq 2$, and if $n = 2$, then either $a_2 = 1$ or $a_2 = 2$, $a_1 \leq 1$. In each case we get $x_1^3 \tau v' = x_1^3 x_2^{a_1} x_3^{a_2} \leq x_1^3 x_2 x_3^2$.

Although Definition 2.8 appears technical, we will soon present a nice interpretation of it that involves leading term cancellation of polynomials. First we verify that it is indeed an ordering.

LEMMA 2.11. *The relation \preceq is an ordering on monomials.*

PROOF. First notice that $w \preceq w$ since we may take $u = 1$ and $\sigma =$ the identity permutation. Next, suppose that $u \preceq v \preceq w$. Then there exist permutations σ, τ in G and monomials u_1, u_2 in X^\diamond such that $v = u_1\sigma u$, $w = u_2\tau v$. In particular, $w = u_2(\tau u_1)(\tau\sigma u)$. Additionally, if $v' \preceq u$, then $u_1\sigma v' \preceq v$, so that $u_2\tau(u_1\sigma v') \preceq w$. It follows that $u_2(\tau u_1)(\tau\sigma v') \preceq w$. This shows transitivity; anti-symmetry of \preceq follows from anti-symmetry of \leq . \square

We offer a useful interpretation of this ordering (which motivates its name). We fix a commutative ring A and let $R = A[X]$ be the ring of polynomials with coefficients from A in the collection of commuting indeterminates X . Its elements may be written uniquely in the form

$$f = \sum_{w \in X^\diamond} a_w w$$

where $a_w \in A$ for all $w \in X^\diamond$, and all but finitely many a_w are zero. We say that a monomial w occurs in f if $a_w \neq 0$. Given a non-zero $f \in R$ we define $\text{lm}(f)$, the *leading monomial* of f (with respect to our choice of term ordering \leq) to be the largest monomial w (with respect to \leq) which occurs in f . If $w = \text{lm}(f)$, then a_w is the *leading coefficient* of f , denoted by $\text{lc}(f)$, and $a_w w$ is the *leading term* of f , denoted by $\text{lt}(f)$. By convention, we set $\text{lm}(0) = \text{lc}(0) = \text{lt}(0) = 0$. We let $R[G]$ be the group ring of G over R (with multiplication given by $f\sigma \cdot g\tau = fg(\sigma\tau)$ for $f, g \in R, \sigma, \tau \in G$), and we view R as a left $R[G]$ -module in the natural way.

LEMMA 2.12. *Let $f \in R, f \neq 0$, and $u, w \in X^\diamond$. Suppose that $\sigma \in G$ witnesses $\text{lm}(f) \preceq w$, and let $u \in X^\diamond$ with $u\sigma \text{lm}(f) = w$. Then $\text{lm}(u\sigma f) = u\sigma \text{lm}(f)$.*

PROOF. Put $v = \text{lm}(f)$. Every monomial occurring in $u\sigma f$ has the form $u\sigma v'$, where v' occurs in f . Hence $v' \leq v$, and since σ witnesses $v \preceq w$, this yields $u\sigma v' \leq w$. \square

Suppose that A is a field, let $v \preceq w$ be in X^\diamond and let f, g be two polynomials in R with leading monomials v, w , respectively. Then, from the definition and the lemma above, there exists a $\sigma \in G$ and a term cu ($c \in A \setminus \{0\}, u \in X^\diamond$) such that all monomials occurring in

$$h = g - cu\sigma f$$

are strictly smaller (with respect to \leq) than w . For readers familiar with the theory of Gröbner bases, the polynomial h can be viewed as a kind of symmetric version of the S -polynomial (see, for instance, [5, Chapter 15]).

EXAMPLE 2.13. In the situation of Example 2.10 above, let $f = x_1x_2^2 + x_2 + x_1^2$ and $g = x_1^3x_2x_3^2 + x_3^2 + x_1^4x_3$. Set $\sigma = (123)$, and observe that

$$g - x_1^3\sigma f = x_1^4x_3 + x_3^2 - x_1^3x_3 - x_1^3x_2^2$$

has a smaller leading monomial than g .

We are mostly interested in the case where our term ordering on X^\diamond is \leq_{lex} , and $G = \mathfrak{S}_X$. Under these assumptions we have:

LEMMA 2.14. *Let $v, w \in X^\diamond$ with $v \preceq w$. Then for every $\sigma \in \mathfrak{S}_X$ witnessing $v \preceq w$ we have $\sigma(X^{\leq|v|}) \subseteq X^{\leq|w|}$. Moreover, if the order type of (X, \leq) is $\leq \omega$, then we can choose such σ with the additional property that $\sigma(x) = x$ for all $x > |w|$.*

PROOF. To see the first claim, suppose for a contradiction that $\sigma x > |w|$ for some $x \in X, x \leq |v|$. We have $\sigma v|w$, so if $x|v$, then $\sigma x|w$, contradicting $\sigma x > |w|$. In particular $x < |v|$, which yields $x <_{\text{lex}} v$ and thus $\sigma x <_{\text{lex}} \sigma v <_{\text{lex}} w$, again contradicting $\sigma x > |w|$. Now suppose that the order type of X is $\leq \omega$, and let σ witness $v \preceq w$. Then $|v| \leq |w|$, and $\sigma X^{\leq|v|}$ can be extended to a permutation σ' of the finite set $X^{\leq|w|}$. We further extend σ' to a permutation of X by setting $\sigma'(x) = x$ for all $x > |w|$. One checks easily that σ' still witnesses $v \preceq w$. \square

2.6. Lovely orderings. We say that a term ordering \leq of X^\diamond is *lovely* for G if the corresponding symmetric cancellation ordering \preceq on X^\diamond is a well-quasi-ordering. If \leq is lovely for a subgroup of G , then \leq is lovely for G .

EXAMPLE 2.15. The symmetric cancellation ordering corresponding to $G = \{1\}$ and a given term ordering \leq of X^\diamond is just

$$v \preceq w \iff v \leq w \wedge v|w.$$

Hence a term ordering of X^\diamond is lovely for $G = \{1\}$ if and only if divisibility in X^\diamond has no infinite antichains; that is, exactly if X is finite.

This terminology is inspired by the following definition from [3] (which in turn goes back to an idea in [2]):

DEFINITION 2.16. Given an ordering \leq of X , consider the following ordering of X :

$$x \sqsubseteq y \iff \begin{cases} x \leq y \text{ and there exists } \sigma \in G \text{ such that } \sigma x = y \\ \text{and for all } x' \leq x, \text{ we have } \sigma x' \leq y. \end{cases}$$

A well-ordering \leq of X is called *nice* (for G) if \sqsubseteq is a well-quasi-ordering.

In [2] one finds various examples of nice orderings, and in [3] it is shown that if X admits a nice ordering with respect to G , then for every field F , the free F -module FX with basis X is Noetherian as a module over $F[G]$. It is clear that the restriction to X of a lovely ordering of X^\diamond is nice. However, there do exist permutation groups (G, X) for which X admits a nice ordering, but X^\diamond does not admit a lovely ordering; see Example 3.4 and Proposition 5.2 below.

EXAMPLE 2.17. Suppose that X is countable. Then every well-ordering of X of order type ω is nice for \mathfrak{S}_X . To see this, we may assume that $X = \mathbb{N}$ with its usual ordering. It is then easy to see that if $x \leq y$ in \mathbb{N} , then $x \sqsubseteq y$, witnessed by any extension σ of the strictly increasing map $n \mapsto n + y - x: \mathbb{N}^{\leq x} \rightarrow \mathbb{N}$ to a permutation of \mathbb{N} .

The following crucial fact (generalizing the last example) is needed for our proof of Theorem 1.1:

THEOREM 2.18. *The lexicographic ordering of X^\diamond corresponding to a cardinal well-ordering of a set X is lovely for the full symmetric group \mathfrak{S}_X of X .*

For the proof, let as above $\text{Fin}(X, \mathbb{N})$ be the set of all sequences in \mathbb{N} indexed by elements in some proper initial segment of X which have finite range, quasi-ordered by \leq_H . For a monomial $w \neq 1$ we define $w^*: X^{\leq |w|} \rightarrow \mathbb{N}$ by

$$w^*(x) := \max \{a \in \mathbb{N} : x^a | w\}.$$

Then clearly $w^* \in \text{Fin}(X, \mathbb{N})$, in fact, $w^*(x) = 0$ for all but finitely many $x \in X^{\leq |w|}$. We also let $1^* :=$ the empty sequence $\emptyset \rightarrow \mathbb{N}$ (the unique smallest element of $\text{Fin}(X, \mathbb{N})$). We now quasi-order $X^\diamond \times \text{Fin}(X, \mathbb{N})$ by the cartesian product of the ordering \leq_{lex} on X^\diamond and the quasi-ordering \leq_H on $\text{Fin}(X, \mathbb{N})$. By Corollary 2.3, Theorem 2.4, and the remark following Proposition 2.1, $X^\diamond \times \text{Fin}(X, \mathbb{N})$ is well-quasi-ordered. Therefore, in order to finish the proof of Theorem 2.18, it suffices to show:

LEMMA 2.19. *The map*

$$w \mapsto (w, w^*): X^\diamond \rightarrow X^\diamond \times \text{Fin}(X, \mathbb{N})$$

is a quasi-embedding with respect to the symmetric cancellation ordering on X^\diamond and the quasi-ordering on $X^\diamond \times \text{Fin}(X, \mathbb{N})$.

PROOF. Suppose that v, w are monomials with $v \leq_{\text{lex}} w$ and $v^* \leq_H w^*$; we need to show that $v \preceq w$. For this we may assume that $v, w \neq 1$. So there exists a strictly increasing function $\varphi: X^{\leq |v|} \rightarrow X^{\leq |w|}$ such that

$$(2.1) \quad v^*(x) \leq w^*(\varphi(x)) \quad \text{for all } x \in X \text{ with } x \leq |v|.$$

By Lemma 2.7 there exists $\sigma \in \mathfrak{S}_X$ such that $\sigma X^{\leq |v|} = \varphi X^{\leq |v|}$. Then clearly $\sigma v | w$ by (2.1). Now let $v' \leq_{\text{lex}} v$; we claim that $\sigma v' \leq_{\text{lex}} \sigma v$. Again we may assume $v' \neq 1$. Then $|v'| \leq |v|$, hence we may write

$$v' = x_1^{a_1} \cdots x_n^{a_n}, \quad v = x_1^{b_1} \cdots x_n^{b_n}$$

with $x_1 < \dots < x_n \leq |v|$ in X and $a_i, b_j \in \mathbb{N}$. Put $y_1 := \varphi(x_1), \dots, y_n := \varphi(x_n)$. Then $y_1 < \dots < y_n$ and

$$\sigma v' = y_1^{a_1} \dots y_n^{a_n}, \quad \sigma v = y_1^{b_1} \dots y_n^{b_n},$$

and therefore $\sigma v' \leq_{\text{lex}} \sigma v$ as required. \square

2.7. The case of countable X . In Section 4 we will apply Theorem 2.18 in the case where X is countable. Then the order type of X is at most ω , and in the proof of the theorem given above we only need to appeal to a special instance (Higman's Lemma) of Theorem 2.4. We finish this section by giving a self-contained proof of this important special case of Theorem 2.18, avoiding Theorem 2.4. Let $\mathfrak{S}_{(X)}$ denote the subgroup of \mathfrak{S}_X consisting of all $\sigma \in \mathfrak{S}_X$ with the property that $\sigma(x) = x$ for all but finitely many letters $x \in X$.

THEOREM 2.20. *The lexicographic ordering of X^\diamond corresponding to a cardinal well-ordering of a countable set X is lovely for $\mathfrak{S}_{(X)}$.*

Let X be countable and let \leq be a cardinal well-ordering of X . Enumerate the elements of X as $x_1 < x_2 < \dots$. We assume that X is infinite; this is not a restriction, since by Lemma 2.14 we have:

LEMMA 2.21. *If the lexicographic ordering of X^\diamond is lovely for $\mathfrak{S}_{(X)}$, then for any n and $X_n := \{x_1, \dots, x_n\}$, the lexicographic ordering of $(X_n)^\diamond$ is lovely for \mathfrak{S}_{X_n} .* \square

We begin with some preliminary lemmas. Here, \preceq is the symmetric cancellation ordering corresponding to $\mathfrak{S}_{(X)}$ and \leq_{lex} . We identify $\mathfrak{S}_{(X)}$ and $\mathfrak{S}_\infty := \mathfrak{S}_{(\mathbb{N})}$ in the natural way, and for every n we regard \mathfrak{S}_n , the group of permutations of $\{1, 2, \dots, n\}$, as a subgroup of \mathfrak{S}_∞ ; then $\mathfrak{S}_n \leq \mathfrak{S}_{n+1}$ for each n , and $\mathfrak{S}_\infty = \bigcup_n \mathfrak{S}_n$.

LEMMA 2.22. *Suppose that $x_1^{a_1} \dots x_n^{a_n} \preceq x_1^{b_1} \dots x_n^{b_n}$ where $a_i, b_j \in \mathbb{N}$, $b_n > 0$. Then for any $c \in \mathbb{N}$ we have $x_1^{a_1} \dots x_n^{a_n} \preceq x_1^c x_2^{b_1} \dots x_{n+1}^{b_n}$.*

PROOF. Let $v := x_1^{a_1} \dots x_n^{a_n}$, $w := x_1^{b_1} \dots x_n^{b_n}$. We may assume $v \neq 1$. Clearly $v \leq_{\text{lex}} w$ and $b_n > 0$ yield $x_1^{a_1} \dots x_n^{a_n} \leq_{\text{lex}} x_1^c x_2^{b_1} \dots x_{n+1}^{b_n}$. Let now $\sigma \in \mathfrak{S}_\infty$ witness $v \preceq w$. Let τ be the cyclic permutation $\tau = (1\ 2\ 3 \dots (n+1))$ and set $\hat{\sigma} := \tau\sigma$. Then $\sigma v|w$ yields $\hat{\sigma}v|\tau w$, hence $\hat{\sigma}v|x_1^c \tau w = x_1^c x_2^{b_1} \dots x_{n+1}^{b_n}$. Next, suppose that $v' \leq_{\text{lex}} v$; then $\sigma v' \leq_{\text{lex}} \sigma v$. By Lemma 2.14 and the nature of τ , the map $\tau\sigma(\{1, \dots, |v|\})$ is strictly increasing, which gives $\hat{\sigma}v' = \tau\sigma v' \leq_{\text{lex}} \tau\sigma v = \hat{\sigma}v$. Hence $\hat{\sigma}$ witnesses $x_1^{a_1} \dots x_n^{a_n} \preceq x_1^c x_2^{b_1} \dots x_{n+1}^{b_n}$. \square

LEMMA 2.23. *If $x_1^{a_1} \dots x_n^{a_n} \preceq x_1^{b_1} \dots x_n^{b_n}$, where $a_i, b_j \in \mathbb{N}$, $b_n > 0$, and $a, b \in \mathbb{N}$ are such that $a \leq b$, then $x_1^a x_2^{a_1} \dots x_{n+1}^{a_n} \preceq x_1^b x_2^{b_1} \dots x_{n+1}^{b_n}$.*

PROOF. As before let $v := x_1^{a_1} \dots x_n^{a_n}$, $w := x_1^{b_1} \dots x_n^{b_n}$. Once again, we may assume $v \neq 1$, and it is clear that $x_1^a x_2^{a_1} \dots x_{n+1}^{a_n} \leq_{\text{lex}} x_1^b x_2^{b_1} \dots x_{n+1}^{b_n}$. Let $\sigma \in \mathfrak{S}_\infty$ witness $v \preceq w$. By Lemma 2.14 we may assume that $\sigma(x_i) = x_i$ for all $i > n$. Let τ be the cyclic permutation $\tau = (1\ 2 \dots (n+1))$. Setting $\hat{\sigma} = \tau\sigma\tau^{-1}$, we have $\hat{\sigma}x_1 = x_1$, hence

$$(2.2) \quad \hat{\sigma}(x_1^a x_2^{a_1} \dots x_{n+1}^{a_n}) = \hat{\sigma}(x_1^a) \hat{\sigma}(x_2^{a_1} \dots x_{n+1}^{a_n}) = x_1^a \tau\sigma v.$$

Since $\sigma v|w$, this last expression divides $x_1^b \tau w = x_1^b x_2^{b_1} \dots x_{n+1}^{b_n}$. Suppose that $v' = x_1^{c_1} \dots x_{n+1}^{c_{n+1}} \leq_{\text{lex}} x_1^a x_2^{a_1} \dots x_{n+1}^{a_n}$, where $c_i \in \mathbb{N}$. Then, since we are using a lexicographic order, we have

$$x_2^{c_2} \dots x_{n+1}^{c_{n+1}} \leq_{\text{lex}} x_2^{a_1} \dots x_{n+1}^{a_n}$$

and therefore

$$\tau^{-1}(x_2^{c_2} \dots x_{n+1}^{c_{n+1}}) = x_1^{c_2} \dots x_n^{c_{n+1}} \leq_{\text{lex}} \tau^{-1}(x_2^{a_1} \dots x_{n+1}^{a_n}) = v.$$

By assumption, this implies that $\sigma\tau^{-1}(x_2^{c_2} \dots x_{n+1}^{c_{n+1}}) \leq_{\text{lex}} \sigma v$ and thus by (2.2)

$$\hat{\sigma}(x_2^{c_2} \dots x_{n+1}^{c_{n+1}}) \leq_{\text{lex}} \tau\sigma v = \hat{\sigma}(x_2^{a_1} \dots x_{n+1}^{a_n}).$$

If this inequality is strict, then since $1 \notin \hat{\sigma}(\{2, \dots, n+1\})$, clearly

$$\hat{\sigma}v' = x_1^{c_1} \hat{\sigma}(x_2^{c_2} \dots x_{n+1}^{c_{n+1}}) <_{\text{lex}} x_1^a \tau\sigma v = \hat{\sigma}(x_1^a x_2^{a_1} \dots x_{n+1}^{a_n}).$$

Otherwise $x_2^{c_2} \dots x_{n+1}^{c_{n+1}} = x_2^{a_1} \dots x_{n+1}^{a_n}$, hence $c_1 \leq a$, in which case we still have $\hat{\sigma}v' \leq_{\text{lex}} \hat{\sigma}(x_1^a x_2^{a_1} \dots x_{n+1}^{a_n})$. Therefore $\hat{\sigma}$ witnesses $x_1^a x_2^{a_1} \dots x_{n+1}^{a_n} \preceq x_1^b x_2^{b_1} \dots x_{n+1}^{b_n}$. This completes the proof. \square

We now have enough to show Theorem 2.20. The proof uses the basic idea from Nash-Williams' proof [13] of Higman's lemma. Assume for the sake of contradiction that there exists a bad sequence

$$w^{(1)}, w^{(2)}, \dots, w^{(n)}, \dots \quad \text{in } X^\diamond.$$

For $w \in X^\diamond \setminus \{1\}$ let $j(w)$ be the index $j \geq 1$ with $|w| = x_j$, and put $j(1) := 0$. We may assume that the bad sequence is chosen in such a way that for every n , $j(w^{(n)})$ is *minimal* among the $j(w)$, where w ranges over all elements of X^\diamond with the property that $w^{(1)}, w^{(2)}, \dots, w^{(n-1)}, w$ can be continued to a bad sequence in X^\diamond . Because $1 \leq_{\text{lex}} w$ for all $w \in X^\diamond$, we have $j(w^{(n)}) > 0$ for all n . For every $n > 0$, write $w^{(n)} = x_1^{a^{(n)}} v^{(n)}$ with $a^{(n)} \in \mathbb{N}$ and $v^{(n)} \in X^\diamond$ not divisible by x_1 . Since \mathbb{N} is well-ordered, there is an infinite sequence $1 \leq i_1 < i_2 < \dots$ of indices such that $a^{(i_1)} \leq a^{(i_2)} \leq \dots$. Consider the monoid homomorphism $\alpha: X^\diamond \rightarrow X^\diamond$ given by $\alpha(x_{i+1}) = x_i$ for all $i > 1$. Then $j(\alpha(w)) = j(w) - 1$ if $w \neq 1$. Hence by minimality of $w^{(1)}, w^{(2)}, \dots$, the sequence

$$w^{(1)}, w^{(2)}, \dots, w^{(i_1-1)}, \alpha(v^{(i_1)}), \alpha(v^{(i_2)}), \dots, \alpha(v^{(i_n)}), \dots$$

is good; that is, there exist $j < i_1$ and k with $w^{(j)} \preceq \alpha(v^{(i_k)})$, or there exist $k < l$ with $\alpha(v^{(i_k)}) \preceq \alpha(v^{(i_l)})$. In the first case we have $w^{(j)} \preceq w^{(i_k)}$ by Lemma 2.22; and in the second case, $w^{(i_k)} \preceq w^{(i_l)}$ by Lemma 2.23. This contradicts the badness of our sequence $w^{(1)}, w^{(2)}, \dots$, finishing the proof.

QUESTION. Careful inspection of the proof of Theorem 2.18 (in particular Lemma 2.7) shows that in the statement of the theorem, we can replace \mathfrak{S}_X by its subgroup consisting of all σ with the property that the set of $x \in X$ with $\sigma(x) \neq x$ has cardinality $< |X|$. In Theorem 2.18, can one always replace \mathfrak{S}_X by $\mathfrak{S}_{(X)}$?

3. Proof of the Finiteness Theorem

We now come to the proof our main result. Throughout this section we let A be a commutative Noetherian ring, X an arbitrary set, $R = A[X]$, and we let G be a permutation group on X . An $R[G]$ -submodule of R will be called a *G-invariant ideal* of R , or simply an *invariant ideal*, if G is understood. We will show:

THEOREM 3.1. *If X^\diamond admits a lovely term ordering for G , then R is Noetherian as an $R[G]$ -module.*

For $G = \{1\}$ and X finite, this theorem reduces to Hilbert's basis theorem, by Example 2.15. We also obtain Theorem 1.1:

COROLLARY 3.2. *The $R[\mathfrak{S}_X]$ -module R is Noetherian.*

PROOF. Choose a cardinal well-ordering of X . Then the corresponding lexicographic ordering of X^\diamond is lovely for \mathfrak{S}_X , by Theorem 2.18. Apply Theorem 3.1. \square

REMARK 3.3. It is possible to replace the use of Theorem 2.18 in the proof of the corollary above by the more elementary Theorem 2.20. This is because if the $R[\mathfrak{S}_X]$ -module R was not Noetherian, then one could find a *countably generated* $R[\mathfrak{S}_X]$ -submodule of R which is not finitely generated, and hence a countable subset X' of X such that $R' = A[X']$ is not a Noetherian $R'[\mathfrak{S}_{X'}]$ -module.

The following example shows how the conclusion of Theorem 3.1 may fail:

EXAMPLE 3.4. Suppose that G has a cyclic subgroup H which acts freely and transitively on X . Then X has a nice ordering (see [2]), but $R = \mathbb{Q}[X^\diamond]$ is not Noetherian. To see this let σ be a generator for H , and let $x \in X$ be arbitrary. Then the $R[G]$ -submodule of $R = \mathbb{Q}[X^\diamond]$ generated by the elements $\sigma^n x \sigma^{-n} x$ ($n \in \mathbb{N}$) is not finitely generated. So by Theorem 3.1, X^\diamond does not admit a lovely term ordering for G .

For the proof of Theorem 3.1 we develop a bit of Gröbner basis theory for the $R[G]$ -module R . For the time being, we fix an arbitrary term ordering \leq (not necessarily lovely for G) of X^\diamond .

3.1. Reduction of polynomials. Let $f \in R$, $f \neq 0$, and let B be a set of non-zero polynomials in R . We say that f is *reducible by B* if there exist pairwise distinct $g_1, \dots, g_m \in B$, $m \geq 1$, such that for each i we have $\text{lm}(g_i) \preceq \text{lm}(f)$, witnessed by some $\sigma_i \in G$, and

$$\text{lt}(f) = a_1 w_1 \sigma_1 \text{lt}(g_1) + \dots + a_m w_m \sigma_m \text{lt}(g_m)$$

for non-zero $a_i \in A$ and monomials $w_i \in X^\diamond$ such that $w_i \sigma_i \text{lm}(g_i) = \text{lm}(f)$. In this case we write $f \xrightarrow{B} h$, where

$$h = f - (a_1 w_1 \sigma_1 g_1 + \cdots + a_m w_m \sigma_m g_m),$$

and we say that f reduces to h by B . We say that f is reduced with respect to B if f is not reducible by B . By convention, the zero polynomial is reduced with respect to B . Trivially, every element of B reduces to 0.

EXAMPLE 3.5. Suppose that A is a field. Then f is reducible by B if and only if there exists some $g \in B$ such that $\text{lm}(g) \preceq \text{lm}(f)$.

EXAMPLE 3.6. Suppose that f is reducible by B as defined (for finite X) in, say, [1, Chapter 4], that is: there exist $g_1, \dots, g_m \in B$ and $a_1, \dots, a_m \in A$ ($m \geq 1$) such that $\text{lm}(g_i) \mid \text{lm}(f)$ for all i and

$$\text{lc}(f) = a_1 \text{lc}(g_1) + \cdots + a_m \text{lc}(g_m).$$

Then f is reducible by B in the sense defined above. (Taking $\sigma_i = 1$ for all i .)

REMARK 3.7. Suppose that $G = \mathfrak{S}_X$, the term ordering \leq of X^\diamond is \leq_{lex} , and the order type of (X, \leq) is $\leq \omega$. Then in the definition of reducibility by B above, we may require that the σ_i satisfy $\sigma_i(x) = x$ for all $1 \leq i \leq m$ and $x > |\text{lm}(f)|$. (By Lemma 2.14.)

The smallest quasi-ordering on R extending the relation \xrightarrow{B} is denoted by \xrightarrow{B}^* . If $f, h \neq 0$ and $f \xrightarrow{B} h$, then $\text{lm}(h) < \text{lm}(f)$, by Lemma 2.12. In particular, every chain

$$h_0 \xrightarrow{B} h_1 \xrightarrow{B} h_2 \xrightarrow{B} \cdots$$

with all $h_i \in R \setminus \{0\}$ is finite. (Since the term ordering \leq is well-founded.) Hence there exists $r \in R$ such that $f \xrightarrow{B}^* r$ and r is reduced with respect to B ; we call such an r a *normal form* of f with respect to B .

LEMMA 3.8. Suppose that $f \xrightarrow{B}^* r$. Then there exist $g_1, \dots, g_n \in B$, $\sigma_1, \dots, \sigma_n \in G$ and $h_1, \dots, h_n \in R$ such that

$$f = r + \sum_{i=1}^n h_i \sigma_i g_i \quad \text{and} \quad \text{lm}(f) \geq \max_{1 \leq i \leq n} \text{lm}(h_i \sigma_i g_i).$$

(In particular, $f - r \in \langle B \rangle_{R[G]}$.)

PROOF. This is clear if $f = r$. Otherwise we have $f \xrightarrow{B} h \xrightarrow{B}^* r$ for some $h \in R$. Inductively we may assume that there exist $g_1, \dots, g_n \in B$, $\sigma_1, \dots, \sigma_n \in G$ and $h_1, \dots, h_n \in R$ such that

$$h = r + \sum_{i=1}^n h_i \sigma_i g_i \quad \text{and} \quad \text{lm}(h) \geq \max_{1 \leq i \leq n} \text{lm}(h_i \sigma_i g_i).$$

There are also $g_{n+1}, \dots, g_{n+m} \in B$, $\sigma_{n+1}, \dots, \sigma_{n+m} \in G$, $a_{n+1}, \dots, a_{n+m} \in A$ and $w_{n+1}, \dots, w_{n+m} \in X^\diamond$ such that $\text{lm}(w_{n+i} \sigma_{n+i} g_{n+i}) = \text{lm}(f)$ for all i and

$$\text{lt}(f) = \sum_{i=1}^m a_{n+i} w_{n+i} \sigma_{n+i} \text{lt}(g_{n+i}), \quad f = h + \sum_{i=1}^m a_{n+i} w_{n+i} \sigma_{n+i} g_{n+i}.$$

Hence putting $h_{n+i} := a_{n+i} w_{n+i}$ for $i = 1, \dots, m$ we have $f = r + \sum_{j=1}^{n+m} h_j \sigma_j g_j$ and $\text{lm}(f) > \text{lm}(h) \geq \text{lm}(h_j \sigma_j g_j)$ if $1 \leq j \leq n$, $\text{lm}(f) = \text{lm}(h_j \sigma_j g_j)$ if $n < j \leq n+m$. \square

REMARK 3.9. Suppose that $G = \mathfrak{S}_X$, $\leq = \leq_{\text{lex}}$, and X has order type $\leq \omega$. Then in the previous lemma we can choose the σ_i such that in addition $\sigma_i(x) = x$ for all i and all $x > |\text{lm}(f)|$. (By Remark 3.7.)

3.2. Gröbner bases. Let B be a subset of R . We let

$$\text{lt}(B) := \langle \text{lc}(g)w : 0 \neq g \in B, \text{lm}(g) \preceq w \rangle_A$$

be the A -submodule of R generated by all elements of the form $\text{lc}(g)w$, where $g \in B$ is non-zero and w is a monomial with $\text{lm}(g) \preceq w$. Clearly for non-zero $f \in R$ we have: $\text{lt}(f) \in \text{lt}(B)$ if and only if f is reducible by B . In particular, $\text{lt}(B)$ contains $\{\text{lt}(g) : g \in B\}$, and for an ideal I of R which is G -invariant, we simply have

$$\text{lt}(I) = \{\text{lt}(f) : f \in I\}.$$

(Use Lemma 2.12.) We say that a subset B of an invariant ideal I of R is a *Gröbner basis* for I (with respect to our choice of term ordering \preceq) if $\text{lt}(I) = \text{lt}(B)$.

LEMMA 3.10. *Let I be an invariant ideal of R and B be a set of non-zero elements of I . The following are equivalent:*

- (1) B is a Gröbner basis for I .
- (2) Every non-zero $f \in I$ is reducible by B .
- (3) Every $f \in I$ has normal form 0. (In particular, $I = \langle B \rangle_{R[G]}$.)
- (4) Every $f \in I$ has unique normal form 0.

PROOF. The implications (1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (4) are either obvious or follow from the remarks preceding the lemma. Suppose that (4) holds. Every $f \in I \setminus \{0\}$ with $\text{lt}(f) \notin \text{lt}(B)$ is reduced with respect to B , hence has two distinct normal forms (0 and f), a contradiction. Thus $\text{lt}(I) = \text{lt}(B)$. \square

Suppose that B is a Gröbner basis for an ideal I of the polynomial ring $R = A[X^\diamond]$, in the usual sense of the word (as defined, for finite X , in [1, Chapter 4]); if I is invariant, then B is a Gröbner basis for I as defined above (by Example 3.6). Moreover, for $G = \{1\}$, the previous lemma reduces to a familiar characterization of Gröbner bases in the usual case of polynomial rings. It is probably possible to also introduce a notion of S -polynomial and to prove a Buchberger-style criterion for Gröbner bases in our setting, leading to a completion procedure for the construction of Gröbner bases. At this point, we will not pursue these issues further, and rather show:

PROPOSITION 3.11. *Suppose that the term ordering \preceq of X^\diamond is lovely for G . Then every invariant ideal of R has a finite Gröbner basis.*

For a subset B of R let $\text{lm}(B)$ denote the final segment of X^\diamond with respect to \preceq generated by the $\text{lm}(g)$, $g \in B$. If A is a field, then a subset B of an invariant ideal I of R is a Gröbner basis for I if and only if $\text{lm}(B) = \text{lm}(I)$. Hence in this case, the proposition follows immediately from the equivalence of (1) and (4) in Proposition 2.1. For the general case we use the following observation:

LEMMA 3.12. *Let S be a well-quasi-ordered set and T be a well-founded ordered set, and let $\varphi: S \rightarrow T$ be decreasing: $s \leq t \Rightarrow \varphi(s) \geq \varphi(t)$, for all $s, t \in S$. Then the quasi-ordering \leq_φ on S defined by*

$$s \leq_\varphi t \quad :\Leftrightarrow \quad s \leq t \wedge \varphi(s) = \varphi(t)$$

is a well-quasi-ordering. \square

PROOF OF PROPOSITION 3.11. Suppose now that our term ordering of X^\diamond is lovely for G , and let I be an invariant ideal of R . For $w \in X^\diamond$ consider

$$\text{lc}(I, w) := \{\text{lc}(f) : f \in I, \text{ and } f = 0 \text{ or } \text{lm}(f) = w\},$$

an ideal of A . Note that if $v \preceq w$, then $\text{lc}(I, v) \subseteq \text{lc}(I, w)$. We apply the lemma to $S = X^\diamond$, quasi-ordered by \preceq , $T =$ the collection of all ideals of A , ordered by reverse inclusion, and φ given by $w \mapsto \text{lc}(I, w)$. Thus by (4) in Proposition 2.1, applied to the final segment X^\diamond of the well-quasi-ordering \leq_φ , we obtain finitely many $w_1, \dots, w_m \in X^\diamond$ with the following property: for every $w \in X^\diamond$ there exists some $i \in \{1, \dots, m\}$ such that $w_i \preceq w$ and $\text{lc}(I, w_i) = \text{lc}(I, w)$. Using Noetherianity of A , for every i we now choose finitely many non-zero elements g_{i1}, \dots, g_{in_i} of I ($n_i \in \mathbb{N}$), each with leading monomial w_i , whose leading coefficients generate the ideal $\text{lc}(I, w_i)$ of A . We claim that

$$B := \{g_{ij} : 1 \leq i \leq m, 1 \leq j \leq n_i\}$$

is a Gröbner basis for I . To see this, let $0 \neq f \in I$, and put $w := \text{lm}(f)$. Then there is some i with $w_i \preceq w$ and $\text{lc}(I, w_i) = \text{lc}(I, w)$. This shows that f is reducible by $\{g_{i1}, \dots, g_{i,n_i}\}$, and hence by B . By Lemma 3.10, B is a Gröbner basis for I . \square

From Proposition 3.11 and the implication (1) \Rightarrow (3) in Lemma 3.10 we obtain Theorem 3.1.

3.3. A partial converse of Theorem 3.1. Consider now the quasi-ordering $|_G$ of X^\diamond defined by

$$v|_G w \quad :\iff \quad \exists \sigma \in G : \sigma v|w,$$

which extends every symmetric cancellation ordering corresponding to a term ordering of X^\diamond . If M is a set of monomials from X^\diamond and F the final segment of $(X^\diamond, |_G)$ generated by M , then the invariant ideal $\langle M \rangle_{R[G]}$ of R is finitely generated as an $R[G]$ -module if and only if F is generated by a finite subset of M . Hence by the implication (4) \Rightarrow (1) in Proposition 2.1 we get:

LEMMA 3.13. *If R is Noetherian as an $R[G]$ -module, then $|_G$ is a well-quasi-ordering.* \square

This will be used in Section 5 below.

3.4. Connection to a concept due to Michler. Let \leq be a term ordering of X^\diamond . For each $\sigma \in G$ we define a term ordering \leq_σ on X^\diamond by

$$v \leq_\sigma w \quad \iff \quad \sigma v \leq \sigma w.$$

We denote the leading monomial of $f \in R$ with respect to \leq_σ by $\text{lm}_\sigma(f)$. Clearly we have

$$(3.1) \quad \sigma \text{lm}(f) = \text{lm}_{\sigma^{-1}}(\sigma f) \quad \text{for all } \sigma \in G \text{ and } f \in R.$$

Let I be an invariant ideal of R . Generalizing terminology introduced in [10], let us call a set B of non-zero elements of I a *universal G -Gröbner basis* for I (with respect to \leq) if B contains, for every $\sigma \in G$, a Gröbner basis (in the usual sense of the word) for the ideal I with respect to the term ordering \leq_σ . If the set X of indeterminates is finite, then every invariant ideal of R has a finite universal G -Gröbner basis. By the remark following Lemma 3.10, every universal G -Gröbner basis for an invariant ideal I of R is a Gröbner basis for I . We finish this section by observing:

LEMMA 3.14. *Suppose that A is field. If B is a Gröbner basis for the invariant ideal I of R , then*

$$GB = \{\sigma g : \sigma \in G, g \in B\}$$

is a universal G -Gröbner basis for I .

PROOF. Let $\sigma \in G$ and $f \in I$, $f \neq 0$. Then $\sigma f \in I$, hence there exists $\tau \in G$ and $g \in B$ such that $w \leq \text{lm}(g) \Rightarrow w \leq_\tau \text{lm}(g)$ for all $w \in X^\diamond$, and $\tau \text{lm}(g) | \text{lm}(\sigma f)$. The first condition implies in particular that $\tau \text{lm}(g) = \text{lm}(\tau g)$, hence $\sigma^{-1} \tau \text{lm}(g) = \text{lm}_\sigma(\sigma^{-1} \tau g)$ and $\sigma^{-1} \text{lm}(\sigma f) = \text{lm}_\sigma(f)$ by (3.1). Put $h := \sigma^{-1} \tau g \in GB$. Then $\text{lm}_\sigma(h) | \text{lm}_\sigma(f)$ by the second condition. This shows that GB contains a Gröbner basis for I with respect to \leq_σ , as required. \square

EXAMPLE 3.15. Suppose that $G = \mathfrak{S}_n$, the group of permutations of $\{1, 2, \dots, n\}$, acting on $X = \{x_1, \dots, x_n\}$ via $\sigma x_i = x_{\sigma(i)}$. The invariant ideal $I = \langle x_1, \dots, x_n \rangle_R$ has Gröbner basis $\{x_1\}$ with respect to the lexicographic ordering; a corresponding (minimal) universal \mathfrak{S}_n -Gröbner basis for I is $\{x_1, \dots, x_n\}$.

4. Invariant Chains of Ideals

In this section we describe a relationship between certain chains of increasing ideals in finite-dimensional polynomial rings and invariant ideals of infinite-dimensional polynomial rings. We begin with an abstract setting that is suitable for placing the motivating problem (described in the next section) in a proper context. Throughout this section, m and n range over the set of positive integers. For each n , let R_n be a commutative ring, and assume that R_n is a subring of R_{n+1} , for each n . Suppose that the symmetric group on n letters \mathfrak{S}_n gives an action (not necessarily faithful) on R_n such that $f \mapsto \sigma f : R_n \rightarrow R_n$ is a ring homomorphism, for each $\sigma \in \mathfrak{S}_n$. Furthermore, suppose that the natural embedding of \mathfrak{S}_n into \mathfrak{S}_m for $n \leq m$ is compatible with the embedding of rings $R_n \subseteq R_m$; that is, if $\sigma \in \mathfrak{S}_n$ and $\hat{\sigma}$ is the corresponding element in \mathfrak{S}_m , then $\hat{\sigma} R_n = \sigma$. Note that there exists a unique action of \mathfrak{S}_∞ on the ring $R := \bigcup_{n \geq 1} R_n$ which extends the action of each \mathfrak{S}_n on R_n . An ideal of R is *invariant* if $\sigma f \in I$ for all $\sigma \in \mathfrak{S}_\infty$, $f \in I$.

We will need a method for lifting ideals of smaller rings into larger ones, and one such technique is as follows.

DEFINITION 4.1. For $m \geq n$, the m -symmetrization $L_m(B)$ of a set B of elements of R_n is the \mathfrak{S}_m -invariant ideal of R_m given by

$$L_m(B) = \langle g : g \in B \rangle_{R_m[\mathfrak{S}_m]}$$

In order for us to apply this definition sensibly, we must make sure that the m -symmetrization of an ideal can be defined in terms of generators.

LEMMA 4.2. *If B is a set of generators for the ideal $I_B = \langle B \rangle_{R_n}$ of R_n , then $L_m(I_B) = L_m(B)$.*

PROOF. Suppose that B generates the ideal $I_B \subseteq R_n$. Clearly, $L_m(B) \subseteq L_m(I_B)$. Therefore, it is enough to show the inclusion $L_m(I_B) \subseteq L_m(B)$. Suppose that $h \in L_m(I_B)$ so that $h = \sum_{j=1}^s f_j \cdot \sigma_j h_j$ for elements $f_j \in R_m$, $h_j \in I_B$ and $\sigma_j \in \mathfrak{S}_m$. Next express each $h_j = \sum_{i=1}^{r_j} p_{ij} g_{ij}$ for $p_{ij} \in R_n$ and $g_{ij} \in B$. Substitution into the expression above for h gives us

$$h = \sum_{j=1}^s \sum_{i=1}^{r_j} f_j \cdot \sigma_j p_{ij} \cdot \sigma_j g_{ij}.$$

This is easily seen to be an element of $L_m(B)$, completing the proof. \square

EXAMPLE 4.3. Let $S = \mathbb{Q}[t_1, t_2]$, $R_n = \mathbb{Q}[x_1, \dots, x_n]$, and consider the natural action of \mathfrak{S}_n on R_n . Let Q be the kernel of the homomorphism induced by the map $\phi: R_3 \rightarrow S$ given by $\phi(x_1) = t_1^2$, $\phi(x_2) = t_2^2$, and $\phi(x_3) = t_1 t_2$. Then, $Q = \langle x_1 x_2 - x_3^2 \rangle$, and $L_4(Q) \subseteq R_4$ is generated by the following 12 polynomials:

$$\begin{aligned} & x_1 x_2 - x_3^2, \quad x_1 x_2 - x_4^2, \quad x_1 x_3 - x_2^2, \quad x_1 x_3 - x_4^2, \\ & x_1 x_4 - x_3^2, \quad x_1 x_4 - x_2^2, \quad x_2 x_3 - x_1^2, \quad x_2 x_3 - x_4^2, \\ & x_2 x_4 - x_1^2, \quad x_2 x_4 - x_3^2, \quad x_3 x_4 - x_1^2, \quad x_3 x_4 - x_2^2. \end{aligned}$$

We would also like a way to project a set of elements in R_m down to a smaller ring R_n ($n \leq m$).

DEFINITION 4.4. Let $B \subseteq R_m$ and $n \leq m$. The n -projection $P_n(B)$ of B is the \mathfrak{S}_n -invariant ideal of R_n given by

$$P_n(B) = \langle g : g \in B \rangle_{R_m[\mathfrak{S}_m]} \cap R_n.$$

We now consider increasing chains I_\circ of ideals $I_n \subseteq R_n$:

$$I_1 \subseteq I_2 \subseteq \dots \subseteq I_n \subseteq \dots,$$

simply called *chains* below. Of course, such chains will usually fail to stabilize since they are ideals in larger and larger rings. However, it is possible for these ideals to stabilize “up to the action of the symmetric group,” a concept we make clear below. For the purposes of this work, we will only consider a special class of chains; namely, a *symmetrization invariant chain* (resp. *projection invariant chain*) is one for which $L_m(I_n) \subseteq I_m$ (resp. $P_n(I_m) \subseteq I_n$) for all $n \leq m$. If I_\circ is both a symmetrization and a projection invariant chain, then it will be simply called an *invariant chain*. We will encounter some concrete invariant chains in the next section. The stabilization definition alluded to above is as follows.

DEFINITION 4.5. A symmetrization invariant chain of ideals I_\circ as above *stabilizes modulo the symmetric group* (or simply *stabilizes*) if there exists a positive integer N such that

$$L_m(I_n) = I_m \quad \text{for all } m \geq n > N.$$

To put it another way, accounting for the natural action of the symmetric group, the ideals I_n are the same for large enough n . Let us remark that if for a symmetrization invariant chain I_\circ , there is some integer N such that $L_m(I_N) = I_m$ for all $m > N$, then I_\circ stabilizes. This follows from the inclusions

$$I_m = L_m(I_N) \subseteq L_m(I_n) \subseteq I_m, \quad n > N.$$

Any chain I_\circ naturally gives rise to an ideal $\mathcal{I}(I_\circ)$ of $R = \bigcup_{n \geq 1} R_n$ by way of

$$\mathcal{I}(I_\circ) := \bigcup_{n \geq 1} I_n.$$

Conversely, if I is an ideal of R , then

$$I_n = \mathcal{J}_n(I) := I \cap R_n$$

defines the components of a chain $\mathcal{J}(I) := I_\circ$. Clearly, for any ideal $I \subseteq R$, we have $\mathcal{I} \circ \mathcal{J}(I) = I$, but, as is easily seen, it is not true in general that $\mathcal{J} \circ \mathcal{I}(I_\circ) = I_\circ$. However, for invariant chains, this relationship does hold, as the following straightforward lemma describes.

LEMMA 4.6. *There is a one-to-one, inclusion-preserving correspondence between invariant chains I_\circ and invariant ideals I of R given by the maps \mathcal{I} and \mathcal{J} .* \square

For the remainder of this section we consider the case where, for a commutative Noetherian ring A , we have $R_n = A[x_1, \dots, x_n]$ for each n , endowed with the natural action of \mathfrak{S}_n on the indeterminates x_1, \dots, x_n . Then $R = A[X^\circ]$ where $X = \{x_1, x_2, \dots\}$. We use the results of the previous section to demonstrate the following.

THEOREM 4.7. *Every symmetrization invariant chain stabilizes modulo the symmetric group.*

PROOF. Given a symmetrization invariant chain, construct the invariant ideal $I = \mathcal{I}(I_\circ)$ of R . One would now like to apply Theorem 1.1, however, more care is needed to prove stabilization. Let \leq be a well-ordering of X of order type ω , and let B be a finite Gröbner basis for I with respect to the corresponding term ordering \leq_{lex} of X° . (Theorem 2.20 and Proposition 3.11.) Choose a positive integer N such that $B \subseteq I_N$; we claim that $I_m = L_m(I_N)$ for all $m \geq N$. Let $f \in I_m$, $f \neq 0$. By the equivalence of (1) and (3) in Lemma 3.10 we have $f \xrightarrow[B]{*} 0$. Hence by Lemma 3.8 there are $g_1, \dots, g_n \in B$, $h_1, \dots, h_n \in R$, as well as $\sigma_1, \dots, \sigma_n \in \mathfrak{S}_\infty$, such that

$$f = h_1 \sigma_1 g_1 + \dots + h_n \sigma_n g_n \quad \text{and} \quad \text{lm}(f) = \max_i \text{lm}(h_i \sigma_i g_i).$$

By Remark 3.9 we may assume that in fact $\sigma_i \in \mathfrak{S}_m$ for each i . Moreover $\text{lm}(h_i) \leq_{\text{lex}} \text{lm}(f)$, hence $|\text{lm}(h_i)| \leq |\text{lm}(f)| \leq m$, for each i . Therefore $h_i \in R_m$ for each i . This shows that $f \in L_m(B) \subseteq L_m(I_N)$ as desired. \square

5. A Chemistry Motivation

We can now discuss the details of the basic problem that is of interest to us. It was brought to our attention by Bernd Sturmfels, who, in turn, learned about it from Andreas Dress.

Fix a natural number $k \geq 1$. Given a set S we denote by $\langle S \rangle^k$ the set of all ordered k -element subsets of S , that is, $\langle S \rangle^k$ is the set of all k -tuples $\mathbf{u} = (u_1, \dots, u_k) \in S^k$ with pairwise distinct u_1, \dots, u_k . We also just write $\langle n \rangle^k$ instead of $\langle \{1, \dots, n\} \rangle^k$. Let K be a field, and for $n \geq k$ consider the polynomial ring

$$R_n = K[\{x_{\mathbf{u}}\}_{\mathbf{u} \in \langle n \rangle^k}].$$

We let \mathfrak{S}_n act on $\langle n \rangle^k$ by

$$\sigma(u_1, \dots, u_k) = (\sigma(u_1), \dots, \sigma(u_k)).$$

This induces an action $(\sigma, x_{\mathbf{u}}) \mapsto \sigma x_{\mathbf{u}} = x_{\sigma \mathbf{u}}$ of \mathfrak{S}_n on the indeterminates $x_{\mathbf{u}}$, which we extend to an action of \mathfrak{S}_n on R_n in the natural way. We also put $R = \bigcup_{n \geq k} R_n$. Note that

$$R = K[\{x_{\mathbf{u}}\}_{\mathbf{u} \in \langle \Omega \rangle^k}],$$

where $\Omega = \{1, 2, 3, \dots\}$ is the set of positive integers, and that the actions of \mathfrak{S}_n on R_n combine uniquely to an action of \mathfrak{S}_∞ on R . Let now $f(y_1, \dots, y_k) \in K[y_1, \dots, y_k]$, let t_1, t_2, \dots be an infinite sequence of pairwise distinct indeterminates over K , and for $n \geq k$ consider the K -algebra homomorphism

$$\phi_n: R_n \rightarrow K[t_1, \dots, t_n], \quad x_{(u_1, \dots, u_k)} \mapsto f(t_{u_1}, \dots, t_{u_k}).$$

The ideal

$$Q_n = \ker \phi_n$$

of R_n determined by such a map is the prime ideal of algebraic relations between the quantities $f(t_{u_1}, \dots, t_{u_k})$. Such ideals arise in chemistry [9, 15, 16]; of specific interest there is when f is a Vandermonde polynomial $\prod_{i < j} (y_i - y_j)$. In this case, the ideals Q_n correspond to relations among a series of experimental measurements. One would then like to understand the limiting behavior of such relations, and in particular, to see that they stabilize up to the action of the symmetric group.

EXAMPLE 5.1. The permutation $\sigma = (123) \in \mathfrak{S}_3$ acts on the elements

$$(1, 2), (2, 1), (1, 3), (3, 1), (2, 3), (3, 2)$$

of $\langle 3 \rangle^2$ to give

$$(2, 3), (3, 2), (2, 1), (1, 2), (3, 1), (1, 3),$$

respectively. Let $f(t_1, t_2) = t_1^2 t_2$. Then the action of σ on the valid relation $x_{12}^2 x_{31} - x_{13}^2 x_{21} \in Q_3$ gives us another relation $x_{23}^2 x_{12} - x_{21}^2 x_{32} \in Q_3$.

It is easy to see that by construction, the chain Q_\circ of ideals

$$Q_k \subseteq Q_{k+1} \subseteq \cdots \subseteq Q_n \subseteq \cdots$$

(which we call the chain of ideals *induced by the polynomial f*) is an invariant chain. As in the proof of Theorem 4.7, we would like to form the ideal $Q = \bigcup_{n \geq k} Q_n$ of the infinite-dimensional polynomial ring $R = \bigcup_{n \geq k} R_n$, and then apply a finiteness theorem to conclude that Q_\circ stabilizes in the sense mentioned above (Definition 4.5). For $k = 1$, Theorem 4.7 indeed does the job. Unfortunately however, this simple-minded approach fails for $k \geq 2$:

PROPOSITION 5.2. *For $k \geq 2$, the $R[\mathfrak{S}_\infty]$ -module R is not Noetherian.*

PROOF. Let us make the dependence on k explicit and denote R by $R^{(k)}$. Then

$$x_{(u_1, \dots, u_k, u_{k+1})} \mapsto x_{(u_1, \dots, u_k)}$$

defines a surjective K -algebra homomorphism $\pi_k: R^{(k+1)} \rightarrow R^{(k)}$ with invariant kernel. Hence if $R^{(k+1)}$ is Noetherian as an $R[\mathfrak{S}_\infty]$ -module, then so is $R^{(k)}$; thus it suffices to prove the proposition in the case $k = 2$. Suppose therefore that $k = 2$. By Lemma 3.13 it is enough to produce an infinite bad sequence for the quasi-ordering $|\mathfrak{S}_\infty$ of X^\diamond , where $X = \{x_i : i \in \langle \Omega \rangle^2\}$. For this, consider the sequence of monomials

$$\begin{aligned} s_3 &= x_{(1,2)}x_{(3,2)}x_{(3,4)} \\ s_4 &= x_{(1,2)}x_{(3,2)}x_{(4,3)}x_{(4,5)} \\ s_5 &= x_{(1,2)}x_{(3,2)}x_{(4,3)}x_{(5,4)}x_{(6,7)} \\ &\vdots \\ s_n &= x_{(1,2)}x_{(3,2)}x_{(4,3)} \cdots x_{(n,n-1)}x_{(n,n+1)} \quad (n = 3, 4, \dots) \\ &\vdots \end{aligned}$$

Now for $n < m$ and any $\sigma \in \mathfrak{S}_\infty$, the monomial σs_n does not divide s_m . To see this, suppose otherwise. Note that $x_{(1,2)}, x_{(3,2)}$ is the only pair of indeterminates which divides s_n or s_m and has the form $x_{(i,j)}, x_{(l,j)}$ ($i, j, l \in \Omega$). Therefore $\sigma(2) = 2$, and either $\sigma(1) = 1, \sigma(3) = 3$, or $\sigma(1) = 3, \sigma(3) = 1$. But since 1 does not appear as the second component j of a factor $x_{(i,j)}$ of s_m , we have $\sigma(1) = 1, \sigma(3) = 3$. Since $x_{(4,3)}$ is the only indeterminate dividing s_n or s_m of the form $x_{(i,3)}$ with $i \in \Omega$, we get $\sigma(4) = 4$; since $x_{(5,4)}$ is the only indeterminate dividing s_n or s_m of the form $x_{(i,4)}$ with $i \in \Omega$, we get $\sigma(5) = 5$; etc. Ultimately this yields $\sigma(i) = i$ for all $i = 1, \dots, n$. But the only indeterminate dividing s_m of the form $x_{(n,j)}$ with $j \in \Omega$ is $x_{(n,n-1)}$, hence the factor $\sigma x_{(n,n+1)} = x_{(n,\sigma(n+1))}$ of σs_n does not divide s_m . This shows that s_3, s_4, \dots is a bad sequence for the quasi-ordering $|\mathfrak{S}_\infty$, as claimed. \square

REMARK 5.3. The construction of the infinite bad sequence s_3, s_4, \dots in the proof of the previous proposition was inspired by an example in [7].

5.1. A criterion for stabilization. Our next goal is to give a condition for the chain Q_\circ to stabilize. Given $g \in R$, we define the *variable size* of g to be the number of distinct indeterminates $x_{\mathbf{u}}$ that appear in g . For example, $g = x_{12}^5 + x_{45}x_{23} + x_{45}$ has variable size 3.

LEMMA 5.4. *A chain of ideals Q_\circ induced by a polynomial $f \in K[y_1, \dots, y_k]$ stabilizes modulo the symmetric group if and only if there exist integers M and N such that for all $n > N$, there are generators for Q_n with variable sizes at most M . Moreover, in this case a bound for stabilization is given by $\max(N, kM)$.*

PROOF. Suppose M and N are integers with the stated property. To see that Q_\circ stabilizes, since Q_\circ is an invariant chain, we need only verify that $N' = \max(N, kM)$ is such that $Q_m \subseteq L_m(Q_n)$ for $m \geq n > N'$. For this inclusion, it suffices that each generator in a generating set for the ideal Q_m of R_m is in $L_m(Q_n)$. Since $m > N$, there are generators B for Q_m with variable sizes at most M . If $g \in B$, then there are at most kM different integers appearing as subscripts of indeterminates in g . We can form a permutation $\sigma \in \mathfrak{S}_m$ such that $\sigma g \in R_{N'}$ and thus in R_n . But then $\sigma g \in P_n(Q_m) \subseteq Q_n$ so that $g = \sigma^{-1}\sigma g \in L_m(Q_n)$ as desired.

Conversely, suppose that Q_\circ stabilizes. Then there exists an N such that $Q_m = L_m(Q_N)$ for all $m > N$. Let B be any finite generating set for Q_N . Then for all $m > N$, $Q_m = L_m(B)$ is generated by elements of bounded variable size, by Lemma 4.2. \square

Although this condition is a very simple one, it will prove useful. Below we will apply it together with a preliminary reduction to the case that each indeterminate y_1, \dots, y_k actually occurs in the polynomial f , which we explain next. For this we let $\pi_k: R^{(k+1)} \rightarrow R^{(k)}$ be the surjective K -algebra homomorphism defined in the proof of Proposition 5.2. We write $Q^{(k)}$ for Q , and considering $f \in K[y_1, \dots, y_k]$ as an element of $K[y_1, \dots, y_k, y_{k+1}]$, we also let $Q^{(k+1)}$ be the kernel of the K -algebra homomorphism

$$R^{(k+1)} \rightarrow K[t_1, t_2, \dots], \quad x_{(u_1, \dots, u_k, u_{k+1})} \mapsto f(t_{u_1}, \dots, t_{u_k}, t_{u_{k+1}}) \quad (= f(t_{u_1}, \dots, t_{u_k})).$$

Note that $\pi_k(Q^{(k+1)}) = Q^{(k)}$, and the ideal $\ker \pi_k$ of $R^{(k+1)}$ is generated by the elements

$$x_{(u_1, \dots, u_k, i)} - x_{(u_1, \dots, u_k, j)} \quad (i, j \in \Omega),$$

in particular $\ker \pi_k \subseteq Q^{(k+1)}$. It is easy to see that as an $R^{(k+1)}[\mathfrak{S}_\infty]$ -module, $\ker \pi_k$ is generated by the single element $x_{(1, \dots, k, k+1)} - x_{(1, \dots, k, k+2)}$. These observations now yield:

LEMMA 5.5. *Suppose that the invariant ideal $Q^{(k)}$ of $R^{(k)}$ is finitely generated as an $R^{(k)}[\mathfrak{S}_\infty]$ -module. Then the invariant ideal $Q^{(k+1)}$ of $R^{(k+1)}$ is finitely generated as an $R^{(k+1)}[\mathfrak{S}_\infty]$ -module. \square*

We let \mathfrak{S}_k act on $\langle \Omega \rangle^k$ by

$$\tau(u_1, \dots, u_k) = (u_{\tau(1)}, \dots, u_{\tau(k)}) \quad \text{for } \tau \in \mathfrak{S}_k, (u_1, \dots, u_k) \in \langle \Omega \rangle^k.$$

This action gives rise to an action of \mathfrak{S}_k on $\{x_{\mathbf{u}}\}_{\mathbf{u} \in \langle \Omega \rangle^k}$ by $\tau x_{\mathbf{u}} = x_{\tau \mathbf{u}}$, which we extend to an action of \mathfrak{S}_k on R in the natural way. We also let \mathfrak{S}_k act on $K[y_1, \dots, y_k]$ by $\tau f(y_1, \dots, y_k) = f(y_{\tau(1)}, \dots, y_{\tau(k)})$. Note that

$$\tau Q_k \subseteq \tau Q_{k+1} \subseteq \dots \subseteq \tau Q_n \subseteq \dots$$

is the chain induced by τf . Using the lemma above we obtain:

COROLLARY 5.6. *Let $f \in K[y_1, \dots, y_k]$. There are $i \in \{0, \dots, k\}$ and $\tau \in \mathfrak{S}_k$ such that $\tau f \in K[y_1, \dots, y_i]$ and each of the indeterminates y_1, \dots, y_i occurs in τf . If the chain of ideals induced by the polynomial τf stabilizes, then so does the chain of ideals induced by f . \square*

5.2. Chains induced by monomials. If the given polynomial f is a monomial, then the homomorphism ϕ_n from above produces a (homogeneous) toric kernel Q_n . In particular, there is a finite set of binomials that generate Q_n (see [17]). Although a proof for the general toric case eludes us, we do have the following.

THEOREM 5.7. *The sequence of kernels induced by a square-free monomial $f \in K[y_1, \dots, y_k]$ stabilizes modulo the symmetric group. Moreover, a bound for when stabilization occurs is $N = 4k$.*

To prepare for the proof of this result, we discuss in detail the toric encoding associated to our problem (see [17, Chapter 14] for more details). By Corollary 5.6, we may assume that $f = y_1 \cdots y_k$. Then $g - \tau g \in Q$ for all $g \in R$. We say that $\mathbf{u} = (u_1, \dots, u_k) \in \langle \Omega \rangle^k$ is *sorted* if $u_1 < \dots < u_k$, and *unsorted* otherwise; similarly we say that $x_{\mathbf{u}}$ is sorted (unsorted) if \mathbf{u} is sorted (unsorted, respectively). For example, x_{135} is a sorted indeterminate, whereas x_{315} is not. Consider the set of vectors

$$\mathcal{A}_n = \{(i_1, \dots, i_n) \in \mathbb{Z}^n : i_1 + \dots + i_n = k, 0 \leq i_1, \dots, i_n \leq 1\}.$$

View \mathcal{A}_n as an n -by- $\binom{n}{k}$ matrix entries with 0 and 1, whose columns are indexed by sorted indeterminates $x_{\mathbf{u}}$ and whose rows are indexed by t_i ($i = 1, \dots, n$). (See Example 5.9 below.) Let $\text{sort}(\cdot)$ denote the operator

which takes any word in $\{1, \dots, n\}^*$ and sorts it in increasing order. By [17, Remark 14.1], the toric ideal $I_{\mathcal{A}_n}$ associated to \mathcal{A}_n is generated (as K -vector space) by the binomials $x_{\mathbf{u}_1} \cdots x_{\mathbf{u}_r} - x_{\mathbf{v}_1} \cdots x_{\mathbf{v}_r}$, where $r \in \mathbb{N}$ and the $\mathbf{u}_i, \mathbf{v}_j$ are sorted elements of $\langle n \rangle^k$ such that $\text{sort}(\mathbf{u}_1 \cdots \mathbf{u}_r) = \text{sort}(\mathbf{v}_1 \cdots \mathbf{v}_r)$. In particular, we have $I_{\mathcal{A}_n} \subseteq Q_n$. Let B be any set of generators for the ideal $I_{\mathcal{A}_n}$.

LEMMA 5.8. *A generating set for the ideal Q_n of R_n is given by*

$$S = B \cup \{x_{\mathbf{u}} - x_{\tau\mathbf{u}} : \tau \in \mathfrak{S}_k, \mathbf{u} \text{ is sorted}\}.$$

PROOF. Elements of Q_n are of the form $g = x_{\mathbf{u}_1} \cdots x_{\mathbf{u}_r} - x_{\mathbf{v}_1} \cdots x_{\mathbf{v}_r}$, in which the \mathbf{u}_i and \mathbf{v}_j are ordered k -element subsets of $\{1, \dots, n\}$ such that $\text{sort}(\mathbf{u}_1 \cdots \mathbf{u}_r) = \text{sort}(\mathbf{v}_1 \cdots \mathbf{v}_r)$. We induct on the number t of \mathbf{u}_i and \mathbf{v}_j that are not sorted. If $t = 0$, then $g \in I_{\mathcal{A}_n}$, and we are done. Suppose now that $t > 0$ and assume without loss of generality that \mathbf{u}_1 is not sorted. Let $\tau \in \mathfrak{S}_k$ be such that $\tau\mathbf{u}_1$ is sorted, and consider the element $h = x_{\tau\mathbf{u}_1} x_{\mathbf{u}_2} \cdots x_{\mathbf{u}_r} - x_{\mathbf{v}_1} \cdots x_{\mathbf{v}_r}$ of Q_n . This binomial involves $t - 1$ unsorted indeterminates, and therefore, inductively, can be expressed in terms of S . But then

$$g = h - (x_{\tau\mathbf{u}_1} - x_{\mathbf{u}_1})x_{\mathbf{u}_2} \cdots x_{\mathbf{u}_r}$$

can as well, completing the proof. □

EXAMPLE 5.9. Let $k = 2$ and $n = 4$. Then

	x_{12}	x_{13}	x_{14}	x_{23}	x_{24}	x_{34}
t_1	1	1	1	0	0	0
t_2	1	0	0	1	1	0
t_3	0	1	0	1	0	1
t_4	0	0	1	0	1	1

represents the matrix associated to \mathcal{A}_4 . The ideal $I_{\mathcal{A}_4}$ is generated by the two binomials $x_{13}x_{24} - x_{12}x_{34}$ and $x_{14}x_{23} - x_{12}x_{34}$. Hence Q_4 is generated by these two elements along with

$$\{x_{12} - x_{21}, x_{13} - x_{31}, x_{14} - x_{41}, x_{23} - x_{32}, x_{24} - x_{42}, x_{34} - x_{43}\}.$$

We are now in a position to prove Theorem 5.7.

PROOF OF THEOREM 5.7. By Lemma 5.4, we need only show that there exist generators for Q_n which have bounded variable sizes. Using [17, Theorem 14.2], it follows that $I_{\mathcal{A}_n}$ has a quadratic (binomial) Gröbner basis for each n (with respect to some term ordering of R_n). By Lemma 5.8, there is a set of generators for Q_n with variable sizes at most 4. This proves the theorem. □

We close with a conjecture that generalizes Theorem 5.7.

CONJECTURE 5.10. *The sequence of kernels induced by a monomial f stabilizes modulo the symmetric group.*

6. Acknowledgment

We would like to thank Bernd Sturmfels for bringing the problem found in Section 5 (originating from Andreas Dress) to our attention and for making us aware of Theorem 14.2 in [17].

References

- [1] W. Adams, P. Lounstaunau, *An Introduction to Gröbner Bases*, Graduate Studies in Mathematics, vol. 3, American Mathematical Society, Providence, RI, 1994.
- [2] G. Ahlbrandt, M. Ziegler, *Quasi-finitely axiomatizable totally categorical theories*, Stability in model theory (Trento, 1984), Ann. Pure Appl. Logic **30** (1986), no. 1, 63–82.
- [3] A. R. Camina, D. M. Evans, *Some infinite permutation modules*, Quart. J. Math. Oxford Ser. (2) **42** (1991), no. 165, 15–26.
- [4] P. Erdős, R. Rado, *A theorem on partial well-ordering of sets of vectors*, J. London Math. Soc. **34** (1959), 222–224.
- [5] D. Eisenbud, *Commutative Algebra with a View Toward Algebraic Geometry*, Graduate Texts in Mathematics, vol. 203, Springer-Verlag, New York, 1995.
- [6] G. Higman, *Ordering by divisibility in abstract algebras*, Proc. London Math. Soc. (3) **2** (1952), 326–336.
- [7] T. A. Jenkyns, C. St. J. A. Nash-Williams, *Counterexamples in the theory of well-quasi-ordered sets*. in: F. Harary (ed.), *Proof Techniques in Graph Theory (Proc. Second Ann Arbor Graph Theory Conf., Ann Arbor, Mich., 1968)*, pp. 87–91, Academic Press, New York, 1969.

FINITE GENERATION OF SYMMETRIC IDEALS

- [8] J. B. Kruskal, *The theory of well-quasi-ordering: A frequently discovered concept*, J. Combinatorial Theory Ser. A **13** (1972), 297–305.
- [9] A. Mead, E. Ruch, A. Schönhofer, *Theory of chirality functions, generalized for molecules with chiral ligands*. Theor. Chim. Acta **29** (1973), 269–304.
- [10] R. Michler, *Gröbner bases of symmetric quotients and applications*, in: C. Christensen et al. (eds.), *Algebra, Arithmetic and Geometry with Applications (West Lafayette, IN, 2000)*, pp. 627–637, Springer-Verlag, Berlin, 2004.
- [11] E. Milner, *Well-quasi-ordering of sequences of ordinal numbers*, J. London Math. Soc. **43** (1968), 291–296.
- [12] C. St. J. A. Nash-Williams, *On well-quasi-ordering finite trees*, Proc. Cambridge Philos. Soc. **59** (1963), 833–835.
- [13] ———, *On well-quasi-ordering transfinite sequences*, Proc. Cambridge Philos. Soc. **61** (1965), 33–39.
- [14] R. Rado, *Partial well-ordering of sets of vectors*, Mathematika **1** (1954), 89–95.
- [15] E. Ruch, A. Schönhofer, *Theorie der Chiralitätsfunktionen*, Theor. Chim. Acta **19** (1970), 225–287.
- [16] E. Ruch, A. Schönhofer, I. Ugi, *Die Vandermondesche Determinante als Näherungsansatz für eine Chiralitätsbeobachtung, ihre Verwendung in der Stereochemie und zur Berechnung der optischen Aktivität*, Theor. Chim. Acta **7** (1967), 420–432.
- [17] B. Sturmfels, *Gröbner Bases and Convex Polytopes*, AMS University Lecture Series, vol. 8, American Mathematical Society, Providence, RI, 1996.

DEPARTMENT OF MATHEMATICS, STATISTICS, AND COMPUTER SCIENCE, UNIVERSITY OF ILLINOIS AT CHICAGO, CHICAGO, IL 60607.

E-mail address: `maschenb@math.uic.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720.

E-mail address: `chillar@math.berkeley.edu`