

FINITE REGULAR BANDS ARE FINITELY RELATED

IGOR DOLINKA

(Received 15 December 2011)

Abstract

An algebra \mathbf{A} is said to be finitely related if the clone $\text{Clo}(\mathbf{A})$ of its term operations is determined by a finite set of finitary relations. We prove that each finite idempotent semigroup satisfying the identity $xyxzx \approx xyzx$ is finitely related.

2010 *Mathematics subject classification*: primary 20M07; secondary 08A40, 05A05.

Keywords and phrases: finite regular band, term operation, variety, finitely related.

1. Introduction

Let $\mathbf{A} = (A, \mathcal{F})$ be an algebra. If $\mathbf{t} = \mathbf{t}(x_1, \dots, x_n)$ is a term of the same similarity type as \mathbf{A} over the alphabet $X_n = \{x_i : 1 \leq i \leq n\}$, then \mathbf{t} induces (by interpretation) an operation $\hat{\mathbf{t}} : A^n \rightarrow A$. Operations on A arising in this way are called the *term operations* of \mathbf{A} . The collection of all term operations of \mathbf{A} is called the *clone* of the algebra \mathbf{A} and denoted by $\text{Clo}(\mathbf{A})$. A k -ary relation $\rho \subseteq A^k$ is said to be *compatible* with \mathbf{A} if ρ is a subalgebra of the direct power \mathbf{A}^k .

For a set \mathcal{R} of finitary relations on a set X , let $\text{Pol}(\mathcal{R})$ denote the set of all *polymorphisms* of \mathcal{R} : these are all finitary operations $f : X^n \rightarrow X$ preserving all relations from \mathcal{R} (so that each relation from \mathcal{R} is compatible with (X, f)). A standard result in clone theory tells us that if $C_{\mathbf{A}}$ is the set of all compatible relations of \mathbf{A} then $\text{Clo}(\mathbf{A}) = \text{Pol}(C_{\mathbf{A}})$; that is, an operation on A arises from a term if and only if it preserves all of the compatible relations. The set $C_{\mathbf{A}}$ is always infinite. It may happen, however, that $\text{Clo}(\mathbf{A})$ is determined by a finite set of relations, so that $\text{Clo}(\mathbf{A}) = \text{Pol}(C')$ holds for a finite $C' \subseteq C_{\mathbf{A}}$. In such a case we say that the algebra \mathbf{A} is *finitely related*. We direct the reader to [6, 12] for an overview of known results on finitely related (finite) algebras; for a general background in universal algebra we refer to [5].

The support of the Ministry of Education and Science of the Republic of Serbia, through grant number 174019, is gratefully acknowledged. The author is also supported by a grant (Contract 114–451–2002/2011) of the Secretariat of Science and Technological Development of the Autonomous Province of Vojvodina.

© 2012 Australian Mathematical Publishing Association Inc. 0004-9727/2012 \$16.00

A major result obtained recently by Aichinger *et al.* [2] implies that any finite algebra having a Mal'cev term operation is finitely related; in particular, this includes all finite groups and rings (see also [1] for a result in a very similar vein). Davey *et al.* [6] presented several classes of finite semigroups with this property: for example, it is enjoyed by a finite semigroup S whenever S is either commutative, or nilpotent. Here we add a new entry to this list of examples. Namely, we prove that each finite *regular band* (idempotent semigroup satisfying $xyxzx \approx xyzx$, see [14]) is finitely related. To achieve this, we apply a general criterion from [6] ensuring the finitely related property for a finite algebra.

Here is how the present note is organised. In the following two sections we gather the prerequisites necessary for our main proof. In more detail, in the next section we revisit the notion (from [6]) of an n -scheme of terms for a given variety \mathcal{V} and formulate the mentioned general result implying that an algebra generating \mathcal{V} is finitely related. These concepts are close relatives of the famous Reconstruction Conjecture of Ulam and Kelly [11, 15] from graph theory, and indeed we prove an auxiliary result of similar flavour about reconstruction of permutations from their maximal proper subpermutations. In Section 3 we quickly recall the lattice of subvarieties of the variety \mathcal{RB} of regular bands and their (well-known) equational theories. Finally, the proof of our main result, whose statement is contained in the title of the paper, is presented in Section 4.

Note added in revision. In mid-February 2012, around two months after this note was submitted, I have learned from a personal communication with Peter Mayr (CAUL, Lisbon) that he independently proved the main result of this paper; this proof is included in the wider study [13] of finite semigroups with respect to the property of being finitely related.

2. Term schemes and reconstructing permutations

Let \mathbf{t} be a term of a given similarity type over the alphabet $X_n = \{x_1, \dots, x_n\}$, $n \geq 1$. Following the terminology related to Ulam's Reconstruction Conjecture (see, for example, [4]) we call a *card* of \mathbf{t} a term $\mathbf{t}^{(ij)}$, $1 \leq i < j \leq n$, obtained from \mathbf{t} by replacing each occurrence of x_i by x_j , so that the letters of $\mathbf{t}^{(ij)}$ are contained in $X_n^{(i)} = X_n \setminus \{x_i\}$. Furthermore, we set $\mathbf{t}^{(ji)}$ to be just $\mathbf{t}^{(ij)}$. The *deck* of \mathbf{t} is the family $\mathcal{D}_{\mathbf{t}} = \{\mathbf{t}^{(ij)} : 1 \leq i < j \leq n\}$ of all cards of \mathbf{t} .

Given a variety \mathcal{V} and an integer $n \geq 2$, the concept of a deck motivates the following abstract definition. Let $\mathcal{S} = \{\mathbf{t}_{ij} : 1 \leq i < j \leq n\}$ be an indexed family of terms (of the same similarity type as \mathcal{V}) over X_n satisfying the following conditions.

- (D) For arbitrary $\mathbf{A} \in \mathcal{V}$, the term operation $\widehat{\mathbf{t}}_{ij}(x_1, \dots, x_n)$ induced on \mathbf{A} by \mathbf{t}_{ij} does not depend on the variable x_i .
- (C1) For any four distinct $1 \leq i, j, p, q \leq n$ such that $i < j$ and $p < q$, \mathcal{V} satisfies $\mathbf{t}_{ij}^{(pq)} \approx \mathbf{t}_{pq}^{(ij)}$.
- (C2) For any three distinct $1 \leq i, j, k \leq n$ such that $i < j < k$, \mathcal{V} satisfies the identities $\mathbf{t}_{ij}^{(jk)} \approx \mathbf{t}_{jk}^{(ik)} \approx \mathbf{t}_{ik}^{(jk)}$.

The condition (D) is called *dependency*, while (C1) and (C2) are *consistency* conditions. The family \mathcal{S} is called an *n-scheme* for the variety \mathcal{V} . It is an easy exercise to show that the notion of an *n-scheme* of terms just defined coincides with the definition of an $(n, n - 1)$ -scheme from [6], since it is straightforward to prove that the previous three properties are equivalent to dependency and consistency conditions (D) and (C) given there. We say that an *n-scheme* \mathcal{S} for \mathcal{V} comes from the term \mathbf{t} if \mathcal{S} and $\mathcal{D}_{\mathbf{t}}$ are equivalent in the sense that $\mathcal{V} \models \mathbf{t}^{(ij)} \approx \mathbf{t}_{ij}$ holds for all $1 \leq i < j \leq n$.

We are now ready to state a general criterion from [6] for a finite algebra to be finitely related. The following constitutes a part of Theorem 2.9 from that paper.

THEOREM 2.1 (Davey *et al.* [6]). *Let \mathbf{A} be a finite algebra generating the variety \mathcal{V} . Then \mathbf{A} is finitely related if and only if there exists an integer n_0 such that for all $n \geq n_0$ every *n-scheme* for \mathcal{V} comes from a term.*

In the case of semigroups, term operations coincide with operations induced by words (elements of the free semigroup X^+ over a suitable alphabet X), and all definitions above remain in place when we introduce words instead of terms, even though words are not terms in the strict sense. (However, they can be made terms for example by left-grouping of parentheses.) So, a finite semigroup S is finitely related if and only if every *n-scheme* of words for the variety generated by S comes from a word whenever n is large enough: that is, every such *n-scheme* is *S-equivalent* to the deck of a single word \mathbf{w} . This is the form of the previous general result that we are going to use in the main course of our proof.

In the following, we provide a ‘combinatorial counterpart’ of the notions just introduced for permutations on finite sets. For $n \geq 1$, we let $[1, n] = \{1, \dots, n\}$ and identify a permutation of $[1, n]$ with a string in which each element of $[1, n]$ occurs precisely once (that is, position i is permuted to the value in position i). Now, for a permutation π of $[1, n]$ and $1 \leq i < j \leq n$ we define its card $\pi^{(ij)}$ as the permutation of $[1, n]^{(i)} = \{1, \dots, i - 1, i + 1, \dots, n\}$ obtained from π by replacing the entry i by j and then deleting the second (from left to right) of the two occurrences of j arising in this way. (For example, $(21354)^{(15)} = 2534$.) The deck of π is just $\{\pi^{(ij)} : 1 \leq i < j \leq n\}$.

A family of permutations $\{\pi_{ij} : 1 \leq i < j \leq n\}$ is a *permutational n-scheme* if the following conditions are satisfied:

- (1) π_{ij} is a permutation of the set $[1, n]^{(i)}$;
- (2) if $1 \leq i, j, p, q \leq n$ are distinct and $i < j, p < q$, then $\pi_{ij}^{(pq)} = \pi_{pq}^{(ij)}$;
- (3) if $1 \leq i, j, k \leq n$ are distinct and $i < j < k$, then $\pi_{ij}^{(jk)} = \pi_{jk}^{(ik)} = \pi_{ik}^{(jk)}$.

PROPOSITION 2.2. *If $n \geq 6$, then each permutational *n-scheme* coincides with the deck of a permutation of $[1, n]$. In more detail, for any scheme $\{\pi_{ij} : 1 \leq i < j \leq n\}$ there exists a permutation π of $[1, n]$ such that $\pi^{(ij)} = \pi_{ij}$ for all $1 \leq i < j \leq n$.*

PROOF. Let p' and p'' be the first two entries in π_{12} (as a string) other than 2. Without loss of generality we may assume $p' < p''$ (up to some ordering on $[1, n]$). Now write $\pi_{p'p''} = p_1 p_2 p_3 q_1 \dots q_{n-6} q' q''$ (we are going to assume that, for example, $q' < q''$,

the other case being different only by notational details); since by (2) we have $\pi_{p'p''}^{(12)} = \pi_{12}^{(p'p'')}$, we conclude that $p'' \in \{p_1, p_2, p_3\}$. On the other hand,

$$\pi_{q'q''}^{(p'p'')} = \pi_{p'p''}^{(q'q'')} = p_1 p_2 p_3 q_1 \dots q_{n-6} q''',$$

so $\pi_{q'q''}$ is just $p_1 p_2 p_3 q_1 \dots q_{n-6} q'''$ with p' inserted either immediately before p'' or after it. In particular, at least one of p' or p'' occur in the first three entries of $\pi_{q'q''}$. We claim that *both* p' and p'' occur among the first four entries of $\pi_{q'q''}$. If $\{q', q''\} = \{1, 2\}$, then p', p'' are already among the first three entries of $\pi_{q'q''}$ (thus $\pi_{q'q''}$ begins with $p' p_1 p_2 p_3, p_1 p' p_2 p_3$ or $p_1 p_2 p' p_3$). Otherwise, consider $\pi_{q'q''}^{(12)}$. If $\{1, 2\} \cap \{q', q''\} = \emptyset$, then $\pi_{q'q''}^{(12)} = \pi_{12}^{(q'q'')}$, and so, since $\{p', p''\}$ is disjoint both from $\{1, 2\}$ and $\{q', q''\}$, it follows that both p', p'' are among the first three entries of $\pi_{q'q''}^{(12)}$, and thus among the first four entries of $\pi_{q'q''}$. On the other hand, if, for example, $q' = 1$ and $q'' \neq 2$, then by (3) we establish $\pi_{1q''}^{(2q'')} = \pi_{12}^{(2q'')}$; as the latter permutation has both p', p'' among its first three entries, these elements must be among the first four entries of $\pi_{1q''}$. The argument is similar if $\{1, 2\} \cap \{q', q''\} = \{2\}$.

We are now prepared to define the permutation π for which we claim that it satisfies the requirements of the proposition. If $r_1 r_2 r_3 r_4$ is the sequence of first four entries of $\pi_{q'q''}$ (which is just $p_1 p_2 p_3$ with p' inserted somewhere) we define

$$\pi = r_1 r_2 r_3 r_4 q_1 \dots q_{n-6} q' q''.$$

In other words, π is obtained by ‘patching’ together the prefix of $\pi_{q'q''}$ of length 4 and the suffix of $\pi_{p'p''}$ of length $n - 4$. The arguments in the previous paragraph show that this is indeed a permutation of $[1, n]$ since it is obtained from $\pi_{q'q''}$ by inserting p' somewhere before q_1 (or before q' if $n = 6$). Note that, in this notation, $\pi_{q'q''} = r_1 r_2 r_3 r_4 q_1 \dots q_{n-6} q' q''$.

It is now immediately clear that $\pi^{(q'q'')} = \pi_{q'q''}$ and it takes only a short reflection to see that

$$\pi^{(p'p'')} = (r_1 r_2 r_3 r_4)^{(p'p'')} q_1 \dots q_{n-6} q' q'' = p_1 p_2 p_3 q_1 \dots q_{n-6} q' q'' = \pi_{p'p''}.$$

Furthermore, let $i < j$ be such that $\{i, j\} \cap \{q', q''\} = \emptyset$. Then $\pi_{ij}^{(q'q'')} = \pi_{q'q''}^{(ij)} = (r_1 r_2 r_3 r_4 q_1 \dots q_{n-6})^{(ij)} q''$, so that $\pi_{ij} = (r_1 r_2 r_3 r_4 q_1 \dots q_{n-6})^{(ij)} \bar{q}$, where \bar{q} is either $q' q''$ or $q'' q'$. However, the second possibility is excluded, because if $\{i, j\} \neq \{p', p''\}$, then $\pi_{ij}^{(p'p'')} = \pi_{p'p''}^{(ij)}$ by (2), or $\pi_{ij}^{(jp'')} = \pi_{p'p''}^{(jp'')}$ by (3) if $i = p'$ or $\pi_{ij}^{(p'p'')} = \pi_{p'p''}^{(ip'')}$ by (3) if $j = p''$; in all three cases we obtain permutations ending with $q' q''$. Hence, $\pi_{ij} = (r_1 r_2 r_3 r_4 q_1 \dots q_{n-6})^{(ij)} q' q'' = \pi_{(ij)}$ in this case. Finally, assume that either $i = q', j \neq q''$ or $i \neq q', j = q''$ (we will consider only the first possibility, the second being similar). Then $\pi_{ij}^{(jq'')} = \pi_{q'q''}^{(jq'')}$, which is the permutation of $[1, n] \setminus \{q', \min\{j, q''\}\}$ obtained from $r_1 r_2 r_3 r_4 q_1 \dots q_{n-6}$ by replacing j by $\max\{j, q''\}$. By a similar argument as above, the card $\pi_{ij}^{(p'p'')}$ ends with q'' , implying that π_{ij} , too, ends with q'' . This, taken together with the available information on $\pi_{ij}^{(jq'')}$, yields that $\pi_{ij} = r_1 r_2 r_3 r_4 q_1 \dots q_{n-6} q'' = \pi^{(ij)}$, as required. Therefore, the given permutational scheme coincides with the deck of π . \square




FIGURE 1. The lattice of regular band varieties.

It is not too difficult to see from the previous proof that, under the given conditions, the permutation π satisfying the above proposition must in fact be unique. Of course, it is an interesting side question whether the bound of 6 from the previous statement can be lowered and by how much. It is likely that this can be verified by computational methods.

3. Varieties of regular bands

In this brief section we review the lattice of all regular band varieties and the description of identities (equational theories) satisfied by these varieties. The lattice $\mathcal{L}(\mathcal{RB})$ of subvarieties of \mathcal{RB} is depicted in Figure 1 (see [3, 8–10, 14]). Here standard notational conventions apply (see, for example, [10]): \mathcal{SL} is the variety of semilattices (commutative bands), \mathcal{ReB} is the variety of rectangular bands (defined by $xyx = x$), \mathcal{NB} is the variety of normal bands (defined by $xyzx = xzyx$), \mathcal{LNB} (\mathcal{RNB}) is the variety of left (right) normal bands (defined by $xyz = xzy$, respectively $yzx = zyx$); finally, \mathcal{LRB} (\mathcal{RRB}) is the variety of left (right) regular bands (defined by $xyx = xy$, respectively $xyx = yx$).

For a word $\mathbf{w} \in X_n^+$, $X_n = \{x_1, \dots, x_n\}$, let $h(\mathbf{w})$ be the *head* of \mathbf{w} , the first letter (from the left) of \mathbf{w} ; dually, the *tail* $t(\mathbf{w})$ of \mathbf{w} is the last letter of \mathbf{w} . Also, let $i(\mathbf{w})$ be the *initial part* of \mathbf{w} , which is obtained by retaining only the first occurrence (from the left) of each letter from the set $c(\mathbf{w})$ of all letters appearing in \mathbf{w} (the *content* of \mathbf{w}). Clearly, to each word \mathbf{w} such that $c(\mathbf{w}) = X_n$ corresponds a permutation $\pi(\mathbf{w})$ of $[1, n]$ simply by recording the indices of letters in $i(\mathbf{w})$; analogously, the *final part* $f(\mathbf{w})$ (obtained by retaining only the last occurrence of each letter appearing in \mathbf{w}) gives rise to the permutation $\bar{\pi}(\mathbf{w})$.

The following provides a summary of the equational theories of ‘interesting’ varieties appearing in Figure 1.

PROPOSITION 3.1. For any $\mathbf{u}, \mathbf{v} \in X_n^+$:

- (i) $\mathcal{LNB} \models \mathbf{u} \approx \mathbf{v}$ if and only if $h(\mathbf{u}) = h(\mathbf{v})$ and $c(\mathbf{u}) = c(\mathbf{v})$;
- (ii) $\mathcal{RNB} \models \mathbf{u} \approx \mathbf{v}$ if and only if $t(\mathbf{u}) = t(\mathbf{v})$ and $c(\mathbf{u}) = c(\mathbf{v})$;
- (iii) $\mathcal{NB} \models \mathbf{u} \approx \mathbf{v}$ if and only if $h(\mathbf{u}) = h(\mathbf{v})$, $t(\mathbf{u}) = t(\mathbf{v})$ and $c(\mathbf{u}) = c(\mathbf{v})$;
- (iv) $\mathcal{LRB} \models \mathbf{u} \approx \mathbf{v}$ if and only if $i(\mathbf{u}) = i(\mathbf{v})$;
- (v) $\mathcal{RRB} \models \mathbf{u} \approx \mathbf{v}$ if and only if $f(\mathbf{u}) = f(\mathbf{v})$;
- (vi) $\mathcal{RB} \models \mathbf{u} \approx \mathbf{v}$ if and only if $i(\mathbf{u}) = i(\mathbf{v})$ and $f(\mathbf{u}) = f(\mathbf{v})$.

We have now gathered all of the necessary prerequisites to start proving that each finite regular band is finitely related.

4. The proof

Throughout, let $\mathcal{W} = \{\mathbf{w}_{ij} : 1 \leq i < j \leq n\}$ be an n -scheme of words for a variety \mathcal{V} from Figure 1 generated by a finite regular band B , where $n \geq 6$. We begin by defining, for each $\alpha \in \{h, t, i, f\}$, a word $\mathbf{u}_{(\alpha)}(\mathcal{W})$, where the case $\alpha = h$ ($\alpha = t$) relies upon the assumption that $\mathcal{LZ} \subseteq \mathcal{V}$ (respectively $\mathcal{RZ} \subseteq \mathcal{V}$), while the case $\alpha = i$ ($\alpha = f$) works under the assumption that $\mathcal{LRB} \subseteq \mathcal{V}$ (respectively $\mathcal{RRB} \subseteq \mathcal{V}$).

First of all, define $\mathbf{u}_{(h)}(\mathcal{W}) = x_r$ if and only if there exist indices $p < q$ such that $r \notin \{p, q\}$ and $h(\mathbf{w}_{pq}) = x_r$.

LEMMA 4.1. The definition of $\mathbf{u}_{(h)}(\mathcal{W})$ is logically correct.

PROOF. Let us first show that there exists an index r with the required properties. If $h(\mathbf{w}_{12}) = x_r$ such that $r \geq 3$, then we are done; otherwise $r \in \{1, 2\}$. Since $n \geq 6$, consider the word \mathbf{w}_{34} . By consistency, the identity $\mathbf{w}_{34}^{(12)} \approx \mathbf{w}_{12}^{(34)}$ holds in \mathcal{V} and thus (by our initial assumptions) in \mathcal{LZ} ; so, $h(\mathbf{w}_{34}^{(12)}) = h(\mathbf{w}_{12}^{(34)}) = [h(\mathbf{w}_{12})]^{(34)} = x_r$. Hence, the first letter of \mathbf{w}_{34} is either x_1 or x_2 ; in any case $h(\mathbf{w}_{34}) = x_r$ such that $r \notin \{3, 4\}$.

It remains to prove that no two distinct indices r, r' satisfy the definition of $\mathbf{u}_{(h)}(\mathcal{W})$. Assume to the contrary: then $h(\mathbf{w}_{pq}) = x_r$ and $h(\mathbf{w}_{p'q'}) = x_{r'}$, while $r \notin \{p, q\}$ and $r' \notin \{p', q'\}$. If $\{p, q\} \cap \{p', q'\} = \emptyset$, then by (C1) we have $\mathcal{V} \models \mathbf{w}_{pq}^{(p'q')} \approx \mathbf{w}_{p'q'}^{(pq)}$, implying

$$x_r^{(p'q')} = h(\mathbf{w}_{pq}^{(p'q')}) = h(\mathbf{w}_{p'q'}^{(pq)}) = x_{r'}^{(pq)}.$$

However, $x_r^{(p'q')} \in \{x_r, x_{q'}\}$ and $x_{r'}^{(pq)} \in \{x_{r'}, x_q\}$, which is a contradiction, as $\{r, q'\} \cap \{r', q\} = \emptyset$. Hence, consider the case $p = p'$ and $q < q'$, the other possibilities being similar. Now, by (C2), \mathcal{V} satisfies $\mathbf{w}_{pq}^{(qq')} \approx \mathbf{w}_{pq'}^{(qq')}$, so

$$x_r = x_r^{(qq')} = h(\mathbf{w}_{pq}^{(qq')}) = h(\mathbf{w}_{pq'}^{(qq')}) = x_{r'}^{(qq')} \in \{x_{r'}, x_{q'}\}.$$

As $r \neq r'$, we must have $r = q'$ and thus $r' = q$. To show this is impossible, choose $s < t \leq n$ such that $s, t \notin \{p, q, q'\}$, which is possible, since $n \geq 6$. Then \mathcal{V} (and, thus, \mathcal{LZ}) satisfies the identities $\mathbf{w}_{pq}^{(st)} \approx \mathbf{w}_{st}^{(pq)}$ and $\mathbf{w}_{st}^{(pq')} \approx \mathbf{w}_{pq'}^{(st)}$. The first of these identities implies

$$x_r = x_r^{(st)} = h(\mathbf{w}_{pq}^{(st)}) = h(\mathbf{w}_{st}^{(pq)}),$$

so that $h(\mathbf{w}_{st}) = x_r$, while the second one yields

$$x_{r'} = x_r^{(st)} = h(\mathbf{w}_{pq'}^{(st)}) = h(\mathbf{w}_{st}^{(pq')}) = x_r^{(pq')} = x_r,$$

a contradiction. □

Dually, we set $\mathbf{u}_{(t)}(\mathcal{W}) = x_r$ if and only if there exist indices $p < q$ such that $r \notin \{p, q\}$ and $t(\mathbf{w}_{pq}) = x_r$. Again, by a dual statement to the above one, this definition is correct, that is, it uniquely determines r .

It is now convenient to take care of the subvarieties of \mathcal{NB} . First we need an auxiliary result.

LEMMA 4.2. *Let $c(\mathcal{W})$ be the set of all letters occurring in some word $\mathbf{w}_{ij} \in \mathcal{W}$, where \mathcal{W} is an n -scheme of words for a variety \mathcal{V} containing \mathcal{SL} . Then for any $p < q$, $c(\mathbf{w}_{pq}) = c(\mathcal{W}) \setminus \{x_p\}$.*

PROOF. By (D), it is immediate that \mathbf{w}_{pq} does not contain x_p (as $\mathcal{SL} \subseteq \mathcal{V}$). Now assume that $x_r \in c(\mathbf{w}_{ij})$ for some $i < j$ and $r \notin \{i, p\}$. If $\{i, j\} \cap \{p, q\} = \emptyset$, then by (C1) and the assumption on \mathcal{V} we have $x_r \in c(\mathbf{w}_{ij}^{(pq)}) = c(\mathbf{w}_{pq}^{(ij)})$, so $x_r \in c(\mathbf{w}_{pq})$. Otherwise, the set $\{i, j, p, q, r\}$ has at most four elements, so there exist at least two indices $k < l$ from $[1, n]$ not belonging to the latter set (as $n \geq 6$). Now we employ (C1) again to conclude that $x_r \in c(\mathbf{w}_{ij}^{(kl)}) = c(\mathbf{w}_{kl}^{(ij)})$, implying $x_r \in c(\mathbf{w}_{kl})$; hence, $x_r \in c(\mathbf{w}_{kl}^{(pq)}) = c(\mathbf{w}_{pq}^{(kl)})$, and so $x_r \in c(\mathbf{w}_{pq})$, as required. □

PROPOSITION 4.3. *Each finite normal band is finitely related.*

PROOF. Since it was proved in [7] that any finite rectangular band is finitely related, and the same holds for any finite semilattice (see, for example, [6, Theorem 3.12]), let us assume first that the finite band B in question generates \mathcal{LNB} . Consider the word

$$\mathbf{w} = \mathbf{u}_{(h)}(\mathcal{W})\mathbf{p}_{c(\mathcal{W})},$$

where $\mathbf{p}_{c(\mathcal{W})}$ denotes the product of the letters from $c(\mathcal{W})$ in the increasing order of their indices. By the previous lemma, for any $p < q$ we have $c(\mathbf{w}^{(pq)}) = c(\mathcal{W}) \setminus \{x_p\} = c(\mathbf{w}_{pq})$. Furthermore, $h(\mathbf{w}^{(pq)}) = \mathbf{u}_{(h)}(\mathcal{W})$ unless $\mathbf{u}_{(h)}(\mathcal{W}) = x_p$, when $h(\mathbf{w}^{(pq)}) = x_q$. On the other hand, Lemma 4.1 shows that if $\mathbf{u}_{(h)}(\mathcal{W}) = x_r$ and $r \notin \{p, q\}$, then $h(\mathbf{w}_{pq}) = x_r$. If $r = p$, then choose $i < j$ such that $\{i, j\} \cap \{p, q\} = \emptyset$; by (C1) we have that $\mathbf{w}_{pq}^{(ij)} \approx \mathbf{w}_{ij}^{(pq)}$ holds in \mathcal{LNB} , so, by Proposition 3.1(i),

$$h(\mathbf{w}_{pq}) = h(\mathbf{w}_{pq}^{(ij)}) = h(\mathbf{w}_{ij}^{(pq)}) = [h(\mathbf{w}_{ij})]^{(pq)} = x_r^{(pq)} = x_q.$$

Finally, if $r = q$ a similar approach would show that $h(\mathbf{w}_{pq}) = x_q = x_r$. This suffices to verify that $h(\mathbf{w}_{pq}) = h(\mathbf{w}^{(pq)})$, so $\mathcal{LNB} \models \mathbf{w}^{(pq)} \approx \mathbf{w}_{pq}$. In other words, \mathcal{W} comes from \mathbf{w} .

The case when B generates \mathcal{RNB} is dual, while the case when it generates \mathcal{NB} is settled along similar lines by using the word

$$\mathbf{w} = \mathbf{u}_{(h)}(\mathcal{W})\mathbf{p}_{c(\mathcal{W})}\mathbf{u}_{(t)}(\mathcal{W});$$

in fact, it suffices to ‘join’ the previous argument for \mathcal{LNB} and its dual. □

Recall that in the process of defining $\mathbf{u}_{(i)}(\mathcal{W})$ we assume that $\mathcal{LRB} \subseteq \mathcal{V}$. Consider the family of permutations $\mathcal{W}^\pi = \{\pi(\mathbf{w}_{ij}) : \mathbf{w}_{ij} \in \mathcal{W}\}$. It is quite easy to see that under the given assumptions \mathcal{W}^π is a permutational n -scheme. Thus, by Proposition 2.2 there is a (unique) permutation $\pi = \pi_1 \dots \pi_n$ of $[1, n]$ such that $\pi^{(ij)} = \pi(\mathbf{w}_{ij})$ for all pairs of indices $i < j$. Now we define $\mathbf{u}_{(i)}(\mathcal{W})$ to be the ‘linear’ (or ‘permutational’) word $x_{\pi_1} \dots x_{\pi_n}$. The word $\mathbf{u}_{(f)}(\mathcal{W})$ is obtained in a dual manner (provided $\mathcal{RRB} \subseteq \mathcal{V}$) from the permutations $\bar{\pi}(\mathbf{w}_{ij})$.

LEMMA 4.4. *For any $1 \leq p < q \leq n$*

$$i([\mathbf{u}_{(i)}(\mathcal{W})]^{(pq)}) = i(\mathbf{w}_{pq}) \quad \text{and} \quad f([\mathbf{u}_{(f)}(\mathcal{W})]^{(pq)}) = f(\mathbf{w}_{pq}).$$

PROOF. By definition, $i([\mathbf{u}_{(i)}(\mathcal{W})]^{(pq)}) = x_{\sigma_1} \dots x_{\sigma_{n-1}}$ where $\sigma = \sigma_1 \dots \sigma_{n-1} = \pi^{(pq)}$, while $i(\mathbf{w}_{pq}) = x_{\tau_1} \dots x_{\tau_{n-1}}$ where $\tau = \tau_1 \dots \tau_{n-1} = \pi(\mathbf{w}_{pq})$. By our construction, we have $\pi^{(ij)} = \pi(\mathbf{w}_{ij})$. Thus $i([\mathbf{u}_{(i)}(\mathcal{W})]^{(pq)}) = i(\mathbf{w}_{pq})$. The second equality follows analogously. \square

The main result of this note is now a consequence of the work done previously and Lemma 4.4.

THEOREM 4.5. *Each finite regular band is finitely related.*

PROOF. If a finite regular band B generates a subvariety of \mathcal{NB} , then the statement follows from Proposition 4.3. If B generates \mathcal{LRB} , then the previous lemma shows that $\mathcal{LRB} \models [\mathbf{u}_{(i)}(\mathcal{W})]^{(pq)} \approx \mathbf{w}_{pq}$ for any $p < q$, implying that the scheme \mathcal{W} comes from the word $\mathbf{u}_{(i)}(\mathcal{W})$. The case when B generates \mathcal{RRB} is dual, while if B generates \mathcal{RB} then \mathcal{W} comes from the word

$$\mathbf{w} = \mathbf{u}_{(i)}(\mathcal{W})\mathbf{u}_{(f)}(\mathcal{W}).$$

Finally, it remains to consider the case when B generates $\mathcal{LRB} \vee \mathcal{RZ}$ (or, dually, $\mathcal{RRB} \vee \mathcal{LZ}$). Then it can be argued that \mathcal{W} comes from the word $\mathbf{w} = \mathbf{u}_{(i)}(\mathcal{W})\mathbf{u}_{(f)}(\mathcal{W})$ (respectively, $\mathbf{w} = \mathbf{u}_{(h)}(\mathcal{W})\mathbf{u}_{(f)}(\mathcal{W})$). Namely, it is immediate that $i(\mathbf{w}^{(pq)}) = i([\mathbf{u}_{(i)}(\mathcal{W})]^{(pq)}) = i(\mathbf{w}_{pq})$; on the other hand, $t(\mathbf{w}^{(pq)}) = t(\mathbf{w}_{pq})$ is established by essentially repeating the arguments dual to those presented in Proposition 4.3, as \mathcal{RNB} is a subvariety of $\mathcal{LRB} \vee \mathcal{RZ}$ (so that the assumption that an identity holds in the latter variety implies the validity of the same identity in the former one). \square

Of course, the following problem naturally presents itself.

QUESTION 4.1. Are all finite bands finitely related?

Acknowledgement

The author is indebted to the anonymous referee for a number of useful comments improving the quality and precision of the exposition.

References

- [1] E. Aichinger, ‘Constantive Mal’cev clones on finite sets are finitely related’, *Proc. Amer. Math. Soc.* **138** (2010), 3501–3507.
- [2] E. Aichinger, P. Mayr and R. McKenzie, ‘On the number of finite algebraic structures’, Preprint, arXiv:1103.2265.
- [3] P. A. Biryukov, ‘Varieties of idempotent semigroups’, *Algebra i Logika* **9** (1970), 255–273, (in Russian).
- [4] R. Brignall, N. Georgiou and R. J. Waters, ‘Modular decomposition and the Reconstruction Conjecture’, Preprint, arXiv:1112.1509.
- [5] S. Burris and H. P. Sankappanavar, *A Course in Universal Algebra*, Graduate Texts in Mathematics, 78 (Springer, New York, 1981).
- [6] B. A. Davey, M. G. Jackson, J. G. Pitkethly and Cs. Szabó, ‘Finite degree: algebras in general and semigroups in particular’, *Semigroup Forum* **83** (2011), 89–110.
- [7] B. A. Davey and B. J. Knox, ‘From rectangular bands to k -primal algebras’, *Semigroup Forum* **64** (2002), 29–54.
- [8] C. F. Fennemore, ‘All varieties of bands I, II’, *Math. Nachr.* **48** (1971), 237–252, 253–262.
- [9] J. A. Gerhard, ‘The lattice of equational classes of idempotent semigroups’, *J. Algebra* **15** (1970), 195–224.
- [10] J. A. Gerhard and M. Petrich, ‘Varieties of bands revisited’, *Proc. Lond. Math. Soc.* (3) **58** (1989), 323–350.
- [11] P. J. Kelly, ‘A congruence theorem for trees’, *Pacific J. Math.* **7** (1957), 961–968.
- [12] P. Marković, M. Maróti and R. McKenzie, ‘Finitely related clones and algebras with cube terms’, *Order*, to appear.
- [13] P. Mayr, ‘On finitely related semigroups’, manuscript (November, 2011), 20 pp.
- [14] M. Petrich, ‘A construction and a classification of bands’, *Math. Nachr.* **48** (1971), 263–274.
- [15] S. M. Ulam, ‘A Collection of Mathematical Problems’, Interscience Tracts in Pure and Applied Mathematics, 8 (Interscience Publishers, New York–London, 1960).

IGOR DOLINKA, Department of Mathematics and Informatics,
University of Novi Sad, Trg Dositeja Obradovića 4, 21101 Novi Sad, Serbia
e-mail: dockie@dmf.uns.ac.rs