

FINITELY RAMIFIED GRAPH-DIRECTED FRACTALS, SPECTRAL ASYMPTOTICS AND THE MULTIDIMENSIONAL RENEWAL THEOREM

B. M. HAMBLY¹ AND S. O. G. NYBERG²

¹*Mathematical Institute, University of Oxford, 24–29 St Giles’,
Oxford OX1 3LB, UK (hambly@maths.ox.ac.uk)*

²*Computas AS, Vollsveien 9, PO Box 482, 1327 Lysaker,
Norway (son@computas.com)*

(Received 28 June 2000)

Abstract We consider the class of graph-directed constructions which are connected and have the property of finite ramification. By assuming the existence of a fixed point for a certain renormalization map, it is possible to construct a Laplace operator on fractals in this class via their Dirichlet forms. Our main aim is to consider the eigenvalues of the Laplace operator and provide a formula for the spectral dimension, the exponent determining the power-law scaling in the eigenvalue counting function, and establish generic constancy for the counting-function asymptotics. In order to do this we prove an extension of the multidimensional renewal theorem. As a result we show that it is possible for the eigenvalue counting function for fractals to require a logarithmic correction to the usual power-law growth.

Keywords: fractal; Dirichlet form; counting-function asymptotics; renewal theory

AMS 2000 *Mathematics subject classification:* Primary 35P20; 58J50
Secondary 28A80; 60K05; 31C25

1. Introduction

The study of the Laplace operator on fractal sets has focused primarily on fractals that are exactly self-similar and have the property of finite ramification. The class of post-critically finite self-similar sets [13, 16] captures most objects with these properties. In this framework there is a reasonable understanding of the Laplace operator, in particular its spectral properties [17] and heat kernel behaviour [12]. It is hoped that these sets will give good insight into more general fractals which may be used in the modelling of physical systems.

The class of graph-directed constructions was introduced in [25] and provides a much wider range of (slightly) more realistic models. Fractals of this type first appeared in [5] and have been called different names, including recurrent iterated function systems [3]. They are no longer exactly self-similar but do consist of self-similar components. An example of a finitely ramified graph-directed fractal, the type of set that we consider, is



Figure 1. The two generators of a graph-directed fractal.




Figure 2. The vector of fractals at stage 3.




Figure 3. The first two stages of construction of a fractal with infinitely ramified generator.

shown in Figure 1. The fractal should be thought of as a pair of fractals, one based on the square and the other on the triangle. The set is formed by recursive application of the maps which replace triangles and squares with the depicted combinations of triangles and squares as shown in Figure 1. The repeated iteration of the maps leads to a pair of fractals as shown in Figure 2. In §6 we will construct a class of fractals that can be decomposed into graph-directed constructions with the finite ramification property. Two examples are the fractal in Figure 3 and the diamond fractal of [18] shown in Figure 5. Note that the generator of the fractal in Figure 3 appears infinitely ramified at stage 1 as the central square is composed of four squares which meet along edges. However, at stage 2 we see that the fractal is a finitely ramified combination of squares and rectangles.

Analysis on fractals has concentrated largely on finitely ramified fractals (a fractal is finitely ramified if any subset of the fractal can be disconnected by removing only a finite number of points) as there are very natural graph approximations to these sets. The

construction of the Laplace operator on the fractal can be reduced to a finite-dimensional fixed-point problem for these graphs.

The framework of graph-directed constructions allows us to extend the class of fractals for which we can construct Laplace operators through quite simple extensions of the techniques developed in [13]. We will assume the existence of a solution to the canonical fixed-point problem and use this to construct a Dirichlet form on the fractal. For some discussion of this existence problem for a homogeneous random version of these fractals see [4]. This paper will then focus on the eigenvalue asymptotics of the associated Laplace operators. As these operators have compact resolvents they have discrete spectra and we can define the associated eigenvalue counting function $N(\lambda)$ as the number of eigenvalues less than λ . We will discuss the leading-order term in the asymptotic expansion of $N(\lambda)$.

There has been considerable interest in studying the behaviour of the eigenvalue counting function for various domains. In the case of a domain with fractal boundary, the Weyl–Berry conjecture has motivated much research. Weyl’s classical result for a bounded open subset $D \subset \mathbb{R}^n$ states that for either the Dirichlet or Neumann Laplacian,

$$\lim_{\lambda \rightarrow \infty} \frac{N(\lambda)}{\lambda^{n/2}} = \frac{B_n |D|}{(2\pi)^n},$$

where $|D|$ denotes the n -dimensional volume of the set D and B_n the volume of the unit ball in \mathbb{R}^n . It was conjectured by Berry that the second-order term would be determined by the Hausdorff dimension of the boundary of D . Subsequent research has shown that the Minkowski dimension of the boundary of D is the right notion of dimension but this does not give the precise behaviour of the second-order term in general (for a discussion see [21]).

In the case where the domain itself is fractal, a number of authors have established the behaviour of the leading term. This is typically different from the Euclidean case and the power which appears in the leading term has become known as the spectral dimension, d_s . In [2, 9–11, 15, 17] it has been shown that the generic situation is the existence of a constant $C > 0$, such that

$$\lim_{\lambda \rightarrow \infty} \frac{N(\lambda)}{\lambda^{d_s/2}} = C.$$

However, if the scale factors in the fractal are related in a certain way, then this limit may not exist and the ratio is asymptotically a periodic function of $\log \lambda$. In the setting of symmetric fractals such as the Sierpinski gasket an explanation for this oscillation is provided by the existence of strictly localized eigenfunctions [2].

We will show in this paper that similar results hold for finitely ramified graph-directed fractals but a wider variety of behaviour is possible. If the graph defining the construction is strongly connected, then the results are given in Theorem 5.4 and are very similar to the case of post-critically finite fractals discussed in [17] with generic constancy and otherwise a possibly periodic function. If the graph is not strongly connected, then there are other possibilities; in particular, the leading-order term in the asymptotic expansion of $N(\lambda)$ may grow faster than $\lambda^{d_s/2}$. Our main result is stated in Theorem 5.6 and shows

that one may have

$$\lim_{\lambda \rightarrow \infty} \frac{N(\lambda)}{(\log \lambda)^m \lambda^{d_s/2}} = C',$$

for a constant $C' > 0$ and an integer $m \geq 1$.

In [27] it was shown that such a logarithmic correction term would be required when studying the spectral asymptotics of the Laplacian on a set called plain mandala. We will see that this set can be considered within the graph-directed framework and we give a more precise limit result than that of [27] in §6.

The outline of the paper is as follows. Firstly, in §2, we state and prove a multidimensional renewal theorem which will be required for proving our main results. Section 3 describes the class of finitely ramified graph-directed fractals in which we work and also uses the renewal theorem to discuss the behaviour of a natural geometric counting function. In §4 we construct Laplace operators on these fractals via their Dirichlet forms. Using some properties of the Dirichlet forms we will derive a multidimensional renewal equation for the eigenvalue counting function and use our renewal theorem in §5. In §6 we discuss some examples, including plain mandala, and construct a class of fractals whose elements may be decomposed into finitely ramified graph-directed fractals. We also discuss some other examples of fractals in this class for which we can prove the existence of a Laplace operator.

2. The multidimensional renewal theorem

The renewal theorem is a powerful probabilistic result that has been used in proving fine results about the behaviour of self-similar sets. Examples include obtaining Hausdorff dimension and Minkowski measurability of fractals [7], studying eigenvalue problems for domains with fractal boundary [23], and studying such problems on fractals themselves [17]. A review of these ideas in the fractal context can be found in [8].

There have been several extensions of the renewal theorem to higher dimensions and we mention two that are related to the one we discuss here. Firstly, in the context of multitype age-dependent branching processes, Ryan [29] derived an extension which could be used to describe the asymptotic growth of the branching process. The second case arose in the study of self-similar measures, where Lau, Wang and Chu [22] obtained a version for separable, metrizable, locally compact, abelian groups, and used it to compute the L^p dimension of such a measure. We will give an extension of the results of [22] in the \mathbb{R}^n case which makes precise all the possible behaviours when the matrix of measures driving the system is reducible. An indication of the behaviour in this case was obtained in [6], when considering the time complexity of certain algorithms.

Firstly we need some notation. Let $M = [m_{ij}]$ be a matrix of Radon measures on \mathbb{R}_+ . We will write F for the matrix of distribution functions of M , that is $F_{ij}(t) = \int_0^t m_{ij}(ds)$ and we will write $F_{ij}(t, t+h) = F_{ij}(t+h) - F_{ij}(t)$. The indices of the matrix will be referred to as states and are the vertices of a graph G . The graph has a directed edge between states i and j if the measure m_{ij} is non-zero.

We denote the operation of convolution of a function a with a measure b by

$$b * a(t) = a * b(t) = \int_0^t a(t - s)b(ds);$$

if both a and b are measures, we convolve the distribution function of a with the measure b . For two matrices A, B of measures we write the ij th element of $C(t) = A * B(t)$ as $c_{ij}(t) = \sum_k a_{ik} * b_{kj}(t)$. Let $\gamma(i, j)$ denote a directed path in the graph G from vertex i to vertex j . We define the measure $m_{\gamma(i,j)}$ by taking the convolution of the measures associated with each given edge in the path. We will also write $m_{i\hat{i}}$ for the i th column of the matrix M with the i th element removed, similarly, $m_{\hat{i}i}$ for the i th row of M with the i th element removed. Finally, we write the matrix of measures with both the i th row and column of M removed as $M_{i\hat{i}}$.

We follow [22] and define the measure

$$\nu_1 = m_{11} + m_{1\hat{1}} * \sum_{k=0}^{\infty} (M_{11})^{*k} * m_{\hat{1}1}. \tag{2.1}$$

It is not difficult to check that, if $F(\infty)$ has maximum eigenvalue 1 and is irreducible, this is a probability measure with support given by $\cup\{\text{supp}(m_{\gamma}) : \gamma \text{ is a simple cycle in } G\}$. If the support is contained in a discrete subgroup of \mathbb{R} , we will call this measure lattice, otherwise we will call it non-lattice. By the irreducibility we see that if ν_1 is non-lattice, then ν_i is non-lattice for all i .

We begin by stating the renewal theorem for the case of irreducible $F(\infty)$. These results appeared as Theorems 4.2 and 4.3 of [22], where they are stated with one slightly stronger condition. The proofs given in [22] use only the assumptions stated here. Firstly, we give the analogue of Blackwell’s Renewal Theorem.

Theorem 2.1. *We assume that $F(t)$ is a matrix of measures in which $F(\infty)$ is irreducible, has maximum eigenvalue 1, $F_{ij}(0-) = 0$, $\int_0^{\infty} t dF_{ij}(t) < \infty$ for all i, j and for each j there is at least one i such that $F_{ij}(0) < F_{ij}(\infty)$. Let $V(t) = \sum_{k=0}^{\infty} F^{*k}(t)$ denote the matrix renewal measure, then if ν_1 is non-lattice,*

$$\lim_{t \rightarrow \infty} V(t, t + h] = Ah,$$

where

$$A = \frac{\mathbf{u}\mathbf{v}^T}{\mathbf{v}^T \mathcal{M} \mathbf{u}},$$

and \mathbf{u}, \mathbf{v} are the unique normalized right and left 1-eigenvectors of $F(\infty)$ and \mathcal{M} is the matrix of first moments of F .

If ν_1 is lattice, with period T , then

$$\lim_{t \rightarrow \infty} [V_{ij}(t + \tau_{ij}, t + \tau_{ij} + T)] = AT,$$

for any $\tau_{ij} \in \text{supp}(m_{\gamma(i,j)})$.

We also state a result concerning the asymptotic behaviour of the solution to the renewal equation.

Theorem 2.2. *Let $\mathbf{z}(t)$ be directly Riemann integrable, and let F be a matrix of measures satisfying the assumptions of Theorem 2.1, then the renewal equation*

$$\mathbf{r}(t) = \mathbf{z}(t) + \mathbf{r} * F(t) \quad (2.2)$$

has a unique solution, bounded on finite intervals. If ν_1 is non-lattice, then

$$\mathbf{r}(t) \rightarrow \int_0^\infty \mathbf{z}(t) dt A, \quad \text{as } t \rightarrow \infty.$$

If ν_1 is lattice with period T , then

$$\mathbf{r}(t) = \lim_{n \rightarrow \infty} [r_i(t + \tau_{1i} + nT)] = \sum_k \mathbf{z}(t + kT)A$$

exists almost surely for every $t \in [0, T]$.

Remark 2.3. Note that if $\text{supp } m_{\gamma(1,j)}$ has period T , then we can remove the vector τ_{1j} from the periodic case. However, this is not true in general, as can be seen from [22, Remark 3.4]. From this point, whenever we encounter a multidimensional periodic function arising as a limit from a renewal equation, we will assume that the initial times were chosen in the support of the underlying measure, as in the previous two theorems.

Remark 2.4. The state 1 is used as a reference state and any other would do.

Remark 2.5. The matrix A has strictly positive entries by the Perron–Frobenius Theorem.

We will concentrate on the case where $F(\infty)$ is reducible and recall some basic facts about non-negative reducible matrices. A state i has *access* to state j if there is a directed path from i to j . The states i and j are said to communicate if they have access to each other. Using the communication relation, we can partition the states into equivalence classes. The spectral radius of a class is the spectral radius of $F(\infty)$ restricted to that class. A class is called *basic* if its spectral radius is the same as that of the matrix itself. If a class is not basic, then it is called *non-basic*. A class J is *final* if J has access to no other class. A *chain* of classes is a collection of classes such that each class has access to or from another in the collection. The *length* of a chain is the number of basic classes that it contains. The *height* of a basic class C is the length of the longest chain of classes which have access to C . Let S_m denote the union of basic classes of height $m + 1$.

Theorem 2.6. *We assume that $F(t)$ is a matrix of measures in which $F(\infty)$ has maximum eigenvalue 1, with $F_{ij}(0-) = 0$, $\int_0^\infty t dF_{ij}(t) < \infty$ for all i, j and for each j there is at least one i such that $F_{ij}(0) < F_{ij}(\infty)$. Let \mathbf{z} be a vector with components that are directly Riemann integrable functions on \mathbb{R}_+ with $z_i \neq 0$ for all $i \in S_0$. If \mathbf{u} is continuous and satisfies the renewal equation (2.2), then $\mathbf{u} = \mathbf{z} * \sum_{k=0}^\infty F^{*k}$ and the components u_i satisfy:*

(1) if $i \in S_0$, then

$$\lim_{t \rightarrow \infty} (u_i(t) - p_i(t)) = 0,$$

where p_i is either periodic or constant depending on whether ν_i is lattice or not;

(2) if $i \in S_m$, for $m > 0$, then

$$\lim_{t \rightarrow \infty} t^{-m} u_i(t) = c_1(i),$$

for some constant $c_1(i) \geq 0$;

(3) if $i \notin S = \cup_m S_m$ and there is no path from S to i , then

$$\lim_{t \rightarrow \infty} u_i(t) = 0;$$

(4) if $i \notin S$ and there is a path from S_0 to i but no path from S_l for any $l > 0$, then

$$\lim_{t \rightarrow \infty} (u_i(t) - p'_i(t)) = 0,$$

for some p'_i which is either constant or periodic;

(5) if $i \notin S$ and there is a path from S_m to i but no path from S_l for any $l > m$, then

$$\lim_{t \rightarrow \infty} t^{-m} u_i(t) = c_2(i),$$

for some constant $c_2(i) \geq 0$.

If $i \in C_k$, an equivalence class in S_m , then if $z_j \neq 0$ for at least one $j \in C_k$, we have $c_1(i), c_2(i) > 0$.

Before we prove this result we need some preliminary lemmas which follow from the irreducible case. Using the above notation we will write $A^{(i)}$ for the limit matrix of Theorem 2.1 corresponding to a matrix renewal measure $V^{(i)}$.

Lemma 2.7. *Let $F(t)$ be a matrix of measures in which $F(\infty)$ has maximum eigenvalue 1, with $F_{ij}(0-) = 0$, $\int_0^\infty t dF_{ij}(t) < \infty$ for all i, j and for each j there is at least one i such that $F_{ij}(0) < F_{ij}(\infty)$ and F is irreducible. For $V = \sum_{k=0}^\infty F^{*k}$ and all $n \geq 0$ we have*

$$\lim_{t \rightarrow \infty} \left[\frac{1}{t} \int_0^t \left(\frac{t-u}{t} \right)^n dV_{ij}(u) \right] = \frac{1}{n+1} A.$$

Proof. We prove this result componentwise. From Theorem 2.1 we can deduce the $n = 0$ case, that is for $\epsilon > 0$ there exists a t_0 such that

$$|t^{-1} V_{ij}(0, t) - A_{ij}| < \epsilon \tag{2.3}$$

for $t > t_0$ for all i, j . For the general case, observe that we can write

$$\frac{1}{n+1} = t^{-n-1} \int_0^t u^n du = t^{-n-1} \int_0^t (t-u)^n du,$$

and hence, using an integration by parts, we have

$$\begin{aligned} \left| t^{-1} \int_0^t \left(\frac{t-u}{t} \right)^n dV_{ij}(u) - \frac{A_{ij}}{n+1} \right| &= \left| \frac{1}{t} \int_0^t \left(\frac{t-u}{t} \right)^n dV_{ij} - \frac{1}{t} \int_0^t \left(\frac{t-u}{t} \right)^n A_{ij} du \right| \\ &\leq \frac{1}{t^{n+1}} \int_0^t nu^{n-1} |V_{ij}(t-u) - (t-u)A_{ij}| du. \end{aligned}$$

Now use the $n = 0$ case, and the fact that the measure V_{ij} is bounded over finite intervals, to obtain

$$\begin{aligned} \left| t^{-1} \int_0^t \left(\frac{t-u}{t} \right)^n dV_{ij}(u) - \frac{1}{n+1} A_{ij} \right| &\leq \frac{\epsilon}{t^{n+1}} \int_0^{t-t_0} nu^{n-1}(t-u) du \\ &\quad + K \frac{1}{t^{n+1}} \int_{t-t_0}^t nu^{n-1}(t-u) du, \end{aligned}$$

for some constant K . If we integrate and choose a value t_n sufficiently large, we see that

$$\left| t^{-1} \int_0^t \left(\frac{t-u}{t} \right)^n dV_{ij}(u) - \frac{1}{n+1} A_{ij} \right| < 2\epsilon, \quad t > t_n,$$

and hence we have the result. □

Corollary 2.8. *If z converges to a constant \bar{z} , as $t \rightarrow \infty$, and F is as above, then*

$$\lim_{t \rightarrow \infty} \left[\frac{1}{t} \int_0^t \left(\frac{t-u}{t} \right)^n \sum_i z_i(t-u) dV_{ij}(u) \right] = \frac{1}{n+1} \bar{z}A.$$

It is also straightforward to show the following result for finite measures.

Lemma 2.9. *Let U be a matrix of finite measures with $U_{ij}(0-) = 0$, then for all $n \geq 0$ we have*

$$\lim_{t \rightarrow \infty} \left[\int_0^t \left(\frac{t-u}{t} \right)^n dU_{ij}(u) \right] = U(\infty).$$

If z converges to a constant \bar{z} , as $t \rightarrow \infty$, then

$$\lim_{t \rightarrow \infty} \left[\int_0^t \left(\frac{t-u}{t} \right)^n \sum_i z_i(t-u) dU_{ij}(u) \right] = \bar{z}U(\infty).$$

A function z will be called asymptotically periodic if there exists a function \bar{z} , satisfying $\bar{z}(t) = \bar{z}(t+kT)$ for all $t > 0, k \in \mathbb{N}$, and for $\epsilon > 0$, there exists a t' such that

$$|z(t) - \bar{z}(t)| < \epsilon, \quad \forall t > t'.$$

Lemma 2.10. *Let z be an asymptotically periodic function with period T and let V be a matrix renewal measure, then*

$$\lim_{t \rightarrow \infty} \frac{1}{t} z * V(t) = \frac{1}{T} \int_0^T \bar{z}(s) dsA.$$

Proof. Let \bar{z} be the periodic function such that for $\epsilon > 0$ there exists a t_0 such that $|\bar{z} - z| < \epsilon$ for $t > t_0$. We first show that $(1/t)(z * V - \bar{z} * V) \rightarrow 0$ as $t \rightarrow \infty$. We prove this componentwise as

$$\begin{aligned} \left| \sum_j \int_0^t (z_j(t-u) - \bar{z}_j(t-u)) dV_{ji}(u) \right| &\leq \sum_j \int_0^t |z_j(t-u) - \bar{z}_j(t-u)| dV_{ji}(u) \\ &\leq \sum_j (\epsilon V_{ji}(t-t_0) + K V_{ji}(t_0)), \end{aligned}$$

where $K = \sup_{0 \leq u \leq t_0, i} |z_i(u) - \bar{z}_i(u)|$. Thus, dividing by t and using the linear growth of V from (2.3), we have the convergence.

As \bar{z} is a bounded function and V is uniformly bounded over finite intervals and converges to a multiple of Lebesgue measure, we have

$$\int_{lT}^{(l+1)T} \sum_j \bar{z}_j(s+u) dV_{ji}(u) \rightarrow \int_0^T \sum_j \bar{z}_j(u) A_{ji} du, \quad \forall 0 \leq s \leq T.$$

Now define

$$a_l(t) = \frac{1}{T} \int_{lT}^{(l+1)T} \sum_j \bar{z}_j(t-u) dV_{ji}(u),$$

and observe that, by the above, for all l we have

$$a_l(t) \rightarrow \frac{1}{T} \int_0^T \sum_j \bar{z}_j(u) A_{ji} du$$

uniformly as $t \rightarrow \infty$. Finally, we can write

$$\begin{aligned} \frac{1}{t} \bar{z} * V(t) &= \frac{T}{t} \sum_{l=0}^{\lfloor t/T \rfloor - 1} \frac{1}{T} \int_{lT}^{(l+1)T} \sum_j \bar{z}_j(t-u) dV_{ji}(u) + \frac{1}{t} \int_{\lfloor t/T \rfloor T}^t \sum_j \bar{z}_j(t-u) dV_{ji}(u) \\ &\leq \frac{T}{t} \sum_{l=0}^{\lfloor t/T \rfloor - 1} a_l(t) + \frac{R_T}{t}, \end{aligned}$$

where R_T is a finite constant. As the terms in the sum converge uniformly to the claimed result, we are done. \square

Write U for a matrix of finite measures.

Lemma 2.11. *If $V^{(i)}$ is as in Lemma 2.7, z is directly Riemann integrable, and $U^{i,i+1}$ are finite matrix measures, then, for $m > 1$,*

$$\begin{aligned} \lim_{t \rightarrow \infty} t^{-m+1} z * V^{(1)} * U^{1,2} * V^{(2)} * \dots * U^{m-1,m} * V^{(m)}(t) \\ = \frac{1}{(m-1)!} \int_0^\infty z * V^{(1)} * U^{1,2}(u) du A^{(2)} U^{2,3}(\infty) \dots U^{m-1,m}(\infty) A^{(m)}. \end{aligned}$$

Proof. Let

$$I^{(m)}(t) = \mathbf{z} * V^{(1)} * U^{1,2} * \dots * V^{(m)}(t).$$

The result for $m = 1$ is the original solution to the renewal equation given in Theorem 2.2. We can write this, distinguishing between the lattice and non-lattice cases, as

$$\lim_{t \rightarrow \infty} (\mathbf{z} * V^{(1)}(t) - p(t)A^{(1)}) = 0, \tag{2.4}$$

where $p(t) = \sum_k \mathbf{z}(t + kT)$ is a periodic function (it may be constant) with period T . We now use induction and begin by proving the result for $m = 2$. First we observe that, as $U^{1,2}$ is a finite measure, $\mathbf{z} * V^{(1)} * U^{1,2}(t)$ is an asymptotically periodic function and we call the associated periodic function \mathbf{h} . Thus we can apply Lemma 2.10 to deduce that

$$\frac{1}{t} \mathbf{z} * V^{(1)} * U^{1,2} * V^{(2)}(t) \rightarrow \frac{1}{T} \int_0^T \mathbf{h}(u) \, du \, A^{(2)}.$$

This constant can then be written in terms of the asymptotically periodic part as

$$\mathbf{h}(u) = \lim_{n \rightarrow \infty} \mathbf{z} * V^{(1)} * U^{1,2}(u + nT) = T \sum_k \mathbf{z} * V^{(1)} * U^{1,2}(u + kT),$$

and hence

$$\frac{1}{T} \int_0^T \mathbf{h}(u) \, du = \int_0^\infty \mathbf{z} * V^{(1)} * U^{1,2}(u) \, du.$$

Now assume the result for $m - 1$ and let the limit vector be $C^{(m-1)}$. We can write

$$I^{(m)}(t) = I^{(m-1)} * U^{m-1,m} * V^{(m)}(t),$$

and hence, in component form,

$$t^{-m+1} I_i^{(m)}(t) = t^{-m+1} \int_0^t \sum_j h_j(t-u) \, dV_{ji}^{(m)}(u), \tag{2.5}$$

where

$$h_j(t) = \int_0^t (t-u)^{(m-2)} \sum_k \frac{I_k^{(m-1)}(t-u)}{(t-u)^{m-2}} \, dU_{kj}^{m-1,m}(u).$$

By our inductive hypothesis and Lemma 2.9, we have

$$t^{-m+2} h_j(t) \rightarrow \sum_i C_i^{(m-1)} U_{ij}^{m-1,m}(\infty),$$

as $t \rightarrow \infty$. Thus we can rewrite (2.5) as

$$t^{-m+1} I_i^{(m)}(t) = t^{-1} \int_0^t \left(\frac{t-u}{t} \right)^{m-2} \sum_j \frac{h_j(t-u)}{(t-u)^{m-2}} \, dV_{ji}^{(m)}(u),$$

and apply Corollary 2.8 to show

$$C^{(m)} = \lim_{t \rightarrow \infty} t^{-m+1} I^{(m)}(t) = \frac{1}{m-1} C^{(m-1)} U^{m-1,m}(\infty) A^{(m)}.$$

Iterating this expression gives the result. □

Proof of Theorem 2.6. This result is an extension of that of [22]. The first part concerning the solution of the renewal equation and Parts (1), (3) and (4) of the limiting result are contained in [22, Theorem 4.5]. We only need to consider Part (2), as Part (5) will follow immediately we have Part (2).

We will write the matrix of measures in canonical form and note that if there are k classes, the renewal equation can be written as

$$[\mathbf{u}_1, \dots, \mathbf{u}_k] = [\mathbf{z}_1, \dots, \mathbf{z}_k] + F * [\mathbf{u}_1, \dots, \mathbf{u}_k],$$

where $\mathbf{u}_i, \mathbf{z}_i$ are the \mathbf{u}, \mathbf{z} vectors restricted to the states in class i . For a non-basic class i , we know that $V^{(i)}$ is a finite measure.

Consider a basic class of height m . For the chain above this class we index the basic classes by i_1, \dots, i_m and can write down a set of renewal equations

$$\begin{aligned} \mathbf{u}_{i_m} &= \mathbf{z}_{i_m} + \mathbf{u}_{i_m} * F_{i_m i_m}, \\ \mathbf{u}_{i_{m-1}} &= \mathbf{z}_{i_{m-1}} + \mathbf{u}_{i_m} * U^{i_m, i_{m-1}} + \mathbf{u}_{i_{m-1}} * F_{i_{m-1} i_{m-1}}, \\ &\vdots \\ \mathbf{u}_{i_1} &= \mathbf{z}_{i_1} + \mathbf{u}_{i_m} * U^{i_m, i_1} + \dots + \mathbf{u}_{i_2} * U^{i_2, i_1} + \mathbf{u}_{i_1} * F_{i_1 i_1}, \end{aligned}$$

where we write $U^{i_j, i_{j+1}}$ for the finite measure which may arise from $F_{i_j i_{j+1}}$ if both i_j and i_{j+1} are basic, or otherwise it is the finite measure obtained from the combination of the measures arising from intermediate non-basic classes in the chain.

We can recursively solve these separate multidimensional renewal equations. If we write $V^{(i_j)} = \sum_{k=0}^\infty F_{i_j i_j}^{*k}$, then

$$\begin{aligned} \mathbf{u}_{i_1} &= (\mathbf{z}_{i_1} + \mathbf{z}_{i_2} * V^{(i_2)}) * U^{i_2, i_1} + \dots + \mathbf{z}_{i_m} * V^{(i_m)} * U^{i_m, i_{m-1}} * V^{(i_{m-1})} * \dots \\ &\quad \dots * U^{i_3, i_2} * V^{(i_2)} * U^{i_2, i_1}) * V^{(i_1)}. \end{aligned}$$

It is clear that each term except the first is an increasing function of t and the only term that will appear in the limit is the last one. As the matrices $U^{i_j, i_{j-1}}$ are finite measures, we can apply Lemma 2.11 to show that there is a constant vector \mathbf{c} such that

$$t^{-m+1} \mathbf{z}_{i_m} * V^{(i_m)} * U^{i_m, i_{m-1}} * V^{(i_{m-1})} * \dots * U^{i_3, i_2} * V^{(i_2)} * U^{i_2, i_1} * V^{(i_1)} \rightarrow \mathbf{c}, \quad \text{as } t \rightarrow \infty.$$

Finally, we must show that $c_1(i) > 0$ if $z_j \neq 0$ for some $j \in C_k(i) \subset S_m$. This follows from the explicit form of the constant and the fact that the classes in the chain have access to one another, ensuring that the matrix $A^{(i_{m-1})} U^{i_{m-1}, i_{m-2}}(\infty) \dots U^{i_2, i_1}(\infty) A^{(i_1)}$ must be positive by Remark 2.5 after Theorem 2.2. \square

In order to apply this result to the renewal equations which occur later, we state the following corollary, a multidimensional version of that found in [16, Appendix B.4].

Corollary 2.12. *Let $t^* > 0$. Let $\mathbf{r}(t)$ be a vector whose components are measurable functions on \mathbb{R} with $r_i(t) = 0, \forall i$, for $t < t^*$, and let $\mathbf{z}(t)$ be a non-negative directly*

Riemann integrable function with $z_i(t) = 0, \forall i$, for $t < t^*$. Assume that \mathbf{r} satisfies the renewal equation

$$\mathbf{r}(t) = \mathbf{z}(t) + (\mathbf{r} * F)(t), \quad t \in \mathbb{R}.$$

The conclusions of Theorem 2.6 then hold.

3. Graph-directed constructions

In [25], graph-directed constructions were defined, extending the class of deterministic fractals. Since then this class has been studied in some detail: see, for example, [28], where fine properties of random recursive graph-directed measures were obtained. We consider here the deterministic case and consider a class of such fractals with the property of finite ramification. Note that we will work in \mathbb{R}^n but it would be possible to define an analogue of post-critically finite (p.c.f.) self-similar sets, which are abstract metric spaces, and indexed, in this setting, by a subshift of finite type.

Let $J = \{J_j\}_{j \in \mathcal{S}}$ be a set of compact and connected subsets of \mathbb{R}^n , indexed by a finite set \mathcal{S} . Let \mathcal{S} be the vertex set for a directed graph $G = (\mathcal{S}, E)$, where E is the set of edges for this graph. Note that, as we will allow multiple edges, the pair (i, j) does not uniquely define the edge (i, j) and we write E_{ij} for the set of edges from i to j . The k th edge in E_{ij} may be denoted $e_k(i, j)$. For each $i \in \mathcal{S}$, let $E_i = \cup_{j \in \mathcal{S}} E_{ij}$. For each edge $e \in E$, let $\phi_e : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be an l_e^{-1} -similitude.

In order to define a graph-directed construction we make the following assumption.

Assumption 3.1.

- (1) For each $i \in \mathcal{S}$, $|E_i| \geq 1$.
- (2) For each $i \in \mathcal{S}$,

$$\bigcup_{j \in \mathcal{S}} \bigcup_{e \in E_{ij}} \phi_e(J_j) \subset J_i.$$

- (3) If a path in the graph G is a cycle $(e_{k_1}(i_1, i_2), \dots, e_{k_{n-1}}(i_{n-1}, i_n))$ with $i_n = i_1$, then

$$\prod_{j=1}^{n-1} l_{e_{k_j}(i_j, i_{j+1})} > 1.$$

An object K constructed from this assumption is called a geometric graph-directed construction, and G is called its *construction graph*. We recall that a graph G is *connected* if for each pair of vertices $i, j \in G$ there is a (non-directed) path between them. We will say that the graph G is *strongly connected* if for each pair of vertices $i, j \in G$ there is a directed path from i to j . A strongly connected component of G is a maximal subgraph H of G such that H is strongly connected. Strongly connected components are pairwise disjoint and do not necessarily cover G . A single vertex may be a strongly connected component if it loops to itself. For our construction graphs, there is at least one

strongly connected component. We construct the fractal as the unique vector of compact sets $(K_1, \dots, K_{|\mathcal{S}|})$ which satisfies

$$K_i = \bigcup_{j \in \mathcal{S}} \bigcup_{e \in E_{ij}} \phi_e(K_j).$$

This result is given as Theorem 1 of [25] and is proved by demonstrating that the map $\psi = \bigcup_{e \in E} \phi_e$ is a contraction on the set of compact subsets of J in the Hausdorff metric.

Let E^n be the set of length- n paths in the graph G . Note that E^n is not the full set of n -sequences of elements of E , but consists of the allowable sequences $\sigma = (e_{k_0}(i_0, i_1), \dots, e_{k_{n-1}}(i_{n-1}, i_n))$ for which $e_{k_j}(i_j, i_{j+1}) \in E$ for $j = 0, \dots, n - 1$. We will write $\sigma(n) = i_n$ for the n th vertex in the path σ where the root of the path is $\sigma(0) = i_0$. Let $E_i^n = \{\sigma \in E^n : \sigma(0) = i\}$. We further define $E^* = \bigcup_{n \in \mathbb{N}} E^n$. Note that E^∞ is a Markov subshift of the full shift space generated by \mathcal{S} . For $\sigma \in E^n$, we write $\phi_\sigma = \phi_{(e_1, \dots, e_n)} = \phi_{e_1} \circ \dots \circ \phi_{e_n}$. For $\sigma \in E_i^n$, we define $K_\sigma = \phi_\sigma(K_{\sigma(n)})$, and call such objects the n -cells of component $\sigma(0)$. The $J_i, i \in \mathcal{S}$, are 0-cells. In this way we can define a canonical projection map from the paths to the fractal $\pi : E^\infty \rightarrow K$, which specifies points in component i as the limit of paths in the graph starting from node i .

In order to determine the Hausdorff dimension of the set, we recall the open-set condition, which ensures that the overlap between sets is sufficiently small.

Definition 3.2. A graph-directed construction $K = \{J_i\}_{i \in \mathcal{S}}$ satisfies the *open-set condition* (OSC) if for all $i \in \mathcal{S}$ there are non-empty open sets $\mathcal{O}_i \subset \mathbb{R}^n$, with $\mathcal{O}_i \subset J_i$, such that $\mathcal{O}_i \supset \bigcup_{e \in E_i} \phi_e(\mathcal{O}_j)$ and $\phi_{e_k(i,j)}(\mathcal{O}_j) \cap \phi_{e_l(i,m)}(\mathcal{O}_m) = \emptyset$ for all $i, j, m \in \mathcal{S}$, $e_k(i, j) \in E_{ij}$, $e_l(i, m) \in E_{im}$.

It is said to satisfy the strong open-set condition if $\mathcal{O}_i \cap K_i \neq \emptyset$ for all $i \in \mathcal{S}$.

It has been shown in [31] that if the graph is not strongly connected, then the OSC may not be equivalent to the strong open-set condition.

The Hausdorff dimension of the fractals K_i was calculated in [25] and we give a brief discussion. Let $l_{\min} = \min_{e \in E} l_e$ and $l_{\max} = \max_{e \in E} l_e$. The length-scale factor can be defined for paths in the graph by setting

$$l_\sigma = \prod_{i=1}^n l_{e_i}$$

for permissible $\sigma = (e_1, \dots, e_n)$. Let \mathbf{H}^s be the non-negative $|\mathcal{S}| \times |\mathcal{S}|$ -matrix whose entries are $h_{ij}^s = \sum_{e \in E_{ij}} l_e^{-s}$. If $s < t$, then $(l_{\min})^{(t-s)} \mathbf{H}^t \geq \mathbf{H}^s \geq (l_{\max})^{(t-s)} \mathbf{H}^t$ elementwise. Letting $\Phi(s)$ be the principal eigenvalue of the matrix \mathbf{H}^s , it is then easily seen by the above that Φ is strictly monotone decreasing and continuous in s . It is also easy to see that the right candidate for a self-similarity dimension would be the t such that $\Phi(t) = 1$. For, if we take any positive vector \mathbf{x} , $\lim_{n \rightarrow \infty} (\mathbf{H}^t)^n \mathbf{x} = c\mathbf{y}$, where \mathbf{y} is \mathbf{H}^t 's principal (right)

eigenvector and c is a constant. If, however, $s \neq t$, this limit would be either 0 or infinity. These heuristics can be formalized to give the following result, which is [25, Theorem 3]. We write $\dim_{H,d}(F)$ for the Hausdorff dimension of a set F with respect to a metric d , and $\mathcal{H}^\alpha(F)$ for the α -dimensional Hausdorff measure of F . If no metric is specified, we mean the Euclidean metric.

Theorem 3.3. *If K satisfies the OSC and if G is strongly connected, then the common Hausdorff dimension of the K_i is given by $\dim_H(K_i) = \alpha = \Phi^{-1}(1)$ and $0 < \mathcal{H}^\alpha(K_i) < \infty$ for all $i \in \mathcal{S}$.*

In [25] the case when the graph G is not strongly connected is also considered. In this situation, if we write $\text{SC}(G)$ for the set of strongly connected components of the graph G , then the Hausdorff dimension is given by

$$\dim_H(K) = \max\{\alpha_F : F \in \text{SC}(G)\}.$$

The Hausdorff measure is finite if and only if $\{F \in \text{SC}(G) : \alpha = \alpha_F\}$ consists of pairwise incomparable elements.

We will be interested in a subset of these fractals consisting of those that are finitely ramified in that the cells only meet at finitely many points. Following [16] we can give an analogue of the definition of p.c.f. self-similar sets in our setting. Let $\mathcal{C} = \cup_{i,j,k \in \mathcal{S}} \cup_{e \in E_{ij}, e' \in E_{ik}} (\phi_e(K_j) \cap \phi_{e'}(K_k))$ be the intersection points between cells. We let the critical set be $\mathcal{C}_\pi = \{\sigma \in E^* : \pi(\sigma) \in \mathcal{C}\}$ and the post-critical set be $\mathcal{P}_\pi = \cup_{n \geq 1} \cup_{\sigma \in \mathcal{C}_\pi} \xi_n(\sigma)$, where ξ is the shift map. Then $F_i^{(0)} = \{\pi(\sigma) : \sigma(0) = i, \sigma \in \mathcal{P}_\pi\}$ is the set of ramification points for the fractal.

We now make an assumption that guarantees that our fractals are connected and finitely ramified.

Assumption 3.4.

- (1) *For each K_i , for each pair of 1-cells A, B , there exists a sequence of 1-cells $\{C_i, i = 0, \dots, n\}$ such that $C_0 = A, C_n = B$ and $C_i \cap C_{i+1} \neq \emptyset$ for $i = 1, \dots, n-1$.*
- (2) *The $F_i^{(0)}$ have finite non-zero cardinality for all $i \in \mathcal{S}$.*

As in [16, Proposition 1.3.5] we can show that $K_a \cap K_b = \phi_a(F_{a(n)}^{(0)}) \cap \phi_b(F_{b(n)}^{(0)})$ for any $a \neq b \in E_n^i$ for all $i \in \mathcal{S}$. By using these ramification points we can define a sequence of graph approximations to the fractal. For $n \geq 1$, we define $F_i^{(n)} = \cup_{\sigma \in E_i^n} \phi_\sigma(F_{\sigma(n)}^{(0)})$. Note that $\{F_i^{(n)}\}_{n \in \mathbb{N}}$ are increasing sequences. Let $F^{(n)} = \cup_{i \in \mathcal{S}} F_i^{(n)}$ and $F^{(\infty)} = F^{(0)} \cup \cup_{\sigma \in E^*} \phi_\sigma(F^{(0)})$.

Definition 3.5. A set K satisfying the above assumptions will be called a *finitely ramified graph-directed fractal set*. We will abbreviate this to *FRGD (fractal) set*. A component K_i of the vector K will be called an *FRGD fractal*.

An FRGD fractal K_i can be a member of many FRGD sets K . We define a map h on subsets \mathcal{S}' of \mathcal{S} by setting $h(\mathcal{S}') = \mathcal{S}' \cup \{i \in \mathcal{S} : \exists e_k(j, i) \in E, \text{ some } j \in \mathcal{S}'\}$. Let $\text{frgd}(i)$

be the smallest set containing i which is closed under h , and let $\text{FRGD}(K_i) = \{K_j : j \in \text{frgd}(i)\}$. If G is strongly connected, we have $\text{FRGD}(K_i) = K, \forall K_i \in K$. From the finite ramification it is easy to see that an FRGD fractal will satisfy the OSC, and hence the dimension results of Theorem 3.3 hold for these fractals.

We conclude this section by discussing a geometric counting function for an FRGD fractal, using the multidimensional renewal theorem. Let $\mathcal{N}(r)$ denote the vector of numbers of sets of radius r required to cover the FRGD fractal K . We can then state a result concerning the limiting behaviour of this function. We begin by defining a suitable matrix renewal measure. Let

$$m_{ij}(\mathrm{d}x) = \sum_{e \in E_{ij}} l_e^{-\alpha} \delta_{\log l_e}(\mathrm{d}x), \tag{3.1}$$

where δ_x is a Dirac measure at x and, as in (2.1), we can define the measure ν_i using this matrix (for simplicity we assume that it is non-lattice for all i). We note that, by choice of α , $F(\infty)$ has maximum eigenvalue 1.

Theorem 3.6. *For an FRGD fractal K , satisfying the OSC, if the measure ν_i is non-lattice for all i , then*

- (1) if $i \in S_m$, for some $m \geq 0$, there exist constants $c_3(i) > 0$ such that

$$\lim_{r \rightarrow 0} \frac{\mathcal{N}_i(r)}{r^{-\alpha} |\log r|^m} = c_3(i);$$

- (2) if $i \notin S$, and there is no path from S to i ,

$$\lim_{r \rightarrow 0} \frac{\mathcal{N}_i(r)}{r^{-\alpha}} = 0;$$

- (3) if $i \notin S$ and there is a path from S_m to i and no path from S_l to i for $l > m$, there exist constants $c_4(i) > 0$ such that

$$\lim_{r \rightarrow 0} \frac{\mathcal{N}_i(r)}{r^{-\alpha} |\log r|^m} = c_4(i).$$

Proof. This is a straightforward application of our renewal theorem using the ideas that can be found in [8, Proposition 7.4]. We decompose the counting function according to the first stage of construction of the set to see that

$$\mathcal{N}_i(r) = \sum_{j \in S} \sum_{e \in E_{ij}} \mathcal{N}_j(r l_e) - a_i(r),$$

where $a_i(r)$ is the remainder term denoting those sets which have been multiply covered. A simple change of variable, setting $r = e^{-t}$, and multiplying through by $e^{\alpha t}$, shows that if we define $\mathcal{N}'_i(t) = e^{\alpha t} \mathcal{N}_i(e^{-t})$ and $a'_i(t) = -e^{\alpha t} a_i(e^{-t})$, then

$$\mathcal{N}'_i(t) = a'_i(t) + \sum_{j \in S} \sum_{e \in E_{ij}} l_e^{-\alpha} \mathcal{N}'_j(t - \log l_e), \quad t \in \mathbb{R}.$$

Using a truncation argument, as in [8], we can transform this into a multidimensional renewal equation with respect to the matrix renewal measure $M = [m_{ij}]$ as defined in (3.1), and with a' directly Riemann integrable. This is an equation of the form considered in § 2 and hence, by suitably shifting the time axis, we can apply Corollary 2.12 to get the results. The constants are strictly positive as the function $a_i(r)$ cannot be 0 for all r . \square

4. Dirichlet forms for FRGD fractals

We will now construct Dirichlet forms on our FRGD fractals. We require a measure and begin by considering any Bernoulli measure on the fractal K . Later we will introduce a ‘natural’ measure on the fractal. We will follow the approach of [13], an essential ingredient of which is to make an assumption about the existence of a suitable harmonic structure, a fixed point of a renormalization map, on the fractal. Note that we will not restrict ourselves to the analogue of Laplace operators derived from a regular harmonic structure.

For each $i \in \mathcal{S}$, let a probability measure μ'_e on E_i be given, with $\mu'_e > 0$ for all $e \in E_i$. This induces a probability measure μ_i on K_i , for $i \in \mathcal{S}$, where, for $\sigma = (e_1, \dots, e_n) \in E_i^n$,

$$\mu_i(K_\sigma) = \mu'_\sigma = \prod_{k=1}^n \mu'_{e_k}. \quad (4.1)$$

Definition 4.1. For a set V , let $\ell(V) = \{f \mid V \rightarrow \mathbb{R}\}$. For V a finite set, define a conductance matrix H to be a symmetric linear map with $H_{ij} \geq 0$, $H_{jj} = -\sum_k H_{jk}$. For $u, v \in \ell(V)$, define a symmetric bilinear form $\mathcal{E}_H(u, v) = -u^T H v$. (V, H) is called a *resistance network* and \mathcal{E}_H is an irreducible Dirichlet form if $\mathcal{E}_H(u, u) \geq 0$ for all $u \in \ell(V)$ with equality if and only if u is constant.

By standard convention, we write $\mathcal{E}(u)$ for $\mathcal{E}(u, u)$ whenever \mathcal{E} is a Dirichlet form. The Dirichlet form can as usual be recovered from knowledge of $\mathcal{E}(u)$ by the polarization identity $\mathcal{E}(u, v) = \frac{1}{4}(\mathcal{E}(u+v) - \mathcal{E}(u-v))$.

Definition 4.2. If (V, H_1) and (U, H_2) are two resistance networks, $U \subset V$, and

$$\mathcal{E}_{H_2}(u) = \min\{\mathcal{E}_{H_1}(v) : v \in \ell(V), v|_U = u\}, \quad (4.2)$$

we say they are *compatible*, and write $(V, H_1) \geq (U, H_2)$.

Remark 4.3. If $U \subset V$ and (V, H_1) is a known resistance network, then (4.2) uniquely defines a resistance network (U, H_2) , called the trace of (V, H_1) on U .

Lemma 4.4. Let (V, H_1) , (U, H_2) be connected resistance networks, $(V, H_1) \geq (U, H_2)$. Then, for each $u \in \ell(U)$, there is a unique function $v \in \ell(V)$ such that $\mathcal{E}_{H_2}(u, u) = \mathcal{E}_{H_1}(v, v)$.

For the proof of this lemma and a discussion of resistance networks in general, see [14].

Remark 4.5. Note that in the above, we can regard $\ell(U)$ as a subset of $\ell(V)$. Further, we see that (4.2) holds when $\ell(U) \subset \ell(V)$, and that it can be used to define a Dirichlet form H_2 on U when the Dirichlet form for (U, H_1) is given.

As an example of this, we can determine the effective resistance between two sets $X, Y \subset V$, given a resistance network (V, H) with Dirichlet form \mathcal{E}_H , as

$$R_H(X, Y) = (\min\{\mathcal{E}_H(u) : u \in \ell(V), u|_X = 1, u|_Y = 0\})^{-1}.$$

In order to construct harmonic structures on the fractal, as in [13], we need to find a discrete Laplace operator which is the fixed point of a renormalization map. For each $i \in \mathcal{S}$ we define the resistivity of a cell K_e in K_i by a choice of a positive finite weight $r_e, e \in E_i$. Let $\mathbf{D} = \{D_i\}_{i \in \mathcal{S}}$, where $(F_i^{(0)}, D_i)$ is a resistance network. Letting $\mathcal{E}_{D_i}^{(0)}$ be the associated Dirichlet forms, we define the Dirichlet forms on $F_i^{(1)}$ as

$$\mathcal{E}_{(r, D_i)}^{(1)}(u) = \sum_{j \in \mathcal{S}} \sum_{e \in E_{ij}} r_e^{-1} \cdot \mathcal{E}_{D_j}^{(0)}(u \circ \phi_e). \tag{4.3}$$

Equation (4.3) defines a coupling mapping \mathcal{T}' , parametrized by r , from the cone of Dirichlet forms on $F^{(0)}$ to the cone of Dirichlet forms on $F^{(1)}$. If we apply the trace map (4.2) to $\mathcal{E}_{(r, D_i)}^{(1)}$, with $H_2 = F^{(0)}$, this gives us a mapping \mathcal{T}'' from the cone of Dirichlet forms on $F^{(1)}$ to the cone of Dirichlet forms on $F^{(0)}$.

Definition 4.6. The renormalization map is defined to be the composition of the two maps $\mathcal{T} = \mathcal{T}'' \circ \mathcal{T}'$, parametrized by r , which takes the cone of Dirichlet forms on $F^{(0)}$ to the cone of Dirichlet forms on $F^{(0)}$.

Assumption 4.7. In this paper, we will assume that the map \mathcal{T} for the FRGD sets with a given fixed set of r , has an eigenform (non-degenerate projective fixed-point) $\mathbf{D} = \{D_i\}_{i \in \mathcal{S}}$, i.e. that there exists a $\lambda > 0$ such that $\mathcal{T}(\mathbf{D}) = \lambda \mathbf{D}$.

From now on we will assume that we have fixed \mathbf{D} and r . In the final section we will provide some non-trivial examples of FRGD fractals for which a fixed point exists for some r . We define the conductivity of a cell K_e in K_i by $\rho_e = \lambda/r_e$ for $e \in E_i$.

We will let $\mathcal{E}_i^{(0)}$, $i \in \mathcal{S}$, be Dirichlet forms on the respective $F_i^{(0)}$ corresponding to this fixed point \mathbf{D} . For $\sigma \in E^n$, let $\rho_\sigma = \prod_{i=1}^n \rho_{e_i}$. For $n \in \mathbb{N}$, define

$$\mathcal{E}_i^{(n)}(u) = \sum_{\sigma \in E_i^n} \rho_\sigma \mathcal{E}_{\sigma(n)}^{(0)}(u|_{\phi_\sigma(F_{\sigma(n)}^{(0)})} \circ \phi_\sigma),$$

and let $H_i^{(n)}$ be the matrices defined by

$$\mathcal{E}_i^{(n)}(f, g) = -f^T H_i^{(n)} g.$$

As is easily seen, $(F^{(n)}, H_i^{(n)}) \leq (F^{(m)}, H_i^{(m)})$ whenever $n \leq m$. This ensures that the sequence of Dirichlet forms is monotone increasing and allows us to construct the limit form.

If the harmonic structure is regular, in that $\rho_e > 1, \forall e \in E$, the effective resistance becomes a metric and the limit form is a local regular Dirichlet form on $L^2(K, \mu)$ for any Borel measure μ . If we do not have regularity of the harmonic structure, we need a further condition. Let $\theta_\sigma = \rho_\sigma/\mu'_\sigma$. We require the following assumption on our FRGD fractals.

Assumption 4.8. $\theta_e > 1, \forall e \in E$.

Remark 4.9. In the light of the Einstein relation (see [1]), we see that θ_σ is the time scaling factor for the approximating random walk on the cell K_σ .

For $m \leq n$, we define the harmonic extension $P_{m,n,i} : \ell(F_i^{(m)}) \rightarrow \ell(F_i^{(n)})$ by setting $P_{m,n,i}(u)$, for $u \in \ell(F_i^{(m)})$, to be $v \in \ell(F_i^{(n)})$ such that $\mathcal{E}_i^{(m)}(u, u) = \mathcal{E}_i^{(n)}(v, v)$. Let $P_{m,*,i} : \ell(F_i^{(m)}) \rightarrow \ell(F_i^{(\infty)})$ be defined by $P_{m,*,i}u(x) = P_{m,n,i}u(x)$ for some $n \in \mathbb{N}$ such that $x \in F_i^{(n)}$. Let $P_{m,i}u$ be the continuous extension of $P_{m,*,i}u$.

As in [13], a function which is in the image of $P_{m,n,i}$ or $P_{m,*,i}$ or $P_{m,i}$ is said to be m -harmonic on K_i .

We define the bilinear form $(\mathcal{E}_i, \mathcal{F}_i)$ by

$$\mathcal{F}_i = \{u \in L^2(K_i, \mu_i) : \sup_n \mathcal{E}_i^{(n)}(u) < \infty\}$$

and

$$\mathcal{E}_i(u) = \lim_{n \rightarrow \infty} \mathcal{E}_i^{(n)}(u), \quad \forall u \in \mathcal{F}_i.$$

Note that $\mathcal{E}_i(P_m u, P_m u) = \mathcal{E}_i^{(m)}(u, u)$.

In order to prove that we have a Dirichlet form and that the corresponding Laplace operator has compact resolvent, we follow [19], which uses ideas of [20].

For a point $x \in K_i$ we write $D_m(x)$ for the points in $\psi_{\sigma|m}(K_{\sigma(m)})$, where $\sigma|m = (\sigma_1, \dots, \sigma_m) \in E_i^m$, for all $\sigma : \pi(\sigma) = x$.

Lemma 4.10. *There exists a constant c_5 such that*

$$\int_{K_i} |f(x) - f(x_m)|^2 \mu(dx) \leq c_5 (\min_e \theta_e)^{-m} \mathcal{E}_i(f, f), \quad \forall x_m \in D_m(x), \quad f \in \mathcal{F}_i, \quad \forall i \in \mathcal{S}.$$

Proof. This can be built up from results on the zeroth level following either [20] or [19]. By definition of \mathcal{E}_i we have

$$\begin{aligned} \mathcal{E}_i(f, f) &= \sup_m \sum_{\sigma \in E_i^m} \rho_\sigma \mathcal{E}_{\sigma(m)}^{(0)}(f \circ \phi_\sigma, f \circ \phi_\sigma) \\ &\geq \sum_{\sigma \in E_i^m} \theta_\sigma \sum_{x, y = \pi(\sigma, e)} (f(x) - f(y))^2 \mu'_\sigma \\ &\geq c \min_{\sigma \in E_i^m} \theta_\sigma \int \sup_{y \in D_m(x)} (f(x) - f(y))^2 \mu_m(dx), \end{aligned} \tag{4.4}$$

for a constant $c > 0$, where $\mu_m(y)$ is the piecewise constant measure defined by $\mu_m(y) = \mu'_\sigma$ for $y \in D_m(x)$ for all $x \in K_i$. We can take a weak limit of this measure to recover

the Bernoulli measure μ_i on the fractal K_i . Let $x_m^a = \pi(\sigma a)$ for $\sigma \in E_i^m$ and a is a path whose projection onto the fractal is a vertex. We can then extend (4.4) to $y \in D_{m+1}(x)$ by modification of the constant c , giving

$$\int \sup_{y \in D_{m+1}(x)} (f(x_m^a) - f(y))^2 \mu(dx) \leq c(\min_{\sigma \in E_i} \theta_\sigma)^{-m} \mathcal{E}_i(f, f).$$

For any point $x \in K_i$ we can find a sequence $y_j \in F^{(j)}$ converging to x so that

$$\begin{aligned} \left(\int_{K_i} |f(x) - f(x_m^a)|^2 \mu(dx) \right)^{1/2} &= \left(\int_{K_i} \left| \sum_{j=m}^\infty f(y_{j+1}) - f(y_j) \right|^2 \mu(dx) \right)^{1/2} \\ &\leq \sum_{j=m}^\infty \left(\int_{K_i} |f(y_{j+1}) - f(y_j)|^2 \mu(dx) \right)^{1/2} \\ &\leq c \sum_{j=m}^\infty \theta_{\min}^{-j/2} \mathcal{E}_i(f, f)^{1/2} \\ &\leq c_5^{1/2} \theta_{\min}^{-m/2} \mathcal{E}_i(f, f)^{1/2}, \end{aligned}$$

as desired. □

This result shows that $\{P_{i,m}f\}_m$ is a Cauchy sequence in $L^2(K_i, \mu_i)$. Following [19] we can show that the limit is an injective map and hence that the form $(\mathcal{F}_i, \mathcal{E}_i)$ is closed in $L^2(K_i, \mu_i)$. We write $\mathcal{E}_{i,\lambda}(f, f) = \mathcal{E}_i(f, f) + \lambda \|f\|_{L^2(K_i, \mu_i)}^2$.

Lemma 4.11. *The bilinear form $(\mathcal{F}_i, \mathcal{E}_i)$ has compact λ -resolvent for all $\lambda > 0$, for each $i \in \mathcal{S}$.*

Proof. Define $T_i^m g(x) = g(x_m^a)$. Lemma 4.10 implies that

$$\|g - T_i^m g\|_{L^2(K_i, \mu_i)}^2 \leq c_5 \theta_{\min}^{-m} \mathcal{E}_i(g, g).$$

Let $I : (\mathcal{F}_i, \mathcal{E}_{i,\lambda}) \rightarrow L^2(K_i, \mu_i)$ denote the inclusion map and observe that therefore $T_i^m \rightarrow I$ in the operator norm as

$$\|I - T_i^m\|_{\mathcal{F}_i \rightarrow L^2(K_i, \mu_i)} \leq c_5 \theta_{\min}^{-m}.$$

As the operators T_i^m are compact, we have that I is compact and hence its adjoint I^* is compact. As I^* is the λ -resolvent we have the result. □

Theorem 4.12. *Let $\mathcal{P}_i = \bigcup_{n \in \mathbb{N}} P_n(l(F_i^{(n)}))$. Then*

- (1) \mathcal{P}_i is dense in $C(K_i, \mathbb{R})$ in the topology of uniform convergence and is a core for \mathcal{F}_i ;
- (2) $(\mathcal{E}_i, \mathcal{F}_i)$ is a regular, local Dirichlet form on $L^2(K_i, \mu_i)$;
- (3) there is a μ_i -symmetric Hunt process W_i , which is almost surely continuous, associated with the Dirichlet form $(\mathcal{E}_i, \mathcal{F}_i)$ on $L^2(K_i, \mu_i)$; and

- (4) there is a self-adjoint operator Δ_i , with compact resolvent, associated with the Dirichlet form $(\mathcal{E}_i, \mathcal{F}_i)$ on $L^2(K_i, \mu_i)$.

Proof. The proof follows the proof of [16, Theorem 3.4.6]. \square

The process W_i will be called the diffusion associated with the triplet $(K_i, \mu_i, \mathcal{E}_i)$. We also introduce a natural measure for the diffusion that could be called Brownian motion on an FRGD fractal. Even though we have not shown uniqueness for our process, we will call this diffusion Brownian motion and the associated generator the Laplacian. Let μ^b be a measure generated by setting $\mu_e^b = \rho_e^{-s}$. Let M be the associated matrix with elements $M_{ij} = \sum_{e \in E_{ij}} \rho_e^{-s}$. We set $\Phi_M(s)$ to be the spectral radius for M . The Hausdorff dimension of the FRGD fractal K in the effective resistance metric is given by $\dim_{H,R}(K) = S = \Phi_M^{-1}(1)$.

5. Spectral asymptotics

In order to calculate the spectral dimension for FRGD fractals we will prove a result about the asymptotics of the spectral counting function. This is a straightforward extension of the ideas of [17], where we use our multidimensional version of the renewal theorem.

For each component of our vector of fractals we have a Laplacian Δ_i , on the fractal K_i , which is the infinitesimal generator of the diffusion on K_i . If we write $(\cdot, \cdot)_{\mu_i}$ for the natural inner product on $L^2(K_i, \mu_i)$, we can also obtain the Laplacian directly from the Dirichlet form by defining $\Delta_i f$, for suitable f , as $\mathcal{E}_i(f, g) = -(\Delta_i f, g)_{\mu_i}$ for all $g \in \mathcal{F}_i$. We are interested in the eigenvalues of the operators, Δ_i , and we define the Dirichlet eigenvalue λ with eigenfunction u for this operator to be the solution to

$$\begin{aligned} \Delta_i u &= -\lambda u, & \text{in } K_i, \\ u(x) &= 0, & x \in \partial K_i. \end{aligned}$$

Note that $\partial K_i = F_i^{(0)}$ is the boundary of the component K_i and it is a finite set.

In order to define a Neumann eigenvalue for Δ_i , we need a notion of normal derivative. Let

$$d_i u(x) = \lim_{n \rightarrow \infty} \sum_{p \in N_n(x)} (u(x) - u(p)) H_i^{(n)}(x, p),$$

where $H_i^{(n)}$ is the finite difference operator defined by the n th Dirichlet form $\mathcal{E}_i^{(n)}$. The Neumann eigenvalue λ with eigenfunction u for the operator Δ_i is then defined by

$$\begin{aligned} \Delta_i u &= -\lambda u, & \text{in } K_i, \\ d_i u(x) &= 0, & x \in \partial K_i. \end{aligned}$$

The eigenvalue problems can be expressed in terms of the Dirichlet form as follows. In the case of the Dirichlet eigenvalue problem, we say that u is an eigenfunction for λ if $u \in \mathcal{F}_i^0 = \{f : f \in \mathcal{F}_i, f|_{\partial K_i=0}\}$, and

$$\mathcal{E}_i(u, v) = \lambda(u, v)_{\mu_i}, \quad \forall v \in \mathcal{F}_i^0.$$

Similarly for the Neumann eigenvalues; u is an eigenfunction for λ if $u \in \mathcal{F}_i$, and

$$\mathcal{E}_i(u, v) = \lambda(u, v)_{\mu_i}, \quad \forall v \in \mathcal{F}_i.$$

By Theorem 4.12 the operator Δ_i has compact resolvent for each $i \in \mathcal{S}$ and hence there is a discrete spectrum of eigenvalues $0 \leq \lambda_1^{(i)} \leq \lambda_2^{(i)} \leq \dots$, whose only accumulation point is ∞ . This enables us to define the spectral counting functions for the Dirichlet and Neumann eigenvalues as

$$N_i^0(x) = \#\{\lambda_k^{(i)} \leq x : \lambda_k^{(i)} \text{ is a Dirichlet eigenvalue of } \Delta_i\},$$

and

$$N_i(x) = \#\{\lambda_k^{(i)} \leq x : \lambda_k^{(i)} \text{ is a Neumann eigenvalue of } \Delta_i\}.$$

These functions are non-decreasing and right continuous with $N_i^0(0) = 0$ and $N_i(0) = 1$.

Our interest is in the large- x asymptotics of the counting functions and we begin with some preliminary results. Firstly there is a natural scaling in the Dirichlet form.

Lemma 5.1. *For all $f, g \in \mathcal{F}_i$, we have*

$$\mathcal{E}_i(f, g) = \sum_{j \in \mathcal{S}} \sum_{e \in E_{ij}} \rho_e \mathcal{E}_j(f \circ \phi_e, g \circ \phi_e).$$

Proof. We prove this by considering the n -level form, which can be decomposed as for $f, g \in l(F_i^{(n)})$:

$$\mathcal{E}_i^{(n+1)}(f, g) = \sum_{j \in \mathcal{S}} \sum_{e \in E_{ij}} \rho_e \mathcal{E}_j^{(n)}(f \circ \phi_e, g \circ \phi_e).$$

Now let $n \rightarrow \infty$. □

As a consequence of this we have

$$\mathcal{E}_i(f, g) = \sum_{\sigma \in E_i^n} \rho_\sigma \mathcal{E}_{\sigma(n)}(f \circ \phi_\sigma, g \circ \phi_\sigma). \tag{5.1}$$

The key relationships for the eigenvalue counting functions are described in the following lemma.

Lemma 5.2. *For all $x \geq 0$ and each $i \in \mathcal{S}$ we have*

$$\sum_{j \in \mathcal{S}} \sum_{e \in E_{ij}} N_j^0(\theta_e^{-1}x) \leq N_i^0(x) \leq N_i(x) \leq \sum_{j \in \mathcal{S}} \sum_{e \in E_{ij}} N_j(\theta_e^{-1}x), \tag{5.2}$$

and there exists a finite constant M such that

$$N_i^0(x) \leq N_i(x) \leq N_i^0(x) + M, \quad \forall x, i. \tag{5.3}$$

Proof. We begin by making a crucial observation on the decimation of the eigenfunctions. We define a new Dirichlet form

$$\tilde{\mathcal{F}}_i = \{f : K_i \setminus F_i^{(1)} \rightarrow \mathbb{R}, f \circ \phi_e = f_e : K_j \setminus F_j^{(0)} \rightarrow \mathbb{R}, f_e \in \mathcal{F}_j, \forall e \in E_{ij}, \forall j \in \mathcal{S}\},$$

and for $f, g \in \tilde{\mathcal{F}}_i$ we define

$$\tilde{\mathcal{E}}_i(f, g) = \sum_{j \in \mathcal{S}} \sum_{e \in E_{ij}} \rho_e \mathcal{E}_j(f \circ \phi_e, g \circ \phi_e).$$

It is easy to see that $\mathcal{F}_i \subset \tilde{\mathcal{F}}_i$ and $\mathcal{E}_i = \tilde{\mathcal{E}}_i$ when restricted to $\mathcal{F}_i \times \mathcal{F}_i$. The form $(\tilde{\mathcal{E}}_i, \tilde{\mathcal{F}}_i)$ is a local regular Dirichlet form on $L^2(K_i, \mu_i)$. The associated resolvent is compact and hence the operator has a spectrum which consists of eigenvalues. Now let u be an eigenfunction for the Dirichlet form $(\tilde{\mathcal{E}}_i, \tilde{\mathcal{F}}_i)$ with eigenvalue k , thus

$$\tilde{\mathcal{E}}_i(u, v) = k(u, v)_{\mu_i}, \quad \forall v \in \tilde{\mathcal{F}}_i.$$

If we rewrite this, using the scaling in the Dirichlet form and in the measure, then

$$\sum_{j \in \mathcal{S}} \sum_{e \in E_{ij}} \rho_e \mathcal{E}_j(u \circ \phi_e, v \circ \phi_e) = k \sum_{j \in \mathcal{S}} \sum_{e \in E_{ij}} \mu'_e(u \circ \phi_e, v \circ \phi_e)_{\mu_j}.$$

Thus we have for any $h \in \mathcal{F}_j$,

$$\mathcal{E}_j(u \circ \phi_e, h) = k \rho_e^{-1} \mu'_e(u \circ \phi_e, h)_{\mu_j}, \quad \forall e \in E_{ij},$$

so that for $e \in E_{ij}$, $u_e = u \circ \phi_e$ is an eigenfunction of $(\mathcal{E}_j, \mathcal{F}_j)$ with eigenvalue $\theta_e^{-1}k$. Clearly, if u_e is an eigenfunction of $(\mathcal{E}_j, \mathcal{F}_j)$ with eigenvalue $\theta_e^{-1}k$ for each $e \in E_i$, then u , defined by

$$u(x) = \begin{cases} u_e(x), & x \in \phi_e(K_i), \\ 0, & \text{otherwise,} \end{cases}$$

is an eigenfunction for $(\tilde{\mathcal{E}}_i, \tilde{\mathcal{F}}_i)$, with eigenvalue k .

Let \tilde{N}_i denote the eigenvalue counting function for $(\tilde{\mathcal{E}}_i, \tilde{\mathcal{F}}_i)$. From the decimation property of the eigenfunctions we have

$$\begin{aligned} \tilde{N}_i(x) &= \#\{k : k \leq x\} \\ &= \sum_{e \in E_i} \#\{k\theta_e^{-1} : k\theta_e^{-1} \leq x\theta_e^{-1}\} \\ &= \sum_{j \in \mathcal{S}} \sum_{e \in E_{ij}} N_j(\theta_e^{-1}x). \end{aligned}$$

Observe that as the domain of $(\tilde{\mathcal{E}}_i, \tilde{\mathcal{F}}_i)$ is larger than $(\mathcal{E}_i, \mathcal{F}_i)$, by a minimax argument the eigenvalues of $(\tilde{\mathcal{E}}_i, \tilde{\mathcal{F}}_i)$ must be less than those for $(\mathcal{E}_i, \mathcal{F}_i)$. Thus we have $N_i \leq \tilde{N}_i$ and we have the left inequality.

We now define another Dirichlet form for the Dirichlet eigenvalues. Let $\tilde{\mathcal{F}}_i^0 = \{f : f \in \mathcal{F}_i^0, f|_{F(1)} = 0\}$ and define $\mathcal{E}_i^0 = \mathcal{E}_i$ restricted to $\tilde{\mathcal{F}}_i^0 \times \tilde{\mathcal{F}}_i^0$. Following the same reasoning

as above we can see that if u is an eigenfunction of $(\tilde{\mathcal{E}}_i^0, \tilde{\mathcal{F}}_i^0)$ with eigenvalue k , then for each $j \in \mathcal{S}$, $e \in E_{ij}$ we have that u_e is an eigenfunction for $(\mathcal{E}_j, \mathcal{F}_j)$ with eigenvalue $\theta_e^{-1}k$. As before we can derive the lower bound for the Dirichlet eigenvalue counting function.

Finally, the middle inequality follows from the minimax argument as $\mathcal{F}_0 \subset \mathcal{F}$.

The second result is a consequence of Dirichlet–Neumann bracketing and the fact that the space of harmonic functions is finite dimensional. For more on this see [17]. \square

In order to state our main result we define the $|\mathcal{S}| \times |\mathcal{S}|$ -matrix R_s to be the one with entries given by $r_s(i, j) = \sum_{e \in E_{ij}} \theta_e^{-s}$. We then let $\Psi(s)$ denote the spectral radius of the matrix R_s .

Theorem 5.3. *If the graph is strongly connected and $d_s^\mu/2$ is the solution to the equation $\Psi(s) = 1$, then*

$$0 < \liminf_{x \rightarrow \infty} N_i^0(x)x^{-d_s^\mu/2} \leq \limsup_{x \rightarrow \infty} N_i^0(x)x^{-d_s^\mu/2} < \infty,$$

and

$$0 < \liminf_{x \rightarrow \infty} N_i(x)x^{-d_s^\mu/2} \leq \limsup_{x \rightarrow \infty} N_i(x)x^{-d_s^\mu/2} < \infty.$$

Proof. As the underlying graph is strongly connected, the matrix R_s is a primitive matrix and hence we can apply the Perron–Frobenius Theorem. There exists an eigenvector u such that

$$u_i = \sum_{j \in \mathcal{S}} \sum_{e \in E_{ij}} \theta_e^{-d_s^\mu/2} u_j.$$

Let $\beta_i(t) = \exp(-td_s^\mu/2)N_i(e^t)$, $\beta_i^0(t) = \exp(-td_s^\mu/2)N_i^0(e^t)$ and rewrite the right-hand side of (5.2) in terms of the function β as

$$\beta_i(t) \leq \sum_{e \in E_i} \theta_e^{-d_s^\mu/2} \beta_j(t - \log \theta_e), \quad t \in \mathbb{R}.$$

By iteration of the inequality we can write this as

$$\beta_i(t) \leq \sum_{\sigma \in E_i^n} (\theta_\sigma)^{-d_s^\mu/2} \beta_{\sigma(n)}(t - \log \theta_\sigma), \quad t \in \mathbb{R}.$$

This will remain true if we alter our definition of E_i^n to be the addresses of variable length, which ensure that $\theta_\sigma = \rho_\sigma/\mu'_\sigma$ is of size e^n ,

$$\hat{E}_i^n = \{\sigma = (\sigma_1, \dots, \sigma_m) : \theta_\sigma \leq e^n \leq \theta_{\sigma, (i_m, i_{m+1})}\}.$$

Note that we still have u as an eigenvector in that

$$u_i = \sum_{\sigma \in \hat{E}_i^n} \theta_\sigma^{-d_s^\mu/2} u_j.$$

Hence we can write

$$\beta_i(t) \leq \sum_{\sigma \in \hat{E}_i^n} (\theta_\sigma)^{-d_s^\mu/2} \beta_{\sigma(m)}(t - \log \theta_\sigma).$$

Let c'_i be such that $\beta_i(t) \leq c'_i$ for $t \in [0, \max_{e \in E} \log \theta_e]$ and choose an n such that for $\sigma \in \hat{E}_i^n$,

$$t - \log(\theta_\sigma) \in [0, \max_{e \in E} \log(\theta_e)].$$

In which case

$$\begin{aligned} \beta_i(t) &\leq \sum_{\sigma \in \hat{E}_i^n} (\theta_\sigma)^{-d_s^\mu/2} \beta_{\sigma(m)}(t - \log \theta_\sigma) \\ &\leq c'_i \sum_{\sigma \in \hat{E}_i^n} (\theta_\sigma)^{-d_s^\mu/2} \\ &\leq \frac{c'_i}{\min_j u_j} \sum_{\sigma \in \hat{E}_i^n} u_{\sigma(m)} (\theta_\sigma)^{-d_s^\mu/2} \\ &= \frac{c'_i u_i}{\min_j u_j}, \end{aligned}$$

which is independent of n and hence holds for all $t > 0$. Similarly for the Dirichlet eigenvalue counting function, we can obtain a lower bound by obtaining a bound over a finite interval and then iterating. This gives a bound of the same form. In order to obtain our result we now use (5.3) to see that there exist constants c, c' such that

$$c\beta_i^0(t) \leq \beta_i(t) \leq c'\beta_i^0(t), \quad \forall t > 0, \quad i \in \mathcal{S}.$$

Using this, and rewriting in terms of the eigenvalue counting function N , we have the result. □

We can provide a finer result using our multidimensional renewal Theorem 2.6. Again we can follow [17] and write equation (5.2) as a renewal type equation.

Let

$$\xi_i(t) = \beta_i^0(t) - \sum_{j \in \mathcal{S}} \sum_{e \in E_{ij}} \theta_e^{-d_s^\mu/2} \beta_j^0(t - \log \theta_e),$$

so that ξ is a non-negative bounded function of t . We also define the measure m_{ij} to be

$$m_{ij}(ds) = \sum_{e \in E_{ij}} \theta_e^{-d_s^\mu/2} \delta_{\log \theta_e}(ds).$$

Let $F(t)$ denote the matrix of measures, expressed as distribution functions. Note that by construction the maximum eigenvalue of $F(\infty)$ is 1. We can now write

$$\beta_i^0(t) = \xi_i(t) + \sum_{j \in \mathcal{S}} \int_0^\infty \beta_j^0(t-s) m_{ij}(ds), \quad t \in \mathbb{R}, \tag{5.4}$$

which can be expressed as

$$\beta^0(t) = \xi(t) + (F * \beta^0)(t), \quad t \in \mathbb{R}.$$

By taking the transpose, this is a matrix renewal equation of the type considered in § 2. Recall that the lattice/non-lattice dichotomy depends upon the probability measure defined by

$$\nu_{11}^1 = m_{11} + m_{1\hat{1}} * \sum_{k=0}^{\infty} M_{11}^{*k} * m_{\hat{1}1}.$$

Thus we will be in the lattice case if, for any path γ in the graph from node 1 to itself, $\log \theta_\gamma$ is supported on a discrete subgroup of \mathbb{R} .

We are now ready to state our main results. These describe the behaviour of the leading-term asymptotics for the spectral counting function and involve a simple application of the multidimensional renewal theorem.

Theorem 5.4. *For an FRGD fractal which is strongly connected, if ν_{11}^1 is not lattice, then there exist constants $c_6(i), c_7(i) > 0$ such that*

$$\lim_{x \rightarrow \infty} N_i(x) x^{-d_s^\mu/2} = c_6(i),$$

and

$$\lim_{x \rightarrow \infty} N_i^0(x) x^{-d_s^\mu/2} = c_7(i).$$

If the measure ν_{11}^1 is lattice, then there exist periodic functions p_i, p'_i such that

$$\lim_{x \rightarrow \infty} (N_i(x) x^{-d_s^\mu/2} - p_i(\log x)) = 0,$$

and

$$\lim_{x \rightarrow \infty} (N_i^0(x) x^{-d_s^\mu/2} - p'_i(\log x)) = 0.$$

Proof. We first observe that the first Dirichlet eigenvalue $\lambda_1 > 0$. Thus there exists an $x_0 \in \mathbb{R}$ such that for $x < x_0$, the counting function $N_i^0(x) = 0$. Thus we can shift our counting function and, as the graph is strongly connected, apply Corollary 2.12. The case of the Neumann counting function follows from (5.3). \square

Remark 5.5. In the case of regular p.c.f. self-similar sets, the spectral dimension d_s can be defined as the maximal value of the spectral exponent d_s^μ over all Bernoulli measures [17]. In our setting, in the corresponding regular case where $\rho_e > 1$ for all $e \in E$, we can maximize over the product measure on the Markov subshift and define $d_s = \max_\mu d_s^\mu$. It is not difficult to see that the measure $\mu_e^b = \rho_e^{-S}$ will achieve the maximum. For this measure, $\theta_e = \rho_e^{S+1}$ and the spectral dimension satisfies $\Psi(d_s/2) = 1$. The remark at the end of § 4 allows us to write $\Psi(u) = \Phi_M(u(S+1))$ and hence $\Psi(u) = 1$ when $u = d_s/2 = S/(S+1)$, where S is the Hausdorff dimension of the set in the effective resistance metric. This relationship between the spectral dimension and the Hausdorff dimension in the effective resistance metric has been observed for other finitely ramified fractals, such as p.c.f. fractals [16] and random recursive Sierpinski gaskets [11].

We now assume that the graph is not strongly connected. As in [25] we write $\text{SC}(G)$ for the set of strongly connected components of the graph G . Each such component defines a fractal within the vector of all fractals. For each $H \in \text{SC}(G)$ we can determine the spectral exponent within the component $d_s^\mu(H)$. There is a partial ordering such that $H_1 \leq H_2$ if for $i \in H_1$, there is a directed path $\sigma \in E_i^\infty$ with $\sigma(m) \in H_2$ for some $m < \infty$. We can classify the strongly connected components, which correspond to classes in the classification of reducible matrices, into basic classes if the component has the same spectral exponent as the whole set. As in our discussion of reducible matrices, we write S_m for the union of the classes of height $m + 1$. Let $\tilde{H}_j = \{H \in \text{SC}(G) \mid \text{there is a path from } j \text{ to some vertex of } H\}$. We state the results for the Neumann counting function as exactly the same results (with different constants, periodic functions) hold for the Dirichlet counting function.

Theorem 5.6. *For a general FRGD fractal the spectral exponent is defined by $d_s^\mu = \max\{d_s^\mu(H) \mid H \in \text{SC}(G)\}$ and $d_s^\mu(K_i) = \max\{d_s^\mu(H) \mid H \in \tilde{H}_i\}$. If $i \in S_0$, then there is a periodic function G_i such that*

$$\lim_{x \rightarrow \infty} \left(\frac{N_i(x)}{x^{d_s^\mu/2}} - G_i(\log x) \right) = 0.$$

If $i \in S_m$, then there are constants $c_7(i) > 0$ such that

$$\lim_{x \rightarrow \infty} \frac{N_i(x)}{(\log x)^m x^{d_s^\mu/2}} = c_7(i).$$

If $i \notin S$ and there is no path from S to i , then

$$\lim_{x \rightarrow \infty} \frac{N_i(x)}{x^{d_s^\mu/2}} = 0.$$

If $i \notin S$ and there is a path from S_0 to i but no path to i from S_l , with $l > 0$, then there is a periodic function G'_i such that

$$\lim_{x \rightarrow \infty} \left(\frac{N_i(x)}{x^{d_s^\mu/2}} - G'_i(\log x) \right) = 0.$$

If $i \notin S$ and there is a path from S_m to i but no path to i from S_l , with $l > m$, then there is a constant $c_8(i) > 0$ such that

$$\lim_{x \rightarrow \infty} \frac{N_i(x)}{(\log x)^m x^{d_s^\mu/2}} = c_8(i).$$

Proof. We observe again that the Dirichlet counting function can be shifted and hence we can apply Corollary 2.12 to give the result. \square

Note that it would be possible to write out an explicit formula for the constants in terms of the function ξ and the matrix of measures.

In the case where the limit is 0, we can obtain a refinement by only considering the components feeding into that class and derive a corresponding eigenvalue counting-function result. We only state here the result for the case of the class of height 1. Let the set of states which have access to a given component H be denoted by $\mathcal{S}(H)$.

Corollary 5.7. *If H is a component with $d_s^\mu(H) < d_s^\mu$, and if i is in a class of height 1 in the $|\mathcal{S}(H)| \times |\mathcal{S}(H)|$ -submatrix, then there is a periodic function $G_{H,i}$ such that*

$$\lim_{x \rightarrow \infty} \left(\frac{N_i(x)}{x^{d_s^\mu(H)/2}} - G_{H,i}(\log x) \right) = 0.$$

6. Examples

6.1. Plain mandala

In [27] a fractal subset of \mathbb{R}^2 , called plain mandala (inspired by the Buddhist image, mandala), was defined. Let L denote the upper semicircle of unit radius with origin at the centre. Set $\xi_1 = (-1, 0)$, $\xi_2 = (1, 0)$ and define for $i = 1, 2$ maps $F_i(x) = \frac{1}{2}(x + \xi_i)$. Also define $F_3, F_4 : L \rightarrow L$ by

$$F_i(x) = \phi\left(\frac{1}{2}(\psi(x) + \psi(\xi_{i-2}))\right), \quad x \in K, \quad i = 3, 4,$$

where $\phi : [0, \pi] \rightarrow L$ is defined by $\phi(\theta) = (\cos(\theta), \sin(\theta))$ and $\psi = \phi^{-1}$. The fractal is defined to be the set K which satisfies

$$K = \bigcup_{i=1,2} F_i(K) \cup L.$$

Though this cannot be directly expressed as a graph-directed fractal, as the maps F_3, F_4 are not similitudes, we can still use the above approach. The two types are given by the semicircle, L and the ‘mandala’ K . Note that the semicircles map into themselves and into the mandalas but the mandalas only map to themselves and the graph describing the fractal will not be strongly connected. The graph-directed formulation for the fractal is

$$\begin{aligned} K &= \bigcup_{i=1,2} F_i(K) \cup \bigcup_{i=3,4} F_i(L), \\ L &= \bigcup_{i=3,4} F_i(L), \end{aligned}$$

and is illustrated in Figure 4. We can regard the construction graph as having two vertices (one for each type) and six edges (one for each map) as shown. We assign a resistivity r_j to each edge of the graph which corresponds to map F_j . In the case where $r_1 + r_2 = 1$ and $r_3 + r_4 = 1$, and $t > 1$, Murai [27] has shown the existence of a regular Dirichlet form with scaling:

$$\begin{aligned} \mathcal{E}_K(f) &= \sum_{i=1,2} (tr_i)^{-1} \mathcal{E}_K(f \circ F_i) + \sum_{i=3,4} (r_i)^{-1} \mathcal{E}_L(f \circ F_i), \\ \mathcal{E}_L(f) &= \sum_{i=3,4} (r_i)^{-1} \mathcal{E}_L(f \circ F_i). \end{aligned}$$

In order to discuss the asymptotic scaling of the eigenvalues we choose a measure. As in [27], we take $\tau, s_i, i = 1, 2, 3, 4$, to be positive real numbers with $\tau \geq t, s_1 + s_2 = 1$,




Figure 4. Plain mandala as an FRGD fractal.

$s_3 + s_4 = 1, s_1s_2 = s_3s_4$. Let (μ_K, μ_L) be probability measures on (K, L) defined as in (4.1) by the measures on the edges $\mu'_j = \tau s_j, j = 1, 2$ and $\mu'_j = s_j, j = 3, 4$.

As there are two strongly connected components we can compute the spectral exponent within each one. Let d_s^1, d_s^2 satisfy

$$\sum_{i=1,2} (tr_i \tau s_i)^{d_s^1/2} = 1, \quad \sum_{i=3,4} (r_i s_i)^{d_s^2/2} = 1.$$

In [27] a result similar to Theorem 5.3 was established showing that the scaling in the spectral counting function could require a logarithmic correction factor, but the existence of the normalized limit was not established. In order to apply our main theorem we let the matrix renewal measure be

$$M = \begin{bmatrix} \sum_{i=1}^2 (tr_i \tau s_i)^{d_s^1/2} \delta_{-\log tr_i \tau s_i} & 0 \\ \sum_{i=3}^4 (r_i s_i)^{d_s^2/2} \delta_{-\log r_i s_i} & \sum_{i=3}^4 (r_i s_i)^{d_s^2/2} \delta_{-\log r_i s_i} \end{bmatrix}.$$

The lattice case occurs for state K if $\log(tr_1 \tau s_1)/\log(tr_2 \tau s_2) \in \mathbb{Q}$ and for state L if $\log(r_3 s_3)/\log(r_4 s_4) \in \mathbb{Q}$. Thus, writing N_K, N_L for either the Dirichlet or Neumann eigenvalue counting function associated with each type K, L , by Theorem 5.6 we have the following.

- (1) If $d_s^1 = d_s^2$, then there exist constants $c_9(i), i = K, L$, such that

$$\lim_{\lambda \rightarrow \infty} \frac{N_i(\lambda)}{\lambda^{d_s^1/2} \log \lambda} = c_9(i), \quad i = K, L.$$




Figure 5. On the left is the generator of the diamond fractal; in the centre the diamond component at level 2 is shown; and on the right are the three component types—the diamond, the chevron and the crown.

(2) If $d_s^1 > d_s^2$, then $d_s = d_s^1$ and there exist periodic functions p_K, p_L such that

$$N_K(\lambda) = \lambda^{d_s/2}(p_K(\log \lambda) + o(1)),$$

and

$$\lim_{\lambda \rightarrow \infty} \frac{N_L(\lambda)}{\lambda^{d_s/2}} = 0,$$

but

$$N_K(\lambda) = \lambda^{d_s^2/2}(p_L(\log \lambda) + o(1)).$$

If state i is in the non-lattice case, then p_i is constant for $i = K, L$.

(3) For $d_s^2 > d_s^1$, we set $d_s = d_s^2$ and there exist periodic functions p'_i such that

$$N_i(\lambda) = \lambda^{d_s/2}(p'_i(\log \lambda) + o(1)), \quad i = K, L.$$

If state i is in the non-lattice case, then p_i is constant for $i = K, L$.

Remark 6.1. We do not have a proof that the periodic function is actually non-constant in the lattice case.

6.2. Fractals with infinitely ramified generator

In this section we discuss a set of fractals in two dimensions whose generators are not finitely ramified but for which we can use graph-directed constructions. We will show the existence of a non-degenerate Dirichlet form on an example fractal from this class in the next subsection. It is an open problem whether the construction of such a form is possible in general.

Our starting point will be a regular polygon P with edge set E_P and vertex set V_P . Let $n_P = |V_P|$. We will let Θ_C be the rotation of P sending $x \in V_P$ to its clockwise nearest neighbour. For $x, y \in V_P$, let Θ_{xy} be the mirror symmetry of P that switches x and y .

Let G be the symmetry group of P . For any line L , let Θ_L be the reflection in L . The Θ_L are not necessarily in G . For a pair of points x, y , let L_{xy} be the line segment from x to y . Example fractals that we consider here can be found in Figures 3 and 5.

Definition 6.2. A set $A \neq \emptyset$ is finitely ramified relative to B if B is a proper superset of A , and $|\overline{A} \cap \overline{A^c}|$ is finite, where the complement is taken with respect to B .

Definition 6.3. Let P be a regular polygon, and consider a set of affine maps $\{\phi_i\}_{i \in I}$, where I is finite. For a subset $J \subset I$, define $\phi_J(\cdot) = \bigcup_{i \in J} \phi_i(\cdot)$ and $\Phi(\cdot) = \phi_I(\cdot)$. Assume the following.

- (1) For all $i \in I$, $\phi_i(P) \subset P$.
- (2) For all $\Theta \in G$, there is a map $\Theta_I : I \rightarrow I$ such that for all $i \in I$, $\Theta(\phi_i(P)) = \phi_{\Theta_I(i)}(P)$. In particular, $\Theta(\Phi(P)) = \Phi(P)$.
- (3) For an $L \in E_P$, for all $i \in I$ such that $\phi_i(P) \cap L \neq \emptyset$ there exists a set A_i with the finite ramification property relative to $\Phi(P)$ and an edge $L' \in E_P$ such that $A_i \cap L' = \emptyset$.
- (4) For all $i, j \in I$, $i \neq j$, either
 - (i) $\phi_i(P) \cap \phi_j(P) = \emptyset$;
 - (ii) there are $x, y \in V_P$ such that $\phi_i(P) \cap \phi_j(P) = \phi_i(x) = \phi_j(y)$;
 - (iii) there are $L, L' \in E_P$ such that $\phi_i(P) \cap \phi_j(P) = \phi_i(L) = \phi_j(L')$ (note that $\Theta_L \circ \phi_i(P) = \phi_j(P)$).

Let K be the unique fixed point of Φ . K is then called a *decomposable nested fractal*.

Observation 6.4. A decomposable nested fractal K is preserved under the symmetries of P , and $K \subset P$. Indeed, $K = \bigcap_{n=1}^{\infty} \Phi^n(P)$.

We will keep a parallel indexing system, so that if X_I is a collection in I , X_Y will be $\{\phi_J(Y) : J \in X_I\}$. Typically, Y will be P or K .

For a decomposable nested fractal as above, let \mathcal{A}_I be the collection of sets $J \subset I$ such that $\phi_J(P)$ has the finite ramification property relative to $\Phi(P)$. Then $\overline{\mathcal{A}}_I = \mathcal{A}_I \cup \{\emptyset, P\}$ is a G -invariant algebra. Let $\mathcal{B}_I = \{A \in \mathcal{A}_I : \neg \exists B \in \mathcal{A}_I [B \subset A]\}$. \mathcal{B}_I is then G -invariant also. Let $\mathcal{A}_P, \mathcal{A}_K, \mathcal{B}_P$ and \mathcal{B}_K be defined accordingly.

Definition 6.5. For a set A , let $B_\varepsilon(A) = \{x \in P : d(x, A) < \varepsilon\}$. $B_\varepsilon(x) = B_\varepsilon(\{x\})$.

Lemma 6.6. If $A \in \mathcal{B}_P$ and $x, y \in A$ are such that $B_\varepsilon(x)$ and $B_\varepsilon(y)$ are contained in A for some $\varepsilon > 0$, there is a curve C and a $\beta > 0$ such that $x, y \in B_\beta(C) \subset A$.

Proof. We easily see that there must be a sequence $S = \{s_1, \dots, s_n\}$ such that $x \in \phi_{s_1}(P)$ and $y \in \phi_{s_n}(P)$ and each pair $\phi_{s_i}(P)$ and $\phi_{s_{i+1}}(P)$ intersect at a line $\phi_i(L)$ for some $L \in E_P$.

Build C piecewise by letting C^0 be the straight line from x to the centre of $\phi_{s_1}(P)$, letting C^i be the straight line from the centre of $\phi_{s_i}(P)$ to the centre of $\phi_{s_{i+1}}(P)$ for

$i = 1, \dots, n - 1$, and letting C^n be the straight line from the centre of $\phi_{s_i}(P)$ to y . We have $C_\varepsilon^0, C_\varepsilon^n \subset A$, and $C_{l \cdot r_{\min}}^n \subset A$, so let $C = \bigcup_{i=0}^n C^i$ and $\beta = \varepsilon \wedge (l \cdot r_{\min})$. Thus $B_\beta(C) \subset A$ as required. \square

Lemma 6.7. *Let $x \in V_P$. If K is connected, there is an $i \in I$ such that $x \in \phi_i(P)$.*

Proof. By symmetry, there is either an $i \in I$ for each $x \in V_P$ so that $x \in \phi_i(P)$, or there does not exist such an $i \in I$ for any $x \in V_P$. So assume such an $i \in I$ does not exist. Then $\Phi(P) \cap V_P = \emptyset$. By Observation 6.4 we get that $\phi_A(K) \cap \phi_{A^c}(K) \subset \phi_A(\phi(P)) \cap \phi_{A^c}(\phi(P))$. But at the same time, Definitions 6.3 (3) and 6.3 (4) give us that $\phi_A(K) \cap \phi_{A^c}(K) \subset \phi_A(V_P) \cap \phi_{A^c}(V_P)$. These two together with $\Phi(P) \cap V_P = \emptyset$ give us that $\phi_A(K) \cap \phi_{A^c}(K) = \emptyset$, contrary to the assumption that K is connected. \square

Observation 6.8. *For $n_P \geq 4$, let C be a curve in P connecting two points x, y , belonging to two non-adjacent edges $L_1, L_2 \in E_P$, respectively. By elementary topology, $C \cap \Theta_C(C) \neq \emptyset$.*

Proposition 6.9. *If $B \subset \mathcal{B}_P$ intersects with two lines $L_1, L_2 \in E_P$, then L_1 and L_2 must be adjacent.*

Proof. This is clearly true for $n_P = 3$. For $n_P > 3$, assume L_1 and L_2 are non-adjacent. Choose $\varepsilon > 0$, $x \in B \cap L_1$ and $y \in B \cap L_2$ such that $B_\varepsilon(\{x, y\}) \subset B$. By Lemma 6.6, there is a curve C connecting x and y and a $\beta > 0$ such that $B_\beta(C) \subset B$. By Observation 6.8, there is a $z \in C \cap \Theta_C(C)$, and $B_\varepsilon(z) \in B \cap \Theta_C(B)$. This means there must be an $i \in I$ such that $\phi_i(P) \subset B \cap \Theta_C(B)$, and so $B_I \cap \Theta_C(B_I)$ is non-empty, and therefore in \mathcal{A}_I . This contradicts the assumption that $B \in \mathcal{B}_P$, unless $B = \Theta_C(B)$. But if $B = \Theta_C(B)$, then Definition 6.3 (3) for decomposable nested fractals gives us an A_i such that $B \cap A$ is a proper subset of B in \mathcal{A} , again contradicting $B \in \mathcal{B}$. \square

Corollary 6.10. *A set $B \subset \mathcal{B}_P$ intersects at most two lines $L_1, L_2 \in E_P$.*

Proof. Assume that B intersects three or more lines in E_P . If $n_P = 3$, B intersects all $L \in E_P$, contradicting Definition 6.3 (3) for decomposable nested fractals. If $n_P \geq 4$, at least two of the lines will be non-adjacent, contradicting Proposition 6.9. \square

Corollary 6.11. *If $B \in \mathcal{B}$ and B intersects L_{xy} on both sides of $z = \frac{1}{2}(x + y)$, then $B = \Theta_{xy}(B)$, and B intersects no $L \in E_P$, $L \neq L_{xy}$.*

Proof. To prove $B = \Theta_{xy}(B)$, we notice that if it was not, then $B \cap \Theta_{xy}(B)$ would be a proper subset of B and at the same time an element of \mathcal{A} , contrary to the assumption that $B \in \mathcal{B}$. From this it follows that if B intersects with an edge $L \neq L_{xy}$, it must also intersect $L' = \Theta_{xy}(L)$, contradicting Corollary 6.10. \square

Let $\angle_x = L_{xy'} \cup L_{xz'}$, where $y' = \frac{1}{2}(x + y)$, $z' = \frac{1}{2}(x + z)$, and y, z are x 's nearest neighbours in V_P . Let α be the angle in \angle_x . Further let $E = \bigcup_{L \in E_P} L$.

Corollary 6.12. *There is, for each $x \in V_P$, a Θ_{yz} -symmetric $B \in \mathcal{B}$ such that $x \in B$ and $B \cap E = B \cap \angle_x$.*

Fix x , set $P_1 = \Phi(P)$ and let P_n be the union of $\Phi(P)$ and a clockwise rotation by α degrees of P_{n-1} around x . $P_1 = \Phi(P)$. We define such P_n for all n smaller than or equal to the maximum number of rotations $360/\alpha$. Let \mathcal{B}_n be the corresponding smallest components, and let $\mathcal{C}' = \cup_n \mathcal{B}_n$. Define an equivalence relation ' \sim ' on \mathcal{C}' by letting $A \sim B$ if and only if there is a bijection $f : A \rightarrow B$ such that $d(x, y) = c \cdot d(f(x), f(y))$. Let $\mathcal{C} = \mathcal{C}' / \sim$, and let $\mathcal{C}_n = \mathcal{B}_n / \sim$.

We want to see how \mathcal{C} differs from \mathcal{C}_1 ; which new element types (\mathcal{C}) have we added from the old types (\mathcal{C}_1) by the procedure above? It is clear that all of the new elements must be unions of old elements. Unions of elements can, again, only happen if two or more old-type elements have intersections of infinite cardinality. This can, again, only happen with \mathcal{C}_1 -elements intersecting E . Let us then divide the old-type elements of \mathcal{C}_1 into three disjoint classes: \mathcal{C}_A , elements which do not intersect with E ; \mathcal{C}_B , elements which intersect exactly one $L \in E_P$; and \mathcal{C}_C , elements which intersect exactly two $L \in E_P$. The previous results describe these sets, and guarantee that \mathcal{C}_C is non-empty.

Proposition 6.13. *The types in the set \mathcal{C} are in one-to-one correspondence with the elements of $\text{FRGD}(K)$.*

Proof. We need only show that each of the equivalence classes of $\Psi^2(P)$ is in one-to-one relation with \mathcal{C} . This, however, is now quite easy. We can divide the elements of $\Psi^2(P)$ into three classes. Firstly, those that are images, by some ϕ_i , of an element of \mathcal{C}_A will correspond exactly to an element of \mathcal{C}_A . Then we have the elements that have been created by a union of two images, by maps ϕ_i and ϕ_j , respectively, of elements of \mathcal{C}_B . They will correspond to elements of \mathcal{C}_2 . Finally, we have the unions of images of elements of \mathcal{C}_C . They are easily seen to correspond to elements in some \mathcal{C}_k for $k \geq 2$. \square

An example of a decomposable nested fractal, called the diamond fractal in [18], with its vector of types is shown in Figure 5.

6.3. Existence results

The key assumption in our construction of the Laplacian on the graph-directed fractals was that there exists a fixed point for the map \mathcal{T} . It is a difficult problem to determine the existence of fixed points for the map, even in the single-component case. For the class of nested fractals, [24] showed that a fixed point would exist by using the symmetry of these fractals. In [30] conditions are given which show the existence of a fixed point for more general fractals, but these are very difficult to check in general.

We will follow [24] and give an existence result for a particular subclass of graph-directed nested fractals which are highly symmetric. We assume that the vector \mathbf{r} has $r_e = 1$, for all $e \in E$.

Let $\{\phi_e : e \in E\}$ denote a family of similitudes in \mathbb{R}^n associated with the edges of a graph G . As before there exists a set of compact subsets $K_i \subset \mathbb{R}^n$ such that

$$K_i = \bigcup_{j \in S} \bigcup_{e \in E_{ij}} \phi_e(K_j).$$

We call K a graph-directed nested fractal if it satisfies the following assumption.

Assumption 6.14.

- (1) K is a finitely ramified graph-directed fractal.
- (2) If $x, y \in F_i^{(0)}$, then reflection in the hyperplane $H_{xy} = \{z : |z - x| = |z - y|\}$ maps $F_i^{(n)}$ to itself.

The second part of the assumption is strong as the symmetry condition is required for each component. With it we can apply the approach of Lindström [24] (or see [1]) to show that the map on transition probabilities has a fixed point. Such a fixed point can be used to construct a fixed point for the \mathcal{T} map.

The fractal which appears in Figure 3 is a graph-directed nested fractal and hence there will be a solution to the fixed-point problem for this set. For the fractal in Figure 2, which is not a graph-directed nested fractal, discussions with Dr V. Metz have shown that, by techniques developed in [26], a fixed point will exist. The existence of the fixed point for the fractal of Figure 5 has also been demonstrated by a direct calculation in [18] and indirectly in [26].

Acknowledgements. The work of B.M.H. was undertaken at BRIMS, Hewlett-Packard Research Laboratories, Bristol, with the aid of a Royal Society Industry Fellowship and also at the Isaac Newton Institute, Cambridge, during the programme ‘Mathematics and Applications of Fractals’. S.O.G.N. was supported at the University of Edinburgh by EPSRC grant GR/L07239. B.M.H. thanks Takashi Kumagai for drawing his attention to the paper [27], Volker Metz for comments on an earlier version of this paper, and a diligent referee for many useful comments.

References

1. M. T. BARLOW, Diffusions on fractals, in *Lectures on probability theory and statistics, École d’été de probabilités de St Flour XXV* (Springer, 1998).
2. M. T. BARLOW AND J. KIGAMI, Localized eigenfunctions of the Laplacian on p.c.f. self-similar sets, *J. Lond. Math. Soc.* **56** (1997), 320–332.
3. M. F. BARNSLEY, J. H. ELTON AND D. P. HARDIN, Recurrent iterated function systems, *Constr. Approx.* **5** (1989), 3–31.
4. J. CROFT, Some generalisations of nested fractal constructions and associated diffusions, PhD thesis, University of Warwick (1999).
5. F. M. DEKKING, Recurrent sets, *Adv. Math.* **44** (1982), 78–104.
6. S. DUBE, Using fractal geometry for solving divide and conquer recurrences, *J. Aust. Math. Soc.* **B37** (1995), 145–171.
7. K. J. FALCONER, On the Minkowski measurability of fractals, *Proc. Am. Math. Soc.* **123** (1995), 1115–1124.

8. K. J. FALCONER, *Techniques in fractal geometry* (Wiley, 1997).
9. M. FUKUSHIMA, Dirichlet forms, diffusion processes and spectral dimensions for nested fractals, in *Ideas and methods in mathematical analysis, stochastics and applications. In memory of R. Høegh-Krohn* (ed. S. Albeverio, J. Fensted, H. Holden and T. Lindstrøm), vol. 1, pp. 151–161 (Cambridge University Press, 1992).
10. M. FUKUSHIMA AND T. SHIMA, On a spectral analysis for the Sierpinski gasket, *Potential Analysis* **1** (1992), 1–35.
11. B. M. HAMBLY, On the asymptotics of the eigenvalue counting function for random recursive Sierpinski gaskets, *Prob. Theory Relat. Fields* **117** (2000), 221–247.
12. B. M. HAMBLY AND T. KUMAGAI, Transition density estimates for diffusion processes on post critically finite self-similar fractals, *Proc. Lond. Math. Soc.* **78** (1999), 431–458.
13. J. KIGAMI, Harmonic calculus on P.C.F. self-similar sets, *Trans. Am. Math. Soc.* **335** (1993), 721–755.
14. J. KIGAMI, Harmonic calculus on limits of networks and its application to dendrites, *J. Funct. Analysis* **128** (1995), 48–86.
15. J. KIGAMI, Distributions of localized eigenvalues of Laplacians on post critically finite self-similar sets, *J. Funct. Analysis* **156** (1998), 170–198.
16. J. KIGAMI, *Analysis on fractals* (Cambridge University Press, 2001).
17. J. KIGAMI AND M. L. LAPIDUS, Weyl’s problem for the spectral distribution of the Laplacian on P.C.F. self-similar fractals, *Commun. Math. Phys.* **158** (1993), 93–125.
18. J. KIGAMI, R. STRICHARTZ AND K. C. WALKER, Constructing a Laplacian on the diamond fractal, *Experimental Math.* **10** (2001), 437–448.
19. T. KUMAGAI, Regularity closedness and spectral dimensions of the Dirichlet forms on p.c.f. self-similar sets, *J. Math. Kyoto Univ.* **33** (1993), 765–786.
20. S. KUSUOKA, Diffusion processes on nested fractals, in *Statistical mechanics and fractals* (ed. R. L. Dobrushin and S. Kusuoka), Lecture Notes in Mathematics, vol. 1567 (Springer, 1993).
21. M. L. LAPIDUS, Vibrations of fractal drums, the Riemann hypothesis, waves in fractal media and the Weyl–Berry conjecture, in *Ordinary and partial differential equations, vol. IV, Proc. 12th Int. Conf. on Theory of Ordinary and Partial Differential Equations, Dundee, 1992*, Research Notes in Mathematics, pp. 126–209 (Longman, London, 1993).
22. K. S. LAU, J. WANG AND C. H. CHU, Vector valued Choquet–Deny theorem, renewal equation and self-similar measures, *Studia Math.* **117** (1995), 1–28.
23. M. LEVITIN AND D. VASSILIEV, Spectral asymptotics, renewal theorem and the Berry conjecture for a class of fractals, *Proc. Lond. Math. Soc.* **72** (1996), 188–214.
24. T. LINDSTRØM, *Brownian motion on nested fractals*, Memoirs of the American Mathematical Society, vol. 83, no. 420 (1990).
25. R. D. MAULDIN AND S. C. WILLIAMS, Hausdorff dimension in graph directed constructions, *Trans. Am. Math. Soc.* **309** (1988), 811–829.
26. V. METZ, ‘Laplacians’ on finitely ramified, graph directed fractals, preprint (2002).
27. J. MURAI, Diffusion processes on mandala, *Osaka J. Math.* **32** (1995), 887–917.
28. L. OLSEN, *Multifractal spectrum for graph directed random recursive self-similar measures* (Longman, London, 1996).
29. T. A. RYAN, A multidimensional renewal theorem, *Ann. Prob.* **4** (1976), 656–661.
30. C. SABOT, Existence and uniqueness of diffusions on finitely ramified self-similar fractals, *Annls Sci. Ec. Norm. Super.* **30** (1997), 605–673.
31. J. L. WANG, The open set condition for graph directed self-similar sets. *Random Comput. Dynam.* **5** (1997), 283–305.