

Fire activity as a function of fire–weather seasonal severity and antecedent climate across spatial scales in southern Europe and Pacific western USA

This content has been downloaded from IOPscience. Please scroll down to see the full text.

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 161.111.180.191

This content was downloaded on 08/11/2016 at 13:22

Please note that [terms and conditions apply](#).

You may also be interested in:

[Modeling very large-fire occurrences over the continental United States from weather and climate forcing](#)

R Barbero, J T Abatzoglou, E A Steel et al.

[Objective identification of multiple large fire climatologies: an application to a Mediterranean ecosystem](#)

J Ruffault, V Moron, R M Trigo et al.

[Future changes in climatic water balance determine potential for transformational shifts in Australian fire regimes](#)

Matthias M Boer, David M J S Bowman, Brett P Murphy et al.

[Controls on interannual variability in lightning-caused fire activity in the western US](#)

John T Abatzoglou, Crystal A Kolden, Jennifer K Balch et al.

[How much global burned area can be forecast on seasonal time scales using sea surface temperatures?](#)

Yang Chen, Douglas C Morton, Niels Andela et al.

[Measurement of inter- and intra-annual variability of landscape fire activity at a continental scale: the Australian case](#)

Grant J Williamson, Lynda D Prior, W Matt Jolly et al.

Environmental Research Letters



LETTER

Fire activity as a function of fire–weather seasonal severity and antecedent climate across spatial scales in southern Europe and Pacific western USA

OPEN ACCESS

RECEIVED
1 August 2015REVISED
19 October 2015ACCEPTED FOR PUBLICATION
20 October 2015PUBLISHED
6 November 2015

Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Itziar R Urbietta¹, Gonzalo Zavala¹, Joaquín Bedia², José M Gutiérrez², Jesús San Miguel-Ayanz³, Andrea Camia³, Jon E Keeley^{4,5} and José M Moreno¹¹ Departamento de Ciencias Ambientales, Universidad de Castilla La Mancha, Av. Carlos III s/n, E-45071 Toledo, Spain² Grupo de Meteorología, Instituto de Física de Cantabria, CSIC-Universidad de Cantabria, Avda. de los Castros, s/n, E-39005 Santander, Spain³ European Commission, Joint Research Centre (JRC), Institute for Environment and Sustainability, I-21027 Ispra Varese, Italy⁴ Sequoia-Kings Canyon Field Station, Western Ecological Research Center, U.S. Geological Survey, Three Rivers, CA 93271, USA⁵ Department of Ecology and Evolutionary Biology, University of California, Los Angeles, CA 90095, USAE-mail: josem.moreno@uclm.es**Keywords:** area burned, climate change, drought, fire weather index, large fires, Mediterranean ecosystemsSupplementary material for this article is available [online](#)**Abstract**

Climate has a strong influence on fire activity, varying across time and space. We analyzed the relationships between fire–weather conditions during the main fire season and antecedent water–balance conditions and fires in two Mediterranean–type regions with contrasted management histories: five southern countries of the European Union (EUMED)(all fires); the Pacific western coast of the USA (California and Oregon, PWUSA)(national forest fires). Total number of fires (≥ 1 ha), number of large fires (≥ 100 ha) and area burned were related to mean seasonal fire weather index (FWI), number of days over the 90th percentile of the FWI, and to the standardized precipitation–evapotranspiration index (SPEI) from the preceding 3 (spring) or 8 (autumn through spring) months. Calculations were made at three spatial aggregations in each area, and models related first-difference (year-to-year change) of fires and FWI/climate variables to minimize autocorrelation. An increase in mean seasonal FWI resulted in increases in the three fire variables across spatial scales in both regions. SPEI contributed little to explain fires, with few exceptions. Negative water–balance (dry) conditions from autumn through spring (SPEI8) were generally more important than positive conditions (moist) in spring (SPEI3), both of which contributed positively to fires. The R^2 of the models generally improved with increasing area of aggregation. For total number of fires and area burned, the R^2 of the models tended to decrease with increasing mean seasonal FWI. Thus, fires were more susceptible to change with climate variability in areas with less amenable conditions for fires (lower FWI) than in areas with higher mean FWI values. The relationships were similar in both regions, albeit weaker in PWUSA, probably due to the wider latitudinal gradient covered in PWUSA than in EUMED. The large variance explained by some of the models indicates that large-scale seasonal forecast could help anticipating fire activity in the investigated areas.

1. Introduction

Climate plays a major role in fires. On a short-term scale, weather closely controls fire ignition and propagation (Rothermel 1972). Daily warnings of fire danger are used by fire agencies across the world to allocate

fire suppression resources based on indices that primarily integrate meteorological variables in ways that reflect the main processes affecting fire ignition, behavior, and difficulty of fire control (Fujioka *et al* 2008, Dowdy *et al* 2010). Numerous studies have related mean fire–weather values of various periods

(the fire season being the most common), alone or combined with other climate variables of the relevant period to explain fire activity (e.g., Preisler *et al* 2008, Koutsias *et al* 2013, Riley *et al* 2013). While many studies emphasize the importance of climate during the fire season in determining fire activity (e.g., Koutsias *et al* 2012, Morton *et al* 2013), antecedent climate (i.e., conditions that occur months or years before the fire season) has been shown to also contribute to fire occurrence (Dimitrakopoulos *et al* 2011), area burned (Trouet *et al* 2006, Archibald *et al* 2010a, Abatzoglou and Kolden 2013), and large fires (Keeley and Zedler 2009, Barbero *et al* 2014), through its influence in fuel load and flammability of live and dead fuels (e.g., Castro *et al* 2003, Dennison *et al* 2008).

On a longer term, climate has a strong but complex influence on fires because it determines primary productivity and vegetation, hence fuels, all of which determine global fire patterns (Krawchuk *et al* 2009, Archibald *et al* 2013). The study of the longer-term climate controls on fires has been recently framed in a context of the fuel amount versus fuel moisture limitations paradigm (Meyn *et al* 2007, Krawchuk and Moritz 2011, Pausas and Ribeiro 2013). According to this, in more arid areas, with less favorable climate for plant growth, low primary productivity more than weather conditions will limit fuels and fires. At the other end of the spectrum, in areas with favorable climate for plant growth, fires may be limited by unfavorable weather conditions for them to ignite and spread owing to excess moisture. Consequently, changes in climate from year to year can differentially affect areas with different controls on fires. Moreover, areas that encompass a variety of vegetation types due to climate could also be differentially responsive to changes in climate variability, depending on what factors most limit fires. Notwithstanding, changes in local conditions (i.e., particular strong winds) (Moritz *et al* 2010) and in the environmental and sociological contexts can occur across the geographic space as well, all of which can modify climate-related controls on fires beyond the fuels/conditions limitations paradigm (Loepfe *et al* 2014, Parks *et al* 2014). As geographic scale increases, the probability of having areas responsive to different climate controls is also likely to increase, hence the relative importance of the various factors affecting fire can vary with the scale (Falk *et al* 2007, Parisien and Moritz 2009, Archibald *et al* 2010b). All of this complicates anticipating how the fire–climate relationships will change when moving from smaller to larger areas or across areas through climate and other gradients.


Here we investigated the relationships between fire activity during the main fire season and climatic conditions during that season and the preceding months for the 1985–2011 period in two Mediterranean-type climate areas that have had different management histories: southern European Union (EUMED) and the

Pacific western USA (Oregon and California, PWUSA). Analyses were done at three spatial scales, from provincial level in Spain/national forests in PWUSA, to subregions in both EUMED/PWUSA, and whole region in each continent. By comparing southern Europe, including fires in all its territory, and PWUSA, focusing on national forest lands, would permit assessing climate controls in two most contrasted territories, sharing a largely-similar climate.

2. Methods

Models of fire activity (total number of fires [≥ 1 ha], number of large fires [≥ 100 ha] and area burned by all fires [≥ 1 ha]) as a function of within-season and antecedent climate conditions were developed for the period 1985–2011 in two regions: the southern countries of the European Union (Portugal, Spain, South France, Italy and Greece; EUMED) (figure 1(a)), and the two southern states of the Pacific western coast of the USA (Oregon and California, PWUSA) (figure 1(b)). EUMED (Latitude: 43.11 to 36.12, Longitude: -9.28 to 28.08 in decimal degrees) covered a wider longitudinal gradient than PWUSA (Latitude: 46.28 to 32.60 , Longitude: -124.0 to -116.27), which represented a longer latitudinal gradient.

Fire–climate models were developed at three geographical scales in each study region: small ($10\,900 \pm 4500$ km² and $11\,850 \pm 3400$ km², (mean \pm SD)), medium ($54\,000 \pm 31\,700$ km² and $23\,700 \pm 6750$ km²), and large ($1188\,010$ km² and $166\,000$ km² for EUMED and PWUSA, respectively) (figure 1). At EUMED, calculations at the smaller scale were made for 47 provinces (i.e., NUTS3—nomenclature of territorial units for statistics—level) (EUROSTAT 2011) at Peninsular Spain based on daily fire statistics (EGIF, Spanish National Forest Fire Statistics, Spanish Ministry of Agriculture, Food and the Environment). In the medium scale, we modeled 21 subregions (proxy of NUTS2) of the different southern EU countries with monthly fire statistics, using the European Fire Database stored at the Joint Research Centre of the European Commission (Camia *et al* 2014), and then aggregated at the larger scale for the whole EUMED (figure 1(a)). At PWUSA, calculations were made using national forests fire data (FAM-WEB Forest Service state data http://fam.nwcg.gov/fam-web/weatherfirecd/state_data.htm, last accessed 5 February 2015). At the smallest scale, we defined 15 areas using single national forests or, when these were too small, aggregations of several of them to reach an approximate equivalent area to the Spanish provinces. For the medium scale, we aggregated national forest data into 7 subregions following US climatic divisions boundaries for Oregon and California (www.esrl.noaa.gov/psd/data/usclimdivs/boundaries.html, last accessed 5 February 2015) (figure 1(b)). Los Padres National Forest was treated as South California



subregion, although one part is located in the Central Coast. Finally, for the larger scale, data were aggregated for the whole PWUSA. Despite similarities in climate between both regions, there were some differences between them that need to be mentioned. At EUMED, the whole territory was used, and thus all fires, the majority of which are caused by people (on average 7% of area burned by lightning, 58% human-caused, and 35% unknown which most probably correspond to human-causes), albeit with some

differences among the various areas (see Vázquez and Moreno 1998 for data from Spain). At PWUSA, only national forests territory, (which amounted to 25% of Oregon and California), and corresponding fires, were analyzed. National forest fires data have been often used in studies assessing large scale patterns of relationships between climate and fire activity (Gedalof *et al* 2005, Trouet *et al* 2006, 2009, Keeley and Syphard 2015). While it has been shown that the inter-annual variation in fire activity in national forests is


correlated with that at the state level for California (Trouet *et al* 2006), differences in climate drivers of fires between forest and non-forest areas have been documented (e.g., Keeley and Syphard 2015). In Oregon, national forests territory covers a diverse range of fire regimes (Agee 1993, Gedalof *et al* 2005), but excludes the grassland fires of the Intermountain West (e.g., Knapp 1998). Thus, while national forests cover a diverse range of ecosystems, yet they may not capture the full range of them and of other factors driving fires in the study area. Forest fires in national forest of PWUSA are caused on average by lightning (36% of area burned) and human causes (64%), with differences between interior mountains, where lightning-ignited fires become more important than along coastal subregions where human-caused fires dominate.

A potential problem when comparing two time series is the possibility of autocorrelation. One means of dealing with such autocorrelations is with the time series technique of differencing, where the value of the variable at time $t - 1$ is subtracted from the variable at time t (Lobell and Field 2007, Trouet *et al* 2009, Turco *et al* 2014, Bedia *et al* 2015). Thus, the relationship between fire activity and climate conditions was ascertained by means of multiple linear regressions between first-difference time series of fire and climate-related variables. The period of study (1985–2011) was common to both regions and for all three scales, as well as the fire season (June to September, from now on abbreviated JJAS) except for South California (Los Padres, Angeles, San Bernardino and Cleveland national forests), in which October fires were included (i.e., JJASO) due to the presence of large fires in some years. This fire season amounted, on average, to 70%, 83% and 82% of number of fires, large fires and area burned, respectively, in EUMED, and 81%, 92% and 97%, respectively, in PWUSA.

To set the modeling framework, we first calculated the yearly change, in percentage, of each response variable (i.e., number of fires (≥ 1 ha), number of large fires (≥ 100 ha) and area burned by all fires ≥ 1 ha). Similarly, we characterized within fire-season conditions using the fire weather index (FWI) of the Canadian Forest Fire Danger Rating System (Van Wagner 1987), which is calculated based on daily meteorological conditions at noon time for air temperature and relative humidity, wind speed, and rainfall accumulated over 24 h. From the FWI time series we calculated the corresponding absolute increment, or yearly delta, of the mean FWI (FWIm) conditions experienced in the fire season. Prior to the modeling exercise, trends in the three metrics of fire activity and FWIm time series data at medium and large scales were analyzed using the Mann–Kendall trend-test, using the R package ‘Kendall v.2.2’ (Hipel and McLeod 2005, R Development Core Team 2013). Furthermore, we also tested the influence of extreme FWI events on fire activity, calculating the delta in the

number of days over the 90th percentile of the FWI historical series for each area (dFWI90). FWI daily values were calculated from WATCH-Forcing-Data-ERA-Interim at 50 km (Weedon *et al* 2010) applied to ERA-Interim data (Dee *et al* 2011), following Bedia *et al* (2012). Finally, in order to test the influence of antecedent climate on fires, we extracted the standardized precipitation-evapotranspiration index (SPEI) (Vicente-Serrano *et al* 2010) from the Global 0.5° gridded SPEI dataset (Beguería *et al* 2014), which indicates the deviations of the current (e.g., period of reference) water-balance (precipitation minus potential evapotranspiration) with respect to the long-term water-balance, with time-scales between 1 and 48 months. Specifically, for each spatial unit of analysis at each scale we sampled the SPEI in May (i.e., to reflect the water-balance conditions accumulated just before the fire season) for two time scales: 3-months SPEI (March to May, SPEI3) to cover spring months, and 8-months SPEI (October to May, SPEI8) to capture the preceding autumn through spring conditions. We calculated the corresponding absolute increment, or yearly delta, for SPEI3 and SPEI8 with positive values indicating an increase in moisture and negative indicating an increase in drought conditions.

A forward stepwise regression analysis was performed with the pool of potential predictors (FWIm, dFWI90, SPEI3, SPEI8), using the R package ‘stats v.3.0.2’ (R Development Core Team 2013). We tested combinations of all possible predictors, maintaining those that were significant ($p < 0.05$) and added variance explained of the fire variables. Goodness of fit of the models was assessed by maximum log-likelihood, explained variance (adjusted R^2), and examining the model residuals. Resulting models for each fire variable were compared using Akaike’s information criterion (AIC) (Akaike 1992). The model with the minimum AIC was selected as the best model. If two models showed a difference in AIC between 0 and 2, these were considered to have equivalent empirical support (Burnham and Anderson 2002), and the simplest one was selected as the best model. The high variability in the fire data in a few small areas led us to carefully examine outliers. These were investigated using the Bonferroni outlier test in ‘car v. 2.0.19’ package (Fox and Weisberg 2011), omitting those values at $p < 0.001$, and after rigorous examination of residual plots. The reasons for such highly deviant years, when present, was beyond our primary interest, and were not further investigated. Additionally, to confirm that results were not sensitive to different assumptions of regressing first-differences, we checked the alternative technique of regressing the detrended residuals (Nicholls 1997) in a number of areas and for various scales, which produced qualitatively similar results (not shown). Before calculations, fire data were transformed to meet linear regression assumptions: number of fires (≥ 1 ha, and ≥ 100 ha) was square-root transformed, and area burned log-transformed.



Finally, to ascertain how climate controls on fires vary in relation to the prevailing climate conditions across areas, for both small and medium scales in both continents, we linearly related the R^2 of the best climate–fire model obtained in each particular area (for each of the three fire variables) with the long-term mean seasonal FWI of the respective area.


3. Results

During the 1985–2011 period, significant decreasing trends were found at the whole EUMED for the number of fires ($\tau = -0.45$, $p < 0.01$), number of large fires ($\tau = -0.40$, $p < 0.01$), and marginally for area burned ($\tau = -0.27$, $p = 0.058$) (figure 2). Similarly, decreasing trends were found for several EUMED subregions, particularly in Spain, Italy and Greece (table A1). Italian Islands were the only subregion showing an increase in the three fire variables ($p < 0.05$). On the other hand, no trend was found at the whole PWUSA (figure 2), and only few subregions showed significant trends in fire activity (table A2). While the number of fires significantly decreased in the South East and South of California ($p < 0.05$), an increasing trend was found

for large fires and area burned in Northern Cascades ($p < 0.05$), and marginally for area burned in the South California subregion. In mean seasonal FWI no significant trends were detected for any of the two regions; at the subregional scales, positive FWIm trends were found in Spain East ($\tau = 0.30$, $p < 0.05$) and Spain South ($\tau = 0.40$, $p < 0.01$) at EUMED, and in the Sacramento Drainage ($\tau = 0.40$, $p < 0.01$) at PWUSA (appendix A).

At the smallest scale of provinces and national forests, fire activity was moderately explained by both, within season FWI and antecedent climate, and for both regions, EUMED and PWUSA. This was true for the three fire variables tested (number of fires, number of large fires and area burned), with minor differences among them within and between regions (figure 3(a)). Only a few (5 provinces and 3 national forests) did not show significant relationships to within season FWI conditions and antecedent climate for any of the three fire variables (tables B1 and B2).

At the medium (i.e., sub-regional) scale, the variance explained by the models increased, more so in EUMED than in PWUSA (figure 3(b)). Differences between fire variables within regions were small. Note the high values obtained in some subregions at



EUMED ($R^2 > 0.70$) (Spain North East, France South West and East, Italy North, Center and South), and at PWUSA ($R^2 > 0.50$) (Northern Cascades, North Coast, South California) (tables C1 and C2).

At the largest scale (i.e., whole region), relationships were still higher and similar in both regions and for the three fire variables, albeit with some differences. All variables were highly predictable (R^2 : 0.46–0.63) in both regions, except for area burned that was less predictable in PWUSA (R^2 : 0.37) (figure 3(c), tables C1 and C2).

Most of the variability at either provinces/national forests or subregions was explained by within-season conditions, with an increase in mean seasonal FWI resulting in increases in the three fire variables (figure 4, appendices B and C). The number of days over the 90th percentile of the FWI (dFWI90) did not enter in any of the best models, and in many instances was highly correlated with mean FWI. Antecedent climate contributed to the models only at a few provinces/national forests or subregions in both continents (appendices B and C). When it did, 3-months SPEI contributed positively (a positive water-balance in spring, i.e., moist conditions, led to more fires and larger burned areas), while 8-months SPEI contributed negatively (i.e., autumn through spring dry conditions led to increased fire activity, with more fires and larger burned areas). While the mean contribution of these variables to the models was small (figure 4), in some cases they contributed quite markedly. This was mainly due to the high contribution of SPEI8, more so at subregional scale (tables C1 and C2): for number of fires (e.g., Spain North), large fires (e.g., Spain East, France South East and West, Central Italy and Greece), and area burned (e.g., France South East and West) in EUMED, and number of fires (e.g., North Coast, Central High Plateau), area burned (e.g., Northern Cascades) and large fires (e.g., southern California) in PWUSA. SPEI3 (spring wet conditions) showed a positive effect on fires only at EUMED, and more so at the small scale (table B1). At the scale of whole region (both, EUMED and PWUSA) no other variable than within-season mean FWI contributed to the models (tables C1 and C2).

Model performance was not homogeneous across the areas studied within each continent. We found a negative relationship, albeit weak, between the R^2 values of the best fire-climate models and the long-term mean FWI of the various areas. This was so for the number of fires at both scales and continents, and for area burned at PWUSA small scale and EUMED subregions (table 1).

4. Discussion

Mean seasonal fire-weather proved critical for fire activity in both regions, with minor influence of longer-term antecedent water-balance climatic conditions. Years with higher fire danger FWI values in summer corresponded to years with increased number of fires (total and large fires) and area burned, and vice versa. The rather strong relationship at the largest scale, with R^2 values that explain almost 60% of the total variance for area burned at EUMED, and above 60% for number of fires and large fires at PWUSA, indicates that fire-climate relationships have been very strong during the nearly three last decades. The variance explained by our models was comparable to that obtained in other studies that used a number of




Figure 4. Average relative contribution (as percentage) of within-season (mean FWI) and antecedent (SPEI3, SPEI8) climatic conditions to number of fires (NF), number of large fires (NLF), and area burned (AB) at small (a), (b) and medium (c), (d) spatial scales, for EUMED and PWUSA.

Table 1. Results of the models fitted between the R^2 values of the best fire–climate models and the long-term mean FWI for EUMED and PWUSA.

	Number of fires			Area burned		
	Intercept	Slope	R^2 (p)	Intercept	Slope	R^2 (p)
Small scale						
EUMED Spanish provinces	0.625	−0.010	0.26 (***)	—	—	n.s
PWUSA national forests	0.615	−0.008	0.21 (*)	0.680	−0.011	0.42 (**)
Medium scale						
EUMED subregions	0.914	−0.015	0.40 (**)	0.782	−0.011	0.15 (*)
PWUSA subregions	1.230	−0.022	0.58 (*)	—	—	n.s

variables for characterizing climate controls on fires (e.g., Trouet *et al* 2009, Amatulli *et al* 2013). The fact that much of the variance in the various fire variables was explained by the mean seasonal FWI implies that a simple, integrated measure of weather conditions during the fire season carries substantial information for anticipating fire activity. These findings are important considering the large socio-economic differences between the two regions, and the contrasted changes over time occurred during the study period, with consequent impacts on landscapes, fuels, ignitions patterns or firefighting, among other. For instance, in EUMED the dominant trend has been rural exodus, abandonment, afforestation and concentration of

population in cities (e.g., Viedma *et al* 2015), while in PWUSA the dominant trend has been increasing urbanization (e.g., Syphard *et al* 2007). National forests may have been indirectly affected by such changes, but not their landscapes on a comparable magnitude to what has happened in EUMED. Nonetheless, in some areas a substantial amount of variance was not explained by climate conditions, thus other factors (e.g., socio-economic, ignition patterns, fire management, etc) are also relevant.

Some studies use conversions of FWI values (i.e., daily, monthly or seasonal severity ratings) to enhance the role of high FWI days (Van Wagner 1970). In our approach, we used raw FWI daily values, without

further additional conversion to daily or monthly severity values. We tested days over the 90th percentile of the FWI to account for possible additional effects of extreme events. Nonetheless, mean FWI seasonal values and days above the 90th percentile were highly correlated, and mean FWI values were the best predictor kept by the models.

Despite previous reports on the significant role of antecedent water-balance conditions on fires in both regions using different approaches (e.g., Keeley 2004, Gedalof *et al* 2005, Riley *et al* 2013, Gudmundsson *et al* 2014, Keeley and Syphard 2015, Marcos *et al* 2015, Sousa *et al* 2015), adding this information did not generally improve our models, except in a few areas in which such variables proved to be relevant. Negative water-balance (i.e., dry) conditions from autumn through spring (SPEI8) were generally more important than positive conditions (i.e., moist) in spring (SPEI3) in contributing to increase fires. The fact that only in a few areas antecedent water-balance was important does not permit any generalization with regard to where (i.e., more or less humid areas) this was more relevant. For example, increasing negative SPEI8 in the antecedent months was important for area burned and number of large fires in southern France, a rather mesic area, but also for number of large fires in southern California, a much drier area. Similarly, while wet conditions in spring (SPEI3) contributed positively to fires in a few areas (e.g., some Spanish provinces) probably due to increasing fine fuel load during the summer months, there was no consistent pattern as to under which dominant climate conditions this variable was more important. Notwithstanding, it must be pointed out that the FWI index also includes drought code as an indicator of seasonal drought effects, which has a 52-day time lag in moderate weather conditions (Lawson and Armitage 2008) and correlates well with long-term live-plant moisture content (e.g., Pellizzaro *et al* 2007). This means that FWI partly carries antecedent water-balance information and that only in a few areas variations in this were sufficiently important or long in time to exceed what was embedded in the FWI index.

Trends in fire danger (mean FWI) during the period analyzed were non-significant, except in a few subregions in both EUMED and PWUSA. However, positive trends in fire danger have been reported in both EUMED (Venäläinen *et al* 2014) and PWUSA (Abatzoglou and Kolden 2013). This is probably due to the shorter period analyzed and different areas of reference, which signals to the sensitivity of trend analysis to the varying database. Trends in fire activity have also been variable among both regions and subregions within them. We found a clear decreasing trend at EUMED (for the three variables and both at large and medium scales), consistent with other studies (San-Miguel-Ayanz *et al* 2013). By contrast, in PWUSA, trends in fires were generally not significant, except in some subregions. While a number of studies

have found an increase in number of fires and the area burned (Keeley *et al* 1999) and in the frequency of large fires in the Western USA (Dennison *et al* 2014), the different periods used and the fact that our data only considered national forests can explain these apparent contradictions.

Despite contrasted trends in both regions, we found significant fire–climate relationships across scales and for the various fire variables. Relationships were detected at the smaller scale investigated, but the R^2 of the relationships generally increased with a growing area of study. We argue that the variability of fire and climate data becomes attenuated as the level of aggregation increases, hence allowing the relationship to emerge more clearly. Aggregating areas has shortcomings since it is possible that trends in the areas aggregated at a higher level might differ from one another, and what is true at the larger level may not necessarily apply at a lower level of aggregation (Parisien *et al* 2011). Therefore, each level of aggregation needs to be evaluated on its own.

Our results also document that, in general, the goodness of fit of the models (the R^2) tended to decrease across areas within a region for some fire variables (i.e., number of fires and area burned) and at certain scales. The pattern was such that followed a climatic gradient, suggesting that areas with lower long-term mean FWI (i.e., milder climatic conditions during the summer) were more responsive to year-to-year changes in climate than those with higher values (i.e., more severe climatic conditions during summer). Similar results have been reported by other authors in both regions (e.g., Vázquez and Moreno 1995, Trouet *et al* 2009, Dimitrakopoulos *et al* 2011). Even if the patterns across the climatic gradient were weak (e.g., not all variables, not all scales), this is partially consistent with the fuels-moisture limitations hypothesis (Pausas and Ribeiro 2013, Parks *et al* 2014). The net result is reduced predictability at the more severe end of the gradient. A saturating performance of FWI index in the upper end of its spectrum of values could also play a role, but that needs further investigation.

The fire–climate relationships that emerged were rather similar, even if weaker in PWUSA at medium scale, and particularly for area burned at the largest scale. While both regions are to a certain degree comparable, the fact that we have covered a wider longitudinal gradient in EUMED (with all fires) and greater latitudinal gradient in PWUSA (only national forest fires) could account for some of the differences. At EUMED we explored more areas that are not that extreme, so the models performed better. The greater latitudinal gradient in PWUSA implies that we covered areas with substantial differences in climate. Several studies have pointed the differential impact of climate on fires across the western US (Gedalof *et al* 2005, Littell *et al* 2009, Stavros *et al* 2014) and between forested and non-forested landscapes (Abatzoglou and Kolden 2013, Keeley and Syphard 2015).

Large scale atmospheric circulation patterns exhibit a pivot point in California, shifting south or north on decadal time-scales that may cause changes in the spatial association of climatic variations and fire activity (Westerling and Swetnam 2003, Trouet *et al* 2006). The longer fire season in southern California, and hence more diverse set of atmospheric drivers, in addition to increased human ignitions compared to the northwest (Miller *et al* 2012), likely contributes to weaken climate–fire linkages (Trouet *et al* 2009, Abatzoglou and Kolden 2013). Moreover, the significant role of local conditions such as Santa Ana winds in area burned in southern California (Keeley and Zedler 2009, Moritz *et al* 2010, Jin *et al* 2014), but not in northern California and Oregon, is causing a much greater inter-annual variability that was not fully accounted for in our models.

Determining the influence of variability and change in climate on fires in the past is important for projecting future climate change impacts (Bowman *et al* 2014, Moritz *et al* 2014). Nonetheless, attributing changes in fires to climate is difficult due to role of confounding factors that affect fire–climate relationships (Cramer *et al* 2014). When using long-term data series for the analysis of climate–fire relationships, studies should use proper analytical procedures or consider the role of other fire-drivers (fuels, landscape, fire management, etc) (Littell *et al* 2009). Relating year-to-year changes in fires to similar changes in the climate-related variables permits arguing that, in such a short period of time and at the spatial scales investigated here, climate with its underlying weather is the main variable that could significantly change. If at the smaller scale it is still conceivable that certain changes from one year to the next could meaningfully affect this relationship (Miller *et al* 2009, Brotons *et al* 2013, Price *et al* 2015), it is most unlikely that significant changes would affect the relationship at the largest scale investigated. For instance, landscapes are dynamic through time, but significant changes in landscape hazard from year to year due to human activities are unlikely. Similar temporal framing can be assumed for other drivers, like changes in ignitions patterns or on fire management, which may take years to unfold. While the effect of the other drivers is difficult to quantify, on such a small temporal window it can be assumed to be much smaller than that of weather/climate. Actually, this implicit assumption is used by the fire services, whose alerts use a scale of indices that only change due to weather, despite trends through time in other non-climate variables affecting fires.

While extrapolating these relationships to a future with more severe regional conditions due to global warming (Westerling *et al* 2011, Bedia *et al* 2014) must be made with caution, it would signal to a clear impact of climate change on fires, at least before changes in fires or other drivers can significantly affect current landscapes and vegetation. Our findings are important in a context of forecasting large scale climate patterns

(Quesada *et al* 2012), as a basis to anticipate fire-seasonal severity. Since the range of FWI studied is quite large, this supports arguing that in some areas an increase in average FWI in the next decades or so, as a consequence of climate change, can result in increased fire activity. Our results also suggest that as the climate becomes more severe due to global warming, fires in these regions are likely to become less predictable, a finding that can have important implications for future fire prevention and planning.

5. Conclusion

Fire activity in EUMED and PWUSA at the three scales investigated was largely explained by within seasonal climate and, in a much lesser extent, by antecedent 3 to 8 months water-balance climatic conditions. Mean seasonal FWI was the most important variable, whereas antecedent water-balance did not contribute much to the models, except in a few areas. In any case, when they did, negative water-balance (i.e., drought) in the antecedent 8 months was more important, and contributed to increase fire activity. Despite differences among the two regions in land-use history, management and type of territory used (whole territory at EUMED, only national forests at PWUSA), the relationships were similar in both regions, albeit weaker, notably for area burned at the whole region level, in PWUSA probably due to the wider latitudinal gradient covered. The relationships tended to increase with the area of aggregation. The rather large variance explained by some of the models indicates that large-scale seasonal forecast could help anticipating the fire season in the investigated areas. The R^2 of the models for some fire-variables decreased in both regions as the long-term mean FWI of the area increased. That is, in areas with lower long-term mean FWI (i.e., less amenable conditions for fires), fires were more susceptible to change with climate variability across the years than in areas with higher long-term mean FWI values (i.e., more amenable to fires). Provided that this trend reflects varied sensitivity to climate under more severe conditions, that would imply a reduced predictability of fire risk with climate change and consequent increases in mean seasonal FWI.

Acknowledgments

We thank Sixto Herrera for his assistance in climate data downloading and pre-processing. We acknowledge the helpful feedback from Julie Yee, Steve Cumming, and two anonymous reviewers to previous versions of the manuscript. This research was funded by the European Union's Seventh Framework Programme (FP7/2007-2013) under grant agreement 243888 (FUME Project). Publication in Open Access was funded by the EC FP7 Post-Grant Access Pilot. Any use of trade, product or firm names is for

descriptive purposes only and does not imply endorsement by the U.S. government.

References

- Abatzoglou J T and Kolden C A 2013 Relationships between climate and macroscale area burned in the western United States *Int. J. Wildland Fire* **22** 1003–20
- Agee J K 1993 *Fire Ecology of Pacific Northwest Forests* (Washington DC: Island Press)
- Akaike H 1992 Information theory and an extension of the maximum likelihood principle *Breakthroughs in Statistics* ed S Kotz and N Johnson (New York: Springer) pp 610–24
- Amatulli G, Camia A and San-Miguel-Ayanz J 2013 Estimating future burned areas under changing climate in the EU-Mediterranean countries *Sci. Total Environ.* **450–451** 209–22
- Archibald S, Lehmann C E R, Gómez-Dans J L and Bradstock R A 2013 Defining pyromes and global syndromes of fire regimes *Proc. Natl Acad. Sci. USA* **110** 6442–7
- Archibald S, Nickless A, Govender N, Scholes R J and Lehsten V 2010b Climate and the inter-annual variability of fire in southern Africa: a meta-analysis using long-term field data and satellite-derived burnt area data *Global Ecol. Biogeogr.* **19** 794–809
- Archibald S, Nickless A, Scholes R J and Schulze R 2010a Methods to determine the impact of rainfall on fuels and burned area in southern African savannas *Int. J. Wildland Fire* **19** 774–82
- Barbero R, Abatzoglou J T, Steel E A and Larkin N K 2014 Modeling very large-fire occurrences over the continental United States from weather and climate forcing *Environ. Res. Lett.* **9** 124009
- Bedia J, Herrera S, Camia A, Moreno J M and Gutiérrez J M 2014 Forest fire danger projections in the mediterranean using ENSEMBLES regional climate change scenarios *Clim. Change* **122** 185–99
- Bedia J, Herrera S, Gutiérrez J M, Benali A, Brands S, Mota B and Moreno J M 2015 Global patterns in the sensitivity of burned area to fire-weather: implications for climate change *Agric. Forest Meteorol.* **214–215** 369–79
- Bedia J, Herrera S, Gutiérrez J M, Zavala G, Urbietta I R and Moreno J M 2012 Sensitivity of fire weather index to different reanalysis products in the Iberian Peninsula *Nat. Hazard Earth Syst.* **12** 699–708
- Beguieria S, Vicente-Serrano S, Reig F and Latorre B 2014 Standardized precipitation evapotranspiration index (SPEI) revisited: parameter fitting, evapotranspiration models, tools, datasets and drought monitoring *Int. J. Climatol.* **34** 3001–23
- Bowman D M J S, Murphy B P, Williamson G J and Cochrane M A 2014 Pyrogeographic models, feedbacks and the future of global fire regimes *Glob. Ecol. Biogeogr.* **23** 821–4
- Brotans L, Aquilué N, de Cáceres M, Fortin M-J and Fall A 2013 How fire history, fire suppression practices and climate change affect wildfire regimes in mediterranean landscapes *Plos One* **8** e62392
- Burnham K P and Anderson D R 2002 *Model Selection and Multimodel Inference: a Practical Information-Theoretic Approach* 2nd edn (New York: Springer)
- Camia A, Durrant T and San-Miguel-Ayanz J 2014 The European fire database: technical specifications and data submission EUR26546 EN Publications Office of the European Union, Luxembourg p 23 (https://ec.europa.eu/jrc/sites/default/files/eudb_tech_spec_final_2register.pdf)
- Castro F X, Tudela A and Sebastià M T 2003 Modeling moisture content in shrubs to predict fire risk in Catalonia (Spain) *Agric. Forest Meteorol.* **116** 49–59
- Cramer W, Yohe G, Auffhammer M, Huggel C, Molau U, Faus da Silva Dias M A, Solow A, Stone D and Tibig L 2014 Detection and attribution of observed impacts *Climate Change 2014: Impacts, Adaptation, and Vulnerability: IPCC Working Group II Contribution to AR5* ch 18, p 94 (https://ipcc-wg2.gov/AR5/images/uploads/WGIAR5-Chap18_FINAL.pdf)
- Dee D P et al 2011 The ERA-Interim reanalysis: configuration and performance of the data assimilation system *Q. J. R. Meteorol. Soc.* **137** 553–97
- Dennison P E, Brewer S C, Arnold J D and Moritz M A 2014 Large wildfire trends in the western United States, 1984–2011 *Geophys. Res. Lett.* **41** 2928–33
- Dennison P E, Moritz M A and Taylor R S 2008 Evaluating predictive models of critical live fuel moisture in the Santa Monica Mountains, California *Int. J. Wildland Fire* **17** 18–27
- Dimitrakopoulos A P, Vlahou M, Anagnostopoulou C G and Mitsopoulos I D 2011 Impact of drought on wildland fires in Greece: implications of climatic change? *Clim. Change* **109** 331–47
- Dowdy A J, Mills G A, Finkele K and de Groot W 2010 Index sensitivity analysis applied to the Canadian forest fire weather index and the McArthur forest fire danger index *Meteorol. Appl.* **17** 298–312
- EUROSTAT 2011 Regions in the European Union Nomenclature of territorial units for statistics *NUTS 2010/EU-27* Publications Office of the European Union, Luxembourg (<http://ec.europa.eu/eurostat/documents/3859598/5916917/KS-RA-11-011-EN.PDF>)
- Falk D, Miller C, McKenzie D and Black A 2007 Cross-scale analysis of fire regimes *Ecosystems* **10** 809–23
- Fox J and Weisberg S 2011 *An R Companion to Applied Regression* 2nd edn (Thousand Oaks CA: Sage) (<http://socserv.socsci.mcmaster.ca/jfox/Books/Companion>)
- Fujioka F M, Gill A M, Viegas D X and Wotton B M 2008 Fire danger and fire behavior modeling systems in Australia, Europe, and North America *Wildland Fires and Air Pollution Developments in Environmental Science* ed A Bytnerowicz et al (Amsterdam: Elsevier) pp 471–97
- Gedalof Z, Peterson D L and Mantua N J 2005 Atmospheric, climatic, and ecological controls on extreme wildfire years in the Northwestern United States *Ecol. Appl.* **15** 154–74
- Gudmundsson L, Rego F C, Rocha M and Seneviratne S I 2014 Predicting above normal wildfire activity in southern Europe as a function of meteorological drought *Environ. Res. Lett.* **9** 084008
- Hipel K W and McLeod A I 2005 *Time Series Modelling of Water Resources and Environmental Systems* (Amsterdam: Elsevier)
- Jin Y, Randerson J T, Faivre N, Capps S, Hall A and Goulden M L 2014 Contrasting controls on wildland fires in Southern California during periods with and without Santa Ana winds *J. Geophys. Res. Biogeogr.* **119** 432–50
- Keeley J E 2004 Impact of antecedent climate on fire regimes in coastal California *Int. J. Wildland Fire* **13** 173–82
- Keeley J E, Fotheringham C J and Morais M 1999 Reexamining fire suppression impacts on brushland fire regimes *Science* **284** 1829–32
- Keeley J E and Syphard A D 2015 Different fire-climate relationships on forested and non-forested landscapes in the Sierra Nevada ecoregion *Int. J. Wildland Fire* **24** 27–36
- Keeley J E and Zedler P H 2009 Large, high-intensity fire events in Southern California shrublands: debunking the fine-grain age patch model *Ecol. Appl.* **19** 69–94
- Knapp P A 1998 Spatio-temporal patterns of large grassland fires in the Intermountain West, USA *Glob. Ecol. Biogeogr. Lett.* **7** 259–72
- Krawchuk M A, Moritz M A, Parisien M A, Van Dorn J and Hayhoe K 2009 Global pyrogeography: the current and future distribution of wildfire *Plos One* **4** e5102
- Krawchuk M A and Moritz M A 2011 Constraints on global fire activity vary across a resource gradient *Ecology* **92** 121–32
- Koutsias N, Arianoutsou M, Kallimanis A S, Mallinis G, Halley J M and Dimopoulos P 2012 Where did the fires burn in Peloponnisos, Greece the summer of 2007? Evidence for a synergy of fuel and weather *Agric. Forest Meteorol.* **156** 41–53
- Koutsias N, Xanthopoulos G, Founda D, Xystrakis F, Nioti F, Pleniou M, Mallinis G and Arianoutsou M 2013 On the relationships between forest fires and weather conditions in Greece from long-term national observations (1894–2010) *Int. J. Wildland Fire* **22** 493–507

- Lawson B D and Armitage O B 2008 *Weather Guide for the Canadian Forest Fire Danger Rating System* (Edmonton, Canada: Canadian Forest Service) p 73
- Littell J S, McKenzie D, Peterson D L and Westerling A L 2009 Climate and wildfire area burned in western US ecoprovinces, 1916–2003 *Ecol. Appl.* **19** 1003–21
- Lobell D B and Field C B 2007 Global scale climate-crop yield relationships and the impacts of recent warming *Environ. Res. Lett.* **2** 014002
- Loepfe L, Rodrigo A and Lloret F 2014 Two thresholds determine climatic control of forest fire size in Europe and northern Africa *Reg. Environ. Change* **14** 1395–404
- Marcos R, Turco M, Bedia J, Llasat M C and Provenzale A 2015 Seasonal predictability of summer fires in a Mediterranean environment *Int. J. Wildland Fire* at press (doi:10.1071/WF15079)
- Meyn A, White P S, Buhk C and Jentsch A 2007 Environmental drivers of large, infrequent wildfires: the emerging conceptual model *Prog. Phys. Geog.* **31** 287–312
- Miller J D, Safford H D, Crimmins M and Thode A E 2009 Quantitative evidence for increasing forest fire severity in the Sierra Nevada and southern Cascade Mountains, California and Nevada, USA *Ecosystems* **12** 16–32
- Miller J D, Skinner C N, Safford H D, Knapp E E and Ramirez C M 2012 Trends and causes of severity, size, and number of fires in northwestern California, USA *Ecol. Appl.* **22** 184–203
- Moritz M A, Moody T J, Krawchuk M A, Hughes M and Hall A 2010 Spatial variation in extreme winds predicts large wildfire locations in chaparral ecosystems *Geophys. Res. Lett.* **37** L04801
- Moritz M A et al 2014 Learning to coexist with wildfire *Nature* **515** 58–66
- Morton D C, Collatz G J, Wang D, Randerson J T, Giglio L and Chen Y 2013 Satellite-based assessment of climate controls on US burned area *Biogeosci.* **10** 247–60
- Nicholls N 1997 Increased Australian wheat yield due to recent climate trends *Nature* **387** 484–5
- Parisien M A and Moritz M A 2009 Environmental controls on the distribution of wildfire at multiple spatial scales *Ecol. Monogr.* **79** 127–54
- Parisien M A, Parks S A, Krawchuk M A, Flannigan M D, Bowman L M and Moritz M A 2011 Scale-dependent controls on the area burned in the boreal forest of Canada, 1980–2005 *Ecol. Appl.* **21** 789–805
- Parks S A, Parisien M A, Miller C and Dobrowski S Z 2014 Fire activity and severity in the western US vary along proxy gradients representing fuel amount and fuel moisture *PloS One* **9** e99699
- Pausas J G and Ribeiro E 2013 The global fire-productivity relationship *Glob. Ecol. Biogeogr.* **22** 728–36
- Pellizzaro G, Cesaraccio C, Duce P, Ventura A and Zara P 2007 Relationships between seasonal patterns of live fuel moisture and meteorological drought indices for Mediterranean shrubland species *Int. J. Wildland Fire* **16** 232–41
- Price O F, Pausas J G, Govender N, Flannigan M, Fernandes P M, Brooks M L and Bird R B 2015 Global patterns in fire leverage: the response of annual area burnt to previous fire *Int. J. Wildland Fire* **24** 297–306
- Preisler H K, Chen S C, Fujioka F, Benoit J W and Westerling A L 2008 Wildland fire probabilities estimated from weather model-deduced monthly mean fire danger indices *Int. J. Wildland Fire* **17** 305–16
- Quesada B, Vautard R, Yiou P, Hirschi M and Seneviratne S I 2012 Asymmetric European summer heat predictability from wet and dry southern winters and springs *Nat. Clim. Change* **2** 736–41
- R Development Core Team 2013 *R: a Language and Environment for Statistical Computing* (Vienna, Austria: R Foundation for Statistical Computing) (<http://R-project.org/>)
- Riley K L, Abatzoglou J T, Grenfell I C, Klene A E and Heinsch F A 2013 The relationship of large fire occurrence with drought and fire danger indices in the western USA, 1984–2008: the role of temporal scale *Int. J. Wildland Fire* **22** 894–909
- Rothermel R C 1972 A mathematical model for predicting fire spread in wildland fuels *Research paper INT-115*, USDA Forest Service, Intermountain Forest and Range Experiment Station p 40
- San-Miguel-Ayanz J, Moreno J M and Camia A 2013 Analysis of large fires in European Mediterranean landscapes: lessons learned and perspectives *Forest Ecol. Manage.* **294** 11–22
- Sousa P M, Trigo R M, Pereira M G, Bedia J and Gutiérrez J M 2015 Different approaches to model future burnt area in the Iberian Peninsula *Agric. Forest Meteorol.* **202** 11–25
- Stavros E N, Abatzoglou J, Larkin N K, McKenzie D and Steel E A 2014 Climate and very large wildland fires in the contiguous western USA *Int. J. Wildland Fire* **23** 899–914
- Syphard A D, Radeloff V C, Keeley J E, Hawbaker T J, Clayton M K, Stewart S I and Hammer R B 2007 Human influence on California fire regimes *Ecol. Appl.* **17** 1388–402
- Trouet V, Taylor A H, Carleton A M and Skinner C N 2006 Fire-climate interaction in forests of the American Pacific coast *Geophys. Res. Lett.* **33** L18704
- Trouet V, Taylor A H, Carleton A M and Skinner C N 2009 Interannual variations in fire weather, fire extent, and synoptic-scale circulation patterns in northern California and Oregon *Theor. Appl. Climatol.* **95** 349–60
- Turco M, Llasat M C, von Hardenberg J and Provenzale A 2014 Climate change impacts on wildfires in a Mediterranean environment *Clim. Change* **125** 369–80
- Van Wagner C E 1970 Conversion of William's severity rating for use with the fire weather index *Information Report PS-X-21* Canadian Forestry Service, Ontario, Canada p 6
- Van Wagner C E 1987 Development and Structure of the Canadian Forest Fire Weather Index System *Forestry Technical Report 35* Canadian Forestry Service, Ottawa, Canada p 37
- Vázquez A and Moreno J M 1995 Patterns of fire occurrence across a climatic gradient and its relationships to meteorological variables in Spain *Global Change and Mediterranean-type Ecosystems* ed J M Moreno and W C Oechel WC (New York: Springer) pp 408–34
- Vázquez A and Moreno J M 1998 Patterns of lightning-, and people-caused fires in peninsular Spain *Int. J. Wildland Fire* **8** 103–15
- Venäläinen A, Korhonen N, Hyvärinen O, Koutsias N, Xystrakis F, Urbietta I R and Moreno J M 2014 Temporal variations and change in forest fire danger in Europe for 1960–2012 *Nat. Hazards Earth Syst.* **14** 1477–90
- Vicente-Serrano S M, Beguería S and López-Moreno J I 2010 A multi-scalar drought index sensitive to global warming: the standardized precipitation evapotranspiration index—SPEI *J. Clim.* **23** 1696–718
- Viedma O, Moity N and Moreno J M 2015 Changes in landscape fire-hazard during the second half of the 20th century: agriculture abandonment and the changing role of driving factors *Agric. Ecosyst. Environ.* **207** 126–40
- Weedon G P, Gomes S, Viterbo P, Österle H, Adam J C, Bellouin N, Boucher O and Best M 2010 The WATCH forcing data 1958–2001: a meteorological forcing dataset for land surface- and hydrological models *WATCH Technical Report 22* p 41
- Westerling A L and Swetnam W 2003 Interannual to decadal drought and wildfire in the Western United States *Eos* **84** 545–60
- Westerling A L, Bryant B P, Preisler H K, Holmes T P, Hidalgo H G, Das T and Shrestha S R 2011 Climate change and growth scenarios for California wildfire *Clim. Change* **109** S445–63