

# First Global Carbon Dioxide Maps Produced from TanSat Measurements

Dongxu YANG<sup>1</sup>, Yi LIU<sup>\*1,2</sup>, Zhaonan CAI<sup>1</sup>, Xi CHEN<sup>1</sup>, Lu YAO<sup>1,2</sup>, and Daren LU<sup>1</sup>

<sup>1</sup>*Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China*

<sup>2</sup>*University of Chinese Academy of Sciences, Beijing 100049, China*

(Received 28 December 2017; revised 1 February 2018; accepted 13 February 2018)

**Citation:** Yang, D. X., Y. Liu, Z. N. Cai, X. Chen, L. Yao, and D. R. Lu, 2018: First global carbon dioxide maps produced from TanSat measurements. *Adv. Atmos. Sci.*, **35**(6), 621–623, <https://doi.org/10.1007/s00376-018-7312-6>.

## 1. The need for global carbon monitoring from space and the TanSat mission

Global warming is a major problem, for which carbon dioxide (CO<sub>2</sub>) is the main greenhouse gas involved in heating the troposphere. However, the poor availability of global CO<sub>2</sub> measurements makes it difficult to estimate CO<sub>2</sub> emissions accurately. Satellite measurements would be very helpful for understanding the global CO<sub>2</sub> flux distribution if the CO<sub>2</sub> column-averaged dry-air mole fraction (XCO<sub>2</sub>) could be measured with a precision of 1–2 ppm (Baker et al., 2010). The Greenhouse Gases Observing Satellite (GOSAT) (Yokota et al., 2009; Yoshida et al., 2013; Kuze et al., 2014) was launched in 2009, followed by the Orbiting Carbon Observatory 2 (OCO-2) (Eldering et al., 2016; Crisp et al., 2017; Bösch et al., 2011) in 2014. Tansat, a Chinese Earth observation satellite dedicated to monitoring CO<sub>2</sub>, was launched in December 2016 and is the third satellite capable of monitoring greenhouse gases by hyper-spectral near-infrared/shortwave infrared (NIR/ SWIR) measurement.

The TanSat mission was supported by the Ministry of Science and Technology of China, the Chinese Academy of Sciences, and the China Meteorological Administration. TanSat is an agile, sun-synchronous satellite that operates in three observation modes—namely, the nadir, sun-glint, and target modes. The line of sight tracks the principal plain in nadir mode and the glint in sun-glint mode, which increases the incident signal level and guarantees high performance of the charge-coupled device (Liu et al., 2013a; Cai et al., 2014). The Atmospheric Carbon dioxide Grating Spectroradiometer (ACGS) was designed to measure near-infrared/shortwave infrared backscattered sunlight in the molecular oxygen A-band (0.76 μm) and two CO<sub>2</sub> bands (1.61 and 2.06 μm) (Wang et al., 2014; Li et al., 2017; Zhang et al., 2017). The Cloud and Aerosol Polarization Imager (CAPI) measures in ultraviolet, visible, and NIR regions to improve the information on aerosol optical properties and the cloud mask for the CDS measurements (Chen et al., 2017a, 2017b; Wang et al., 2017).

\* Corresponding author: Yi LIU  
Email: liuyi@mail.iap.ac.cn


## 2. In-orbit CO<sub>2</sub> measurements from TanSat

After TanSat was launched in December 2016, in-orbit and calibration tests were completed in the summer of 2017, and the performance of the instrument has since been evaluated in test sessions. TanSat XCO<sub>2</sub> retrieval algorithm was developed based on Institute of Atmospheric Physics Carbon dioxide retrieval Algorithm for Satellite remote sensing, referred to as IAPCAS (Yang et al., 2015). IAPCAS also informed the development of ATANGO (Application of TanSat XCO<sub>2</sub> Retrieval Algorithm in GOSAT Observations). Its retrieval accuracy and precision have been validated by Total Carbon Column Observing Network (TCCON) measurements (Liu et al., 2013b; Yang et al., 2015), and the retrieval product has been applied in estimations of carbon flux inversion in China (Yang et al., 2017). The retrieval relies on mathematical and physical models to approach XCO<sub>2</sub> from hyper-spectrum measurements restored in the L1B data. The result is the best estimate after comparison between the simulated satellite-received spectrum and measurements. The first global XCO<sub>2</sub> maps based on TanSat measurements show the global distribution over land in April and July 2017 (Fig. 1).

Based on the maps, a seasonal decrease in CO<sub>2</sub> concentration from spring to summer in the Northern Hemisphere is obvious, and results from a change in the rate of photosynthesis. This effect is also reflected in the XCO<sub>2</sub> gradient between the Northern Hemisphere and Southern Hemisphere shown in Fig. 1a. Emission hotspots due to anthropogenic activity, such as industrial activity and fossil fuel combustion, are clearly evident in eastern China, the eastern United States, and Europe, as reflected by the relatively high levels of XCO<sub>2</sub>.

## 3. Outlook

There are still gaps in the TanSat measurements between the footprints of each orbit, and these missing measurements also continue to impact on the carbon flux inversion estimation (Feng et al., 2009). The gaps can be filled by using both OCO-2 and TanSat XCO<sub>2</sub> measurements because the footprint tracks are almost parallel and interlaced, such that



**Fig. 1.** Global XCO<sub>2</sub> maps produced from TanSat in nadir mode in (a) April and (b) July 2017. The colored marks indicate the XCO<sub>2</sub> values and the color scale bar is shown at the bottom of each figure.

OCO-2 provides an additional measurement track between two TanSat tracks. This improves the spatial coverage significantly when compared with the use of a single satellite (i.e., either OCO-2 or TanSat). Accuracy and precision of XCO<sub>2</sub> data are essential for the joint application of OCO-2 and TanSat data. Hence, research focusing on the validation of satellite in-orbit calibration and retrieval algorithms is re-

quired to evaluate their precision and reduce the bias associated with their use.

**Acknowledgements.** This work was supported by the National Key R&D Program of China (2016YFA0600203), the National High-Tech Research and Development Program (2011AA12A104), and External Cooperation Program of the Chinese Academy of Sci-

ences (Grant No. GJHZ1507). The people working on the TanSat mission are highly appreciated. The authors would also like to thank the science teams of GOSAT, SCHIAMACHY, and OCO-2 for valuable discussions, as well as groups at the University of Leicester and RemoTeC (SRON and KIT) for their valuable suggestions.

## REFERENCES

- Baker, F., H. Bösch, C. Doney, D. O'Brien, D. S. Schimel, 2010: Carbon source/sink information provided by column CO<sub>2</sub> measurements from the Orbiting Carbon Observatory. *Atmos. Chem. Phys.*, **10**, 4145–4165, <https://doi.org/10.5194/acp-10-4145-2010>.
- Bösch, H., D. Baker, B. Connor, D. Crisp, and C. Miller, 2011: Global characterization of CO<sub>2</sub> column retrievals from shortwave-infrared satellite observations of the orbiting carbon observatory-2 mission. *Remote Sensing*, **3**, 270–304, <https://doi.org/10.3390/rs3020270>.
- Cai, Z. N., Y. Liu, and D. X. Yang, 2014: Analysis of XCO<sub>2</sub> retrieval sensitivity using simulated Chinese carbon satellite (TanSat) measurements. *Science China Earth Sciences*, **57**, 1919–1928, <https://doi.org/10.1007/s11430-013-4707-1>.
- Chen, X., D. X. Yang, Z. N. Cai, Y. Liu, and R. Spurr, 2017a: Aerosol retrieval sensitivity and error analysis for the cloud and aerosol polarimetric imager on board TanSat: The effect of multi-angle measurement. *Remote Sensing*, **9**, 183, <https://doi.org/10.3390/rs9020183>.
- Chen, X., J. Wang, Y. Liu, X. G. Xu, Z. N. Cai, D. X., Yang, C. X. Yan, and L. Feng, 2017b: Angular dependence of aerosol information content in capi/tansat observation over land: Effect of polarization and synergy with a-train satellites. *Remote Sensing of Environment*, **196**, 163–177, <https://doi.org/10.1016/j.rse.2017.05.007>.
- Crisp, D., and Coauthors, 2017: The on-orbit performance of the orbiting carbon observatory-2 (OCO-2) instrument and its radiometrically calibrated products. *Atmospheric Measurement Techniques*, **10**, 59–81, <https://doi.org/10.5194/amt-10-59-2017>.
- Eldering, A., and Coauthors, 2016: The orbiting carbon observatory-2: First 18 months of science data products. *Atmospheric Measurement Techniques Discussions*, **10**, 549–563, <https://doi.org/10.5194/amt-10-549-2017>.
- Feng, L., P. I. Palmer, H. Bösch, S. Dance, 2009: Estimating surface CO<sub>2</sub> fluxes from space-borne CO<sub>2</sub> dry air mole fraction observations using an ensemble kalman filter. *Atmos. Chem. Phys.*, **9**, 2619–2633, <https://doi.org/10.5194/acp-9-2619-2009>.
- Kuze, A., and Coauthors, 2014: Long-term vicarious calibration of GOSAT short-wave sensors: Techniques for error reduction and new estimates of radiometric degradation factors. *IEEE Trans. Geosci. Remote Sens.*, **52**, 3991–4004, <https://doi.org/10.1109/TGRS.2013.2278696>.
- Li, Z. G., and Coauthors, 2017: Prelaunch spectral calibration of a carbon dioxide spectrometer. *Measurement Science and Technology*, **28**, 065801, <https://doi.org/10.1088/1361-6501/aa6507>.
- Liu, Y., Z. N. Cai, D. X. Yang, M. Z. Duan, and D. Lü, 2013a: Optimization of the instrument configuration for TanSat CO<sub>2</sub> spectrometer. *Chinese Science Bulletin*, **58**, 2787–2789. (in Chinese)
- Liu, Y., D. X. Yang, and Z. N. Cai, 2013b: A retrieval algorithm for TanSat XCO<sub>2</sub> observation: Retrieval experiments using GOSAT data. *Chinese Science Bulletin*, **58**, 1520–1523, <https://doi.org/10.1007/s11434-013-5680-y>.
- Wang, Q., Z. D. Yang, and Y. M. Bi, 2014: Spectral parameters and signal-to-noise ratio requirement for TanSat hyper spectral remote sensor to measure atmospheric CO<sub>2</sub>. *Remote Sensing of the Atmosphere, Clouds, and Precipitation*, <https://doi.org/10.1117/12.2067572>.
- Wang, X., Z. Guo, Y. P. Huang, H. J. Fan, and W. B. Li W, 2017: A cloud detection scheme for the Chinese carbon dioxide observation satellite (TANSAT). *Adv. Atmos. Sci.*, **34**(1), 16–25, <https://doi.org/10.1007/s00376-016-6033-y>.
- Yang, D. X., Y. Liu, Z. N. Cai, J. B. Deng, J. Wang, and X. Chen, 2015: An advanced carbon dioxide retrieval algorithm for satellite measurements and its application to GOSAT observations. *Science Bulletin*, **60**, 2063–2066, <http://doi.org/10.1007/s11434-015-0953-2>.
- Yang, D. X., H. F. Zhang, Y. Liu, B. Z. Chen, Z. N. Cai, and D. R. Lü, 2017: Monitoring carbon dioxide from space: Retrieval algorithm and flux inversion based on GOSAT data and using carbontracker-china. *Adv. Atmos. Sci.*, **34**, 965–976, <https://doi.org/10.1007/s00376-017-6221-4>.
- Yokota, T., Y. Yoshida, N. Eguchi, Y. Ota, T. Tanaka, H. Watanabe, and S. Maksyutov, 2009: Global concentrations of CO<sub>2</sub> and CH<sub>4</sub> retrieved from GOSAT: First preliminary results. *Sola*, **5**, 160–163, <https://doi.org/10.2151/sola.2009-041>.
- Yoshida, Y., and Coauthors, 2013: Improvement of the retrieval algorithm for GOSAT SWIR XCO<sub>2</sub> and XCO<sub>4</sub> and their validation using TCCON data. *Atmospheric Measurement Techniques*, **6**, 1533–1547, <https://doi.org/10.5194/amt-6-1533-2013>.
- Zhang, H., Y. Q. Zheng, C. Lin, W. Q. Wang, Q. Wang, and S. Li, 2017: Laboratory spectral calibration of TanSat and the influence of multiplex merging of pixels. *Int. J. Remote Sens.*, **38**, 3800–3816, <https://doi.org/10.1080/01431161.2017.1306142>.