

 Open access • Journal Article • DOI:10.1136/VR.101868

First isolation of “*Brachyspira hampsonii*” from pigs in Europe — Source link

Maxime Mahu, E. de Jong, N. De Pauw, L. Vande Maele ...+7 more authors

Institutions: Ghent University

Published on: 11 Jan 2014 - Veterinary Record (British Medical Journal Publishing Group)

Topics: Brachyspira hyodysenteriae, Brachyspira pilosicoli and Brachyspira

Related papers:

- [Phenotypic and molecular characterization of a novel strongly hemolytic *Brachyspira* species, provisionally designated "*Brachyspira hampsonii*".](#)
- ["Brachyspira hampsonii" clade I isolated from Belgian pigs imported to Germany.](#)
- [Isolation and characterization of *Brachyspira* spp. including "*Brachyspira hampsonii*" from lesser snow geese \(*Chen caerulescens caerulescens*\) in the Canadian Arctic](#)
- [First Identification of “*Brachyspira hampsonii*” in Wild European Waterfowl](#)
- [Reproduction of Mucohaemorrhagic Diarrhea and Colitis Indistinguishable from Swine Dysentery following Experimental Inoculation with “*Brachyspira hampsonii*” Strain 30446](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/first-isolation-of-brachyspira-hampsonii-from-pigs-in-europe-28wshzzm06>

1 **First isolation of “*Brachyspira hamptonii*” from pigs in Europe**

2 Maxime Mahu^{1*}, Ellen de Jong², Nele De Pauw¹, Lien Vande Maele^{1,3}, Virginie
3 Vandenbroucke², Tamara Vandersmissen², Cora Miry², Frank Pasmans¹, Freddy
4 Haesebrouck¹, An Martel¹ and Filip Boyen¹

5 1 Department of Pathology, Bacteriology and Poultry Diseases, Faculty of Veterinary
6 Medicine, Ghent University, Belgium

7 2 Animal Health Care Flanders, Deinse Horsweg 1, B-9031, Drongen, Belgium

8 3 Institute for Agricultural and Fisheries Research (ILVO), Brusselsesteenweg 370, B-9090
9 Melle, Belgium

10 *Corresponding author: Maxime Mahu, Salisburylaan 133, 9820 Merelbeke, Belgium.

11 maxime.mahu@ugent.be

12

13

14

15

16

17

18

19

20

21 **Abstract.** Swine dysentery in Europe is classically attributed to *Brachyspira hyodysenteriae*.
22 However, other *Brachyspira* species have been increasingly associated with intestinal
23 disorders in pigs. This case report describes the first diagnosis of a “*Brachyspira hampsonii*”
24 infection in European pigs. In a routine quarantine monitoring protocol, two gilts were
25 presented for necropsy, in which soft watery non-haemorrhagic colonic content was found.
26 Microbial culture from the colonic content and from faecal samples revealed the presence of
27 strongly haemolytic, ring-phenomenon positive spirochetes indicative for *Brachyspira*
28 *hyodysenteriae*. A diagnostic commercial PCR could not confirm the presence of *B.*
29 *hyodysenteriae*. Phenotypic characterisation and PCRs targeting the 16S rRNA, 23S rRNA,
30 *nox*, *hlyA* and *tlyA* genes of different swine-related *Brachyspira* spp. were performed.
31 Phylogenetic analysis of sequences of the partial *nox* and 16S rRNA genes and multi locus
32 sequence typing demonstrated that the isolates in this case were “*B. hampsonii*” isolates. This
33 case report shows that the diagnosis of infections caused by new, emerging *Brachyspira*
34 species is not self-evident and that the combination of microbial culture and PCR is
35 recommended, completed with more extensive genotyping if necessary.

36

37

38

39

40

41

42 **Key words:** *Brachyspira*; swine dysentery; haemolysis; “*Brachyspira hampsonii*”

43 **Case report: first isolation of “*Brachyspira hampsonii*” from pigs in Europe**

44

45 Infections with *Brachyspira* spp. in swine occur in most swine-rearing countries and
46 can result in substantial economic losses. Of all swine-related *Brachyspira* spp. infections
47 classical swine dysentery, caused by *Brachyspira hyodysenteriae*, results in the most severe
48 clinical symptoms (eg. mucohaemorrhagic diarrhea, weight loss, poor feed conversion). *B.*
49 *hyodysenteriae* was first recognized as the cause of swine dysentery in 1971 (Taylor and
50 Alexander 1971). At that time, the strong haemolysis of *B. hyodysenteriae* appeared indicative
51 for pathogenicity since other, weakly haemolytic *Brachyspira* (formerly *Serpulina*, *Serpula*
52 and *Treponema*) appeared to be commensal and were therefore named *Brachyspira innocens*
53 (Kinyon and Harris 1979). Several reports of clinical disease caused by weakly hemolytic
54 *Brachyspira* indicated that not all weakly hemolytic *Brachyspira* spp. were non-pathogenic
55 for pigs (Taylor and others 1980, Neef and others 1994). Further research of these weakly
56 haemolytic isolates including DNA-DNA hybridisation, resulted in the designation of three
57 more weakly haemolytic species namely *B. intermedia*, *B. murdochii* and *B. pilosicoli* (Trott
58 and others 1996, Stanton and others 1997).

59 These weakly haemolytic species of *Brachyspira* diverge in the severity of clinical
60 symptoms they cause. *B. pilosicoli* is pathogenic and causes spirochetel colitis in pigs, which
61 is marked by non-haemorrhagic diarrhea and a poor feed conversion. For *B. intermedia* and
62 *B. murdochii* the pathogenic potential is less clear-cut. Although both species have been
63 isolated from clinical cases of diarrhea, the clinical symptoms are mild or absent in
64 experimental infections and yet high numbers of spirochetes are necessary to cause an effect
65 (Jensen and others 2004, Jensen and others 2010).

66 Recently, a new type of *Brachyspira* infection has been described. Outbreaks of
67 mucohaemorrhagic diarrhea, caused by strongly haemolytic *Brachyspira* strains inconsistent

68 with *B. hyodysenteriae*, were reported in the USA and Canada. Phylogenetic analysis of these
69 strains showed such a large genetic divergence between those isolates and all other
70 *Brachyspira* spp. that these isolates likely represent a novel species, for which the name
71 “*Brachyspira hampsonii*” has been proposed (Chander and others 2012). The current case
72 report describes, to the best of our knowledge, the first confirmed “*B. hampsonii*” infection in
73 pigs outside North-America.

74 Two gilts, imported from the Czech Republic, were presented for necropsy in a routine
75 quarantine monitoring protocol. General macroscopic findings consisted of a low body weight
76 and dilated large intestines in which soft watery non-haemorrhagic colonic content was
77 present. Histological examination of these large intestines was not performed. Microbial
78 culture of the colonic content was performed on Tryptic Soy Agar (BD, Heidelberg,
79 Germany) supplemented with 5% sheep blood (IMP, Brussels, Belgium), 0.1% yeast extract
80 (Oxoid, Aalst, Belgium) and following antimicrobials: spectinomycin (200µg/ml), spiramycin
81 (25 µg/ml), rifampin (12.5 µg/ml), colistin (6.25 µg/ml), and vancomycin (6.25µg/ml)
82 (Hommez and others 1998). The microbial cultures revealed strongly haemolytic, ring
83 phenomenon-positive spirochetes, indicative for *B. hyodysenteriae* (Fellström and others
84 1995, Hommez and others 1998). Some of the pigs, housed in the same group as the two gilts
85 presented for necropsy, showed mild diarrhea. From the next batch of gilts from the same
86 origin, additional faecal samples were taken in the quarantine. Strongly haemolytic
87 *Brachyspira* isolates, with ring phenomenon, were again found on microbial culture, whereas
88 commercial diagnostic PCR analysis (Adiavet Brachy, Paris, France) did not confirm the
89 presence of *B. hyodysenteriae* in these samples. All faecal samples were negative for
90 *Salmonella*.

91 Phenotypic characterisation tests were performed on pure cultures which were
92 obtained by at least three subcultures on Tryptic Soy Agar (TSA) plates supplemented with

93 5% defibrinated sheep blood and 0,1% yeast extract (Jenkinson and Wingar, 1981).
94 Phenotypic characterisation was performed on 4-day old cultures and was based on beta
95 haemolysis, indole production, hippurate hydrolysis and the presence or absence of α -
96 galactosidase, α -glucosidase and β -glucosidase (Fellström and others 1995). Indole
97 production was determined using a spot-indole test (Remel BactiDrop, Dartford, UK) and for
98 the other biochemical characteristics commercial discs were used according to the
99 manufacturer's instructions (Rosco Diatabs, Taastrup, Denmark). Type strains of *B.*
100 *hyodysenteriae* (ATCC 27164), *B. pilosicoli* (ATCC 51139) and *B. innocens* (ATCC 29796)
101 were included to provide positive controls for all the phenotypic characteristics that were
102 examined.

103 Several species-specific PCRs were performed, based on the following genes: *tlyA*
104 (Råsbäck and others 2006), 23S rRNA (Leser and others 1997) and *nox* (Phillips and others
105 2006) for *B. hyodysenteriae*, *nox* (Phillips and others 2010) and 23s rRNA (Leser and others
106 1997) for *B. intermedia*, 16S rRNA (Phillips and others 2006) for *B. pilosicoli* and *nox*
107 (Ateyo and others 1999) for *B. murdochii/B. innocens*. Additionally, PCR's were performed for
108 the haemolysis related genes *hlyA* and *hlyA-ACP* (Barth and others 2012).

109 Forward primer 5' TAGCYTGCGGTATYGCWCTTT 3' and reverse primer 5'
110 GCMTGWATAGCTTCRGCATGRT 3' were used to partially sequence the *nox* gene
111 (Weissenböck and others 2005). A product of 1014 base pairs was obtained. Forward primer
112 5' GTTTGATYCTGGCTCAGARCKAACG 3' and reverse primer 5'
113 CTTCCGGTACGGMTGCCTTGTTACG 3' were used to partially sequence the 16S rRNA
114 gene of which a 1044 base pair product was obtained (Johansson and others 2004).
115 Sequencing reactions were performed on purified PCR-product with the same primers as for
116 PCR. *Nox* and 16S rRNA sequences from other *Brachyspira* isolates were retrieved from

117 GenBank and compared with the sequences of the described field case isolate (D52) by
118 BLAST analysis.

119 The sequences of the *nox* gene of the strain retrieved in this case report (D52), of *B.*
120 *hyodysenteriae*, *B. intermedia*, *B. murdochii* and *B. innocens* ATCC type strains and of 42
121 additional strains of several *Brachyspira* spp. retrieved from GenBank were aligned using
122 ClustalW. Sequences of clade I strain 30599 and clade II strain 30446 of *B. "hampsonii"*
123 were also included (Rubin and others, 2013b). Phylogenetic analysis was performed with an
124 alignment sequence fragment of 540 bp and Kimura distance calculation and neighbour-
125 joining method were used.

126 For multilocus sequence typing (MLST) primers and PCR conditions as described by
127 Råsbäck and others in 2007(b) were used to analyse genes encoding alcohol dehydrogenase
128 (*adh*), esterase (*est*), glutamate dehydrogenase (*gdh*), glucose kinase (*glpK*),
129 phosphoglucomutase (*pgm*) and acetyl-coA acetyltransferase (*thi*). For each locus the
130 sequence obtained from the D52 isolate was matched with the online MLSTdatabase
131 (www.pubmlst.org/brachyspira).

132 The phenotypic characteristics of isolate D52 corresponded to those of *B. "hampsonii"*
133 as described by Chander and others (2012). The isolate was strongly beta haemolytic, indole
134 negative, hippurate negative, negative for α -galactosidase and α -glucosidase, and positive for
135 β -glucosidase. Although not exclusively, most isolates of clade I are positive for β -
136 glucosidase as compared to clade II, in which most isolates are negative for β -glucosidase.

137 Table 1 shows the PCR results. Isolate D52 generated a positive result in the two
138 species-specific PCRs for *B. intermedia* based on the 23S rRNA and *nox* gene respectively.
139 Interestingly, the PCR targeting *tlyA*, presumed typical for *B. hyodysenteriae*, also generated a
140 positive result. The PCRs for several haemolysis associated genes, *hlyA* and ACP(*fabF-fabG*),
141 were positive as well.

142 The *nox* sequence of isolate D52 (GenBank accession nr KF202498) showed a
143 similarity of 100% over 547 basepairs with *B. "hampsonii"* isolate NSH-16, which is
144 described by Chander and others as the reference strain of *B. "hampsonii"* clade I (Chander
145 and others 2012). Besides, the *nox* sequence of isolate D52 showed a similarity of more than
146 99% over 874 basepairs with previously described isolates KC35 en EB106 (JX197410.1 and
147 JX197409.1) (Burrough and others 2012b). These isolates, originally described as strongly
148 haemolytic *B. intermedia* are recently referred to as "*B. hampsonii* clade I" (GenBank). With
149 the type strain *B. hampsonii* 30599 (clade I, NZ_AOMM01000255.1) as described by Rubin
150 and others (2013b), the *nox* sequence of our isolate showed a similarity of 99% over 1014 bp
151 (difference of 2 nucleotides). The 16S rRNA sequence of isolate D52 (GenBank accession nr
152 KF586484) showed a sequence similarity of 99% over 1044 bp with "*B. hampsonii*" isolate
153 NSH-16.

154 Phylogenetic analysis of the *nox* sequence of isolate D52 and *nox* sequences of other
155 *Brachyspira* spp. clearly place isolate D52 in the cluster of isolates comprising clade I of "*B.*
156 *hampsonii*" (fig. 1)

157 As described for "*B. hampsonii*" in previous studies (Chander and others 2012) three
158 of the seven loci for MLST could not be amplified. From the sequences of the 4 loci that
159 could be amplified (*est*, *pgm*, *glp* and *thi*), none of them gave an exact match with known
160 alleles in the MLST database. The *thi* sequence matched closest with allele 24 of "*Serpulina*
161 sp. P280/1" (difference of 21 nucleotides) in accordance with the findings of Chander and
162 others, 2012, for "*B. hampsonii*".

163 The results of the phenotypic characteristics, sequence comparisons, MLST and
164 phylogenetic analysis based on the *nox* sequence, identify the D52 isolate as "*B. hampsonii*"
165 clade I. To the best of our knowledge it is the first time that "*B. hampsonii*" isolates from

166 porcine origin are described in Europe, although the isolate *Serpulina* sp. P280/1 (Neef and
167 others 1994) in retrospect also may belong to “*B. hampsonii*”.

168 The isolates of strain D52 obtained from the current field case, all contained the *hlyA*,
169 *tlyA* and ACP(*fabF*,*fabG*) genes. HlyA is the protein responsible for the strong haemolysis in
170 *B. hyodysenteriae* (Hsu and others 2001). In order to adequately perform its actions, the *hlyA*
171 gene has to be correctly placed between the accompanying *fab-F* and *fab-G* genes, coding for
172 an ACP-reductase and –synthetase (Zuerner and others 2004). Although the presence of *hlyA*
173 has been reported in some weakly haemolytic *Brachyspira* spp. isolates, the *fabF* and *fabG*
174 genes were in those cases absent (Barth and others 2012), probably rendering the *hlyA* gene
175 functionally inactive. Another hemolysin, namely *tlyA* is consistently found in *B.*
176 *hyodysenteriae*. Although it has also been twice reported in weakly haemolytic species (Pati
177 and others 2010, Wanchanthuek and others 2010), these sequences show low sequence
178 similarity (82-83%) with *tlyA* of *B. hyodysenteriae* (Barth and others 2012). The presence of
179 both these hemolysin encoding genes in the isolates in the current field case may be
180 responsible for the strong haemolysis displayed by these isolates.

181 Rubin and others (2013a) could experimentally induce mucohaemorrhagic diarrhea in
182 swine when infected with a “*B. hampsonii*” strain 30446. The clinical signs were
183 indistinguishable from swine dysentery. It should, however, be noted that the strain 30446
184 clearly falls into the cluster II isolates of “*B. hampsonii*” whereas the strain from this case
185 falls into cluster I as shown in the phylogenetic tree in figure 1. Although experimentally “*B.*
186 *hampsonii*” strain 30599, which belongs to clade I, can induce severe clinical symptoms
187 (Harding and others, 2013), the symptoms in this case report were rather mild. This could be
188 due to difference in pathogenic potential between strains of clade I or be related to
189 *Brachyspira colitis* being a multifactorial disease. Environmental or nutritional factors may
190 alter the severity of clinical signs.

191 This case report and the recent case reports from Canada and USA (Burrough and
192 others 2012a, Rubin and others 2013a) indicate that new, emerging species of *Brachyspira*
193 can be important in swine-rearing countries. The results of the species-specific PCRs show
194 that diagnosis of infections caused by these emerging species can be confusing. When
195 diagnosis is solely based on microbial culture, all strongly haemolytic isolates will be reported
196 as *B. hyodysenteriae*, whereas they could belong to the provisionally named species “*B.*
197 *hampsonii*”, “*B. suanatina*” (Chander and others 2012, Burrough and others 2012a, Råsbäck
198 and others 2007a, Rubin and others 2013a) or even other emerging *Brachyspira* species. On
199 the other hand, when diagnosis is entirely based on PCR, strongly haemolytic isolates
200 inconsistent with *B. hyodysenteriae* could easily be missed. For now, the combination of
201 microbial culture and PCR, complemented with sequencing if necessary, is presumably the
202 most complete method for diagnosis of *Brachyspira* spp. infections.

203

204 **Declaration of conflicting interests**

205 Sources of financial support have been acknowledged and the authors declare that they have
206 no competing interests.

207

208

209 **Funding**

210 This work was supported by the Institute for the Promotion of Innovation by Science and
211 Technology in Flanders (IWT Vlaanderen), Brussels, Belgium (grant IWT Landbouw
212 100850) and by ‘Veepeiler Varken’, the Fund for Animal Health, Belgium.

213

214 **References**

- 215 ATYEO, R. F., STANTON, T. B., JENSEN, N. S., SURIYAARACHICHI, D. S. &
216 HAMPSON, D. J. (1999) Differentiation of *Serpulina* species by NADH oxidase gene
217 (nox) sequence comparisons and nox-based polymerase chain reaction tests. *Veterinary*
218 *microbiology* **67**, 47–60.
- 219 BARTH, S., GÖMMEL, M., BALJER, G. & HERBST, W. (2012) Demonstration of
220 genes encoding virulence and virulence life-style factors in *Brachyspira* spp. isolates
221 from pigs. *Veterinary microbiology* **155**, 438–43.
- 222 BURROUGH, E., STRAIT, E., KINYON, J., BOWER, L., MADSON, D.,
223 SCHWARTZ, K., FRANA, T. & SONGER, G. (2012a) Comparison of atypical
224 *Brachyspira* spp. clinical isolates and classic strains in a mouse model of swine
225 dysentery. *Veterinary microbiology* **160**, 387–94.
- 226 BURROUGH, E. R., STRAIT, E. L., KINYON, J. M., BOWER, L. P., MADSON, D.
227 M., WILBERTS, B. L., SCHWARTZ, K. J., FRANA, T. S. & SONGER, G. S.
228 (2012b) Comparative virulence of clinical *Brachyspira* spp. isolates in inoculated pigs.
229 *Journal of veterinary diagnostic investigation: official publication of the American*
230 *Association of Veterinary Laboratory Diagnosticians, Inc* **24**, 1025–34.
- 231 CHANDER, Y., PRIMUS, A., OLIVEIRA, S. & GEBHART, C. J. (2012) Phenotypic
232 and molecular characterization of a novel strongly hemolytic *Brachyspira* species,
233 provisionally designated “*Brachyspira hampsonii*”. *Journal of veterinary diagnostic*
234 *investigation: official publication of the American Association of Veterinary*
235 *Laboratory Diagnosticians, Inc* **24**, 903–10.
- 236 FELLSTRÖM, C., PETTERSSON, B., UHLÉN, M., GUNNARSSON, A. &
237 JOHANSSON KE. (1995) Phylogeny of *Serpulina* based on sequence analyses of the
238 16S rRNA gene and comparison with a scheme involving biochemical classification.
239 *Research in Veterinary Science* **59**:5-9.
- 240 HAMPSON, D. J. (2012) *Brachyspira* infections in pigs: an update on pathogenesis
241 and control. Proceedings of the 4th European Symposium of Porcine Health
242 Management, pp 86-91, April 25-27, Brugge, Belgium
- 243 HARDING, J. C. S., LEMIEUX, H., FERNANDO C., DETMER, S.E., COSTA, M.O.,
244 RUBIN, J. AND HILL, J.E. (2013) Experimental reproduction of mucohaemorrhagic
245 diarrhoea following inoculation with “*Brachyspira hampsonii*” clade I (Canadian
246 isolate 30599) Proceedings of the 6th Conference on colonic spirochaetal infections in
247 animals and humans, p 60, September 5-6, Surrey, UK
- 248 HOMMEZ, J., CASTRYCK, F., HAESBROUCK, F. & DEVRIESE, L. A. (1998)
249 Identification of porcine *Serpulina* strains in routine diagnostic bacteriology.
250 *Veterinary microbiology* **62**, 163–9.
- 251 HSU, T., HUTTO, D. L., MINION, F. C., ZUERNER, R. L. & WANNEMUEHLER,
252 M. J. (2001) Cloning of a beta-hemolysin gene of *Brachyspira* (*Serpulina*)
253 *hyodysenteriae* and its expression in *Escherichia coli*. **69**, 706–711.

- 254 JENKINSON, S. R. & WINGAR C.R. (1981) Selective medium for the isolation of
255 *Treponema hyodysenteriae*. *The Veterinary record* **109**, 384-385
- 256 JENSEN, T. K., BOYE, M. & MØLLER, K. (2004) Extensive intestinal spirochaetosis
257 in pigs challenged with *Brachyspira pilosicoli*. *Journal of medical microbiology* **53**,
258 309-312
- 259 JENSEN, T. K., CHRISTENSEN, A. S. & BOYE. M. (2010) *Brachyspira murdochii*
260 colitis in pigs. *Veterinary Pathology* **47**, 334-338
- 261 JOHANSSON, K.E., DUHAMEL, G. E., BERGSJÖ; B., ENGVALL, E. O.,
262 PERSSON, M., PETTERSON, B. & FELLSTRÖM, C. (2004) Identification of three
263 clusters of canine intestinal spirochaetes by biochemical and 16S rDNA sequence
264 analysis. *Journal of medical microbiology* **53**, 345-350.
- 265 KINYON, J. & HARRIS, D. (1979) *Treponema innocens*, a new species of intestinal
266 bacteria, and emended description of the type strain of *Treponema hyodysenteriae*.
267 *International Journal of Systematic Bacteriology* **29**,102-109.
- 268 LESER, T. D., MØLLER, K., JENSEN, T. K. & JORSAL, S. E. (1997) Specific
269 detection of *Serpulina hyodysenteriae* and potentially pathogenic weakly beta-
270 haemolytic porcine intestinal spirochetes by polymerase chain reaction targeting 23S
271 rDNA. *Molecular and Cellular Probes* **11**, 363-372.
- 272 NEEF, N. LYSONS, R. J., TROTT, D. J., HAMPSON, D. J., JONES, P. W. &
273 MORGAN, J. H.(1994) Pathogenicity of porcine intestinal spirochetes in gnotobiotic
274 pigs. *Infection and immunity* **62**, 2395-403.
- 275 PATI, A., SIKORSKI, J., GRONOW, S., MUNK, C., LAPIDUS, A., *et al.* (2010)
276 Complete genome sequence of *Brachyspira murdochii* type strain (56-150^T). *Standards*
277 *in Genomic Sciences* **2**, 260-269
- 278 PHILLIPS, N. D., LA, T. & HAMPSON, D. J. (2006) Development of a two-step
279 nested duplex PCR assay for the rapid detection of *Brachyspira pilosicoli* and
280 *Brachyspira intermedia* in chicken faeces. *Veterinary Microbiology* **116**, 239-245.
- 281 PHILLIPS, N. D., LA, T., AMIN, M.M. & HAMPSON, D.J. (2010) *Brachyspira*
282 *intermedia* strain diversity and relationships to the other indole-positive *Brachyspira*
283 species. *Veterinary Microbiology* **143**, 246-254
- 284 RÅSBÄCK, T., FELLSTRÖM, C., GUNNARSSON, A & ASPÁN, A. (2006)
285 Comparison of culture and biochemical tests with PCR for detection of *Brachyspira*
286 *hyodysenteriae* and *Brachyspira pilosicoli*. *Journal of Microbiological Methods* **66**,
287 347-353.
- 288 RÅSBÄCK, T., JANSSON, D. S., JOHANSSON, K. & FELLSTRÖM, C. (2007a) A
289 novel enteropathogenic , strongly haemolytic spirochaete isolated from pig and
290 mallard, provisionally designated “*Brachyspira suanatina*” sp . nov . *Environmental*
291 *Microbiology* **9**, 983-991.

292 RÅSBÄCK, T., JOHANSSON, K., JANSSON, D. S., FELLSTRÖM, C., ALIKHANI,
293 M.Y., LA, T., DUNN, D. S. & HAMPSON, D.J.(2007b) Development of a multilocus
294 sequence typing scheme for intestinal spirochaetes within the genus *Brachyspira*.
295 *Microbiology* **153**, 4074-4084

296 RUBIN, J. E., COSTA, M. O., HILL, J. E., KITTRELL, H. E., FERNANDO, C.,
297 HUANG, Y., O'CONNOR, B. & HARDING, J. C. S. (2013a) Reproduction of
298 mucohaemorrhagic diarrhea and colitis indistinguishable from swine dysentery
299 following experimental inoculation with “*Brachyspira hampsonii*” Strain 30446. *PloS*
300 *one* **8**, e57146. doi:10.1371/journal.pone.0057146

301 RUBIN, J. E., HARMS, J. N., FERNANDO, C., SOOS, C., DETMER, S. E.,
302 HARDING, J. C. S. & HILL, J. (2013b) Isolation and characterization of *Brachyspira*
303 spp. including *Brachyspira “hampsonii”* from Lesser snow geese (*Chen caerulescens*
304 *caerulescens*) in the Canadian arctic. *Microbial ecology* doi:10.1007/s00248-013-
305 0273-5

306 TAYLOR, D. J. & ALEXANDER, T. J. (1971) The production of dysentery in swine
307 by feeding cultures containing a spirochaete. *The British Veterinary Journal* **127**, 58–
308 61.

309 TAYLOR, D. J., SIMMONS, J. R. & LAIRD, H.M. (1980) Production of diarrhoea
310 and dysentery in pigs by feeding pure cultures of a spirochete differing from
311 *Treponema hyodysenteriae*. *The veterinary record* **106**:324-332

312 TROTT, D. & STANTON, T. (1996) *Serpulina pilosicoli* sp. nov., the agent of porcine
313 intestinal spirochetosis. *International journal of Systematic bacteriology* **46**, 206–215.

314 WANCHANTHUEK, P., BELLGARD, M. I., LA, T., RYAN, K., MOOLHUIJZEN,
315 P., CHAPMAN, B., BLACK, M., SCHIBECI, D., HUNTER, A., BARRERO, R.,
316 PHILLIPS, N. D. & HAMPSON, D. J. (2010) The complete genome sequence of the
317 pathogenic intestinal spirochete *Brachyspira pilosicoli* and comparison with other
318 *Brachyspira* genomes. *PloS one* **5**, e11455. doi:10.1371/journal.pone.0011455

319 WEISSENBOCK, H., MADERNER, A., HERZOG, A. M., LUSSY, H. &
320 NOWOTNY, N. (2005) Amplification and sequencing of *Brachyspira* spp. specific
321 portions of nox using paraffin-embedded tissue samples from clinical colitis in
322 Austrian pigs shows frequent solitary presence of *Brachyspira murdochii*. *Veterinary*
323 *microbiology* **111**, 67–75.

324 ZUERNER, R. L., STANTON, T. B., MINION, C., LI, C., CHARON, N. W., TROTT,
325 D. J. & HAMPSON, D. J. (2004) Genetic variation in *Brachyspira*: chromosomal
326 rearrangements and sequence drift distinguish *B. pilosicoli* from *B. hyodysenteriae*.
327 *Anaerobe* **10**, 229–237.

328

329

330 **Tables and figures**

331 *Table 1:* Primers used in PCRs and results for *B.hydysenteriae* reference strain ATCC 27164, *B. intermedia*

332 reference strain ATCC 51140 and the field case isolate D52.

Target Gene	Species-specificity	Primer name	Primer Sequence (5'-3')	Result D 52	Result ATCC 27164 <i>B.hydysenteriae</i>	Result ATCC 51140 <i>B. intermedia</i>
<i>hlyA</i>	Non- specific	hlyAFo	TCG ATG AAA TTA AAG ATG TTG TT	positive	positive	positive
		hlyARe	TTT TTC TTG ATC TTC TTG AGG A			
ACP(fabF-fabG)	Non-specific	ACPFo	AGG IGA AGT IAT AGC IGT TGA CG	positive	positive	positive
		ACPRE	GAA ACA CCA TTA AGI AIA TTA TCC CA			
23S	<i>B. hydysenteriae</i>	Hyo23SFo	CGG TAA GTG ATG TAC TTG	negative	positive	negative
		Hyo23SRe	AGC CTC AAC CTT AAA GA			
<i>nox</i>	<i>B. hydysenteriae</i>	HyonoxFo	ACT AAA GAT CCT GAT GTA TTT G	negative	positive	negative
		HyonoxRe	CTA ATA AAC GTC TGC TGC			
<i>tlyA</i>	<i>B. hydysenteriae</i>	tlyAFo	GCA GAT CTA AAG CAC AGG AT	positive	positive	negative
		tlyARE	GCC TTT TGA AAC ATC ACC TC			
<i>nox</i>	<i>B. intermedia</i>	IntnoxFo	AGA GTT TGA AGA CAC TTA TGA C	positive	negative	positive
		IntnoxRe	ATA AAC ATC AGG ATC TTT GC			
23S	<i>B. intermedia</i>	Int23SFo	CCG TTG AAG GTT TAC CGT G	positive	negative	positive
		Int23SRe	CGC CTG ACA ATG TCC GG			
16S	<i>B. pilosicoli</i>	Pilo16SFo	AGA GGA AAG TTT TTT CGC TTC	negative	negative	negative
		Pilo16SRe	GCA CCT ATG TTA AAC GTC CTT G			
<i>nox</i>	<i>B. innocens/ B. murdochii</i>	Innmurdnox Fo	CCT GAA AGT TTA AAA GCT G	negative	negative	negative
		Innmurdnox Re	CGA TGT ATT CTT CTT TTC C			

333

334 *Figure 1:* Phylogenetic tree based on the alignment (540bp) of the *nox* gene of *Brachyspira* spp. The
335 alignment was created using CLUSTALw, distance calculation (Kimura) and neighbour joining using PHYLIP.
336 Bootstrap values are indicated. Scale bar indicates 0,02 substitutions per site.

