

Fish biodiversity in coral reefs and lagoon at the Maratua Island, East Kalimantan

HAWIS H. MADDUPPA¹, SYAMSUL B. AGUS¹, AULIA R. FARHAN², DEDE SUHENDRA³,
BEGINER SUBHAN¹

¹Department of Marine Science and Technology, Faculty of Fisheries and Marine Sciences, Bogor Agricultural University. Jl. Agatis No.1, Bogor 16680, West Java, Indonesia. Tel./Fax +62 251 8623644, ✉email: hawis@ipb.ac.id

²Agency for Marine and Fisheries Research and Development, Ministry of Marine Affairs and Fisheries, Republic of Indonesia. Kompleks Bina Samudera, Jl. Pasir Putih I, East Ancol, North Jakarta 14430, Indonesia

³Fish Quarantine Centre Class II, Tanjung Priok. Jl. Enggano Raya No. 16 Tanjung Priok, Jakarta 14320, Indonesia

Manuscript received: 10 June 2012. Revision accepted: 22 July 2012.

ABSTRACT

Madduppa HH, Agus SB, Farhan AR, Suhendra D, Subhan B. 2012. Fish biodiversity in coral reefs and lagoon at the Maratua Island, East Kalimantan. Biodiversitas 13: 145-150. Fishes are one of the most important biotic components in the aquatic environment. They are filling different habitats, including coral reef and lagoon. This study aims to (i) assess biodiversity in coral reef and lagoon in Maratua Island, East Kalimantan, and (ii) compare the fish community indices (Shannon-Wiener diversity, Evenness, and Dominance) between the coral reef and lagoon. A total of 159 fish species of belonging to 30 families were observed during five visual census of the study period. The number of species on coral reefs is higher (121 species) than in the lagoons (47 species). Relative abundance (%) of each species also varied and did not form a specific pattern. However, a clear cluster between the coral reef and lagoon habitats from fish relative abundance based on multivariate analysis and dendrogram Bray-Curtis Similarity was revealed. The Evenness index value (E) ranged from 0.814 to 0.874, the dominance index (C) ranged from 0.023 to 0.184, and the Shannon-Wiener diversity index (ln base, H') ranged from 1.890 to 4.133. Fish biodiversity in coral reefs was higher ($H' = 3.290 \pm 0.301$) than in the lagoon ($H' = 2.495 \pm 0.578$).

Key words: diversity index, Maratua Island, lagoon, fish visual census, reef fishes

INTRODUCTION

Fishes are one of the most important biotic components in the aquatic environment. They fill a very specific habitat by meeting a variety of waters substratum. Several studies have mentioned the importance of glittering fish communities in ecosystem processes through trophic relationships with other biotic components (e.g. Carrasson and Cartes 2002). Coral reefs are used by fishes as a territorial (Robertson et al. 1976), a feeding ground (Reese 1981), as a place to hide (Hixon 1991), and as a reproduction and spawning ground (Wootton 1992). Fish reach their high biodiversity in coral reef ecosystem (Allen and Werner 2002), especially in Indonesia (Allen and Adrim 2003). Indonesian coral reefs harbour more than 2000 fish species (Allen and Adrim 2003). Consequently, Indonesia has been declared as the centre of marine biodiversity (Allen 2008; Allen and Werner 2002).

Lagoon ecosystem is one of shallow water ecosystems that are separated by barrier islands or coral from larger water systems, which is characterized by predominant sand substratum (Hutomo and Moosa 2005). Sedberry and Carter (1993) states that due to differences in characteristics of the substrate between coral reef and lagoon area, the fish community will be different in terms of composition, relative abundance and biomass. In addition, lagoon surrounded by a coral reef may have

similar to the biota of coral reef ecosystem, because corals may grow on hard substrate at its leeseide with a variety of coral lifeforms (Hutomo and Moosa 2005).

Maratua Island, one of the islands in Derawan Islands, has a land area of 384.36 km² and sea area of 3735.18 km², which is fringed with coral reefs, lagoon and uplifted atoll (Tomascik et al. 1997). Geographically the island is situated on a peninsula north of Berau seas (02°15'12" N and 118°38'41" E). Maratua's climatic conditions in the region are affected by the rainy season (October-May) and dry season (July to September). Oceanographic factors influenced the seasonal movement of currents and Indonesian Trough Flow (Arlindo) from the Pacific to the Indian Ocean through the Straits of Makassar (Wyrcki 1961). These conditions create an environment that supports a high biodiversity of marine organisms, especially in fish.

However, documentations of the biodiversity of marine organisms such as fish in Maratua Island are poorly known. The island's coral reefs are threatened by many anthropogenic pressures like other Indonesian reef system that potentially lead many organisms to extinction. Therefore, this study aims to (i) assess fish biodiversity at the coral reefs and lagoons in Maratua Island, East Kalimantan, and (ii) compare the fish community indices between the reef and lagoon.

MATERIAL AND METHODS

Study sites and periods

The reef fish communities on the island of Maratua were observed at the two stations on the reef slope (Maratua 1, and Maratua 2) and three stations for the lagoon (Tanjung Duwata, Karang Bentukan, and Buar) (Figure 1). The study was conducted on 27-28 November 2005 and November 13 to 14, 2006 (Table 1).

Table 1. Information on data collection for each site

Sites	Geographical position		Study period
	N	E	
Coral reefs:			
Maratua 1	2° 16' 51.806"	118° 33' 45.003"	13-11-2006
Maratua 2	2° 15' 42.705"	118° 33' 26.999"	14-11-2006
Lagoons:			
Tanjung Duwata	2° 9' 50.809"	118° 38' 45.014"	27-11-2005
Karang Bentukan	2° 7' 8.309"	118° 42' 23.888"	27-11-2005
Buar	2° 12' 14.387"	118° 37' 12.515"	28-11-2005

Data collection

The location of the study sites were determined initially by snorkeling and observing the coral reef conditions and representative areas (3-7 meters depth). Data obtained by the method of fish visual census along 50 m transect line (English et al. 1997). A total of two transects were laid on coral reefs, and three transects on lagoon. The fish transect lines are straight and follow the depth contour. The basic unit of data collection for the fish visual census was 50 m x 5 m (250 m²). The procedure was to wait for at least 10 minutes before surveying reef fish species (Halford and Thompson 1994), and the approximate time of fish census was up to 60 minutes for each transect. The SCUBA observer swims slowly along transect and recording fish encountered within 2.5 m on both side and 5 m above transect. After data collection, reef fish identification was confirmed by using several fish identification books, i.e. Allen (2000) and Lieske and Myers (2001).




Figure 1. Map of study sites: the reef slope (station 1 and 2) and the lagoon (stations 3, 4 and 5) on the island Maratua within Derawan Islands, East Kalimantan. Insert: Maratua Island in relation to Indonesia indicated by a red box.

Data analysis

Shannon-Wiener Diversity Index (H' ; ln basis), Evenness Index (E), and Simpson Dominance Index (C) (Odum 1971; Magurran 1988; Ludwig and Reynolds 1988; Smith 2002), as well as species richness and their relative abundance were compared among sites. In addition, functional categories (target, major and indicator) were analyzed. The target fish is individual fish that has economic value and food resources for fishermen. This group is also known as the economically important fish or fish consumption. The indicator fish is group of fish that determines the health of coral reefs. This is caused by the strong relationship between the fish and coral, such as Family Chaetodontidae. Member of chaetodontids used coral reefs as a source of food. The major fish is fish groups that are not included in the targets and indicators. This group is generally found in large quantities and many of them are traded as marine ornamental fish. Non-metric Multidimensional Scaling (MDS) and similarity dendrogram were used to visualize the differences in fish communities of the two different habitats (coral reefs and lagoons) (Shepard 1962; Kruskal 1964) by using PRIMER-5 software (Clarke and Gorley 2001). MDS was based on Bray-Curtis similarities. The quality of the MDS plot is indicated by the stress value. Values <0.2 give a potentially useful 2-dimensional picture, stress <0.1 corresponds to a good ordination and stress <0.05 gives an excellent representation (Kruskal 1964; Field et al. 1982; Clarke 1993). This approach has been widely applied to multivariate analysis of the various communities (Field et al. 1982; Clarke and Green 1988).

RESULTS AND DISCUSSION

Fish community structure

A total of 2145 individuals from 160 species of fish belonging into 30 families were observed in this study (Table 2). The number of species on coral reefs was higher (121 species) than in the lagoons (47 species). List of families that have three top species composition in coral reef areas were Pomacentridae, Labridae and Acanthuridae, while in the lagoon were Pomacentridae, Chaetodontidae, and Nemipteridae (Figure 2). These families are also mainly observed on the other coral reefs around Indonesia (Estradivari et al. 2007; Ferse 2008).

Fish abundance in the coral reef was higher (690 ± 163 individual/ 250 m^2) than in the lagoon (255 ± 92 individual/ 250 m^2), and relative abundance (%) of each species also varied over study sites (Table 2). A clear difference between the reef and lagoon habitats based on MDS and dendrogram was revealed (Figure 3). This is similar with other study that showed differences in fish community based on the abundance of species from the results of the multivariate analysis (Madduppa et al. 2012). The presence of reef fish in the waters depends on coral health indicated by the percentage of live coral cover. It is very possible because of the live reef fish associated with the shape and type of coral as a shelter, protection and

places to look for food (Madduppa 2006). Besides the health of coral, reef structure complexities have enriched reef fishes (Nybakken 1992).

Reef fish communities observed in this study were grouped into three functional categories (Table 2). The target fish families were observed in the study sites as follows: Haemulidae, Nemipteridae, Serranidae and Pomacanthidae. The target fish in the coral reef stations (40 ± 8 species/ 250 m^2) were greater than in the lagoons (7 ± 2 species/ 250 m^2). The presence of the target fish in the coral reef ecosystem are due to searching for food (feeding ground) or spawning and nursery. In the Philippines, many coral reef fishes are caught for the small-scale fisheries including surgeonfish, groupers, and snappers (Amar et al. 1996).

A total of 16 species of fish were found in all indicators of the study sites. Groups of fish indicators also showed a similar pattern in the area where the coral reef was higher (7 ± 1 species/ 250 m^2) than in the lagoon (3 ± 2 species/ 250 m^2). Due to their greatest biodiversity in coral reef ecosystems (Allen and Werner 2002) and their interdependence to the health of coral reef ecosystem (Hourigan et al. 1988), chaetodontid has been considered as a reliable way to indirectly assess the changes of a coral reef and monitor it through time (Tanner et al. 1994; Markert et al. 2003). Many members of Chaetodontidae are highly dependent on live coral polyps, and multiple studies have proven that they are corallivorous (e.g. Harmelin-Vivien and Bouchon-Navaro 1983; Pratchett 2005; Reese 1981; Birkeland and Neudecker 1981; Alwany et al. 2003; Madduppa 2006).

The major groups of fish on the coral reefs stations were higher (39 ± 7 species/ 250 m^2) than in the lagoon (13 ± 4 species/ 250 m^2). Function and role of the fish were not yet clear but could be as one link in the ecological system and food webs in coral reef ecosystems. Major fish species are found mainly from the family Pomacentridae.

Fish biodiversity

The Evenness index (E) value ranging from 0.814 to 0.874, the dominance index (C) 0.023-0.184 range, and the value of diversity index (H') ranged from 1.890-4.133. Fish communities in reef areas were more diverse ($H' = 3.290 \pm 0.301$) than in the lagoon ($H' = 2.495 \pm 0.578$) (Figure 4). The value of diversity can indicate the level of stress or pressure received by the species from the environment (Lardicci et al. 1997). In the lagoon, reef fish communities in the lagoon which is located outside near reef edge (Station 3, see Figure 1) was higher in comparison with two other lagoon stations which are located in the inside (Station 4 and 5). This is similar to Hutomo and Moosa (2005) that fish communities in the lagoon near coral reefs is highly influenced by the benthic community (including corals).

Diversity of species of reef fish have a close relationship with the characteristics of the substrate in the area, such as the existence of the herbivorous fishes of the family Scaridae, because of dead coral covered with macroalgae (Madduppa et al. 2012). The fish will tend to




Figure 2. Comparison of the number of species of fish from each family were found among the coral reefs and lagoons




Figure 3. Multivariate analysis: (left) Bray-Curtis Similarity and (right) non-metric MDS (multidimensional scaling) of the fish community based on the relative abundance of each species per station on coral reefs (1 and 2) and in the lagoon (3, 4 and 5)




Figure 4. Comparison of reef fish community index values between the lagoon and reef slope are seen from the value of Shannon-Wiener diversity (H'), Evenness (E), and Dominance (C)

cluster in certain forms of corals and generally have limited movement compared to other invertebrates that have the same size cite. Because the reef fish communities have a close relationship with the coral reefs as a habitat, so that if a high percentage of dead coral will cause a significant decrease in the number of fish species and individuals associated with coral reefs cite. The current study observed that some fishes found in a specific habitat, but many species were found in more than one habitat. Generally each species has specific habitat preferences (Hutomo 1986), and they have two different interaction mode in coral reefs: direct interaction (e.g. as a refuge from predators or prey, interactions in search of food the relationship between corals and fish that live on the reef biota, including algae) and indirect interaction due to reef structure and hydrology and sediment (Choat and Bellwood 1991).

Table 2. List of taxa, number of individual (\pm SE individual/250 m²), species richness (\pm SE species/250 m²), fish categorical function (\pm SE species/250 m², T=Target, M=Major, I=Indicator), and relative abundance (%) for each species at each site (1 and 2 = reef slope, 3-5 = lagoon).

Taxa (family/species)	Cate- gory	Relative abundance (%)				
		1	2	3	4	5
Acanthuridae						
<i>Acanthurus auranticavus</i>	T	1.3	-	-	-	-
<i>Acanthurus gahhm</i>	T	0.8	2.3	-	-	-
<i>Acanthurus leucocheilus</i>	T	0.8	-	-	-	-
<i>Acanthurus leucosternon</i>	T	0.1	1.1	-	-	-
<i>Acanthurus lineatus</i>	T	-	0.4	-	-	-
<i>Acanthurus thompsoni</i>	T	0.8	2.1	-	-	-
<i>Naso annulatus</i>	T	0.2	0.2	-	-	-
<i>Naso brevirostris</i>	T	-	0.2	-	-	-
<i>Naso lituratus</i>	T	0.4	-	-	-	-
<i>Naso sp.</i>	T	-	-	1.9	6.4	-
<i>Paracanthurus hepatus</i>	T	-	-	11.3	-	-
<i>Zebrasoma scopas</i>	T	2.2	2.1	-	-	-
Apogonidae						
<i>Apogon cavities</i>	M	-	-	-	21.3	-
<i>Apogon cookie</i>	M	-	-	5.8	-	-
<i>Apogon doederleini</i>	M	2.7	4.0	-	-	-
<i>Apogon kauderni</i>	M	-	1.3	-	-	-
<i>Sphaeramia nematoptera</i>	M	0.5	0.8	-	-	-
Balistidae						
<i>Balistapus undulatus</i>	M	-	0.2	-	-	-
<i>Balistoides viridescens</i>	M	0.1	-	-	-	-
<i>Odonus niger</i>	M	-	-	3.0	-	-
<i>Pseudobalistes flavimarginatus</i>	M	-	5.1	-	-	-
<i>Pseudobalistes fuscus</i>	M	1.6	0.2	-	-	-
Belontiidae						
<i>Strongylura incisa</i>	M	1.1	-	-	-	-
<i>Meiacanthus vittatus</i>	M	0.1	-	0.3	-	-
Caesionidae						
<i>Caesio cunning</i>	T	0.1	0.2	-	-	-
<i>Caesio lunaris</i>	T	0.1	0.2	-	-	-
<i>Caesio teres</i>	T	0.1	-	-	-	-
<i>Caesio xanthonota</i>	T	-	-	0.3	-	-
Carangidae						
<i>Gnathanodon speciosus</i>	T	-	0.2	-	-	-
Chaetodontidae						
<i>Chaetodon auriga</i>	I	-	-	0.6	-	-
<i>Chaetodon collare</i>	I	-	-	0.8	-	-
<i>Chaetodon collare</i>	I	-	-	0.3	0.9	-
<i>Chaetodon decussatus</i>	I	-	-	-	0.6	-
<i>Chaetodon decussatus</i>	I	-	-	0.8	-	-
<i>Chaetodon ephippium</i>	I	-	-	2.2	-	-
<i>Chaetodon guttatissimus</i>	I	0.4	0.2	0.6	-	-
<i>Chaetodon melapterus</i>	I	0.2	0.2	-	-	-
<i>Chaetodon meyeri</i>	I	0.6	0.6	-	-	-
<i>Chaetodon rafflesi</i>	I	-	0.2	-	-	-
<i>Chaetodon trifascialis</i>	I	0.1	-	-	-	-
<i>Chaetodon trifasciatus</i>	I	0.2	2.3	-	-	-
<i>Chaetodon vagabundus</i>	I	0.1	-	-	-	-
<i>Chelmon rostratus</i>	I	-	-	0.3	-	-
<i>Hemitaurichthys polylepis</i>	I	1.3	0.4	-	-	-
<i>Parachaetodon ocellatus</i>	I	-	-	12.9	-	-
Cirrhitidae						
<i>Paracirrhites arcatus</i>	M	-	-	-	9.7	-
<i>Paracirrhites forsteri</i>	M	-	-	6.3	2.7	31.5
<i>Paracirrhites nesus</i>	M	-	-	-	2.4	-
Diploprionidae						
<i>Diploprion bifasciatum</i>	M	-	0.6	-	-	-
<i>Platax orbicularis</i>	M	4.1	-	-	-	-

<i>Platax pinnatus</i>	M	0.4	-	-	-	-	<i>Amphiprion ocellaris</i>	M	-	-	-	-	9.6
<i>Platax teira</i>	M	0.1	-	-	-	-	<i>Amphiprion sandaracinos</i>	M	-	-	10.2	-	-
Gobiidae							<i>Chrysiptera unimaculata</i>	M	-	0.2	-	-	-
<i>Gobi</i> sp.	M	1.1	-	-	-	-	<i>Chromis analis</i>	M	-	1.3	-	-	-
<i>Istigobius nigroocellatus</i>	M	-	-	1.4	-	2.7	<i>Chromis atripectoralis</i>	M	0.4	0.2	-	-	-
Haemulidae							<i>Chromis delta</i>	M	0.1	0.2	-	-	1.4
<i>Plectorhinchus sordidus</i>	T	0.2	-	-	-	-	<i>Chromis dimidiata</i>	M	0.4	0.2	-	1.2	-
<i>Plectorhinchus flavomaculatus</i>	T	0.7	-	-	-	-	<i>Chromis nitida</i>	M	0.2	-	-	-	-
<i>Plectorhinchus chaetodonoides</i>	T	0.2	7.8	-	-	-	<i>Chromis verater</i>	M	0.6	1.7	-	-	-
Holocentridae							<i>Chromis viridis</i>	M	-	-	0.6	1.5	4.1
<i>Myripristis amaena</i>	T	-	-	-	3.3	-	<i>Chrysiptera cyanea</i>	M	-	-	-	1.2	-
<i>Sargocentron caudimaculatus</i>	T	1.5	0.4	-	-	-	<i>Chrysiptera hemicyanea</i>	M	-	-	1.4	2.1	-
Labridae							<i>Chrysiptera rollandi</i>	M	-	-	1.9	0.3	-
<i>Bodianus mesothorax</i>	T	0.1	0.4	-	-	-	<i>Chrysiptera springeri</i>	M	0.2	-	-	-	-
<i>Cheilinus fasciatus</i>	T	-	0.6	-	-	-	<i>Dascyllus aruanus</i>	M	0.2	-	-	-	-
<i>Cheilinus oxycephalus</i>	T	0.2	-	-	-	-	<i>Dascyllus carneus</i>	M	0.4	-	-	-	-
<i>Cheilinus trilobatus</i>	T	0.1	-	-	-	-	<i>Dascyllus melanurus</i>	M	0.9	-	-	-	-
<i>Choerodon zamboanga</i>	T	0.1	-	-	-	-	<i>Dascyllus trimaculatus</i>	M	0.1	-	-	-	-
<i>Choerodon anchorago</i>	T	0.1	-	-	-	-	<i>Dischistodus melanotus</i>	M	0.1	-	-	-	-
<i>Gomphosus caeruleus</i>	T	0.1	0.9	-	-	-	<i>Dischistodus perspicillatus</i>	M	0.1	-	-	-	-
<i>Gomphosus varius</i>	T	0.8	-	-	-	-	<i>Dischistodus prosopotaenia</i>	M	2.2	4.7	-	-	-
<i>Halichoeres hortulanus</i>	T	1.6	0.2	-	-	-	<i>Hemiglyphidodon plagiometopon</i>	M	0.4	-	1.7	-	-
<i>Halichoeres melanurus</i>	T	0.8	-	-	-	-	<i>Neopomacentrus azysron</i>	M	-	-	2.8	7.0	-
<i>Halichoeres ornatissimus</i>	T	0.5	0.2	-	-	-	<i>Neopomacentrus cyanomos</i>	M	-	-	8.5	-	-
<i>Halichoeres purpurascens</i>	T	-	-	1.9	0.9	-	<i>Neopomacentrus xanthurus</i>	M	-	-	0.6	-	-
<i>Labroides dimidiatus</i>	M	1.4	-	0.8	-	-	<i>Plectroglyphidodon lacrymatus</i>	M	0.1	1.7	-	-	-
<i>Thalassoma grammaticum</i>	M	0.8	4.2	-	-	-	<i>Pomacentrus alexandrae</i>	M	-	1.3	-	-	-
<i>Thalassoma lunare</i>	M	2.9	-	-	-	-	<i>Pomacentrus amboinensis</i>	M	0.2	0.4	-	-	-
Lethrinidae							<i>Pomacentrus brachialis</i>	M	0.5	-	-	-	-
<i>Gymnocranius griseus</i>	T	0.1	0.2	-	-	-	<i>Pomacentrus lepidogenys</i>	M	0.4	2.1	-	-	-
<i>Monotaxis grandoculis</i>	T	0.1	-	0.3	-	-	<i>Pomacentrus milleri</i> (juv.)	M	-	0.6	-	-	-
Lutjanidae							<i>Pomacentrus nigromarginatus</i>	M	0.7	-	-	-	-
<i>Lutjanus decussatus</i>	T	2.6	-	-	-	-	<i>Pomacentrus tripunctatus</i>	M	1.3	0.4	-	-	-
<i>Lutjanus ehrenbergii</i>	T	-	0.2	-	-	-	<i>Stegastes aureus</i>	M	1.4	5.5	-	-	-
<i>Lutjanus fulviflamma</i>	T	-	0.4	-	-	-	Scaridae						
<i>Lutjanus kasmira</i>	T	-	0.9	-	-	-	<i>Chlorurus microrhinos</i>	T	0.1	-	-	-	-
<i>Lutjanus quinquelineatus</i>	T	1.9	0.2	-	-	-	<i>Scarus globiceps</i>	T	1.2	2.7	-	-	-
<i>Lutjanus semicinctus</i>	T	0.1	-	-	-	-	<i>Scarus melanurus</i>	T	-	1.7	-	-	-
<i>Lutjanus vitta</i>	T	-	0.2	-	-	-	<i>Scarus niger</i>	T	0.5	-	-	-	-
Microdesmidae							<i>Scarus rivulatus</i>	T	3.4	-	-	-	-
<i>Ptereleotris evides</i>	M	0.1	0.4	-	-	-	<i>Scarus rubroviolaceus</i>	T	0.1	0.2	-	-	-
Mullidae							<i>Scarus</i> sp.	T	-	-	-	0.3	-
<i>Parupeneus bifasciatus</i>	M	-	-	4.4	4.0	-	Serranidae						
<i>Upeneus</i> sp.	M	0.6	0.4	-	-	-	<i>Cephalopholis cyanostigma</i>	T	-	4.6	-	-	-
<i>Upeneus vittatus</i>	M	0.5	0.4	-	-	-	<i>Epinephelus fasciatus</i>	T	1.1	-	-	-	-
Nemipteridae							<i>Epinephelus spilotoceps</i>	T	-	-	0.3	-	17.8
<i>Parascalopsis eriomma</i>	T	-	-	1.1	1.5	9.6	<i>Plectropomus leopardus</i>	T	0.4	-	-	-	-
<i>Pentapodus caninus</i>	T	-	-	7.2	0.6	-	<i>Pseudanthias bimaculatus</i>	T	1.9	-	-	-	-
<i>Pentapodus trivittatus</i>	T	1.5	4.7	-	-	-	<i>Pseudanthias pleurotaenia</i>	T	2.6	-	-	-	-
<i>Scolopsis bilineatus</i>	T	-	-	-	3.3	16.4	<i>Pseudanthias randalli</i>	T	0.2	-	-	-	-
<i>Scolopsis ghanam</i>	T	-	-	-	0.6	-	<i>Pseudanthias squamipinnis</i>	T	1.6	-	-	-	-
<i>Scolopsis lineatus</i>	T	0.1	-	-	-	-	<i>Pseudanthias tuka</i>	T	1.1	0.4	-	-	-
<i>Scolopsis margaritifera</i>	T	2.1	2.5	-	-	-	Siganidae						
<i>Scolopsis trilineatus</i>	T	1.6	-	-	-	-	<i>Siganus puellus</i>	T	-	-	0.3	-	-
Platycephalidae							<i>Siganus virgatus</i>	T	0.1	1.9	-	-	-
<i>Thysanophrys arenicola</i>	M	5.6	-	-	-	-	Tetraodontidae						
Plesiopidae							<i>Canthigaster compressa</i>	T	0.2	0.2	-	-	-
<i>Calloplepsiops altivelis</i>	M	-	0.2	-	-	-	Theraponidae						
Pomacanthidae							<i>Terapon jarbua</i>	T	0.8	0.4	-	-	-
<i>Centropyge eibli</i>	M	5.5	4.0	-	-	-	Zanclidae						
<i>Centropyge multispinis</i>	M	6.3	8.2	-	-	-	<i>Zanclus cornutus</i>	M	2.5	0.2	2.5	2.1	-
<i>Centropyge nox</i>	M	3.4	-	-	-	-	Number of individual		690	±163	255	±92	
<i>Pygoplites diacanthus</i>	T	3.6	3.0	-	-	-	Species richness		86	±16	22	±7	
Pomacentridae							Target (T)		40	±8	7	±2	
<i>Abudefduf sexfasciatus</i>	M	-	-	2.5	21.3	-	Major (M)		39	±7	13	±4	
<i>Abudefduf vaigiensis</i>	M	0.9	-	2.5	4.6	6.8	Indicator (I)		7	±1	3	±2	
<i>Amblyglyphidodon curacao</i>	M	0.5	2.1	-	-	-							

CONCLUSION

The study observed that the diversity of fish species in the reef slope was higher than in the lagoon. This shows that the characteristics of the habitat were instrumental in shaping the fish community. Habitats in a tropical lagoon are an important area for various fish species of coral reef ecosystems cite. Despite the complexity of the lagoon habitat characteristics are not as high in coral reef ecosystems, but also provides a location for a few juvenile fish to grow and evolve, which is also vital for the conservation of species. Therefore, management strategies and conservation of coral reef ecosystems should include a lagoon habitat as an important and integrated part.

ACKNOWLEDGEMENTS

The research is part of the project 'Action plan for the outer islands of East Kalimantan' by BAPPEDA East Kalimantan Province year 2006 and 'Inventory of Lagoon ecosystem' by the Ministry of Marine Affairs and Fisheries year 2005.

REFERENCES

- Allen GR. 2000. Marine fishes of South-East Asia. Periplus. Singapore.
- Allen GR. 2008. Conservation hotspots of biodiversity and endemism for Indo-Pacific coral reef fishes. *Aquatic Conservation: Mar Freshw Ecosyst* 18: 541-556.
- Allen GR, Adrim M. 2003. Coral reef fishes of Indonesia. *Zool Stud* 42: 1-72.
- Allen GR, Werner TB. 2002. Coral reef fish assessment in the 'Coral Triangle' of Southeastern Asia. *Environ Biol Fish* 65: 209-214
- Alwany M, Thaler E, Stachowitsch M. 2003. Food selection in two coralivorous butterfly fishes, *Chaetodon austriacus* and *C. trifascialis*, in the Northern Red Sea. *Mar Ecol* 24: 165-167.
- Amar EC, Cheong RMT, Cheong MVT. 1996. Small-scale fisheries of coral reefs and the need for community-based resource management in Malalison Island, Philippines. *Fish Res* 25: 265-277
- Birkeland C, Neudecker S. 1981. Foraging behavior of two Caribbean chaetodontids: *Chaetodon capistratus* and *C. aculeatus*. *Copeia* 1: 169-178
- Carrasson M, Cartes JE. 2002. Trophic relationships in a Mediterranean deep-sea fish community: partition of food resources, dietary overlap and connections within the benthic boundary layer. *Mar Ecol Prog Ser* 241: 41-55
- Choat JH, Bellwood DR. 1991. Reef fishes: Their history and evolution. In: Sale PF (ed). *The ecology of the fishes on coral reefs*. Academic Press, California.
- Clarke KR. 1993. Non-parametric multivariate analyses of changes in community structure. *Australian J Ecol* 18: 117-143.
- Clarke KR, Gorley NR. 2001. PRIMER v5: Unser Manual/Tutorial. PRIMER-E Ltd., Plymouth, UK
- Clarke KR, Green RH. 1988. Statistical design and analysis for a 'biological effects' study. *Mar Ecol Prog Ser* 46: 213-226
- English S, Wilkinson C, Baker V. 1997. Survey Manual for Tropical Marine Resources. 2nd ed. Australian Institute of Marine Science. Australia (AIMS), Townsville.
- Estradivari, Syahrir M, Susilo N, Yusri S, Timotius S. 2007. Jakarta coral reefs: Long-term monitoring of the Kepulauan Seribu coral reefs (2004-2005). Yayasan Terangi, Jakarta. [Indonesia]
- Ferse SCA. 2008. Artificial reef structures and coral transplantation: fish community responses and effects on coral recruitment in North Sulawesi/Indonesia. [Dissertation]. University of Bremen, Bremen, Germany.
- Field JG, Clarke KR, Warwick RM. 1982. A practical strategy for analysing multispecies distribution patterns. *Mar Ecol Prog Ser* 8: 37-52
- Halford AR, Thompson AA. 1994. Visual Census Surveys of Reef Fish. Long-term monitoring of the Great Barrier Reef. Australian Institute of Marine Science (AIMS), Townsville, Australia
- Harmelin-Vivien ML, Bouchon-Navaro Y. 1983. Feeding diets and significance of coral feeding among chaetodontid fishes in Moorea (French Polynesia). *Coral Reefs* 2: 119-127.
- Hixon MA. 1991. Predation as a process structuring coral reef fish communities. In: Sale PF (ed). *The ecology of fishes on coral reefs*. Academic Press, San Diego.
- Hutomo M. 1986. Coral reef fish resources and their relation to reef condition: Some case studies in Indonesian waters. *Biotrop spec. publ* 19: 67-78.
- Hutomo M, Moosa MK. 2005. Indonesian marine and coastal biodiversity: Present status. *Indian J Mar Sci* 34: 88-97.
- Hourigan TF, Tricas TC, Reese ES. 1988. Coral reef fishes as indicators of environmental stress in coral reefs. In: Soule DF, Kleppel GS (Eds.), *Marine Organisms as Indicators*. Springer-Verlag, New York City, NY, USA, pp. 107-135
- Kruskal JB. 1964. Multidimensional scaling by optimizing goodness of fit to a nonmetric hypothesis. *Psychometrika* 29: 1-27
- Lardicci C, Rossi F, Castelli A. 1997. Analysis of macrozoobenthic community structure after severe dystrophic crises in a Mediterranean coastal lagoon. *Mar Poll Bull* 34: 536-547
- Lieske E, Myers R. 2001. Reef fishes of the world: Indo Pacific and Caribbean. (Revised edition). Periplus. Singapore.
- Ludwig JA, Reynolds JF. 1988. Statistical ecology: A primer on methods and computing. John Wiley and Sons, New York.
- Madduppa HH. 2006. Bioecological study on eight band butterfly fish (*Chaetodon octofasciatus*, Bloch 1787) to detect condition of coral reef ecosystem in East Petondan Island, Seribu Islands, Jakarta. [MSc Thesis]. Bogor Agricultural University, Bogor
- Madduppa HH, Ferse SCA, Aktani U, Palm HW. 2012. Seasonal trends and fish-habitat associations around Pari Island, Indonesia: setting a baseline for environmental monitoring. *Environ Biol Fish*. DOI 10.1007/s10641-012-0012-7
- Magurran AE. 1988. Ecological diversity and its measurements. Princeton University Press, Princeton.
- Markert BA, Breure AM, Zechmeister HG. 2003. Bioindicators and biomonitoring: Principles, concepts, and applications. Elsevier, Amsterdam
- Nybakken JW. 1992. Marine biology: An ecology approach. Gramedia, Jakarta.
- Odum EP. 1971. Fundamental of ecology. 3rd ed. Toppan Co, Ltd. London.
- Pratchett MS. 2005. Dietary overlap among coral-feeding butterfly fishes (Chaetodontidae) at Lizard Island, northern Great Barrier Reef. *Mar Biol* 148: 373-382
- Reese ES. 1981. Predation on corals by fishes of the family Chaetodontidae: Implication for conservation and management of coral reef ecosystems. *Bull Mar Sci* 31: 594-604.
- Robertson DR, Sweatman HPA, Fletcher EA, Cleland MG. 1976. Schooling as a mechanism for circumventing the territoriality of competitors. *Ecology* 57: 1208-1220.
- Sedberry GR, Carter J. 1993. The fish community of a shallow tropical lagoon in Belize, Central America. *Estuaries* 16: 198-215
- Shepard R. 1962. The analysis of proximities: Multidimensional scaling with an unknown distance function. II. *Psychometrika* 27: 219-246
- Smith EP. 2002. Ecological statistics. Vol 2. In: Abdel HE, Walter WP (eds). John Wiley and Sons, Chichester.
- Tanner JE, Hughes TP, Connell JH. 1994. Species coexistence, keystone species, and succession - A sensitivity analysis. *Ecology* 75: 2204-2219
- Tomascik T, Mah AJ, Nontji A, Moosa MK. 1997. The ecology of the Indonesian seas. Part Two. Periplus. Singapore.
- Wootton RJ. 1992. Fish ecology (Tertiary Level Biology). Blackie and Son, New York.
- Wyrtki K. 1961. Physical oceanography of Southeast Asian waters. In: NAGA report. The University of California, Scripps Institution of Oceanography. LaJolla, California.