

Five new sucrose esters from the whole plants of *Phyllanthus cochinchinensis*

Jian-Qiang ZHAO,^{a,b,†} Yan-Ming WANG,^{a,b,†} Dong WANG,^a Chong-Ren YANG,^a Min XU,^{a,*} and Ying-Jun ZHANG^{a,*}

^aState Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, China

^bUniversity of Chinese Academy of Sciences, Beijing 100049, China

[†]These authors contributed equally to this work.

Received 21 March 2013; Accepted 5 April 2013

© The Author(s) 2013. This article is published with open access at Springerlink.com

Abstract: Chemical investigation of the whole plants of *Phyllanthus cochinchinensis* (Euphorbiaceae) led to the isolation of five new sucrose benzoyl esters, 3,6'-di-*O*-benzoylsucrose (**1**), 3,6'-di-*O*-benzoyl-2'-*O*-acetylsucrose (**2**), 3,6'-di-*O*-benzoyl-4'-*O*-acetylsucrose (**3**), 3,6'-di-*O*-benzoyl-3'-*O*-acetylsucrose (**4**) and 3-*O*-benzoyl-6'-*O*-(*E*)-cinnamoylsucrose (**5**), together with two known secoiridoid glycosides, jasminoside (**6**) and jaslaceoside B (**7**). Their structures were established on the basis of detailed spectroscopic analysis and chemical method.

Keywords: *Phyllanthus cochinchinensis*, sucrose esters, secoiridoid glycosides

Introduction

Phyllanthus, comprising about 600 species, is the largest genus in family Euphorbiaceae, of which most species are important medicinal plants having been used for the treatment of infectious diseases. Previous chemical studies on this genus have revealed the occurrence of flavonoids, alkaloids, sesquiterpenoids, triterpenoids, lignans, and tannins.^{1–5} Among them, some sesquiterpenoids exhibited potent antiviral activity against coxsackie virus B3 (CVB3),⁴ and inhibition of the growth of murine P-388 lymphocytic leukemia cell line.⁵

P. cochinchinensis, a shrub up to 3 m height, is mainly growing in the montane sparse forests, forest margins, scrub on slopes, and wastelands of the southern part of China. It is also widely distributed in Cambodia, India, Laos, and Vietnam. So far, no chemical study was reported on this species. As a part of our continuing study on bioactive compounds from *Phyllanthus* species,^{1,2,4} five new sucrose benzoyl esters, 3,6'-di-*O*-benzoylsucrose (**1**), 3,6'-di-*O*-benzoyl-2'-*O*-acetylsucrose (**2**), 3,6'-di-*O*-benzoyl-4'-*O*-acetylsucrose (**3**), 3,6'-di-*O*-benzoyl-3'-*O*-acetylsucrose (**4**) and 3-*O*-benzoyl-6'-*O*-(*E*)-cinnamoylsucrose (**5**), were isolated from the whole plants of *P. cochinchinensis*, together with two known secoiridoid glycosides, jasminoside (**6**) and jaslaceoside B (**7**). Their structures were established by means of MS and extensive




Figure 1. Structure of compounds 1–7

NMR spectroscopic analysis and chemical method.

Results and Discussion

The air-dried and powdered whole plants of *P. cochinchinensis* were extracted with MeOH under reflux. Further column chromatography (CC) over Diaion HP20SS, Sephadex LH-20 and silica gel, followed with semi-preparative HPLC purification of the MeOH extract yielded five new compounds (**1–5**), together with two known ones. The known compounds were elucidated as jasminoside (**6**)⁶ and jaslaceoside B (**7**)⁷ by comparison of their spectroscopic data with reported literature values.

Compound **1** was isolated as a white amorphous powder. Its

*To whom correspondence should be addressed. E-mail: xumin@mail.kib.ac.cn (M. Xu); zhangyj@mail.kib.ac.cn (Y.J. Zhang)

Table 1. ^1H NMR spectroscopic data of compounds 1–5


Pos.	1 ^a	2 ^b	3 ^b	4 ^b	5 ^c
1	3.62, d (12.0); 3.67 ^d	3.45, d (11.8); 3.63, d (11.8)	3.66, d (12.3); 3.71 ^d	3.63, d (12.5); 3.72 ^d	3.39 ^e ; 3.34 ^e
3	5.63, d (8.0)	5.65, d (8.2)	5.63, d (7.1)	5.71, d (8.3)	5.53, d (7.9)
4	4.43, t (8.1)	4.41, t (8.4)	4.44, t (7.2)	4.45, t (8.3)	4.24, br. s
5	3.96, m	3.94, ddd (8.8, 6.6, 2.5)	4.03, td (7.0, 3.5)	4.00, ddd (8.4, 6.9, 2.7)	3.85, td (7.6, 3.2)
6	3.68 ^d ; 3.83 ^d	3.78 ^d ; 3.69, dd (12.1, 2.5)	3.74, dd (12.0, 3.5); 3.84, dd (12.0, 6.8)	3.71 ^d ; 3.86, m	3.61, d (11.4); 3.73, br. s
1'	5.46, d (3.6)	5.58, d (3.6)	5.52, d (3.6)	5.55, d (3.6)	5.23, d (3.3)
2'	3.47 ^d	4.61, dd (10.1, 3.6)	3.57, dd (9.8, 3.6)	3.72 ^d	3.29 ^e
3'	4.17, dd (9.9, 2.5)	3.76 ^d	3.71 ^d	5.36, t (9.7)	3.44 ^e
4'	3.49 ^e	4.15, d (3.6)	4.98 ^e	3.69, t (8.5)	3.15, t (9.3)
5'	3.66, m	3.56, t (9.5)	4.32 ^d	4.29 ddd (10.0, 4.2, 2.0)	4.03, t (8.4)
6'	4.50, dd (12.0, 4.5)	4.50, dd (12.1, 2.1)	4.45 ^d	4.55, dd (12.1, 4.3)	4.18, dd (11.8, 7.1)
	4.60, dd (12.0, 1.9)	4.46, dd (12.1, 4.1)	4.34, dd (12.2, 3.9)	4.63, dd (12.1, 2.0)	4.43, d (11.4)
2'',6''	8.12, d (7.4)	8.14, d (7.3)	8.15, d (7.2)	8.19, d (7.2)	8.04, d (7.6)
3'',5''	7.50, t (7.4)	7.52, t (7.4)	7.51, t (7.2)	7.53, t (7.2)	7.42 ^d
4''	7.61, t (7.4)	7.63, t (7.3)	7.64, t (7.2)	7.62, t (7.2)	7.70 ^d
2''',6'''	8.05, d (7.4)	8.05, d (7.3)	8.07, d (7.2)	8.07, d (7.2)	7.42 ^d
3''',5'''	7.48, t (7.4)	7.50, t (7.3)	7.49, t (7.2)	7.50, t (7.2)	7.56, t (7.7)
4'''	7.61, t (7.4)	7.61, t (7.3)	7.62, t (7.2)	7.60, t (7.2)	7.70 ^d
7'''					7.67, d (15.6)
8'''					6.68, d (15.6)
AcO		2.04, s	2.06, s	2.11, s	

^aRecorded in CD_3OD at 500 MHz; ^bRecorded in CD_3OD at 600 MHz; ^cRecorded in $\text{DMSO}-d_6$ at 600 MHz; ^dOverlapping ^1H NMR signals; ^eOverlapped by solvent

molecular formula was determined to be $\text{C}_{26}\text{H}_{30}\text{O}_{13}$, on the basis of HRESIMS (m/z 573.1578 [$\text{M} + \text{Na}$]⁺). The IR spectrum showed the presence of hydroxyl (3432 cm^{-1}) and carbonyl (1721 cm^{-1}) groups. The ^1H NMR spectrum of **1** (Table 1) displayed characteristic signals of two benzoyl groups [δ_{H} 8.12, 8.05 (each 2H, d, $J = 7.4$ Hz), 7.50, 7.48 (each 2H, t, $J = 7.4$ Hz), 7.61 (2H, t, $J = 7.4$ Hz)]. The ^{13}C NMR (DEPT) spectra of **1** (Table 2) gave 26 carbon signals, including 12 aromatic (δ_{C} 129.8–134.6) and two carbonyl carbons (δ_{C} 167.5 and 168.1) arising from two benzoyl units, and 12 oxygen-bearing aliphatic carbons (δ_{C} 63.8–104.9) due to two hexosyl moieties. Alkaline hydrolysis of **1** with 0.5% NaOH in MeOH yielded sucrose ($[\alpha]_{\text{D}}^{16} + 28.5$),⁸ indicating that **1** is a bisbenzoyl sucrose ester. The ^1H and ^{13}C NMR signals of the sugar units were assigned unambiguously by ^1H - ^1H COSY, HSQC, and HMBC analysis (Tables 1 and 2). In the HMBC spectrum of **1**, a correlation of glucosyl anomeric proton at δ_{H} 5.46 (H-1') with the fructosyl C-2 (δ_{C} 104.9) confirmed the sucrose moiety in **1**. Furthermore, HMBC correlations of δ_{H} 5.63 (H-3) with δ_{C} 167.5 (C-7''), δ_{C} 74.3 (C-4) and δ_{C} 65.3 (C-1), and δ_{H} 4.50 (H-6'a) with δ_{C} 168.1 (C-7'') and δ_{C} 74.9 (C-5') indicated that the two benzoyl units were linked to C-3 and C-6' of sucrose moiety, respectively. Accordingly, compound **1** was determined to be 3,6'-di-*O*-benzoylsucrose.

Compound **2**, a white amorphous powder, gave an [$\text{M} + \text{Na}$]⁺ peak at m/z 615.1675 ($\text{C}_{28}\text{H}_{32}\text{O}_{14}\text{Na}$) in HRESIMS, which was 42 Da more than that of **1**. The ^1H and ^{13}C NMR spectra (Tables 1 and 2) of **2** showed high similarity to those of **1**, except for the appearance of an additional acetyl group [δ_{H} 2.04 (3H, s) and δ_{C} 172.7 and 21.1]. In the HMBC spectrum of **2**, the correlation of δ_{H} 4.61 (H-2') with the acetyl carbonyl carbon at δ_{C} 172.7 was observed, allowing the assignment of the acetyl group located at C-2' of sucrose moiety. This also resulted in the higher field shifted carbon resonances of C-1' and C-3' of **2** as compared with those of **1** (Table 2). Thus, the structure of **2** was established as 3,6'-di-*O*-benzoyl-2'-*O*-acetylsucrose.

Compounds **3** and **4** had the same molecular formula

**Figure 2.** Key HMBC correlations of **1**

$\text{C}_{28}\text{H}_{32}\text{O}_{14}$, on the basis of their HRESIMS (m/z 615.1692 and 615.1689 [$\text{M} + \text{Na}$]⁺ for **3** and **4**, respectively), which exhibited the same molecular weight with that of **2**. The extensive comparison of 1D and 2D NMR data with those of compound **2** suggested that both **3** and **4** had the same 3,6'-di-*O*-benzoylsucrose skeleton, while the difference among **2**, **3** and **4** was the position of the acetyl group. The ^1H and ^{13}C NMR signals of compounds **3** and **4** could be assigned unambiguously by HSQC, and HMBC analysis (Tables 1 and 2), respectively. In the HMBC experiment, H-4' (δ_{H} 4.98) in **3** and H-3' (δ_{H} 5.36) in **4** were correlated with the acetyl carbonyl carbon at δ_{C} 172.0 (**3**) and δ_{C} 172.8 (**4**), respectively, indicating that the acetyl groups in **3** and **4** were located at C-4' and C-3', respectively. Other HMBC correlations confirmed the structures of **3** and **4** as shown in Fig 1. Consequently, compounds **3** and **4** were determined to be 3,6'-di-*O*-benzoyl-4'-*O*-acetylsucrose (**3**) and 3,6'-di-*O*-benzoyl-3'-*O*-acetylsucrose (**4**), respectively.

Compound **5**, a white amorphous powder, possessed a molecular formula $\text{C}_{28}\text{H}_{32}\text{O}_{13}$, as deduced from the HRESIMS (m/z 599.1737 [$\text{M} + \text{Na}$]⁺) and ^{13}C NMR (DEPT) spectra. The ^{13}C NMR spectrum of **5** (Table 2) exhibited the presence of 12 oxygen-bearing carbon signals relating to two hexosyl units, two carbonyl carbons at δ_{C} 165.1 and 166.4, in addition to 12 aromatic methines (δ_{C} 118 to 145) and two aromatic quaternary carbons at δ_{C} 129.8 and 134.1, suggesting the

Table 2. ^{13}C NMR spectroscopic data of compounds 1–5

Pos.	1 ^a	2 ^b	3 ^b	4 ^b	5 ^c
1	65.3, CH ₂	64.8, CH ₂	65.5, CH ₂	65.0, CH ₂	63.2, CH ₂
2	104.9, C	105.4, C	105.5, C	104.9, C	103.2, C
3	80.3, CH	79.3, CH	80.6, CH	80.4, CH	78.1, CH
4	74.3, CH	74.0, CH	74.8, CH	74.3, CH	73.0, CH
5	84.3, CH	84.2, CH	85.0, CH	84.3, CH	83.1, CH
6	63.8, CH ₂	63.7, CH ₂	63.7, CH ₂	63.9, CH ₂	62.6, CH ₂
1'	93.5, CH	91.0, CH	93.3, CH	93.5, CH	91.3, CH
2'	73.1, CH	74.5, CH	73.0, CH	71.3, CH	71.3, CH
3'	72.6, CH	72.2, CH	72.8, CH	76.8, CH	72.9, CH
4'	71.5, CH	72.3, CH	72.5, CH	69.7, CH	70.2, CH
5'	74.9, CH	71.5, CH	70.1, CH	72.6, CH	70.7, CH
6'	65.3, CH ₂	65.0, CH ₂	64.4, CH ₂	65.0, CH ₂	64.6, CH ₂
1''	131.4, C	131.4, C	131.3, C	131.3, C	129.8, C
2'',6''	131.1, CH	131.1, CH	131.1, CH	131.2, CH	129.5, CH
3'',5''	129.9, CH	130.0, CH	129.9, CH	130.0, CH	128.9, CH
4''	134.6, CH	134.8, CH	134.7, CH	134.6, CH	133.6, CH
7''	167.5, C	167.6, C	167.1, C	167.8, C	165.1, C
1'''	131.3, C	131.0, C	131.2, C	131.3, C	134.1, C
2''',6'''	130.8, CH	130.8, CH	130.0, CH	130.8, CH	128.5, CH
3''',5'''	129.8, CH	129.8, CH	129.8, CH	129.8, CH	129.0, CH
4'''	134.5, CH	134.5, CH	134.6, CH	134.5, CH	130.6, CH
7'''	168.1, C	168.1, C	167.8, C	168.0, C	144.8, CH
8'''					118.0, CH
9'''					166.4, C
AcO		21.1, CH ₃ 172.7, C	21.2, CH ₃ 172.0, C	21.3, CH ₃ 172.8, C	


^aRecorded in CD₃OD at 125 MHz; ^bRecorded in CD₃OD at 150 MHz; ^cRecorded in DMSO-*d*₆ at 150 MHz.

presence of two mono-substituted benzene rings and one double bond. The ^1H NMR spectrum (Table 1) of **5** showed the presence of two *trans*-coupled olefinic protons at δ_{H} 6.68 and 7.67 (each 1H, d, $J = 15.6$ Hz). The above data revealed that compound **5** was also an analogue of **1**, except for the additional double bond with *trans* configuration. All the proton and carbon signals in **5** were assigned unambiguously by HSQC and HMBC analysis (Tables 1 and 2). In the HMBC spectrum of **5** (Figure 3), correlations of the olefinic proton at δ_{H} 6.68 (H-8''') with one aromatic quaternary carbon at δ_{C} 134.1 (C-1''') and another olefinic proton at δ_{H} 7.67 (H-7''') with one carbonyl carbon at δ_{C} 166.4 (C-9''') indicated the additional *trans* double bond belonging to a *trans* cinnamoyl moiety. The HMBC correlations of the glucosyl H-6' (δ_{H} 4.43 and 4.18) with C-9''' (δ_{C} 166.4) and glucosyl C-4' (δ_{C} 70.2) revealed the linkage of the (*E*)-cinnamoyl moiety with C-6'. Moreover, the HMBC correlations of the fructosyl H-3 (δ_{H} 5.53) with δ_{C} 165.1 (C-7'''), 63.2 (C-1) and 73.0 (C-4) suggested the benzoyl unit located at C-3. Other HMBC correlations (Figure 3) further confirmed the structures of **5** as shown in Fig 1. Therefore, compound **5** was determined to be 3-*O*-benzoyl-6'-*O*-(*E*)-cinnamoylsucrose.

The isolated compounds were evaluated for their cytotoxicities against five human cancer cell lines (breast cancer MCF-7, hepatocellular carcinoma SMMC-7721, human myeloid leukemia HL-60, colon cancer SW480, and lung cancer A-549). All of them showed no cytotoxic activity against the five human cancer cell lines at a concentration of 40 μM .

Experimental Section

General Experimental Procedures. Optical rotations were measured with a HORIBA SEPA-300 high-sensitive polarimeter. IR spectra were measured on a Bio-Rad FTS-135 series spectrometer. UV spectra were recorded on a Shimadzu UV2401A ultraviolet-visible spectrophotometer. ESIMS and HRESIMS were run on an API QSTAR Pular-1 spectrometer.

**Figure 3.** Key HMBC correlations of **5**

NMR spectra measured in methanol-*d*₄ or DMSO-*d*₆ solution and recorded on a Bruker AV-400, DRX-500 or AV III-600 spectrometer, using TMS as an internal standard. Chemical shifts were reported in units of δ (ppm) and coupling constants (J) were expressed in Hz. Column chromatography (CC) were carried out over silica gel (200–300 mesh, Qingdao Haiyang Chemical Co., Ltd., Qingdao, China), Diaion HP20SS (Mitsubishi Chemical Industry, Ltd., Tokyo, Japan), and Sephadex LH-20 (25–100 μm , Pharmacia Fine Chemical Co., Ltd., Uppsala, Sweden). Pre-coated silica gel plates (Qingdao Haiyang Chemical Co., Ltd., Qingdao, China) were used for TLC. Detection was done under UV light (254 and 365 nm) and by spraying the plates with 10% sulfuric acid followed by heating. An Agilent series 1260 (Agilent Technologies) were used for HPLC. An Agilent ZORBAX SB-C18 column 5 μm 143 \AA column (250 mm \times 9.4 mm) were used for semi-preparative HPLC separations.

Plant Material. The whole plant of *P. cochinchinensis* was collected from Guangdong Province, China, on December 2011. A voucher specimen (KUN-1215860) was deposited at the State Key Laboratory of Phytochemistry and Plant

Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, and was identified by Prof. Xiao-Min Fang.

Extraction and Isolation. The air-dried and powdered whole plants of *P. cochinchinensis* (3.2 kg) were extracted with MeOH (3 times, 3h each time) under reflux at 60 °C. Evaporation of the solvent under vacuum gave a residue (270 g), which was suspended in water and then extracted sequentially with chloroform and butanol. The butanol extract (90 g) was chromatographed on Diaion HP20SS eluting with a gradient of MeOH-H₂O (1:9 → 9:1, finally MeOH), to give 6 fractions F1–F6. F6 (16 g) was subjected to Sephadex LH-20 CC eluted with a gradient MeOH-H₂O (1:9 → 6:4) to attain 3 fractions (F601–F603). F602 was fractionated through silica gel CC, using CHCl₃-MeOH-H₂O (90:10:1 and 80:20:2) as solvent to afford 4 subfractions (F0201–F0204). F0201 (60 mg) was purified by semi-preparative HPLC (16%, MeCN-H₂O) to furnish **6** (*t_R* 20 min, 17 mg). F0202 was fractionated through semi-preparative HPLC (17%, MeCN-H₂O) to afford two fractions at 15 min and at 18 min. The fraction at 15 min was purified through semi-preparative HPLC (17%, MeCN-H₂O) to attain **5** (*t_R* 14 min, 2 mg), and the other fraction at 18 min was further purified by semi-preparative HPLC (18%, MeCN-H₂O) to afford **1** (*t_R* 15 min, 13 mg), **2** (*t_R* 16 min, 4 mg), **3** (*t_R* 17 min, 6 mg), and **4** (*t_R* 18 min, 3 mg). F5 (2.0 g) was chromatographed through Sephadex LH-20 CC eluted with a gradient MeOH-H₂O (1:9 → 6:4) to attain 2 fractions (F501–F502). F502 (315 mg) was purified through silica gel CC, using CHCl₃-MeOH-H₂O (90:10:1 and 80:20:2) as solvent, and then subjected to semi-preparative HPLC (18%, MeCN-H₂O) to afford **7** (88 mg).

3,6'-Di-O-benzoylsucrose (1): white amorphous powder; $[\alpha]_D^{22} + 30.2$ (*c* 0.18, MeOH). UV (MeOH) λ_{\max} (log ϵ) 200 (4.24), 229 (4.33), 273 (3.26) nm; IR (KBr) ν_{\max} 3432, 2924, 1721, 1603, 1453, 1379, 1279, 1124, 1070, 994 cm⁻¹; ¹H and ¹³C NMR data, see Tables 1 and 2; negative ESIMS *m/z* 585 [M + Cl]⁻; positive HRESIMS *m/z* 573.1578 [M + Na]⁺ (calcd for C₂₆H₃₀O₁₃Na, 573.1578).

3,6'-Di-O-benzoyl-2'-O-acetylsucrose (2): white amorphous powder; $[\alpha]_D^{22} + 30.9$ (*c* 0.08, MeOH). UV (MeOH) λ_{\max} (log ϵ) 200 (4.28), 229 (4.36), 273 (3.31) nm; IR (KBr) ν_{\max} 3427, 2924, 2854, 1721, 1631, 1603, 1453, 1278, 1123, 1070, 1052, 994 cm⁻¹; ¹H and ¹³C NMR data, see Tables 1 and 2; negative ESIMS *m/z* 627 [M + Cl]⁻; positive HRESIMS *m/z* 615.1675 [M + Na]⁺ (calcd for C₂₈H₃₂O₁₄Na, 615.1684).

3,6'-Di-O-benzoyl-4'-O-acetylsucrose (3): white amorphous powder; $[\alpha]_D^{22} + 26.0$ (*c* 0.17, MeOH). UV (MeOH) λ_{\max} (log ϵ) 200 (4.06), 229 (4.18), 273 (3.09) nm; IR (KBr) ν_{\max} 3425, 2926, 1721, 1603, 1453, 1376, 1279, 1119, 1052, 939 cm⁻¹; ¹H and ¹³C NMR data, see Tables 1 and 2; positive ESIMS 615 [M + Na]⁺; positive HRESIMS *m/z* 615.1692 [M + Na]⁺ (calcd for C₂₈H₃₂O₁₄Na, 615.1684).

3,6'-Di-O-benzoyl-3'-O-acetylsucrose (4): white amorphous powder; $[\alpha]_D^{22} + 29.3$ (*c* 0.07, MeOH). UV (MeOH) λ_{\max} (log ϵ)

200 (4.23), 229 (4.28), 273 (3.20) nm; IR (KBr) ν_{\max} 3427, 2925, 1721, 1603, 1452, 1376, 1279, 1120, 1054, 1000 cm⁻¹; ¹H and ¹³C NMR data, see Tables 1 and 2; positive ESIMS 615 [M + Na]⁺; positive HRESIMS *m/z* 615.1689 [M + Na]⁺ (calcd for C₂₈H₃₂O₁₄Na, 615.1684).

3-O-Benzoyl-6'-O-(E)-cinnamoylsucrose (5): white amorphous powder; $[\alpha]_D^{22} + 25.6$ (*c* 0.07, MeOH). UV (MeOH) λ_{\max} (log ϵ) 201 (4.28), 220 (4.26), 278 (4.21) nm; IR (KBr) ν_{\max} 3439, 2924, 1709, 1635, 1452, 1281, 1179, 1071, 1056, 1000 cm⁻¹; ¹H and ¹³C NMR data, see Tables 1 and 2; negative ESIMS *m/z* 611 [M + Cl]⁻; positive HRESIMS *m/z* 599.1737 [M + Na]⁺ (calcd for C₂₈H₃₂O₁₃Na, 599.1737).

Alkaline Hydrolysis of 1. A mixture of **1** (6 mg), 0.5% NaOH (0.5 mL), and MeOH (3 mL) was stirred at room temperature for 6 h. The reaction mixture was neutralized with 1 N HCl and extracted with CHCl₃ (3 × 10 mL). The aqueous layer was applied to silica gel CC, eluting with CHCl₃-MeOH-H₂O (6:4:1), to give sucrose (1.0 mg): $[\alpha]_D^{16} + 28.5$ (*c* 0.8, H₂O); positive ESIMS *m/z* 365 [M + Na]⁺. The alkaline hydrolysis of **2–5** was not performed due to the limited amount of samples. The disaccharides of **2–5** were determined to be sucrose on the basis of biogenetic arguments of **1**.⁸

Cytotoxicity Assay. Five human cancer cell lines, human myeloid leukemia HL-60, hepatocellular carcinoma SMMC-7721, lung cancer A-549 cells, breast cancer MCF-7, and colon cancer SW480, were used in the cytotoxic assay. All the cells were cultured in RPMI-1640 or DMEM medium (Hyclone, USA), supplemented with 10% fetal bovine serum (Hyclone, USA). The cytotoxicity assay was performed according to the MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) method in 96-well microplates.⁹ Briefly, adherent cells (100 μ L) was seeded into each well of 96-well cell culture plates and allowed to adhere for 12 h before drug addition, while suspended cells were seeded just before drug addition with an initial density of 0.5 × 10⁵–1 × 10⁵ cells/mL. Each tumor cell line was exposed to the test compound dissolved in DMSO in triplicates for 48 h at 37 °C, with DDP and taxol (Sigma, USA) as positive controls. Then, MTT (50 μ L) was added to each well, and the tumor cells were incubated for another 4 h at 37 °C. After the supernatant liquor was removed, SDS (200 μ L) was added to each well. The optical density was measured at 595 nm on a microplate reader. Cell viability was detected and a cell growth curve was graphed. IC₅₀ values were calculated by Reed and Muench's method.¹⁰

Electronic Supplementary Material

Supplementary material is available in the online version of this article at <http://dx.doi.org/10.1007/s13659-013-0026-7> and is accessible for authorized users.

Acknowledgments

The authors are grateful to the members of the Analytical Group in State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, for measurements of all spectra. We would like to acknowledge

Prof. Yan Li for cytotoxicity assay. This work was supported by the NSFC 21002105, the 973 Program of Ministry of Science and Technology of China (2011CB915503), the Fourteenth Batch Candidates of the Young Academic Leaders of Yunnan Province (Min XU, 2011CI044) and by West Light Foundation of the Chinese Academy of Sciences.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] Zhang, Y. J.; Tanaka, T.; Iwamoto, Y.; Yang, C. R.; Kouno, I. *J. Nat. Prod.* **2000**, *63*, 1507–1510.
- [2] Zhang, Y. J.; Abe, T.; Tanaka, T.; Yang, C. R.; Kouno, I. *J. Nat. Prod.* **2001**, *64*, 1527–1532.
- [3] Ratnayake, R.; Covell, D.; Ransom, T. T.; Gustafson, K. R.; Beutler, J. A. *Org. Lett.* **2009**, *11*, 57–60.
- [4] Liu, Q.; Wang, Y. F.; Chen, R. J.; Zhang, M. Y.; Wang, Y. F.; Yang, C. R.; Zhang, Y. J. *J. Nat. Prod.* **2009**, *72*, 969–972.
- [5] Pettit, G. R.; Schaufelberger, D. E.; Nieman, R. A.; Dufresne, C.; Saenz-Renauld, J. A. *J. Nat. Prod.* **1990**, *53*, 1406–1413.
- [6] Shen, Y. C.; Lin, S. L. *Planta Med.* **1996**, *62*, 515–518.
- [7] Shen, Y. C.; Lin, S. L.; Chen, C. C. *Phytochemistry* **1997**, *44*, 891–895.
- [8] Dong, L. B.; He, J.; Li, X. Y.; Wu, X. D.; Deng, X.; Xu, G.; Peng, L. Y.; Zhao, Y.; Li, Y.; Gong, X.; Zhao, Q. S. *Nat. Prod. Bioprospect.* **2011**, *1*, 41–47.
- [9] Chang, S. L.; Chang, C. L.; Chiang, Y. M.; Hsieh, R. H.; Tzeng, C. R.; Wu, T. K.; Sytwu, H. K.; Shyur, L. F.; Yang, W. C. *Planta Med.* **2004**, *70*, 1045–1051.
- [10] Reed, L. J.; Muench, H. *Am. J. Hyg.* **1938**, *27*, 493–497.