

## Flash droughts present a new challenge for subseasonal-to-seasonal prediction

— [Source link](#) 





Angeline G. Pendergrass, Gerald A. Meehl, Roger S. Pulwarty, Michael T. Hobbins ...+19 more authors

**Institutions:** National Center for Atmospheric Research, Earth System Research Laboratory,  
Cooperative Institute for Research in Environmental Sciences, University of California, Irvine ...+10 more institutions

**Published on:** 01 Mar 2020 - Nature Climate Change (Nature Publishing Group)

### Related papers:

- [Flash Droughts: A Review and Assessment of the Challenges Imposed by Rapid-Onset Droughts in the United States](#)
- [A Multiscalar Drought Index Sensitive to Global Warming: The Standardized Precipitation Evapotranspiration Index](#)
- [A Methodology for Flash Drought Identification: Application of Flash Drought Frequency across the United States](#)
- [Examining Rapid Onset Drought Development Using the Thermal Infrared–Based Evaporative Stress Index](#)
- [The drought monitor](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/flash-droughts-present-a-new-challenge-for-subseasonal-to-ytl0iqv2om>



## Flash droughts present a new challenge for subseasonal-to-seasonal prediction

Item Type	Article
Authors	Pendergrass, Angeline G.; Meehl, Gerald A.; Pulwarty, Roger; Hobbins, Mike; Hoell, Andrew; AghaKouchak, Amir; Bonfils, Céline J. W.; Gallant, Ailie J. E.; Hoerling, Martin; Hoffmann, David; Kaatz, Larna; Lehner, Flavio; Llewellyn, Dagmar; Mote, Philip; Neale, Richard B.; Overpeck, Jonathan T.; Sheffield, Amanda; Stahl, Kerstin; Svoboda, Mark; Wheeler, Matthew C.; Wood, Andrew W.; Woodhouse, Connie A.
Citation	Pendergrass, A.G., Meehl, G.A., Pulwarty, R. et al. Flash droughts present a new challenge for subseasonal-to-seasonal prediction. Nat. Clim. Chang. 10, 191–199 (2020). <a href="https://doi.org/10.1038/s41558-020-0709-0">https://doi.org/10.1038/s41558-020-0709-0</a>
DOI	<a href="https://doi.org/10.1038/s41558-020-0709-0">10.1038/s41558-020-0709-0</a>
Publisher	NATURE PUBLISHING GROUP
Journal	NATURE CLIMATE CHANGE
Rights	This is a U.S. government work and not under copyright protection in the U.S.; foreign copyright protection may apply 2020.
Download date	30/05/2022 15:10:08
Item License	<a href="https://creativecommons.org/publicdomain/mark/1.0/">https://creativecommons.org/publicdomain/mark/1.0/</a>
Version	Final published version

Link to Item	<a href="http://hdl.handle.net/10150/640941">http://hdl.handle.net/10150/640941</a>
--------------	---



# Flash droughts present a new challenge for subseasonal-to-seasonal prediction

Angeline G. Pendergrass<sup>1</sup>✉, Gerald A. Meehl<sup>1</sup>, Roger Pulwarty<sup>2</sup>, Mike Hobbins<sup>2,3</sup>, Andrew Hoell<sup>2</sup>, Amir AghaKouchak<sup>4</sup>, Céline J. W. Bonfils<sup>5</sup>, Ailie J. E. Gallant<sup>6,7</sup>, Martin Hoerling<sup>2</sup>, David Hoffmann<sup>6,7</sup>, Larna Kaatz<sup>8</sup>, Flavio Lehner<sup>1</sup>, Dagmar Llewellyn<sup>9</sup>, Philip Mote<sup>10</sup>, Richard B. Neale<sup>1</sup>, Jonathan T. Overpeck<sup>11</sup>, Amanda Sheffield<sup>12</sup>, Kerstin Stahl<sup>13</sup>, Mark Svoboda<sup>14</sup>, Matthew C. Wheeler<sup>15</sup>, Andrew W. Wood<sup>1</sup> and Connie A. Woodhouse<sup>16</sup>

**Flash droughts are a recently recognized type of extreme event distinguished by sudden onset and rapid intensification of drought conditions with severe impacts. They unfold on subseasonal-to-seasonal timescales (weeks to months), presenting a new challenge for the surge of interest in improving subseasonal-to-seasonal prediction. Here we discuss existing prediction capability for flash droughts and what is needed to establish their predictability. We place them in the context of synoptic to centennial phenomena, consider how they could be incorporated into early warning systems and risk management, and propose two definitions. The growing awareness that flash droughts involve particular processes and severe impacts, and probably a climate change dimension, makes them a compelling frontier for research, monitoring and prediction.**

Drought is perhaps the most complex and least understood of all “weather and climate extremes”<sup>1</sup>. Droughts can span timescales from a few weeks to decades, and spatial scales from a few kilometres to entire regions. Their impacts usually develop slowly, are often indirect and can linger for long after the end of the drought itself. The drought risk, therefore, is often underestimated and continues to remain a ‘hidden’ hazard<sup>2</sup>. A comprehensive overview of traditional drought characteristics, processes, mechanisms and impacts is provided in ref. <sup>3</sup>.

In a future warmer climate, droughts are likely to increase in duration and intensity in many regions of the world<sup>4,5</sup>. A better understanding of drought phenomena, especially of the physical processes leading to drought, their propagation through the hydrological cycle, and the societal and environmental vulnerability to drought and its wide-ranging impacts is more important than ever. The key challenge is to move from a reactive society, responding to impacts, to a proactive society that is resilient and adapted to drought risk—that is, adopts proactive risk management strategies<sup>3,6</sup>.

Droughts whose impacts arise in part from their long duration, such as the Dust Bowl and the 2011–2015 California drought, have formed strong imagery in the United States, and ‘megadroughts’ lasting more than 20 years have also been documented in tree-ring records. Much research has been conducted on aspects of drought that play out over multiple years, but more recently attention has been drawn to the rapid development of some drought events, in

the space of a few weeks: ‘flash droughts’, a specific definition for which we will provide below. These events, distinguished by their sudden onset and rapid intensification, can have severe impacts<sup>7,8</sup>. Flash droughts develop on the subseasonal-to-seasonal (S2S) timescale (weeks to months) and present a new challenge for prediction efforts on that timescale, which are currently surging in interest<sup>9</sup>.


One flash drought that brought attention to the phenomenon occurred in the US Midwest in 2012<sup>8,10</sup> (Fig. 1). The areal extent of abnormally dry conditions expanded from 30% of the continental United States (CONUS) in May 2012 to over 60% by August. This event had considerable impacts on agriculture and water-borne transportation in the region. Although other rapidly developing droughts had been identified before<sup>11</sup>, the widespread impacts of the 2012 event caught the attention of the US public and leadership.

Flash drought is not confined to the United States<sup>12</sup>. For example, processes that can produce flash droughts are foci of research in China<sup>13,14</sup>. In southern Queensland, Australia, a flash drought in early 2018 de-vegetated the landscape and drove livestock numbers to their lowest level in a century, a significant impact for agriculture<sup>15</sup>.

A drought monitoring and early-warning system is the foundation of effective, proactive drought policy because it enables notice of potential and impending drought conditions. It identifies climate and water-resource trends, and detects the emergence or probability of occurrence and the likely severity of droughts and their impacts. Reliable information must be communicated in a timely manner

<sup>1</sup>National Center for Atmospheric Research, Boulder, CO, USA. <sup>2</sup>NOAA/Earth System Research Laboratory/Physical Sciences Division, Boulder, CO, USA.

<sup>3</sup>Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder, Boulder, CO, USA. <sup>4</sup>Department of Civil and Environmental Engineering, and Department of Earth System Science, University of California, Irvine, CA, USA. <sup>5</sup>Program for Climate Model Diagnosis and Intercomparison, Lawrence Livermore National Laboratory, Livermore, CA, USA. <sup>6</sup>School of Earth, Atmosphere and Environment, Monash University, Clayton, Victoria, Australia. <sup>7</sup>ARC Centre of Excellence for Climate Extremes, Monash University, Clayton, Victoria, Australia. <sup>8</sup>Denver Water, Denver, CO, USA. <sup>9</sup>US Department of Interior, Bureau of Reclamation, Albuquerque, NM, USA. <sup>10</sup>Oregon Climate Change Research Institute, and Graduate School, Oregon State University, Corvallis, OR, USA. <sup>11</sup>School for Environment and Sustainability, University of Michigan, Ann Arbor, MI, USA. <sup>12</sup>NOAA/NIDIS, Scripps Institution of Oceanography, La Jolla, CA, USA. <sup>13</sup>University of Freiburg, Freiburg, Germany. <sup>14</sup>National Drought Mitigation Center, University of Nebraska–Lincoln, Lincoln, NE, USA. <sup>15</sup>Australian Bureau of Meteorology, Melbourne, Victoria, Australia. <sup>16</sup>School of Geography and Development, University of Arizona, Tucson, AZ, USA. ✉e-mail: [apgrass@ucar.edu](mailto:apgrass@ucar.edu)



**Fig. 1 | Evolution of a flash drought across the US Midwest in 2012.**

**a–d**, Evaporative drought (ED) categories based on 2-week Evaporative Demand Drought Index (EDDI) at 5-week intervals during the drought onset. **e–h**, US Drought Monitor (USDM). **i**, Percent of High Plains region in USDM categories from 1 June to 3 July 2012. Adapted with permission from: **e–h**, ref. <sup>85</sup>; **i**, US National Drought Mitigation Center.

to water and land managers, policy makers and the public through appropriate communication channels to trigger actions documented in a drought plan, which is particularly critical for flash droughts. That information, if used effectively, can form the basis for reducing vulnerability and improving mitigation and response capacities of people and systems at risk.

In this Perspective, we build on a recent review of flash droughts<sup>8</sup> and discuss the observational and predictive skill of key processes with an eye towards impact assessment and early warning of flash drought. We highlight the current understanding of the physical processes that can drive flash droughts, the existing capabilities of

predicting them, and what is needed to make progress to establish the predictability and effective early warning of flash droughts on S2S timescales. Following earlier suggestions of possible definitions for flash droughts<sup>8,16</sup>, we propose, for consideration by the community, two quantitative definitions for flash drought that can be used for applications related to operations, analysis of observations, model simulations of present and future climate, and assessing S2S initialized model predictions.

### Physical processes that produce flash drought


To illustrate the physical processes involved with producing a flash drought, we consider another recent flash drought, in the US northern Great Plains in 2017. This event shows some recurring flash-drought characteristics, including precipitation deficit and above-average temperatures preceding or coinciding with a rapid soil-moisture decline (Fig. 2). Precipitation deficits began before April, when precipitation would climatologically increase. Soil moisture was nonetheless high in April, but continued precipitation deficits throughout the month eroded it slowly at first, before a rapid decline in May.

We can examine the physical processes driving the moisture balance of the land surface to understand mechanisms that can lead to rapid drought intensification<sup>17,18</sup>. Moisture flux into the surface is driven by precipitation. Like other aspects of drought, precipitation deficit often plays an important role<sup>19</sup>. Moisture flux away from the surface—evapotranspiration (ET)—can also play an important role in flash drought, driving feedbacks between the land and atmosphere. An important concept is the demand for moisture from the atmosphere—evaporative demand, which is the amount of evaporation that would occur given an unlimited supply of moisture. Evaporative demand can be thought of as the ‘thirst’ of the atmosphere. It both drives and responds to ET. Starting from a state with sufficient soil moisture (energy-limited conditions; Fig. 3), evaporative demand and evaporation vary together—when evaporative demand increases, evaporation follows. With enough evaporation and no replenishment, surface moisture eventually becomes insufficient to supply further water for evaporation; water becomes the limiting factor. Under water-limited conditions, further increases in evaporation can no longer continue, and evaporation decreases. If the same factors that had been driving increases in evaporative demand persist, then evaporative demand will diverge from evaporation. Meanwhile, sensible heat flux increases instead of evaporation, which increases near-surface air temperature and vapour pressure deficit, and thus also evaporative demand—an amplifying feedback<sup>20–22</sup>.

Although much of the focus on flash droughts has been in humid regions, flash droughts and their impacts are also a concern in semi-arid and arid regions where evaporative demand usually exceeds evapotranspiration (locations that start on the right side of Fig. 3; see ‘Impacts-based early warning’ below). Starting from a dry, moisture-limited state, flash droughts in arid regions can be driven by precipitation deficits, and amplification of warm air temperatures by sensible heat flux feedbacks is also of concern.

The local moisture imbalance during flash drought is conditioned by large-scale atmospheric circulation. The large-scale circulation can modify the frequency and intensity of precipitation, and it can increase evaporative demand by reducing cloud cover (which increases incoming solar radiation at the surface), increasing wind speeds and/or increasing temperatures<sup>16,23,24</sup>. In the midlatitudes in summer, this can involve a persistent ‘blocking’ pattern, with a strong quasi-stationary ridge of positive geopotential height anomalies and associated anomalously high surface pressure<sup>16</sup>.

Large-scale atmospheric circulation associated with flash droughts can vary from one event to the next and between different regions. While moisture-bearing storms were largely absent during the 2012 US Midwest flash drought, the atmospheric circulation during the event varied from one month to the next<sup>23</sup>. For the



**Fig. 2 | US Northern Great Plains flash drought in May 2017.** **a**, Soil-moisture percentile (top 1 m) from University of Washington simulation of the variable infiltration capacity (VIC) model<sup>86</sup>, forced by an estimate of the time-varying meteorology<sup>87,88</sup>. **b**, **c**, Precipitation (**b**) and daily maximum temperature (**c**) in 2017 depicted as departures from the long-term climatology (solid black lines) from a collection of Global Historical Climatology Network (GHCN-D) stations. **d**, Rank of accumulated May–July 2017 precipitation relative to the 1895–2017 climatology. The time series are data averaged over eastern Montana (demarcated by the dotted line in the right panel). Adapted with permission from ref. <sup>73</sup>.

southern US Great Plains, the atmospheric circulation associated with rapid declines in soil moisture in conjunction with precipitation deficits can be different from that associated with such declines in conjunction with heat waves<sup>24</sup>.

Flash droughts may be triggered or exacerbated by compound extreme events—extremes of multiple factors that occur simultaneously<sup>25</sup>. A classic example would be an extreme deficit of precipitation coinciding with a heat wave, such as occurred in southern Queensland in January 2018<sup>15</sup>. If these are superimposed on more slowly evolving factors, such as a building soil-moisture deficit, rapid onset or intensification of drought conditions can result.

Vegetation type can also influence flash drought through its mediating role in transpiration. Trees become moisture-stressed over the course of long-term drought, whereas crops and pasture can be moisture-stressed much more quickly and might be more sensitive to moisture in the upper soil layer.


### The challenge of drought for S2S prediction

Compared with slowly evolving droughts, the relatively fast development timescale of flash droughts requires different approaches to monitoring and prediction. Many drought monitoring and prediction products are updated at monthly or at most weekly timescales. Given a flash drought's onset timescale of only a few weeks, these are not sufficient. Instead, products that update daily are required. This provides an opportunity to leverage synoptic weather forecasts in combination with seasonal forecasting efforts that have recently become available at shorter timescales, such as the SubX system<sup>26</sup>.

Prediction efforts focused on flash drought are currently in their infancy. One key challenge is skilfully forecasting precipitation deficit on the S2S timescale. However, for a successful flash-drought prediction, more is needed than just a forecast of deficient precipitation. Prediction skill is also required for other potential ingredients of rapid drought onset and intensification: high temperatures, low humidity, strong winds and excess insolation. In the 2012 US Midwest event, high temperatures and precipitation deficits may have been driven by a blocking high, whereas the substantial soil-moisture deficit may have been due to anomalous seasonal circulation associated with La Niña<sup>27</sup>. For other flash-drought events and locations, different processes and phenomena probably contribute to or affect development, such as land–atmospheric interaction, the Madden–Julian Oscillation, the Southern and Northern Annular Modes, and the Indian Ocean Dipole. Each of these has been argued to provide or influence predictability of surface-climate variables on timescales relevant for flash drought<sup>28–31</sup>, and so these processes and phenomena are fundamental to the prospects of S2S prediction<sup>32</sup>.

Global coupled prediction systems show some S2S skill for precipitation<sup>33–35</sup> and temperature<sup>35–38</sup>. Seasonal forecasts of evaporative demand are more skilful than for precipitation over the CONUS<sup>39</sup>, and at least as skilful globally<sup>40</sup>, although skill for extreme conditions on subseasonal timescales, which may be more relevant for flash drought, has not been established. Predictions are only as accurate as the models that make them. In the case of global climate models, which are the primary tool for S2S prediction systems, there are significant biases.





**Fig. 3 | The response of evaporative demand and evapotranspiration to feedbacks from drying land.** Schematic evolution of evaporative demand ( $E_0$ ), evapotranspiration (ET) and surface moisture availability, starting from a wet (energy-limited, left side) state, which has enough water for ET to further increase if additional energy is added to the surface, and developing into a dry (water-limited, right side) state. Adapted with permission from ref. <sup>89</sup>.

For example, a challenge for US flash-drought prediction is a summertime dry and warm bias over the central United States in many models<sup>41</sup>. Another key factor for prediction is the fidelity of teleconnections; some models have biased Madden–Julian Oscillation teleconnections<sup>42</sup> that could play a role in flash-drought predictions. Furthermore, land surface models underestimate characteristics of evaporative drought<sup>43</sup>.

Establishing predictability and credibility of predictions present considerable challenges. One aspect is the number of past flash droughts needed to build up a large enough set of samples to test hindcast efficacy. But a property of a flash drought is that it is an unusual event. If the expected return period were more than a year, then testing predictability using hindcasts would require over 20 years of hindcasts; this is more than is available for some current operational S2S prediction systems<sup>44</sup>. Achieving credible predictions will be a continuing challenge with limited computing resources that face competing demands from increased model resolution, ensemble size, and the number and complexity of physical processes.

Other challenges for flash-drought prediction lie in our ability to monitor the current state of the land surface and soil, and to use this information to initialize forecast models. The initial state of the soil-moisture profile is expected to have greater impact on S2S predictions than on shorter or longer timescales<sup>45</sup>. Despite recent improvement, accurate monitoring of soil moisture is still poor compared with many meteorological variables. Perhaps even further afield from operational systems, but still of potential importance, are interactions between vegetation and the land surface. Dynamic vegetation models (such as eco-demographic models<sup>46</sup>) are becoming available, but initializing them in an operational context will present another challenge.

### Context within longer droughts and climate change

The factors driving flash drought can change with climate variability and change on longer timescales, but only a few studies have examined observed regional trends in flash drought (using varied definitions)<sup>47–49</sup> so how they could be affected by different climatic background states remains unclear. In this section, we consider the context within which flash droughts occur, and how climate

background states, multidecadal variability and climate change can influence flash droughts.

Human influence has been identified on various aspects of hydroclimate, including droughts<sup>50,51</sup>; external forcing that drives anthropogenic climate change will significantly change the background climate state as we move further into the twenty-first century. Future changes to precipitation, temperature and atmospheric circulation will all induce changes to surface water availability and evaporative demand<sup>52</sup> and would thus affect flash drought. Aridity, defined in terms of evaporative demand, increases in many drought-prone regions in climate-change projections<sup>53</sup> and also influences soil moisture<sup>54</sup>. But how evaporative demand is formulated—via temperature-dependent measures like the Palmer Drought Severity Index, versus more comprehensive measures—can alter its projected response<sup>55–58</sup>. Actual evaporation and its changes are mediated by vegetation and growing season length, which can counter or exacerbate increasing evaporative demand<sup>59–62</sup>. How these changes in aridity, evaporation and land–atmosphere feedbacks<sup>63</sup> affect flash drought should be a research priority.

Flash droughts can manifest as discrete drought episodes (for example the 2012 US Midwest drought<sup>23,64</sup>), but they may also manifest as a rapid increase in severity from a longer-term drought already in progress. If they are not terminated, they may continue into a period of longer-lasting drought (for example the 2018 eastern Australian drought<sup>15</sup>). Flash droughts can be embedded within climate variability occurring at decadal and longer timescales; the characteristics of the more slowly varying climate will influence the impact of a flash drought. Centuries-long records of climate from palaeoclimatic data are useful<sup>52</sup> for understanding how short, severe droughts that might have developed rapidly are distributed over longer timescales and under a variety of climate conditions.

Although temporal resolution of even the highest-quality palaeoclimatic data is insufficient to capture subseasonal timescales, these records can nonetheless provide insights on the frequency and distribution of extreme single-year or multi-year drought events. Annually resolved tree-ring reconstructions of streamflow for the Colorado River document these extreme occurrences over the past 1,200 years under varying baseline climates<sup>65</sup>. If a ‘palaeo flash drought’ is defined as a year with flow less than two standard deviations below average, then it is possible to identify and characterize periods during which palaeo flash droughts occur. For example, in the medieval period (900–1300 CE), characterized by persistent droughts and temperatures warmer than during any period until the past few decades in southwestern North America<sup>66</sup>, the mid-twelfth-century period of persistent drought had no occurrences of flash-drought years in Colorado River streamflow<sup>65</sup>. The thirteenth century, which was also dry but less persistently so, contained two flash-drought years. However, the most notable cluster of flash-drought years occurred between 1495 and 1506, a 12-year period during which four flash droughts occurred; this was not a period of particularly persistent drought. Similar behaviour can be found in streamflow reconstruction of the Upper Rio Grande<sup>67</sup>, where only one of three flash-drought clusters in a record over 500 years long was associated with persistent drought conditions. Furthermore, palaeo reconstructions contain years in which runoff efficiency is lower than expected from annual streamflow alone<sup>68,69</sup>. Notwithstanding reconstruction uncertainties, such years could have harboured flash droughts that affected runoff efficiency while leaving little imprint on annual streamflow.

Slowly varying or changing background states present an additional challenge for S2S prediction of flash drought since the climatic base state can alter S2S predictive skill. For example, a decline in predictability of the Eastern Pacific El Niño in the early twenty-first century has been attributed to a change in the background state of the tropical Pacific<sup>70</sup>. Potential changes in S2S skill as the climate baseline evolves need further investigation.

### Proposed definition

We have seen that the physical processes leading to flash drought are a matter of ongoing research; we will see that specific impacts of flash drought are too. To aid in the identification of flash droughts, we adopt three principles to describe them that are broadly consistent with previously proposed definitions<sup>8,16</sup> and that lend themselves to analysis, yet remain useful for monitoring and prediction. We then apply these principles to propose two specific quantitative definitions of flash drought: one for US operations with the US Drought Monitor (<https://droughtmonitor.unl.edu/AboutUSDM/WhatIsTheUSDM.aspx>), and another that can be used globally for analysis of observations and climate models.

The first principle is that the event should involve a rapid onset and intensification, as emphasized previously<sup>8</sup>. To adequately reflect the rapid onset rate, the onset period should be short enough to distinguish flash droughts from the general population of droughts, encompassing the upper tail of the distribution of events. The second principle is that the intensification rate should be high, as advocated previously<sup>8</sup>. The third principle is that the event should end in a state severe enough to qualify as drought. These principles should apply across drought types, sectors, regions and seasons, and not only be adapted to definitions based on different variables (precipitation or drought indices) but also offer broad guidelines for the development of specific flash-drought definitions. An additional principle that would be desirable is that the event should have impacts to qualify as a flash drought, but this requires more work to quantitatively document past and potential future impacts.

Next, we propose two quantitative definitions of subseasonal flash drought that encompass some of the principles outlined above, using the 2012 US Midwest event as guidance. These definitions follow from recommendations made previously<sup>8</sup> and are designed to be quantitative measures that can be evaluated in the context of past flash droughts, used operationally, and also applied to model simulations analyses and forecast evaluations. Their purpose is not to make further prescriptions, but rather to provide concrete definitions for scrutiny and analysis by the community. The goal is to arrive at quantitative, usable definitions—whether these or a revision.

The first definition is based on Evaporative Demand Drought Index (EDDI; <https://www.esrl.noaa.gov/psd/eddi/>), which is an experimental drought monitoring and early-warning guidance tool based on how anomalous the evaporative demand is for a given location. The caveat accompanying EDDI is that for a flash drought to develop, the enhanced atmospheric demand should not be compensated by increased precipitation<sup>8</sup>. The second definition, useful only for US operations and based on a previous proposal<sup>8</sup>, relies on the US Drought Monitor (USDM) and can be applied in near real-time for early warning applications:

- **Flash drought definition 1** (applications: international operations, prediction, research): 50% increase in EDDI (toward drying) over two weeks, sustained for at least another two weeks
- **Flash drought definition 2** (application: US operations): two-category change in the USDM in 2 weeks, sustained for at least another 2 weeks

Regarding the second definition above, the USDM is a weekly operational product based on multiple inputs from observations (for example weather, climate, hydrology) and empirical input from regional observers and expert judgement evaluations from the team of scientists (authors) who curate the USDM<sup>7</sup>. A caveat limiting application of this definition to the United States is that the USDM involves expert judgement, beyond raw input of observational data, and hence it cannot be directly applied outside the US operational setting, although drought monitors in other countries could also be used (for example <https://droughtwatch.eu>). Even with that caveat,

as the USDM is familiar and widely referenced by stakeholders and other users, basing a flash-drought index on the USDM categories would have a readily applicable operational utility not possible from other indices. Following from conditions experienced in recent events (Figs. 1 and 2), the rapidity of onset of flash-drought conditions is reflected by requiring a two-category change in the drought monitor in a 2-week period. Impacts can emerge on the timescale of weeks during a flash drought, so this definition requires the two-category change in the drought monitor index to be sustained for at least another 2 weeks after it is established. This definition includes no prescription beyond this 4-week period; the event could persist beyond that time or it could terminate. For example, during the US Midwest 2012 event (Fig. 1), 45% of the High Plains went from D0 ('Abnormally dry') to D2 ('Severe drought') between 12 June and 26 June, a two-category change in the USDM in 2 weeks.


A more general flash-drought definition is the first one listed above, which can be used for international operations, prediction, analysis of observations and climate model output, research into future projections, and applications to periods prior to the USDM. This general definition is based on EDDI, which is multiscalar, can be calculated at 1-week to 12-month timescales, and can capture drying dynamics that operate at the timescales of flash droughts. EDDI provides information on the emergence and persistence of anomalous evaporative demand in a region. The rapid onset characteristic is reflected in the EDDI-based definition by requiring an increase in EDDI of 50 percentiles (toward drying) over 2 weeks, which must then be sustained for at least the next 2 weeks. Again returning to the guidance for this definition provided by the US 2012 event (Fig. 1, left), there are large areas of the United States that experienced at least a 50-percentile change in the EDDI from 5 June to 3 July.

Related to changes in EDDI are changes in soil moisture. The spatial patterns of the frequency of occurrence of 40-, 50- and 60-percentile decreases in soil moisture during a 20-day period over approximately the past 100 years are shown in Fig. 4. Large variations in soil moisture are common over the wettest areas (east of the Mississippi River). Ideally, EDDI would see a comparable change over these periods. In cases in which the anomalously dry conditions persist beyond the initial week, and result in sufficiently dry conditions, these events would qualify as flash drought, although the threshold for 'sufficiently dry' remains to be assessed.

A phenomenon related to but separate from flash drought is rapid-intensification snow drought, which occurs when snowpack has a sudden and fast decline. These are of particular concern for regions that rely on snowpack for water supply and power generation. A rapid-intensification snow drought can be induced, for example, by dust-on-snow, rain-on-snow, or anomalously warm temperatures<sup>71</sup>. Other processes that drive a rapid decrease in snowpack could also include advection or sublimation events, for example due to high winds. Because of the cross-timescale interactions between snowpack loss and impacts, and the substantial differences in processes from the flash droughts discussed above, we propose that rapid-intensification snow drought should be considered separately from flash drought. Nonetheless, because of its impacts, it does also require attention<sup>72</sup>.

The next steps are to test and apply these definitions retrospectively, to verify that they appropriately encompass events generally described as flash-drought events, that they are sufficiently rare that they describe unusual events, and that they describe events that are impactful in one or more dimensions. Extending the definition to require that impacts occur would require quantifying those impacts; this could be addressed by extending the definitions. Further refinement of flash-drought definitions may also be useful for specific regions, seasons, sectors and drought types, using criteria of sufficient intensification rate, impact and rarity<sup>8</sup>. That said, these definitions are designed as proposals to elicit discussion





**Fig. 4 | Frequency of different drought intensification rates.** Frequency of soil-moisture decreases exceeding 40-, 50- and 60-percentile thresholds over four pentads for a 100-year period (1916–2017). Soil moisture is from the same VIC simulations as Fig. 2.

in the community over their appropriateness and applicability. It is expected that they would be fine-tuned in the future.

### Impacts-based early warning

Impacts particular to flash drought arise from its rapid and intense development. Because drought response plans developed by communities and governments are often designed around slower-onset events that unfold over the course of months, rapid onset and intensification have the potential to inhibit the initial response: there may be less time than is allocated to prepare or implement mitigation measures. In the 2012 US Midwest flash drought, during the May–July growing season, dry weather dominated the agricultural areas in the Central Plains and Midwest. Several states had record dry seasons: Arkansas (April–June and other seasons), Kansas (May–July), Nebraska (June–August and other seasons) and South Dakota (July–September). Impacts included, but were not limited to, the reduction in crop yields and commerce-related activities on major river systems. The Mississippi River had water levels that went below 2 m depth and was closed to navigation three times with less load carried, barges running aground, slower speeds and increased dredging costs. The US summer drought of 2012 also contributed to unusually high acreage burned by wildfires. The 2017 Northern Great Plains flash drought also brought wildfire and affected water resources and agriculture<sup>73</sup>.

Severity of drought impacts is not only aggravated by other climatic factors, such as high temperatures, high winds and low relative humidity, but also by the timing (that is, principal season of occurrence, delays in the start of the rainy season, and occurrence of rains in relation to principal crop-growth stages) and effectiveness of the rains (rainfall intensity and number of rainfall events)<sup>3</sup>. Other impacts may be associated with hazards that compound drought, such as heatwaves, wildfires and soil erosion. These may induce public-health effects of heat stress or air-quality degradation due to forest fires. Water quality may degrade, affecting aquatic habitats. Depletion of water storage, low river flows and associated consequences for water supply systems and hydropower production can occur with flash drought, though perhaps with some delays. The recreation sector could feel impacts from wildfire as well as low river flows. This impact- and sector-specific vulnerability to flash droughts requires more in-depth investigation, especially as buffers against drought impacts (such as water storage, or grain/feed stores for livestock agriculture) are used up more quickly than for slower-onset drought.

Even though much of the focus of flash drought has been on humid regions where agriculture is a primary activity, impacts are

also keenly felt in arid and semi-arid regions. A baseline environment that is already water-stressed renders arid regions more vulnerable to drought with less buffer until impacts are felt. For example, a flash drought could deplete reservoirs, affecting both water availability and hydropower generation capacity in places like the southwest United States, where water is highly managed. Because some physical mechanisms (see above) and affected systems will differ from humid regions, understanding and predicting flash droughts to provide early warning in arid regions presents an additional challenge.

Overall, some types of flash-drought-related impacts will present different challenges from slower-onset drought. An accelerated ‘time to impact’ from the onset of a meteorological drought also means that forecasting gains importance compared with monitoring (which remains important, but not sufficient) in operational early warning and risk management of drought. Furthermore, translating drought development into mitigation action, and predicting the likelihood of termination versus continuation into long-term drought, are also important. A systematic assessment of where and when (in terms of seasonal timing) vulnerability to flash drought is highest is needed to guide efforts on where prediction and early warning would be most useful.

Early warning can enable communities to prepare for impacts. The United Nations office for Disaster Risk Reduction has established four key areas of people-centred early warning: risk knowledge, monitoring and warning, communication, and response capability. Early warning systems in such contexts are needed not only for event onset, at which a threshold above some socially acceptable or safe level is exceeded, but also for intensification and duration<sup>74</sup>. The phrase ‘early warning information system’ can be used to describe an integrated process of risk assessment, communication and decision support, of which an early warning is a central output. An early warning information system involves much more than development and dissemination of a forecast; it is the systematic collection and analysis of relevant information about, and coming from, areas of impending risk that informs the development of strategic responses to anticipate crises and crisis evolution; provides capabilities for generating problem-specific risk assessments and scenarios; and effectively communicates options to critical actors for the purposes of decision-making, preparedness and mitigation<sup>74</sup>.

In summary, with improved monitoring and credible S2S timescale predictions, drought early warning could include flash drought. For risk management before, during and after flash-drought events, improvements in monitoring and also predicting not just onset of flash drought but termination of events would be beneficial.

## Ethics in drought research and applications

The ultimate goal of research on flash drought, as for many impactful environmental phenomena, is to avoid or decrease the negative effects of drought on individuals and communities. Inequalities influence the ability of communities to cope and adapt to disasters<sup>75</sup>. Across the early warning and response continuum lie three cross-cutting elements: capacity-building, governance, and gender and social inclusion. These elements are best served through a focus on procedural justice and the resulting ethics of participation<sup>76,77</sup>. Effective information-based services engage affected people and multiple perspectives in the development of knowledge, in decision-making, and as recipients of policies<sup>78,79</sup>. Identifying and understanding how flash drought and other climate impacts affect communities and individuals requires integrating local knowledge about impacts. This is knowledge that is inclusive of many different types of individuals in each community, including people who can successfully and meaningfully engage with those affected in the research and monitoring process. People from many different identities are underrepresented in the environmental science workforce; a well-documented example is women. Women in many parts of the world are at greater risk of harm due to climate-related disasters<sup>75</sup>, and yet they remain underrepresented among one influential set of climate scientists—IPCC authors<sup>80</sup>. Improving diversity of the scientific workforce and taking an inclusive approach to engaging with stakeholders, while remaining mindful of those that are not included, is essential to ethical research on weather and climate in general and droughts, including flash drought, in particular.

The following objectives are suggested to support the ethical practice of drought research:

- Enhance engagement between users and researchers.
- Develop capacity in the segment of the work dedicated to being an interface with stakeholders and users.
- Support individual actions to improve scientific culture.
- Make institutional efforts to change the culture of science and its reward system.
- Collaborate on interdisciplinary work.
- Share research outcomes with society, users and stakeholders.

## Future directions in flash drought

Key areas where progress could be made on flash droughts include improved understanding of events in the recent and more distant past and their impacts; establishing predictability and improving prediction of flash-drought events; applying these predictions to improve early warning systems for impending events as well as responding to events as they unfold; and understanding how flash drought will respond to climate variability and change. We identify some key challenges and directions for achieving this below.

To identify developing flash-drought events, monitoring systems must attend to shorter timescales and more frequent updates than are needed to capture slower, longer-term drought events. Products that are only updated monthly (such as the North American Multi-Model Ensemble, NMME<sup>81</sup>) are not very useful for flash-drought monitoring and prediction. Some countries have drought monitoring and drought early warning systems. In countries with less monitoring and prediction infrastructure, there is also potential to leverage systems that provide global hydrological information, such as the Global Flood Awareness System (GloFAS)<sup>82</sup>, World-Wide Hydrological Predictions for the Environment (HYPE)<sup>83</sup>, experimental Global Drought Information Systems (GDIS), Global Drought Observatory (GDO) and Integrated Drought Management Programme (IDMP).

There remain open questions about how to define flash drought. One challenge for identifying flash-drought events is their wide variation in spatial scale. What areal extent is sufficient to assert that a flash drought is occurring? Assessment of regions and times of year with

high sensitivity to or preponderance for flash drought should also be factored into its identification; model representation of land use and its change can play a role as well. A better understanding of flash droughts requires more in-depth research on relevant compound and cascading physical processes that can trigger or increase the likelihood of a flash drought. These include relationships among soil moisture, land–atmosphere interactions, their connections to large-scale meteorological conditions (and precursor conditions), and how these are forced by remote SST patterns and influenced by internal atmospheric variability. Furthermore, research is needed into how these conditions will change as the climate base state changes<sup>52,84</sup>, and to incorporate the changing climate into the definition of flash drought so that flash-drought definitions remain meaningful in the future.

Prediction systems focus mostly on physical quantities such as precipitation, but the motivation and ultimate goal of flash-drought monitoring and prediction is to provide as much anticipatory information as possible of impending impacts of flash-drought events, and aid response during and after those events. This requires engagement with relevant stakeholders, building capacity, establishing ethical practices of research to document potential impacts of flash drought and when and where these are a concern, and identifying relationships between flash-drought indicators and impacts. Such efforts should cross disciplines and engage researchers and decision makers at all stages that bridge the weather–climate continuum.

## Data availability

EDDI is available for the CONUS at [http://ftp.cdc.noaa.gov/Projects/EDDI/CONUS\\_archive](http://ftp.cdc.noaa.gov/Projects/EDDI/CONUS_archive) and for the globe at [http://ftp.cdc.noaa.gov/Projects/EDDI/global\\_archive](http://ftp.cdc.noaa.gov/Projects/EDDI/global_archive). Figure 1i is generated from the USDM (droughtmonitor.unl.edu). The data analysed in Figs. 2 and 4 are available from <http://ftp.cdc.noaa.gov/pub/Public/jeischeid/andy/>. The data to generate Fig. 4 are available at [github.com/apendergrass/flashdroughtperspectivefigure](https://github.com/apendergrass/flashdroughtperspectivefigure).

## Code availability

Figure 1i is generated from the USDM (droughtmonitor.unl.edu). Figures 2 and 4 were generated following the protocol [http://192.12.137.7/pub/dcp/archive/OBS/livneh2014.1\\_16deg/](http://192.12.137.7/pub/dcp/archive/OBS/livneh2014.1_16deg/). The code to generate Fig. 4 is available at [github.com/apendergrass/flashdroughtperspectivefigure](https://github.com/apendergrass/flashdroughtperspectivefigure).

## Online content

Any methods, additional references, Nature Research reporting summaries, source data, extended data, supplementary information, acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41558-020-0709-0>.

Received: 10 June 2019; Accepted: 21 January 2020;

Published online: 2 March 2020

## References

1. Pulwarty, R. S. & Sivakumar, M. V. K. Information systems in a changing climate: early warnings and drought risk management. *Weather Clim. Extrem.* **3**, 14–21 (2014).
2. *Global Assessment Report on Disaster Risk Reduction* (UNDRR, 2019).
3. Wilhite, D. A. & Pulwarty, R. S. in *Drought and Water Crises: Integrating Science, Management, and Policy* (eds Wilhite, D. & Pulwarty, R. S.) Ch. 25 (CRC, 2017).
4. Christensen, J. et al. in *Climate Change 2007: The Physical Science Basis* (eds Solomon, S. et al.) Ch. 11 (IPCC, Cambridge Univ. Press, 2007).
5. Seneviratne, S. I. et al. in *Special Report on Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation* (eds Field, C. B. et al.) 109–230 (IPCC, Cambridge Univ. Press, 2012).
6. Wilhite, D. A., Sivakumar, M. V. K. & Pulwarty, R. Managing drought risk in a changing climate: the role of national drought policy. *Weather Clim. Extrem.* **3**, 4–13 (2014).

7. Svoboda, M. et al. The Drought Monitor. *Bull. Am. Meteorol. Soc.* **83**, 1181–1190 (2002).
8. Otkin, J. A. et al. Flash droughts: a review and assessment of the challenges imposed by rapid-onset droughts in the United States. *Bull. Am. Meteorol. Soc.* **99**, 911–919 (2018).
9. Robertson, A. W. et al. Improving and promoting subseasonal to seasonal prediction. *Bull. Am. Meteorol. Soc.* **96**, ES49–ES53 (2015).
10. Hoerling, M. P. et al. Is a transition to semipermanent drought conditions imminent in the U.S. Great Plains? *J. Clim.* **25**, 8380–8386 (2012).
11. Namias, J. Anatomy of Great Plains protracted heat waves (especially the 1980 U.S. summer drought). *Mon. Weather Rev.* **110**, 824–838 (1982).
12. Yuan, X., Wang, L. & Wood, E. F. Anthropogenic intensification of southern African flash droughts as exemplified by the 2015/16 season. *Bull. Am. Meteorol. Soc.* **99**, S86–S90 (2018).
13. Yuan, X., Ma, Z., Pan, M. & Shi, C. Microwave remote sensing of short-term droughts during crop growing seasons. *Geophys. Res. Lett.* **42**, 4394–4401 (2015).
14. Li, Y. et al. Mechanisms and early warning of drought disasters: experimental drought meteorology research over China. *Bull. Am. Meteorol. Soc.* **100**, 673–687 (2019).
15. Nguyen, H. et al. Using evaporative stress index to monitor flash drought in Australia. *Environ. Res. Lett.* <https://doi.org/10.1088/1748-9326/ab2103> (2019).
16. Ford, T. W. & Labosier, C. F. Meteorological conditions associated with the onset of flash drought in the eastern United States. *Agric. Meteorol.* **247**, 414–423 (2017).
17. Hobbins, M. T., Ramirez, J. A. & Brown, T. C. Trends in pan evaporation and actual evapotranspiration across the conterminous U.S.: paradoxical or complementary? *Geophys. Res. Lett.* **31**, <https://doi.org/10.1029/2004GL019846> (2004).
18. Ramirez, J. A., Hobbins, M. T. & Brown, T. C. Observational evidence of the complementary relationship in regional evaporation lends strong support for Bouchet's hypothesis. *Geophys. Res. Lett.* **32**, L15401 (2005).
19. Koster, R. D. et al. Flash drought as captured by reanalysis data: disentangling the contributions of precipitation deficit and excess evapotranspiration. *J. Hydrometeorol.* <https://doi.org/10.1175/JHM-D-18-0242.1> (2019).
20. Seneviratne, S. I., Lüthi, D., Lietschi, M. & Schär, C. Land–atmosphere coupling and climate change in Europe. *Nature* **443**, 205–209 (2006).
21. Fischer, E. M., Seneviratne, S. I., Lüthi, D. & Schär, C. Contribution of land–atmosphere coupling to recent European summer heat waves. *Geophys. Res. Lett.* **34**, L06707 (2007).
22. Su, H., Yang, Z.-L., Dickinson, R. E. & Wei, J. Spring soil moisture–precipitation feedback in the Southern Great Plains: how is it related to large-scale atmospheric conditions? *Geophys. Res. Lett.* **41**, L283–L289 (2014).
23. Hoerling, M. et al. Causes and predictability of the 2012 Great Plains drought. *Bull. Am. Meteorol. Soc.* **95**, 269–282 (2014).
24. Mo, K. C. & Lettenmaier, D. P. Precipitation deficit flash droughts over the United States. *J. Hydrometeorol.* **17**, 1169–1184 (2016).
25. Chiang, F., Mazdiyasi, O. & AghaKouchak, A. Amplified warming of droughts in southern United States in observations and model simulations. *Sci. Adv.* **4**, eaat2380 (2018).
26. Pegion, K. et al. The Subseasonal Experiment (SubX): a multi-model subseasonal prediction experiment. *Bull. Am. Meteorol. Soc.* <https://doi.org/10.1175/BAMS-D-18-0270.1> (2019).
27. Chen, L. G. et al. Flash drought characteristics based on U.S. Drought Monitor. *Atmosphere (Basel)* **10**, 498 (2019).
28. Dirmeyer, P. A., Gentile, P., Ek, M. B. & Balsamo, G. *Sub-Seasonal to Seasonal Prediction: The Gap Between Weather and Climate Forecasting* (Robertson, A. W. & Vitart, F.) 165–181 (Elsevier, 2019).
29. Waliser, D. E. et al. Potential predictability of the Madden–Julian oscillation. *Bull. Am. Meteorol. Soc.* **84**, 33–50 (2003).
30. Hendon, H. H. et al. Australian rainfall and surface temperature variations associated with the Southern Hemisphere annular mode. *J. Clim.* **20**, 2452–2467 (2007).
31. Zhao, M. & Hendon, H. H. Representation and prediction of the Indian Ocean dipole in the POAMA seasonal forecast model. *Q. J. R. Meteorol. Soc.* **135**, 337–352 (2009).
32. *Next Generation Earth System Prediction: Strategies for Subseasonal to Seasonal Forecasts* (National Academies Press, 2016).
33. Zhu, H. et al. Seamless precipitation prediction skill in the tropics and extratropics from a global model. *Mon. Weather Rev.* **142**, 1556–1569 (2014).
34. Wheeler, M. C., Zhu, H., Sobel, A. H., Hudson, D. & Vitart, F. Seamless precipitation prediction skill comparison between two global models. *Q. J. R. Meteorol. Soc.* **143**, 374–383 (2017).
35. Wang, L. & Robertson, A. W. Week 3–4 predictability over the United States assessed from two operational ensemble prediction systems. *Clim. Dyn.* **52**, 5861–5875 (2019).
36. Hudson, D. et al. Forewarned is forearmed: extended-range forecast guidance of recent extreme heat events in Australia. *Weather Forecast.* **31**, 697–711 (2016).
37. Vitart, F. & Robertson, A. W. The sub-seasonal to seasonal prediction project (S2S) and the prediction of extreme events. *npj Clim. Atmos. Sci.* **1**, 3 (2018).
38. Lehner, F. et al. Mitigating the impacts of climate nonstationarity on seasonal streamflow predictability in the U.S. Southwest. *Geophys. Res. Lett.* **44**, 12208–12217 (2017).
39. McEvoy, D. J. et al. The Evaporative Demand Drought Index. Part II: CONUS-wide assessment against common drought indicators. *J. Hydrometeorol.* **17**, 1763–1779 (2016).
40. Shukla, S. et al. Examining the value of global seasonal reference evapotranspiration forecasts to support FEWS NET's food insecurity outlooks. *J. Appl. Meteorol. Climatol.* **56**, 2941–2949 (2017).
41. Zhang, C. et al. CAUSES: diagnosis of the summertime warm bias in CMIP5 climate models at the ARM southern Great Plains site. *J. Geophys. Res. Atmos.* **123**, 2968–2992 (2018).
42. Vitart, F. Madden–Julian Oscillation prediction and teleconnections in the S2S database. *Q. J. R. Meteorol. Soc.* **143**, 2210–2220 (2017).
43. Ukkola, A. M. et al. Land surface models systematically overestimate the intensity, duration and magnitude of seasonal-scale evaporative droughts. *Environ. Res. Lett.* **11**, 104012 (2016).
44. Vitart, F. et al. The Subseasonal to Seasonal (S2S) Prediction Project Database. *Bull. Am. Meteorol. Soc.* **98**, 163–173 (2017).
45. Koster, R. D. et al. Contribution of land surface initialization to subseasonal forecast skill: first results from a multi-model experiment. *Geophys. Res. Lett.* **37**, <https://doi.org/10.1029/2009GL041677> (2010).
46. Fisher, R. A. et al. Vegetation demographics in Earth System Models: a review of progress and priorities. *Glob. Change Biol.* **24**, 35–54 (2018).
47. Mo, K. C. & Lettenmaier, D. P. Heat wave flash droughts in decline. *Geophys. Res. Lett.* **42**, 2823–2829 (2015).
48. Wang, L., Yuan, X., Xie, Z., Wu, P. & Li, Y. Increasing flash droughts over China during the recent global warming hiatus. *Sci. Rep.* **6**, 30571 (2016).
49. Zhang, Y., You, Q., Chen, C. & Li, X. Flash droughts in a typical humid and subtropical basin: a case study in the Gan River Basin, China. *J. Hydrol.* **551**, 162–176 (2017).
50. Barnett, T. P. et al. Human-induced changes in the hydrology of the Western United States. *Science* **319**, 1080–1083 (2008).
51. Marvel, K. et al. Twentieth-century hydroclimate changes consistent with human influence. *Nature* **569**, 59–65 (2019).
52. Cook, B. I., Mankin, J. S. & Anchukaitis, K. J. Climate change and drought: from past to future. *Curr. Clim. Change Rep.* **4**, 164–179 (2018).
53. Seager, R. et al. Model projections of an imminent transition to a more arid climate in southwestern North America. *Science* **316**, 1181–1184 (2007).
54. Zhou, S. et al. Land–atmosphere feedbacks exacerbate concurrent soil drought and atmospheric aridity. *Proc. Natl Acad. Sci. USA* **116**, 18848–18853 (2019).
55. Roderick, M. L., Greve, P. & Farquhar, G. D. On the assessment of aridity with changes in atmospheric CO<sub>2</sub>. *Water Resour. Res.* **51**, 5450–5463 (2015).
56. Milly, P. C. D. & Dunne, K. A. Potential evapotranspiration and continental drying. *Nat. Clim. Change* **6**, 946–949 (2016).
57. Feng, S. et al. Why do different drought indices show distinct future drought risk outcomes in the U.S. Great Plains? *J. Clim.* **30**, 265–278 (2017).
58. Lehner, F. et al. Projected drought risk in 1.5 °C and 2 °C warmer climates. *Geophys. Res. Lett.* **44**, 7419–7428 (2017).
59. Swann, A. L. S., Hoffman, F. M., Koven, C. D. & Randerson, J. T. Plant responses to increasing CO<sub>2</sub> reduce estimates of climate impacts on drought severity. *Proc. Natl Acad. Sci. USA* **113**, 10019–10024 (2016).
60. Bonfils, C. et al. Competing influences of anthropogenic warming, ENSO, and plant physiology on future terrestrial aridity. *J. Clim.* **30**, 6883–6904 (2017).
61. Yang, Y., Roderick, M. L., Zhang, S., McVicar, T. R. & Donohue, R. J. Hydrologic implications of vegetation response to elevated CO<sub>2</sub> in climate projections. *Nat. Clim. Change* **9**, 44–48 (2019).
62. Mankin, J. S., Seager, R., Smerdon, J. E., Cook, B. I. & Williams, A. P. Mid-latitude freshwater availability reduced by projected vegetation responses to climate change. *Nat. Geosci.* <https://doi.org/10.1038/s41561-019-0480-x> (2019).
63. Dirmeyer, P. A. et al. Evidence for enhanced land–atmosphere feedback in a warming climate. *J. Hydrometeorol.* **13**, 981–995 (2012).
64. Otkin, J. A. et al. Assessing the evolution of soil moisture and vegetation conditions during the 2012 United States flash drought. *Agric. Meteorol.* **218–219**, 230–242 (2016).
65. Meko, D. M. et al. Medieval drought in the upper Colorado River Basin. *Geophys. Res. Lett.* **34**, L17075 (2007).
66. Woodhouse, C. A., Meko, D. M., MacDonald, G. M., Stahle, D. W. & Cook, E. R. A 1,200-year perspective of 21st century drought in southwestern North America. *Proc. Natl Acad. Sci. USA* **107**, 21283–21288 (2010).
67. Woodhouse, C., Stahle, D. & Villanueva Diaz, J. Rio Grande and Rio Conchos water supply variability over the past 500 years. *Clim. Res.* **51**, 147–158 (2012).



68. Woodhouse, C. A. & Pederson, G. T. Investigating runoff efficiency in Upper Colorado River streamflow over past centuries. *Water Resour. Res.* **54**, 286–300 (2018).
69. Lehner, F., Wahl, E. R., Wood, A. W., Blatchford, D. B. & Llewellyn, D. Assessing recent declines in Upper Rio Grande runoff efficiency from a paleoclimate perspective. *Geophys. Res. Lett.* **44**, 4124–4133 (2017).
70. Zhao, M. et al. Weakened eastern Pacific El Niño predictability in the early twenty-first century. *J. Clim.* **29**, 6805–6822 (2016).
71. Huning, L. S. & AghaKouchak, A. Mountain snowpack response to different levels of warming. *Proc. Natl Acad. Sci. USA* **115**, 10932–10937 (2018).
72. Harpold, A., Dettinger, M. & Rajagopal, S. Defining snow drought and why it matters. *Eos* <https://doi.org/10.1029/2017EO068775> (2017).
73. Hoell, A., Perlwitz, J. & Eischeid, J. *Drought Assessment Report: The Causes, Predictability, and Historical Context of the 2017 US Northern Great Plains Drought* (NOAA/NIDIS/CIRES, 2019).
74. Pulwarty, R. S. & Verdin, J. P. in *Measuring Vulnerability to Natural Hazards: Towards Disaster Resilient Societies* 2nd edn (ed. Birkmann, J.) 124–147 (United Nations Univ. Press, 2013).
75. Cutter, S. et al. in *Special Report on Managing the Risks of Extremes and Disaster to Advance Climate Change Adaptation* (eds Field, C. B. et al.) 291–338 (IPCC, Cambridge Univ. Press, 2012).
76. Shrader-Frechette, K. S. *Environmental Justice: Creating Equality, Reclaiming Democracy. Environmental Ethics and Science Policy* (Oxford Univ. Press, 2002).
77. Jamieson, D. *Ethics and the Environment: An Introduction* (Cambridge Univ. Press, 2008).
78. Pulwarty, R. S. et al. in *Mapping Vulnerability: Disasters, Development and People* (eds Bankoff, G. & Frerks, G.) Ch. 6 (Routledge, 2004).
79. Allis, E. et al. The future of climate services. *World Meteorol. Organ. Bull.* **68**, <https://public.wmo.int/en/resources/bulletin/future-of-climate-services> (2019).
80. Gay-Antaki, M. & Liverman, D. Climate for women in climate science: women scientists and the Intergovernmental Panel on Climate Change. *Proc. Natl Acad. Sci. USA* <https://doi.org/10.1073/pnas.1710271115> (2018).
81. Kirtman, B. P. et al. The North American Multimodel Ensemble: Phase-1 seasonal-to-interannual prediction; Phase-2 toward developing intraseasonal prediction. *Bull. Am. Meteorol. Soc.* **95**, 585–601 (2014).
82. Alfieri, L. et al. GloFAS—global ensemble streamflow forecasting and flood early warning. *Hydrol. Earth Syst. Sci.* **17**, 1161–1175 (2013).
83. Arheimer, B. et al. Global catchment modelling using World-Wide HYPE (WWH), open data and stepwise parameter estimation. *Hydrol. Earth Syst. Sci. Discuss.* <https://doi.org/10.5194/hess-2019-111> (2019).
84. Yuan, X. et al. Anthropogenic shift towards higher risk of flash drought over China. *Nat. Commun.* **10**, 4661 (2019).
85. Hobbins, M. T., McEvoy, D. J. & Hain, C. R. in *Drought and Water Crises: Integrating Science, Management, and Policy* (eds Wilhite, D. A. & Pulwarty, R. S.) Ch. 11 (CRC, 2017).
86. Liang, X., Lettenmaier, D. P., Wood, E. F. & Burges, S. J. A simple hydrologically based model of land surface water and energy fluxes for general circulation models. *J. Geophys. Res.* **99**, 14415 (1994).
87. Livneh, B. et al. A long-term hydrologically based dataset of land surface fluxes and states for the conterminous United States: update and extensions. *J. Clim.* **26**, 9384–9392 (2013).
88. Livneh, B. et al. A spatially comprehensive, hydrometeorological data set for Mexico, the U.S., and Southern Canada 1950–2013. *Sci. Data* **2**, 150042 (2015).
89. Lukas, J., Hobbins, M. T., Rangwala, I. & EDDI Team. *The EDDI User Guide* (NOAA, 2017); [https://www.esrl.noaa.gov/psd/eddi/pdf/EDDI\\_UserGuide\\_v1.0.pdf](https://www.esrl.noaa.gov/psd/eddi/pdf/EDDI_UserGuide_v1.0.pdf)

## Acknowledgements

The perspectives in this manuscript emerged from an Aspen Global Change Institute (AGCI) workshop in September 2018; we thank all participants (<https://www.agci.org/event/18s4>). This material is based on work supported by the National Center for Atmospheric Research, which is a major facility sponsored by the National Science Foundation (NSF) under Cooperative Agreement no. 1947282. Portions of this study were supported by the Regional and Global Model Analysis (RGMA) component of the Earth and Environmental System Modeling Program of the US Department of Energy's Office of Biological & Environmental Research (BER) via NSF IA 1844590. M.H. was supported by a National Oceanic and Atmospheric Administration (NOAA) Joint Technology Transfer Initiative (JTII) award and a US Agency for International Development (USAID)—Famine Early Warning Systems Network (FEWS NET) award (NA17OAR4320101). C.J.W.B. was supported by an Early-Mid Career LLNL Laboratory Directed Research and Development award (tracking code 17-ERD-115) under the auspices of the US Department of Energy by Lawrence Livermore National Laboratory under contract DE-AC52-07NA27344. A.J.E.G. is supported by Australian Research Council Discovery Early Career Researcher Award DE150101297. M.C.W. was partially supported by the Northern Australia Climate Program (NACP), funded by Meat and Livestock Australia, the Queensland Government and the University of Southern Queensland. F.L. is also supported by NSF AGS-0856145 Amendment 87 and by the Bureau of Reclamation under Cooperative Agreement R16AC00039. The US Drought Monitor is jointly produced by the National Drought Mitigation Center at the University of Nebraska-Lincoln, the United States Department of Agriculture, and NOAA. Map (Fig. 1i) courtesy of NDMC. Opinions expressed by D.L. represent professional opinions of the co-author, not official positions of the government.

## Author contributions

A.G.P., G.A.M. and R.P. coordinated the text and organized the workshop. M.H. and A.H. provided figures. All authors except A.A. participated in discussions at the workshop, and all authors contributed to the writing.

## Competing interests

The authors declare no competing interests.

## Additional information

**Correspondence** should be addressed to A.G.P.

**Peer review information** *Nature Climate Change* thanks Laura Ferranti and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This is a U.S. government work and not under copyright protection in the U.S.; foreign copyright protection may apply 2020