

FLAVONE C-GLYCOSIDES AND CUCURBITACIN GLYCOSIDES FROM *CITRULLUS COLOCYNTHIS*

¹ABBAS DELAZAR, ²SIMON GIBBONS, ¹ALI REZA KOSARI, ¹HOSSEIN NAZEMIYEH, ¹MASOUD MODARRESI, ³LUTFUN NAHAR, ⁴SATYAJIT D. SARKER

¹Faculty of Pharmacy and Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran., ²Centre for Pharmacognosy and Phytotherapy, The School of Pharmacy, University of London, 29-39 Brunswick Square, London WC1N 1AX, UK., ³Department of Chemistry, University of Aberdeen, Aberdeen, Scotland, UK., ⁴School of Biomedical Sciences, University of Ulster at Coleraine, Cromore Road, Coleraine BT52 1SA, Co. Londonderry, Northern Ireland, UK

ABSTRACT

Citrullus colocynthis (L.) Schrad. (Cucurbitaceae) is an Iranian medicinal plant that has traditionally been used as an abortifacient and to treat constipation, oedema, bacterial infections, cancer and diabetes. As part of our on-going studies on Iranian medicinal plants, thorough phytochemical investigation was carried out on this plant. The reversed-phase preparative HPLC was employed to isolate compounds from the butanol fraction of the hydro-methanolic (70%) extract of the fruits of the locally grown *C. colocynthis*. Structures of the isolated compounds [1-5] were elucidated by spectroscopic means. The antioxidant property of the flavonoids 1-3 was determined by the 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay. Three flavone glucosides, isosaponarin [1], isovitexin [2] and isoorientin 3'-O-methyl ether [3] and two cucurbitacin glucosides, 2-O-β-D-glucopyranosylcucurbitacin I [4] and 2-O-β-D-glucopyranosylcucurbitacin L [5] were isolated and identified. Flavonoids 1-3 showed significant antioxidant properties. Since reactive oxygen species are important contributors to tissue injury, inflammation, cancer and many other ailments, the antioxidant properties of 1-3 probably contribute, at least to some extent, to the pharmacological and traditional medicinal uses of the *C. colocynthis*.

Keywords: *Citrullus colocynthis*, Cucurbitaceae, Isosaponarin, Isovitexin, Isoorientin 3'-O-methyl ether, Cucurbitacin I, Cucurbitacin L, DPPH.

INTRODUCTION

Citrullus colocynthis (L.) Schrad. (Cucurbitaceae), commonly known as 'bitter apple', 'colosynth', 'vine-of-Sodom' 'tumba' or 'wild gourd', is a tropical plant that grows abundantly in the south of Iran, and widely in other parts of the world (1). In the traditional medicine of Iran, this plant has been used to treat constipation, oedema, bacterial infections, cancer and diabetes, and as an abortifacient (2). The ethnobotanical uses of this plant include its use as an abortifacient, cathartic, purgative and vermifuse, and for the treatment of fever, cancer, amenorrhoea, jaundice, leukemia, rheumatism and tumour (3). In Syria, it has also been used as an insect repellent (3). A number of plant secondary metabolites including cucurbitacins, flavonoids, caffeic acid derivatives and terpenoids have previously been reported from this plant (4-10). As part of our on-going studies on Iranian medicinal plants (11-15), the isolation and identification of the chemical constituents,

namely, isosaponarin [1], isovitexin [2], isoorientin 3'-O-methyl ether [3], 2-O-β-D-glucopyranosylcucurbitacin I [4], 2-O-β-D-glucopyranosylcucurbitacin L [5] of the endemic Iranian species *C. colocynthis*, and the antioxidant properties of the flavonoids [1-3] are described.

MATERIALS AND METHODS

General procedures

NMR spectra were recorded on a Bruker NMR Spectrometer (200 MHz for ¹H and 50 MHz for ¹³C NMR) using DMSO-d₆ for the compounds 1 – 3, and those of 4 and 5 were obtained (500 MHz for ¹H and 125 MHz for ¹³C NMR) on Bruker AVANCE 500 spectrometer using CD₃OD as solvent. The residual solvent peaks were used as internal standards. ESIMS analysis was performed on Finnigan MAT95 spectrometer.

Plant material

Fruits of *Citrullus colocynthis* were collected




Figure 1. Structure of compounds [1-5] isolated from *C. colocynthis*

during August and September, 2003, from Ahvaz in Khozestan province (Iran) and the identity was confirmed by anatomical examination in comparison with the herbarium specimen retained in the School of Pharmacy, Tabriz University of Medical Sciences, Iran. A voucher specimen (TUM-ADA 87) for this collection has been deposited in the herbarium of the Faculty of Pharmacy, Tabriz University of Medical Science, and also in the herbarium of the Plant and Soil Science Department, University of Aberdeen, Scotland (ABD).

Extraction, isolation and structure elucidation

The pulp was separated from the seeds. The ground pulp (500 g) was macerated in water/methanol (30/70) for 24 hours (three times), and the hydro-methanolic solution was subsequently concentrated by evaporation up to 1/3 of the initial volume. The extract was fractionated by the following solvents with increasing polarity: petroleum ether, chloroform, ethyl acetate, butanol and water fractions. The fractions were concentrated to dryness by rotary evaporator. The butanol fraction (2 g) was subjected to Sep-Pack (ODS) fractionation using a step gradient of MeOH-water mixture (10:90, 20:80, 40:60, 60:40, 80:20 and 100:0). The preparative reversed-phase HPLC analysis (Shim-Pak ODS column 20 μ m, 250 mm \times 20 mm; mobile phase: 0 to 50 min gradient 25 to 45% MeOH in water; flow-rate: 20 mL/min, detection at 267 nm) of the 40% methanolic Sep-Pack fraction (2.5 g) resulted in the isolation of compounds **1** (6.4 mg, t_R = 11.3 min), **2** (4.7 mg, t_R = 27.3 min), **3** (7.2 mg, t_R = 29.2 min), **4** (49.8 mg, t_R = 45.8 min) and **5** (35.3 mg, t_R = 51.3 min). The structures of the isolated compounds [1-5] were determined by spectroscopic means.

Isosaponarin [**1**]. Gum, UV λ_{max} (MeOH): Table 1; ESIMS m/z 595 [M+H]⁺; ¹H NMR (200 MHz, DMSO-d₆): Table 2; ¹³C NMR (50.0 MHz, DMSO-d₆): Table 2.

Isovitexin [**2**]. Gum, UV λ_{max} (MeOH): Table 1; ESIMS m/z 433 [M+H]⁺; ¹H NMR (200 MHz, DMSO-d₆): Table 2; ¹³C NMR (50.0 MHz, DMSO-d₆): Table 2.

Isoorientin 3'-O-methyl ether [**3**]. Gum, UV λ_{max} (MeOH): Table 1; ESIMS m/z 463 [M+H]⁺; ¹H NMR (200 MHz, DMSO-d₆): Table 2; ¹³C NMR (50.0 MHz, DMSO-d₆): Table 2.

2-O- β -D-Glucopyranosylcurbitacin I [**4**]. Gum, ESIMS m/z 677 [M+H]⁺; ¹H NMR (500 MHz, CD₃OD): Table 3; ¹³C NMR (125 MHz, CD₃OD): Table 3.

2-O- β -D-Glucopyranosylcurbitacin L (bryoamaride) [**5**]. Gum, ESIMS m/z 679 [M+H]⁺; ¹H NMR (500 MHz, CD₃OD): Table 3; ¹³C NMR (125 MHz, CD₃OD): Table 3.

Antioxidant assay

2,2-Diphenyl-1-picrylhydrazyl radical (DPPH), molecular formula C₁₈H₁₂N₅O₆, was obtained from Fluka Chemie AG, Bucks. The method used by Takao *et al.* (16) was adopted with suitable modifications (17). A solution of DPPH (0.08 mg/mL) in MeOH was used. The compounds **1-3** were dissolved in MeOH to obtain a concentration of 1 \times 10⁻¹ mg/mL. Dilutions were made to obtain concentrations of 5.00 \times 10⁻², 2.5 \times 10⁻², 1.25 \times 10⁻², 6.25 \times 10⁻³, 3.13 \times 10⁻³, 1.56 \times 10⁻³ mg/mL. Diluted solutions (5 mL each) were mixed with DPPH (5 mL) and allowed half hour for any reaction to occur. The UV absorbance was recorded at 517 nm. The experiment was performed in duplicate and average absorption was noted for each concentration. Data were processed using EXCEL and the concentration

that caused a 50% reduction in absorbance (RC_{50}) was calculated. The same procedure was followed for the standard (quercetin).

RESULTS AND DISCUSSION

The reversed-phase preparative HPLC analysis of the butanol fraction of the methanol extract of *C. colocynthis* fruits afforded three flavonoid glycosides, isosaponarin [1], isovitexin [2] and isoorientin 3'-*O*-methyl ether [3] and two cucurbitacin glucosides, 2-*O*- β -D-glucopyranosylcucurbitacin I [4] and 2-*O*- β -D-glucopyranosylcucurbitacin L [5]. While the chemical structures of 1-3 were elucidated by extensive UV analyses (Table 1), ESIMS, and NMR spectroscopic analyses (Table 2), and also by comparing experimental data with respective literature data (4, 18-25), the identity of cucurbitacins 4 and 5 was confirmed unambiguously by a series of 1D and 2D NMR analyses, including ^{13}C DEPT 135, 1H - 1H COSY, 1H - 1H NOESY, 1H - ^{13}C HMBC and 1H - ^{13}C HMQC. The 1H and ^{13}C NMR data of 5 and 6 were in good agreement with the published data (8).

Compounds 1-3 displayed characteristic UV absorption maxima for a flavone skeleton (Table 1) (26). The 1H and ^{13}C NMR data (Table 2) for these compounds also confirmed the presence of a flavone nucleus in these molecules (26). The results of the UV analyses (Table 1) on these compounds using various shift reagents were in good agreement with the substitution patterns depicted in the structures of 1-3 (26). An ESIMS mass spectrum of 1 revealed $[M+H]^+$ (positive ion mode) ion peak at m/z 595, suggesting $M_r = 594$ and solving for $C_{27}H_{30}O_{15}$. In the 1H and ^{13}C NMR spectra (Table 2), there were signals for a 6-substituted 5,7, 4'-trihydroxy flavone nucleus and also for two glucose moieties. The 1H and ^{13}C chemical shift values were comparable to the published data for the C-glucoside, isosaponarin or isovitexin 4'-*O*- β -D-glucopyranoside [1] (18-22). While isovitexin derivatives have previously been isolated from *C. colocynthis*, to our knowledge, this is the first report on the occurrence of isosaponarin [1] in this plant. However, compound 1 has previously been reported from a number of other genera including *Gentiana*, *Gypsophila*, *Psoralea* and *Vaccaria* (10).

An ESIMS mass spectrum of 2 displayed the $[M+H]^+$ (positive ion mode) ion peak at m/z 433, suggesting $M_r = 432$ and solving for $C_{21}H_{20}O_{10}$. The 1H and ^{13}C NMR spectra (Table 2) displayed the signals similar to those of 1, excepting that there were signals for only one glucose unit in 2 (instead of two in 1). Thus compound 2 was

identified as isovitexin, and the 1H and ^{13}C chemical shift values were comparable to the published data for isovitexin [2] (23, 24). Isovitexin [2] is widely distributed in a number of genera including *Citrullus*, *Geranium*, *Fagopyrum*, *Vitex*, *Polygonum*, *Swertia* and *Passiflora* (10).

An ESIMS mass spectrum of 3 exhibited the $[M+H]^+$ (positive ion mode) ion peak at m/z 463, suggesting $M_r = 462$ and solving for $C_{22}H_{22}O_{11}$. The 1H and ^{13}C NMR spectra (Table 2) displayed, in addition to the signals corresponding to isoorientin (25), a singlet at δ 3.90, integrating for 3H, which could be assigned to a methoxyl group. Comprehensive UV analyses using shift reagents (Table 1) confirmed the attachment of this methoxyl group at C-3'. The spectroscopic data of 3 were in good agreement with those published for isoorientin 3'-*O*-methyl ether (4). While isoorientin is well distributed in the plant kingdom, compound 3 has only been reported from *C. colocynthis* (10).

An ESIMS mass spectrum of 4 exhibited the $[M+H]^+$ (positive ion mode) ion peak at m/z 677, suggesting $M_r = 676$ and solving for $C_{36}H_{52}O_{12}$. The 1H and ^{13}C NMR spectra (Table 3) displayed signals comparable to the published data for 2-*O*- β -D-glucopyranosylcucurbitacin I (8). The extensive 2D NMR analyses including COSY, NOESY, HMQC and HMBC confirmed the identity of this compound. Similarly, an ESIMS mass spectrum of 5 exhibited the $[M+H]^+$ (positive ion mode) ion peak at m/z 679, suggesting $M_r = 678$ and solving for $C_{36}H_{54}O_{12}$. In the 1H NMR spectrum of 5 (Table 2), the two distinct signals for *trans* olefinic protons (as in 4) were absent. The ^{13}C NMR spectrum (Table 3) also revealed that instead of two olefinic methine signals (as in 4), signals for two methylene carbons were present. Thus compound 5 was identified as 2-*O*- β -D-glucopyranosylcucurbitacin L, and all data were in good agreement with the published data for this compound (8). Both 4 and 5, and their aglycones as well as various other cucurbitacins are well distributed in the genus *Citrullus* and the family Cucurbitaceae (10).

The DPPH assay measures the free radical scavenging property of a compound (16). DPPH is a molecule containing a stable free radical. In the presence of an antioxidant which can donate an electron to DPPH, the purple color, which is typical to free DPPH radical, decays, and the change in absorbance at 517 nm is followed spectrophotometrically. In the cases where the structure of the electron donor is not known (e.g., a plant extract), this method can afford data on the reduction potential of the sample, and hence can be helpful in comparing the reduction potential of

Table 1. UV-visible absorption peaks (in nm) of **1-3** in MeOH and their shifts with different reagents

Compounds	MeOH	Shift reagents				
		NaOMe	AlCl ₃	AlCl ₃ /HCl	NaOAc	NaOAc/ H ₃ BO ₃
1	326.7 sh	364.5	375.5(sh)	375.5(sh)	366.8(sh)	329.0(sh)
	274	279.5	334.5	334.5	279.5	276.4
			302.6(sh)	302.6(sh)		
2	333.6	398.5	383.0	380.7	392.8	340.94
	272.0	328.0(sh)	349.0	340.9	303.6(sh)	273.6
		278.5	302.8(sh)	301.9(sh)	278.5	
			278.8	279.3		
		339.5	406.0	389.0	384.3	400.5
3	272.5	329.6	356.0	348.0	348.9	342.3
		280.0	297.5(sh)	294.0(sh)	278.5	272.5
			276.0	279.0		

Table 2. ¹H (200 MHz, coupling constant *J* in Hz in parentheses) and ¹³C NMR (50.0 MHz) data of **1-3** in DMSO-d₆

Carbon no.	Chemical shifts δ in ppm					
	δ _H			δ _C		
	1	2	3	1	2	3
2	-	-	-	164.2	164.2	164.3
3	6.52 s	6.53 s	6.55 s	103.6	103.6	104.0
4	-	-	-	182.8	182.7	182.9
5	-	-	-	161.5	161.5	161.5
6	-	-	-	109.8	109.8	109.7
7	-	-	-	164.4	164.4	164.3
8	6.92 s	6.80 s	6.92 s	94.5	94.5	94.6
9	-	-	-	157.1	157.1	157.1
10	-	-	-	104.2	104.2	104.2
1'	-	-	-	121.9	122.0	122.3
2'	8.06 d (8.6)	7.95 d (8.7)	7.56 d (1.8)	129.3	129.2	109.7
3'	7.20 d (8.6)	6.94 d (8.7)	-	116.8	116.6	148.9
4'	-	-	-	162.1	162.4	151.6
5'	7.20 d (8.6)	6.94 d (8.7)	6.95 d (8.0)	116.8	116.6	116.6
6'	8.06 d (8.6)	7.95 d (8.7)	7.50 dd (1.8, 8.0)	129.3	129.2	121.2
C-glucose						
1''	4.60 d (9.5)	4.60 d (9.7)	4.56 d (9.7)	73.9	73.9	73.9
2''	3.20-3.90*	3.20-3.90*	3.20-3.90*	71.5	71.5	71.4
3''	3.20-3.90*	3.20-3.90*	3.20-3.90*	79.8	79.8	78.8
4''	3.20-3.90*	3.20-3.90*	3.20-3.90*	71.1	71.1	71.1
5''	3.20-3.90*	3.20-3.90*	3.20-3.90*	80.0	80.0	80.1
6''	3.20-3.90*	3.20-3.90*	3.20-3.90*	62.3	62.3	62.4
O-Glucose						
1'''	5.05 d (7.2)	-	-	102.0	-	-
2'''	3.20-4.00*	-	-	74.1	-	-
3'''	3.20-4.00*	-	-	77.1	-	-
4'''	3.20-4.00*	-	-	70.5	-	-
5'''	3.20-4.00*	-	-	77.0	-	-
6'''	3.20-4.00*	-	-	62.1	-	-
OCH ₃	-	-	3.90 s	-	-	56.8

* Overlapped peaks.

Table 3. ^1H (500 MHz, coupling constant J in Hz in parentheses) and ^{13}C NMR (125.0 MHz) data and key ^1H - ^{13}C long-range correlation observed in the HMBC spectra of **4** and **5**

Carbon no.	Chemical shift δ in ppm				HMBC correlations	
	δ_{H}		δ_{C}		2J	3J
	4	5	4	5		
1	6.11 d (2.6)	6.12 d (2.6)	123.8	123.8	C-2	C-3, C-5, C-9
2	-	-	147.4	147.4	-	-
3	-	-	199.9	199.9	-	-
4	-	-	50.7	50.7	-	-
5	-	-	137.6	137.6	-	-
6	5.84 br s	5.84 br s	122.5	122.5	C-5	C-4, C-8
7	2.40 m, 2.04 m	2.40 m, 2.05 m	24.6	24.6	-	C-9, C-14
8	2.10 m	2.09 m	43.3	43.3	C-9	C-13, C-15
9	-	-	50.5	50.4	-	-
10	3.69 br s	3.69 br s	38.2	38.2	-	C-6, C-11
11	-	-	216.6	216.6	-	-
12	3.41 d (14.8)	3.42 d (14.8)	50.1	50.1	C-11	C-14, C-17
	2.65 d (14.8)	2.65 d (14.8)				
13	-	-	50.4	50.4	-	-
14	-	-	50.2	50.2	-	-
15	1.96 dd (12.8, 8.8)	1.95 dd (12.8, 8.8)	46.7	46.7	C-16	C-13, C-17
	1.44 m	1.47 m				
16	4.48 m	4.47 m	71.6	71.7	C-15	C-20, C-13
17	2.59 d (00)	2.60 m	59.8	59.8	C-13	C-21, C-22
18	0.95 s	0.97 s	20.8	20.8	C-13	C-14, C-17
19	1.01 s	1.01 s	20.6	20.6	C-9	C-10, C-8, C-11
20	-	-	80.9	81.0	-	-
21	1.40 s	1.42 s	25.5	25.5	C-20	C-17, C-22
22	-	-	205.2	218.0	-	-
23	6.84 d (15.5)	2.90 ddd (17.4, 9.6)	121.4	33.6	C-22,	C-25
		2.78 ddd (6.2, 9.6)			C-24	
24	6.97 d (15.5)	1.74 m	155.5	38.2	C-25,	C-22
					C-23	
25	-	-	71.8	70.9	-	-
26	1.33 s	1.21 s	29.5	29.3	C-25	C-24, C-27
27	1.31 s	1.21 s	29.5	29.1	C-25	C-24, C-26
28	1.29 s	1.30 s	28.3	28.3	C-4	C-3, C-5, C-29
29	1.27 s	1.28 s	20.8	20.8	C-4	C-3, C-5, C-28
30	1.41 s	1.42 s	18.8	18.8	C-14	C-13, C-15
1'	4.65 d (7.5)	4.66 d (7.5)	101.3	101.3	C-2'	C-2, C-5'
2'	3.42 *	3.42 *	71.9	71.9	C-1'	C-4'
3'	3.44 *	3.44 *	78.3	78.3	-	C-1', C-5'
4'	3.53 *	3.53 *	70.9	70.9	C-5'	-
5'	3.38 *	3.38 *	80.1	80.1	C-6'	-
6'	4.04 dd (12.0, 3.5)	4.06 dd (12.1, 3.5)	62.1	62.1	C-5'	C-4'
	3.86 dd (12.0, 2.5)	3.87 dd (12.1, 2.5)				

*Overlapped peaks; Spectra obtained in CD_3OD

unknown materials. The antioxidant activity of **1-3** was determined by this method and the RC_{50} values were found to be 7.13×10^{-2} , 5.62×10^{-4} and 3.47×10^{-3} mg/mL, respectively. The RC_{50} value of the positive control, quercetin, was 2.78×10^{-5} mg/mL.

CONCLUSION

Since reactive oxygen species are important contributors to tissue injury, inflammation, cancer

and many other ailments, the antioxidant properties of **1-3** probably contribute, at least to some extent, to the pharmacological and traditional medicinal uses of the *C. colocynthis*.

ACKNOWLEDGEMENTS

We thank EPSRC National Mass Spectrometry Service Centre (Department of Chemistry, University of Wales, Swansea, Swansea, Wales, UK) for MS analysis.

REFERENCES

- GRIN Database, USDA, ARS, National Genetic Resources Program, Germplasm Resources Information Network (GRIN) online database, National Germplasm Resources Laboratory, Beltsville, Maryland. 2006; Available on-line at: <http://www.ars-grin.gov/cgi-bin/npgs/html/taxon.pl?10674>
- Madari, H., Jacobs, R. S., An analysis of cytotoxic botanical formulations used in the traditional medicine of ancient Persia as abortifacient, *J. Nat. Prods.* 2004; 67: 1204-1210.
- Dr. Duke's Phytochemical and Ethnobotanical Databases, Ethnobotanical uses of *Citrullus* (Cucurbitaceae). 2006; Available on-line at: <http://www.ars-grin.gov/cgi-bin/duke/ethnobot.pl>
- Maatooq, G. T., El-Sharkawy, S. H., Afifi, M. S., Rosazza P. N., C-*p*-Hydroxybenzoylglycoflavanones from *Citrullus colocynthis*, *Phytochemistry* 1997; 44: 187-190.
- Hatam, N. A. R., Whiting, D. A., Yousif, N. J., Cucurbitacin glycosides from *Citrullus colocynthis*, *Phytochemistry* 1989; 28: 1268-1271.
- Nmila, R., Gross, R., Rchid, H., Roye, M., Manteghetti, M., Petit, P., Tijane, M., Ribes, G., Sauvaire, Y., Insulinotropic effect of *Citrullus colocynthis* fruit extract, *Planta Medica* 2000; 66: 418-423.
- Yankov, L. K., Hussein, S. M., Fatty acid from the oil of the seeds of *Citrullus colocynthis*, *Dokl. Bolg. Akad. Nouk* 1975; 28: 209-12.
- Seger, C., Sturm, S., Mair, M.-E., Ellmerer, E. P., Stuppner, H., ¹H and ¹³C NMR signal assignment of cucurbitacin derivatives from *Citrullus colocynthis* (L.) Schrader and *Ecballium elaterium* L. (Cucurbitaceae), *Magnetic Resonance in Chemistry* 2005; 43: 489-491.
- Dr. Duke's Phytochemical and Ethnobotanical Databases, Chemical in *Citrullus colosynthis* (Cucurbitaceae). 2006; Available on-line at: <http://www.ars-grin.gov/cgi-bin/duke/farmacy2.pl>
- ISI Database, ISI Web of Knowledge Service for UK Education, supported by MIMAS at The University of Manchester and hosted by Thomson Scientific. 2006; Available on-line at: <http://portal.isiknowledge.com/portal.cgi?DestApp=WOS&Func=Frame>
- Delazar, A., Modarresi, M., Shoeb, M., Nahar, L., Reid R. G., Majinda, R. R. T., Sarker, S. D. Eremostachiin: A new furanolabdane diterpene glycoside from *Eromostachys glabra*, *Natural Product Research* 2006; 20: 167-172.
- Delazar, A., Celik, S., Gokturk, R. S., Unal, O., Nahar, L., Sarker, S. D., Two acylated flavonoids from *Stachys bombycina* and their free radical scavenging activity, *Die Pharmazie* 2005; 60: 878-880.
- Delazar, A., Byres, M., Gibbons, S., Kumarasamy, Y., Nahar, L., Modarresi, M., Shoeb, M., Sarker, S. D. (2004) Iridoid glycosides from *Eromostachys glabra*, *J. Nat. Prod.* 67: 1584-1587.
- Delazar, A., Reid, R. G., Sarker, S. D., GC-MS analysis of essential oil of the oleoresin from *Pistacia atlantica* var *mutica*, *Chemistry of Natural Compounds* 2004; 40: 24-27.
- Delazar, A., Shoeb, M., Kumarasamy, Y., Byres, M., Nahar, L., Modarresi, M., Sarker, S. D., Two bioactive ferulic acid derivatives from *Eremostachys glabra*, *DARU* 2004; 12: 49-53.
- Takao, T., Watanabe, N., Yagi, I., Sakata, K., A simple screening method for antioxidants and isolation of several antioxidants produced by marine bacteria from fish and shellfish, *Biosci. Biotech. Biochem* 1994; 58:1780-1783.
- Kumarasamy, Y., Fergusson, M., Nahar, L., Sarker, S. D., Bioactivity of moschamindole from *Centarea moschata*. *Pharmaceutical Biology.* 2002; 40:307-310.
- Bock, K., Jensen, S. R., Neilsen, B. J., Secogalioside, an iridoid glucoside from *Galium album* Mill and C-13 nmr-spectra of some seco-iridoid glucosides, *Acta Chemica Scandinavica* 1976; 30: 743-748.
- Davoust, D., Massias, M., Molho, D., C-13 nmr investigation of flavonoid C-β-D-glucosides - detection of a conformational equilibrium, *Organic Magnetic Resonance* 1980; 13: 218-219.
- Markham, K. R., Ternai, B., Stanley, R., Geiger, H., Mabry, T. J., Carbon-13 NMR studies of flavonoids—III : Naturally occurring flavonoid glycosides and their acylated derivatives, *Tetrahedron* 1978; 34: 1389-1397.
- Hostettmann, K., Bellmann, G., Tabacchi, R., Jacot-Guillarmod, A., *Phytochemistry of genus Gentiana* .3. study on flavone and xanthone compounds in leaves of *Gentiana lutea* L. 2. *Helvetica Chimica Acta* 1973; 56: 3050-3054.
- Goetz, M., Hostettmann, K., Jacot-Guillarmod, A., A new C-glycosylflavone from *Gentiana asclepiadea*, *Phytochemistry* 1976; 15: 2014.
- Wagner, H., Horhammer, L., Kiraly, C., Flavon-C-glykoside in *Croton zambezicus*, *Phytochemistry* 1970; 9: 897.
- Harborne, J. B. and Mabry, T. J., in 'The Flavonoids: Advances in Research'. University Press, Cambridge, pp. 19, 1982.
- Kumarasamy, Y., Byres, M., Cox, P. J., Delazar, A., Jaspars, M., Nahar, L., Shoeb, M., Sarker, S. D., Isolation, structure elucidation and biological activity of flavone C-glycosides from the seeds of *Alliaria petiolata*, *Chemistry of Natural Compounds* 2004; 40: 122-128.
- Mabry, T. J., Markham, K. R., Thomas, K. R., *The Systematic Identification of Flavonoids*, Springer-Verlag, New York, USA, 1970.