

Floral abundance and resource quality influence pollinator choice

Article (Accepted Version)

Fowler, Robert E, Rotheray, Ellen L and Goulson, Dave (2016) Floral abundance and resource quality influence pollinator choice. *Insect Conservation and Diversity*, 9 (6). pp. 481-494. ISSN 1752-458X

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/63437/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

1 Title: Floral abundance and resource quality influence pollinator choice

2

3 Authors: R. E. Fowler¹, E. L. Rotheray*¹, and D. Goulson¹

4 Affiliations:

5 ¹School of Life Sciences, University of Sussex, Falmer, BN1 9QG,

6 *Co-first author

7 Correspondence author: R.E.Fowler@sussex.ac.uk, Tel: +44 1273 872757

8

9 Key words: pollen, nectar, protein, sugar, bees

10 Running Title: Floral rewards drive pollinator choice

11

12

13

14

15

16

17

18

19

20

21

23

24 1. Pollinator declines caused by forage habitat loss threaten insect pollination services. Pollinating insects
25 depend on adequate floral resources, and their ability to track these resources. Variability of these resources
26 and the effect on insect foraging choice is poorly understood.

27 2. We record patterns of visitation to six wildflower species and test the hypotheses that: pollinators
28 preferentially visit the most rewarding flowers; nectar diurnal variations affect foraging preferences;
29 pollinators respond most strongly to nectar rewards.

30 3. Nectar volume and sugar concentration were negatively correlated within plant species over time of day
31 where greater concentration and lower volume was evident in the afternoon, but this did not correspond to
32 pollinator visitation. Both floral abundance and nectar quality (total sugar per inflorescence) positively affect
33 insect visitation. For some foragers the positive effects of high quality rewards were only evident when floral
34 abundance was high (>50 inflorescences per patch), perhaps reflecting the low probability of pollinators
35 detecting scarce rewards. Pollen quality (total protein per inflorescence) was negatively related to visitation
36 of *Apis mellifera* and *Bombus pascuorum*.

37 4. Fewer pollinators visiting flowers of higher pollen quality could reflect plant allocation trade-offs or the
38 presence of secondary metabolites in pollen, meaning pollen foraging is likely affected by factors other than
39 protein concentration. Nectar rather than pollen appeared to be the main driver of floral choice by insects in
40 this system.

41 5. Conservation schemes for bees in farmland or gardens might benefit from ensuring that rewarding plant
42 species are present at high density and/or are aggregated in space.

43

44

45

46

47

48

49

51

52 Over the past 80 years, local and UK-wide changes in farming practice and agricultural intensification
53 have led to a reduction in diversity of crop and non-crop plants (Robinson & Sutherland, 2002; Öckinger *et al.*,
54 2007; Geiger *et al.*, 2010). This includes the loss of meadow plants (Goulson *et al.*, 2005), arable weeds and
55 hedgerows (Hanley & Wilkins, 2015), which provide valuable foraging resources for flower visiting insects. This
56 habitat degradation has been identified as the primary reason for population declines in insects reliant on flowers
57 to provide nectar and/or pollen as a main food source, including adult butterflies (Brereton *et al.*, 2011) and adult
58 and offspring honey bees (Potts *et al.*, 2010), bumblebees (Goulson *et al.*, 2008) and solitary bees (Ollerton *et al.*,
59 2014). Nesting and hibernation resources aside, it is imperative that bees and other pollinating insects are able to
60 forage effectively for nutritional resources in increasingly fragmented landscapes in order to survive and
61 reproduce, particularly as they are facing other pressures such as diseases and pathogens (Cox-Foster *et al.*, 2007),
62 global environmental change (Tylianakis *et al.*, 2008) and pesticide use (Goulson *et al.*, 2015), all of which impact
63 on their survival. Population decline in flower visiting insects could jeopardise the pollination services provided
64 to entomophilous crops (Kremen *et al.*, 2002; Klein *et al.*, 2007).

65 Nectar provides sugars (mainly sucrose, glucose and fructose), which energise pollinators to continue
66 foraging (on nectar or other sources of nutrients), undertake nesting activities, find mates, and provide for
67 offspring. It also contains ions, water and small amounts of amino acids, which may contribute to nutrition (Kim
68 & Smith, 2000). Pollen comprises proteins, carbohydrates, lipids, vitamins and minerals (Roulston & Cane, 2000;
69 Nicolson, 2011). Although many flower visiting insects consume pollen for sustenance (e.g. beetles, flies), bees
70 (Hymenoptera: Apoidea) also collect pollen to feed larvae, and many simultaneously collect both nectar and pollen
71 from flowers, depending on what requirements they have and what flower species they are feeding on (Heinrich,
72 1979a; Heinrich, 1979b; Goulson *et al.*, 2005).

73 Survival and reproductive success of pollinating insects is dependent on them successfully gathering
74 adequate protein and sugar to provide for their energy and nutritional needs. For example, many Lepidoptera,
75 Diptera and Hymenoptera rely on energy gained from nectar to undertake mating flights, for dispersal and/or
76 migration, and to find suitable places to lay their eggs. Butterflies tend to exhibit lower fecundity when nectar
77 limited (Boggs & Ross, 1993), and the hoverfly *Episyrphus balteatus* has greater longevity when fed a sugar and
78 protein rich diet (Pinheiro *et al.*, 2015). Bees require both pollen and nectar to feed to their offspring. When fed

79 on protein rich diets, bumblebee colonies are more reproductively successful (Génissel *et al.*, 2002; Kitaoka &
80 Nieh, 2009), and the solitary bee *Lasioglossum zephyrum* (Roulston & Cane, 2002) and honey bees (Basualdo *et*
81 *al.*, 2012) produce larger adults. Larger bees generate and retain heat faster leading to earlier and more frequent
82 forage flights (Stone, 1993), they have greater diapause survival (Strohm and Linsenmair, 1997) and are better
83 able to cope with adverse conditions such as parasitism and disease (Alaux *et al.*, 2010; Di Pasquale *et al.*, 2013).

84 In order to forage effectively, pollinators use olfactory cues to enable detection of non-depleted nectar
85 resources (Goulson *et al.*, 1998a; Howell & Alarcón, 2007) and greater nectar volume and sugar content (Pyke,
86 1978; Heinrich, 1979a; Wolff, 2006). Likewise, insect foragers tend to select pollen with greater protein and
87 essential amino acid content (Levin & Bohart, 1955; Schmidt, 1982; Robertson *et al.*, 1999; Cook *et al.*, 2003;
88 Arenas & Farina, 2012), and obligate insect pollinated plant species produce pollen that is richer in protein and
89 amino acids compared to facultative species (Hanley *et al.*, 2008, but see Roulston *et al.*, 2000).

90 Few studies have investigated how eusocial or solitary bees integrate information on nectar and pollen
91 quantity and quality simultaneously, and those that have, indicate that nectar is the primary factor influencing
92 foraging preference for bees, and pollen a secondary consideration (Konzmann & Lunau, 2014; but see Somme
93 *et al.*, 2015). In addition, studies focusing on flower selection by pollinators tend to look only at individual species
94 of bee and specific pollinators for individual plant species (e.g. Robertson *et al.*, 1999), or solely bees as a group
95 (e.g. Heinrich, 1979b; Pernal & Currie, 2001). Floral choice tests are frequently undertaken in controlled
96 conditions (e.g. Konzmann & Lunau, 2014), but the limited number of field studies investigating the influence of
97 rewards on the foraging decision of pollinators makes it difficult for the conclusions gained by laboratory studies
98 to be applied, especially as there are external factors in the natural environment likely to influence the results,
99 such as the spatial distribution and reward phenology of flowers. For example, Hanley & Wilkins (2015) describe
100 greater food plant abundance in roadside, compared with field facing hedgerows, and noted a corresponding
101 increase in bumblebee abundance. In addition, several studies indicate that some flower species offer less nectar
102 in the middle of the day and afternoon compared to the morning and evening (Mačukanović-Jocić *et al.*, 2004;
103 Silva *et al.*, 2004; Mačukanović-Jocić & Đurđević, 2005), providing a further challenge to efficient foraging by
104 pollinators since the relative rewards provided by different flower species may alter hour by hour through the day.

105 In this study, we examine the foraging choices made by all flower visiting insects in relation to the
106 relative nectar and pollen quantity and quality of six common plant species under natural conditions. We tested
107 the hypothesis that nectar quantity or quality shows diurnal variation and that nectar or pollen quantity or quality,

108 or a combination of reward metrics, predicts insect visits. We record how nectar volume and sugar concentration
109 changes through the day, and also the pollen weight and protein concentration for each test plant species, and
110 relate these values to insect flower choices through the day. Specifically, this study's objectives were to; (i) record
111 nectar and pollen quantity and quality estimates for six test plant species, (ii) investigate the diurnal variation in
112 nectar quantity and quality for the test plant species, and (iii) assess how nectar and pollen quantity and quality
113 influence foraging choices by flower visiting insects.

114

115 Methods

116

117 *Test Plant Species*

118

119 Six species of flowering plant that are common in southern UK were selected for nectar and pollen
120 quantity and quality estimation, and insect visitation surveys in the field. These were *Lamium album* L. and
121 *Glechoma hederacea* L. (Lamiaceae), *Crataegus monogyna* Jacq. and *Rubus fruticosus* L. agg. (Rosaceae),
122 *Symphytum officinale* L. (Boraginaceae) and *Ranunculus repens* L. (Ranunculaceae). These spring or early
123 summer flowering plants considered to be beneficial foraging plants for insects in previous studies (Lack, 1982;
124 Goulson *et al.*, 1998b; Kipling & Warren, 2013) (further species details are listed in Table S1 in the Electronic
125 Supplementary Material). Previously, *R. fruticosus* and *S. officinale* were included in studies of pollen protein
126 quality and its effects on bee foraging preferences (Hanley *et al.*, 2008), but to our knowledge the other test plant
127 species have not been studied in this context.

128

129 *Site Selection*

130

131 Sites were chosen to be included in this study if they contained at least three of the test plant species
132 within 50 m of each other, were on chalk soil, were easy to access and were subject to intermediate levels of
133 disturbance (e.g. mowed once a year, or grazed intermittently by sheep). Sites could be divided into two groups,

134 road verges and semi-improved grassland. In terms of management, road verge sites were mowed once a year and
135 semi-improved grassland were grazed several times throughout the year. However, none of the survey sites were
136 either mown or grazed whilst this study was taking place. Once sites were identified, nectar and pollen samples
137 were collected from test plant species and visitation sampling was conducted.

138

139 *Nectar Volume and Sugar Concentration*

140

141 Nectar sampling was undertaken on 30 inflorescences of each test plant species, collected from at least
142 three different survey sites. Nectar was sampled from inflorescences of plant species that had recently come in to
143 bloom and were easily accessible. For each inflorescence nectar production was estimated from morning,
144 afternoon and evening periods for each plant species tested. For each species then flowers were emptied of nectar
145 and bagged by 09:00 (GMT) using a fine-mesh cotton fabric and masking tape. At 15:00, 21:00 and 09:00 the
146 following morning flowers were emptied of nectar using 5 µl micropipettes (BLAUBRAND® intraMARK,
147 Wertheim, Germany). The nectar from each inflorescence was then expelled onto a field refractometer
148 (Bellingham and Stanley Ltd, Basingstoke, Hants, UK) to measure the proportion of sugar in each sample (Bolten
149 *et al.*, 1979). This produced both mean nectar volume and mean sugar concentration for each species in each time
150 period, and over a 24 hour period when totalled.

151

152 *Pollen Weight and Protein Concentration*

153

154 Pollen was collected from 10 inflorescences from at least three different sites for each plant species. Standing
155 crop of pollen was taken just once for each replicate so variation in pollen quantity or quality was not measured
156 over time of day. Flowers were collected in the field before anthers had dehisced, and placed in water in an
157 unheated and well ventilated laboratory until anthers opened. Pollen was collected systematically until anthers
158 were empty. It is conceivable that placing flowers in water affects pollen quality, but all plant species were
159 treated in the same way. Pollen was either stored in a – 20 °C freezer for drying at a later date, or immediately
160 dried in an oven for 24 hours at 40 °C. After drying, pollen was weighed to measure total pollen weight per

161 inflorescence. Protein extraction and detection was undertaken using the Bradford assay (Bradford, 1976). The
162 assay binds protein to Coomassie Brilliant Blue G-250 dye (4.7% [weight: volume] ethanol, and 8.5% [weight:
163 volume] phosphoric acid dissolved in water). Light absorbance is then measured against known protein
164 standards. From each of the 10 inflorescence pollen from each species, 1 mg pollen were dusted with aluminium
165 powder, wetted with 20 µl 0.1 mol/L NaOH and ground with a micro-pestle. Ground pollen was reanimated
166 with 480 µl 0.1 mol/L NaOH and placed in a refrigerator for 24 hours before analysis, but used within 1 week.
167 Prior to absorbance measurement samples were placed in boiling water for 5 minutes and centrifuged for 5
168 minutes. Then 10 µl of supernatant was slowly vortexed with 300 µl of dye reagent. This was repeated in
169 triplicate for each sample and left to incubate at room temperature for 15 minutes.

170 Protein standards were made up each time samples were run, using pre-mixed concentrations of Bovine
171 Serum Albumin (BSA) from the BIO-RAD (Hertfordshire, UK) Quick-Start™ Bradford Protein Assay kit. Once
172 samples and standards were created, they were measured for absorbance within an hour of mixing at 595 nm using
173 a Thermo Scientific (Paisley, UK) Nanodrop 2000 UV-Vis Spectrophotometer. This produced the standing crop
174 of both mean pollen weight and a crude mean protein concentration for each test plant species.

175

176 *Visitation Surveys*

177

178 Visitation surveys took place at seven sites across Sussex, UK between May and June 2014 (for site
179 details see Table S2). Surveys were undertaken in the morning (08:00—10:00), afternoon (13:00—15:00) and
180 evening (19:00—21:00). Each survey consisted of a 10 minute standing observation of a 4 m² area of each test
181 plant species at each site. The number and species of all visiting insects to that plant species were recorded. All
182 observed flower visiting insects appeared to collect nectar and, in most cases, pollen from plant species, therefore
183 we did not attempt to distinguish which resources insects were collecting. The number of inflorescences of the
184 plant species, ambient temperature, and wind speed were recorded for each survey. Surveys were only conducted
185 when air temperature was > 14°C and average wind speeds was < 20 mph.

186 Common species of bumblebee (genus *Bombus*) and European honeybees (*Apis mellifera* L.) were
187 identified on the wing. Due to the unreliability of morphological characters for separating *Bombus terrestris* L.
188 and *B. lucorum* agg. workers in the field, these two species were grouped. Although others have recorded flower

189 visiting bumblebees into groups based on colour type (Haghton *et al.*, 2003), in this study the bumblebee species
190 observed were easily separated to species in the field. However, faded or suspected cleptoparasitic bees (*Psithyrus*
191 spp.) were caught and checked using a hand magnifying glass. Other visiting insects too small to be identified on
192 the wing were collected and identified to species or genus. Although unlikely given the short period of each survey,
193 the possibility of double counting insects was the same for all surveys due to equal sampling time and therefore
194 assumed to be unbiased across the study.

195

196 *Statistical Analysis*

197

198 Means tests of variance were used to assess differences in floral resources available from the plant species,
199 as well as the number of insect visits, between periods of the day. Based on whether resource and insect visit
200 metrics showed homogeneity of variance between species or periods of the day using Levene's test, either
201 ANOVA (parametric) or Kruskal-Wallis (non-parametric) tests were applied to assess variance for all groups
202 before either Tukey HSD tests or pairwise Wilcox tests were applied to assess differences between groups,
203 respectively.

204 Generalized Linear Mixed Models (GLMMs) were used to test the effects of resource metrics on insect
205 visitation (i.e. counts per 10 minute survey). Prior to applying models, proposed explanatory variables were
206 checked for multicollinearity using Variance Inflation Factors (VIFs), if variables had VIFs greater than 3 or
207 correlation coefficients more than 0.6 with other variables, they were excluded from models (Zuur *et al.*, 2010).
208 These analyses indicated nectar volume and pollen weight were correlated with sugar and protein concentration,
209 respectively. To enable modelling of resource quality and quantity without violating model assumptions, nectar
210 volume and pollen weight were multiplied by sugar and protein concentration to create total sugar and protein in
211 milligrams per inflorescence as measurements of quality in the plant species. To account for the change in density
212 of the solution based on the amount of sugar recorded, percentage sugar was multiplied by the mass density (g/cm³)
213 of sugar at that concentration. These measurements also allowed the interpretation of the nutritional gain available
214 to foraging insect pollinators, and is referred to as total sugar and total protein from here on.

215 GLMMs were initially run with the environmental variables temperature and wind speed included as
216 explanatory variables, however these had no effect on visitation or the outcome of the models so were removed.

217 We modelled five visitation response variables (including: all insects; all bumblebees; the three most recorded
218 species, *B. pascuorum* Scopoli, *B. pratorum* L., and *A. mellifera*) in response to the number of inflorescences (log
219 transformed) for each test plant species surveyed in each observation area (4 m²) (floral abundance), and total
220 sugar and protein. We did not analyse data on other groups of flower visiting insects alone as they were recorded
221 in such small numbers that data analysis would have been unreliable. However, they were included in the data
222 analysis as part of ‘all insects’ in statistical models. We included interactions between floral abundance and total
223 sugar and protein explanatory variables. Sample site was fitted as a random factor because observations were
224 nested within sample sites and contained different combinations of test plant species. Visitation rates were count
225 data, therefore models were applied with Poisson errors (O’Hara & Kotze, 2010). Model residuals were assessed
226 for normality and heteroscedasticity. Model simplification was carried out using likelihood ratio tests, and
227 sequentially deleting terms that did not significantly decrease model deviance, beginning with higher order
228 interactions.

229 To further investigate how particular plant species affect insect visitation, linear mixed effects (LME)
230 models were used to test the effects of floral abundance and species on the same response variables as the GLMMs,
231 with site as a random factor. Models were tested for significance using likelihood ratio tests, with and without
232 species as an explanatory variable.

233 Already published protein estimates for each plant species were compared to confirm the extraction
234 results and accuracy of our values. We found that only *Lamium album* (4.48%) differed from the literature for
235 *Lamium* sp. (22.8% in Roulston *et al.*, 2000), so to assess whether this estimate affected our findings, we
236 substituted it in the two models that included total protein (for *A. mellifera* and *B. pascuorum* visitation).

237 Statistical analysis was conducted using R v3.1.2 (R Core Team, 2014) within RStudio (RStudio, 2012)
238 using packages ‘lme4’ (Bates *et al.*, 2015) and ‘usdm’ (Naimi, 2013), and plots were generated using ‘ggplot2’
239 (Wickham, 2009).

240

241 Results

242

243 *Nectar and Pollen Quantity and Quality*

244

245 A large amount of between-inflorescence variation was found in nectar and pollen resources for each test
246 plant species (Table 1). Nevertheless, nectar volume (Kruskal-Wallis $H = 211.64$, d.f. = 5, $P < 0.001$), nectar sugar
247 concentration ($H = 112.61$, d.f. = 5, $P < 0.001$) and pollen protein concentration (ANOVA $F_{5, 54} = 20.6$, $P < 0.001$)
248 varied significantly between plant species, with only pollen weight ($F_{5, 54} = 1.25$, $P = 0.3$) showing no significant
249 differences.

250 Over 24 hours, *Rubus fruticosus* produced the greatest volume of nectar, which contained the lowest
251 sugar concentration (Table 1). In contrast, *Ranunculus repens* produced the least nectar, whilst *Crataegus*
252 *monogyna* offered the highest sugar concentration. *Lamium album* pollen offered the greatest mass per
253 inflorescence but had the lowest protein concentration of all species. *C. monogyna* produced the least pollen mass
254 whilst *G. hederacea* offered the highest protein concentration. We found no correlation between nectar and pollen
255 quantity (Pearson $R = -0.46$, $P = 0.26$) or quality (Pearson $R = -0.33$, $P = 0.52$).

256 Nectar volume and pollen weight were multiplied by sugar and protein concentration, respectively, to
257 create total sugar and protein in milligrams per inflorescence (Table 1). Over 24 hours, *Symphytum officinale*
258 produced the greatest total sugar per inflorescence whilst *R. repens* produced the least. *C. monogyna* contained
259 the most total protein per inflorescence and *S. officinale* the least (Table 1).

260

261 *Diurnal Variation in Nectar Quantity and Quality*

262

263 Mean nectar volume and sugar concentration averaged between plant species (regardless of time period)
264 were not significantly correlated (Pearson $R = -0.49$, $P = 0.32$, Fig. 1A). When mean for each species at each time
265 period was taken, there was a significant negative correlation between nectar volume and sugar concentration
266 (Pearson $R = -0.52$, $P = 0.02$, Fig. 1B).

267 Volumes of nectar produced differed markedly between species, and between morning, afternoon and
268 evening periods. *R. fruticosus* produced the greatest volume of nectar of all species in the evening and morning,
269 yet recorded the lowest sugar concentration of all species in these periods (Table 1). The lowest volume of nectar
270 was detected in *R. repens* in the evening, and the greatest sugar concentration was found in *C. monogyna* in the

271 afternoon. When averaged across species, mean nectar volume did not differ significantly between periods of the
272 day ($F_{2, 15} = 1.18$, $P = 0.35$). However, mean nectar sugar concentration across species increased significantly in
273 the afternoon compared to the morning ($F_{2, 15} = 4.01$, $P = 0.04$) (Fig. 2).

274 Once sugar and protein concentrations were calculated between time periods, total sugar was
275 significantly lower in the afternoon period compared to the morning or evening for *R. fruticosus*, *G. hederacea*,
276 *L. album* and *R. repens* (Fig. 3). The exceptions were *C. monogyna* which produced comparable amounts of total
277 sugar in each time period sampled, and *S. officinale* which had marginally more total sugar in the afternoon
278 compared to morning and evening periods.

279

280 *Foraging choices of flower visiting insects*

281

282 In total, we made 112 ten-minute observations (38, 39 and 35 in morning, afternoon and evening periods,
283 respectively), between May and June 2014 at seven sites in Sussex, during which 574 insects were recorded
284 visiting test plant species. Proportionately, of all insect visits, 93% were by bees (Hymenoptera: Apidae), 6% by
285 Diptera (< 1% of which were hoverflies [Syrphidae]), < 1% by Lepidoptera and < 1% by Vespidae (Table S3).
286 Of all flower visiting insects, the three most numerous were the bee species *Bombus pascuorum*, *B. pratorum*, and
287 *Apis mellifera*, which represented 33%, 30% and 16% of all insect visits, respectively (Table S3). The majority
288 of insects were recorded visiting *R. fruticosus* (47%), *S. officinale* (24%) and *L. album* (19%) (Table S3). LME
289 models showed significant species-specific preferences for the most commonly recorded flower visiting insects
290 (Table S4). *B. pascuorum* mainly visited *L. album* and *S. officinale* (49% and 31% of this species' visits,
291 respectively), *B. pratorum* preferred *R. fruticosus* and *S. officinale* (57% and 36%), and *A. mellifera* mainly visited
292 *R. fruticosus* (87%) (Fig. S1 & Table S3).

293 Fewer insects were recorded in the evening (122) than the morning (221) and afternoon periods (231),
294 and differences in abundance were also evident within and between recorded insect and test plant species (Table
295 S3). However, these differences were not significant when tested between surveys for all insect abundance ($F_{2, 109}$
296 = 2.59, $P = 0.07$). No significant difference was found between any response variable between different periods
297 of the day when all test plant species were grouped or tested separately, except *G. hederacea* which had
298 significantly more insects visiting during the afternoon compared to morning and evening periods ($F_{2, 24} = 9.03$, P

299 = 0.001; Fig. 4). This however does not correspond with the timing of maximum sugar availability, which
300 appeared highest in the morning or evening for most species (Fig. 3). As standing crop of pollen was taken just
301 once for each replicate, we could not quantify variation in pollen quantity or quality over the day.

302 Overall, flower abundance had a positive effect on the total number of bumblebee (bees within the genera
303 *Bombus*) and all insect visitation (Table 2). Total sugar predicted insect visitation, with this relationship being
304 positive for all response categories apart from *B. pascuorum* which was negative (Table 2). There were also a
305 number of significant higher order interactions between floral abundance and total sugar and protein. For *Apis*
306 *mellifera*, all bumblebees and all insects, the relationship between insect visitation and total sugar was weak or
307 absent at low floral abundance (10 – 50 inflorescences per 4 m²) but positive at high floral abundance (51 – 1000
308 inflorescences per 4 m²) (Fig. 5). Total protein had a significant negative effect on *A. mellifera* visitation (Table
309 2). In addition, a negative interaction between total protein and floral abundance was found for *B. pascuorum*
310 (Table 2), with the relationship weak or absent at low floral abundance (10 – 50 inflorescences per 4 m²) but
311 negative at high floral abundance (51 – 1000 inflorescences per 4 m²). However, the three plant species most
312 visited by insects (*R. fruticosus*, *S. officinale* and *L. album*) were also recorded as producing the lowest total pollen
313 protein. When the protein value for *L. album* was replaced with the higher value reported in Roulston *et al.* (2000)
314 there was no longer a negative significant effect of total protein on *B. pascuorum* visitation, and we found a
315 positive significant effect of total sugar (Table S5).

316

317 Discussion

318

319 Many insects forage on flowering plants to gain key nutritional resources, largely nectar and pollen. What
320 factors determine when plants secrete nectar is still largely unknown, as both internal and external factors can
321 affect the rate of secretion (Heil, 2011). In our study, floral rewards of nectar and pollen (apart from pollen weight)
322 significantly differed between test plant species. Similarly, nectar resources varied between periods of the day,
323 whilst the single measure of pollen for each replicate meant this resource could not be tested for diurnal differences.
324 Nectar volume and sugar concentration were negatively correlated in test plant species, with sugar concentration
325 greater in the afternoon, which corresponds with previous studies showing similar trends (Mačukanović-Jocić *et al.*
326 *al.*, 2004; Silva *et al.*, 2004; Mačukanović-Jocić & Đurđević, 2005). This may be due to decreased water

327 availability or flowers reabsorbing nectar in the afternoon (Silva *et al.*, 2004) when humidity decreases and
328 temperatures rise (Silva *et al.*, 2004; Mačukanović-Jocić & Đurđević, 2005). While our results suggests that nectar
329 resources vary in quantity and quality across the day, insect visitation did not track nectar availability (with the
330 exception of *G. hederacea* where the opposite was found: more insects were recorded visiting this species in the
331 afternoon when lower total sugar was recorded). This may be an indication that some pollinators are not capable
332 of accurately responding to changes in nectar production throughout the day, or it may be that their activity is
333 constrained by other factors such as temperature. It is important to note that our methods meant that afternoon and
334 evening nectar sampling occurred at 6-hour intervals, compared with morning sampling which was undertaken
335 12 hours after the flowers had been emptied and bagged the previous evening. We considered this as providing a
336 fair estimate of how much nectar is likely available to foraging insects by the morning, however, the effect of
337 early morning foraging between 05:00 and 09:00 GMT was not represented by our data and this should be taken
338 into account in interpretations.

339 Due to the high energy cost of foraging, successfully selecting the most rewarding flowers is predicted
340 to have a large impact on survival and reproductive success, especially when floral resources are fragmented. The
341 most documented resource offered to insects by plants as an attractant for pollination is nectar. Foraging insects
342 are capable of learning nectar rewards gained from visited flowers (Pyke, 1978), preferring to forage on flowers
343 with significantly more nectar (Heinrich, 1979a; Wolff, 2006) and with greater sugar concentration (Hendriksma
344 *et al.*, 2014). After testing for relationships between nectar and pollen resources and pollinator visitations our
345 results support this, as typically, we found that greater nectar resources had a positive effect on insect and bee
346 visitation. Single species specialism in insect-plant mutualisms is rare (Waser *et al.*, 1996), and the majority of
347 flower visiting insects have flexible foraging choices. Foraging bumblebees show flower constancy (‘majoring’
348 on one particular species of known reward) and flower infidelity (‘minoring’ on other flowers checking reward
349 change over time) (Heinrich, 1979b). This behaviour allows foragers to track resources in multiple flower species
350 in a habitat. In our study, the positive effects of high floral rewards i.e. sugar were often only apparent when the
351 floral abundance of test plant species was high (> 50 inflorescences), with scarce flowers tending to be visited
352 less frequently even when comparatively highly rewarding. Our results appear to support Heinrich (1979b); if a
353 flower species is both abundant and rewarding then insects are very likely to have discovered its value and
354 preferentially visit it. In addition, where there are more flowers in a patch, there is a greater total quantity of nectar.
355 Hence, although quality has an influence on forager choice, the abundance of floral rewards in the local
356 environment is important in insect selection of resources.

357 There is evidence that bees show preferences for pollen with higher protein (Levin & Bohart, 1955;
358 Robertson *et al.*, 1999) and essential amino acid content (Cook *et al.*, 2003), and these preferences are supported
359 by obligate insect pollinated plant species producing pollen that is richer in protein and amino acids (Hanley *et al.*
360 *et al.*, 2008). However, other studies describe contrasting results where protein-rich pollen seems to be preferred in
361 some cases, but not in others (Wille *et al.*, 1985 in Praz *et al.*, 2008; Roulston & Cane, 2002). Our results suggest
362 greater total protein was negatively related to visitation by *A. mellifera*, and for *B. pascuorum* the negative
363 relationship between protein content and visitation was more apparent when the floral abundance of test plant
364 species was high (> 50 inflorescences). There are several possible explanations. Firstly, although not significant,
365 there was a negative relationship between nectar and pollen quantity per inflorescence (correlation coefficient -
366 0.46), so if bees are basing decisions primarily on nectar rewards they will tend to visit flowers with less pollen.
367 We did not attempt to discern whether bees were collecting pollen only, nectar only, or both, but previous studies
368 suggest that the majority of visits are for nectar (e.g. Goulson *et al.*, 2005). Second, bees may have been responding
369 to other nutritional compounds present in pollen. Plant species may be making trade-offs between protein and
370 other nutritional elements that drive foraging preferences such as sterols (Somme *et al.*, 2014), lipids and starch
371 (Roulston & Cane, 2000) or pollen-specific odours (Dötterl & Vereecken, 2010) not addressed in this study. Third,
372 some plant species protect their pollen with defensive secondary compounds that may affect bee foraging choices
373 (Gosselin *et al.*, 2013). For example, *Echium vulgare* has high protein content and also high concentrations of the
374 hepatotoxins 1,2-dehydropyrrolizidine alkaloids and their N-oxides (Boppré *et al.*, 2005), which can be toxic to
375 insects (e.g. Boppré *et al.*, 2005; Sedivy *et al.*, 2012) and affect flower selection (Kessler & Halitschke, 2009). In
376 this study the species recorded with the greatest amount of pollen protein were *Glechoma hederacea* and
377 *Ranunculus repens*, and pollen of *Ranunculus* spp. is known to contain the toxic lactone protoanemonin
378 ranunculin with reportedly negative effects on honeybees (Sedivy *et al.*, 2012; Jurgens & Dötterl, 2004); and *G.*
379 *hederacea* is toxic to some species of herbivorous insects as it produces a defensive insecticide protein (Hutchings
380 & Price, 1999; Van Damme, 2008). It is also important to note that the negative effect we found did not persist
381 for *B. pascuorum* once our values for *Lamium album* were replaced with higher values found in the literature. Our
382 methods for extracting and measuring protein content are crude, and results can be variable so should be treated
383 with caution. More detailed investigations are needed in which the full range of compounds present in pollen are
384 quantified if we are to fully understand how bees choose which flowers to visit when collecting pollen.

385 Most studies do not measure both nectar and pollen rewards in relation to insect visit frequency, and in
386 studies that do, conflicting results have been found. Konzmann & Lunau (2014) found that, in bumblebees, nectar

387 rewards appear more important than pollen quality, whereas Somme *et al.*, (2014) found when pollen loads are
388 analysed in conjunction with nectar from forage plants, both nectar and pollen quality appear important. In this
389 study, while pollen and nectar are not negatively correlated, total nectar production appears to influence the
390 visitation of insects to a greater extent than pollen. This could mean that flower visiting insects are more concerned
391 with the quality of nectar, with pollen as a secondary consideration.

392 Insect visitation to test plant species appears to be species-specific, which can go further to explain our
393 results. *B. pascuorum* mainly visited *L. album* and *S. officinale*, most likely due to its longer tongue allowing
394 access to their deeper corollas. Contrary to other visiting species, we detected a negative effect of total sugar on
395 *B. pascuorum* visitation. This may be because *B. pascuorum* had little competition from other insect foragers for
396 *L. album* (only 16 other individuals recorded foraging on this species, which produced comparatively low amounts
397 of sugar), or because this species was at the start of its life cycle when this study's sampling was undertaken and
398 newly emerged queens were focusing foraging efforts on pollen collection. *Bombus pratorum*, which as a short-
399 tongued bee may have exhibited restricted foraging choices, tending to visit flowers with greater total sugar.
400 Although this species has a short tongue, it is a secondary nectar robber; 93% of recorded visits to *S. officinale*
401 were via robbing (behaviour previously reported in Goulson *et al.*, 1998b). *Apis mellifera*, although also a short
402 tongued species, mainly visited *R. fruticosus* and was not recorded robbing in this study, though it has been
403 recorded acting as a secondary robber elsewhere (Darwin, 1872). Differences between foraging behaviour of
404 bumblebees and honeybees suggest bumblebees (*B. terrestris* and *B. pascuorum*) show less fidelity when
405 collecting pollen than honeybees, which have a highly flower-constant strategy (Leonhardt & Blüthgen, 2012).

406 Promoting and developing resources for pollinating insects is predominantly conducted through agri-
407 environmental schemes promoting flower-rich field edges (Carvell *et al.*, 2007), or through targeted planting in
408 urban spaces or private gardens (Hanley *et al.*, 2014). However, our understanding of the way in which pollinating
409 insects respond to differences in the quality of resources offered by managed planting is limited. Our results
410 suggest more consideration should be given when selecting plants for conservation management efforts, notably
411 in terms of differing insect species requirements for pollen and nectar quality. Differences in flower selection
412 between pollinator species may relate to the variation of life histories and may reduce competition for resources.
413 Nectar resource quality appears to be the main driver of flower selection by most insect foragers in this study but,
414 importantly, the benefits of greater resource quality in plants is dependent on local floral abundance. One practical
415 conclusion to be drawn from this is that bees may benefit more from plantings of flowers (be they in farmland,

416 parks or gardens) where species are presented in large clumps rather than in heterogeneous mixtures. More broadly,
417 it is clear that there is still much that we do not understand about the role of sugars, proteins and other compounds
418 in nectar and pollen in determining the foraging preferences of pollinators under field conditions.

419

420 Acknowledgments

421

422 We thank Rebecca Crowe and Charles Linton for help with observation surveys, and the Biotechnology
423 and Biological Sciences Research Council (BBSRC) (BB/J014753/2) for funding this project. We also thank Mick
424 Hanley and three anonymous reviewers for their comments on the original manuscript, which helped improve it
425 considerably.

426

427 References

428

429 Alaux, C., Ducloz, F., Crauser, D. & Le Conte, Y. (2010) Diet effects on honeybee immunocompetence.

430 *Biology Letters*, **6**, 562–565.

431 Arenas, A. & Farina, W.M. (2012) Learned olfactory cues affect pollen-foraging preferences in honeybees, *Apis*

432 *mellifera*. *Animal Behaviour*, **83**, 1023–1033.

433 Basualdo, A.M., Barragán, S., Vanagas, L., García, C., Solana, H., Rodríguez, E. & Bedascarrasbure, E. (2012)

434 Conversion of high and low pollen protein diets into protein in worker honey bees (Hymenoptera:

435 Apidae). *Economic Entomology*, **106**, 1553–1558.

436 Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal*

437 *of Statistical Software*, **67**, 1–48.

438 Brereton, T.M., Roy, D.B., Middlebrook, I., Botham, M. & Warren, M. (2011) The development of butterfly

439 indicators in the United Kingdom and assessments in 2010. *Journal of Insect Conservation*, **15**, 139–151.

- 440 Boggs, C.L. & Ross, C.L. (1993) The effect of adult food limitation on life history traits in *Speyeria mormonia*
441 (Lepidoptera: Nymphalidae). *Ecology*, **74**, 433–441.
- 442 Bolten, A.B., Feinsinger, P., Baker, H.G. & Baker, I. (1979) On the calculation of sugar concentration in flower
443 nectar. *Oecologia*, **41**, 301–304.
- 444 Boppré, M., Colegate, S.M. & Edgar, J.A. (2005) Pyrrolizidine alkaloids of *Echium vulgare* honey found in
445 pure pollen. *Journal of Agricultural and Food Chemistry*, **53**, 594–600.
- 446 Bradford, M.M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein
447 utilizing the principle of protein-dye binding. *Analytical Biochemistry*, **72**, 248–254.
- 448 Carvell, C., Meek, W.R., Pywell, R.F., Goulson, D. & Nowakowski, M. (2007) Comparing the efficacy of agri-
449 environment schemes to enhance bumblebee abundance and diversity on arable field margins. *Journal of*
450 *Applied Ecology*, **44**, 29–40.
- 451 Cook, S.M., Awmack, C.S. & Williams, I.W. (2003) Are honey bees' foraging preferences affected by pollen
452 amino acid composition? *Ecological Entomology*, **28**, 622–627.
- 453 Cox-Foster, D.L., Conlan, S., Holmes, E.C., Palacios, G., Evans, J.D., Moran, N.A., Quan, P.-L., Briese, T.,
454 Hornig, M., Geiser, D.M., Martinson, V., vanEngelsdorp, D., Kalkstein, A.L., Drysdale, A., Hui, J., Zhai,
455 J., Cui, L., Hutchison, S.K., Simons, J.F., Egholm, M., Pettis, J.S. & Lipkin, W.I. (2007) A metagenomic
456 survey of microbes in honey bee colony collapse disorder. *Science*, **318**, 283–287.
- 457 Darwin, C. (1872) *The effects of cross and self fertilisation in the vegetable kingdom*. Murray, London, UK.
- 458 Di Pasquale, G., Salignon, M., Le Conte, Y., Belzunces, L.P., Decourtye, A., Kretzschmar, A., Suchail, S.,
459 Brunet, J. & Alaux, C. (2013) Influence of pollen nutrition on honey bee health: do pollen quality and
460 diversity matter? *PLoS ONE*, **8**, e72016.
- 461 Dötterl, S. & Vereecken, N.J. (2010) The chemical ecology and evolution of bee–flower interactions: a review
462 and perspectives. *Canadian Journal of Zoology*, **88**, 668–697.
- 463 Geiger, F., Bengtsson, J., Berendse, F., Weisser, W.W., Emmerson, M., Morales, M.B., Ceryngier, P., Liira, J.,
464 Tschardtke, T., Winqvist, C., Eggers, S., Bommarco, R., Pärt, T., Bretagnolle, V., Plantegenest, M.,
465 Clement, L.W., Dennis, C., Palmer, C., Oñate, J.J., Guerrero, I., Hawro, V., Aavik, T., Thies, C., Flohre,

- 466 A., Hänke, S., Fischer, C., Goedhart, P.W. & Inchausti, P. (2010) Persistent negative effects of pesticides
467 on biodiversity and biological control potential on European farmland. *Basic and Applied Ecology*, **11**,
468 97– 105.
- 469 Génissel, A., Aupinel, P., Bressac, C., Tasei, J.-N. & Chevrier, C. (2002) Influence of pollen origin on
470 performance of *Bombus terrestris* micro-colonies. *Entomologia Experimentalis et Applicata*, **104**, 329–
471 336.
- 472 Gosselin, M., Michez, D., Vanderplanck, M., Roelants, D., Glauser, G. & Rasmont, P. (2013) Does *Aconitum*
473 *septentrionale* chemically protect floral rewards to the advantage of specialist bumblebees? *Ecological*
474 *Entomology*, **38**, 400–407.
- 475 Goulson, D., Hawson, S.A. & Stout, J.C. (1998a) Foraging bumblebees avoid flowers already visited by
476 conspecifics or by other bumblebee species. *Animal Behaviour*, **55**, 199–206.
- 477 Goulson, D., Stout, J.C., Hawson, S.A. & Allen, J.A. (1998b) Floral display size in comfrey, *Symphytum*
478 *officinale* L. (Boraginaceae): Relationships with visitation by three bumblebee species and subsequent
479 seed set. *Oecologia*, **113**, 502–508.
- 480 Goulson, D., Hanley, M.E., Darvill, B., Ellis, J.S. & Knight, M.E. (2005) Causes of rarity in bumblebees.
481 *Biological Conservation*, **122**, 1–8.
- 482 Goulson, D., Lye, G. & Darvill, B. (2008) Decline and conservation of bumble bees. *Annual Review of*
483 *Entomology*, **53**, 191–208.
- 484 Goulson, D., Nicholls, B., Botías, C. & Rotheray, E.L. (2015) Bee declines driven by combined stress from
485 parasites, pesticides, and lack of flowers. *Science*, **347**, 6229.
- 486 Hanley, M.E., Franco, M., Pichon, S., Darvill, B. & Goulson, D. (2008) Breeding system, pollinator choice and
487 variation in pollen quality in British herbaceous plants. *Functional Ecology*, **22**, 592–598.
- 488 Hanley, M.E., Awbi, A.J. & Franco, M. (2014) Going native? Flower use by bumblebees in English urban
489 gardens. *Annals of Botany*, **113**, 799–806.
- 490 Hanley, M.E. & Wilkins, J.P. (2015) On the verge? Preferential use of road-facing hedgerow margins by
491 bumblebees in agro-ecosystems. *Journal of Insect Conservation*, **19**, 67–74.

492 Haughton, A. J., Champion, G. T., Hawes, C., Heard, M. S., Brooks, D. R., Bohan, D. A., Clark, S. J., Dewar,
493 A. M., Firbank, L. G., Osborne, J. L., Perry, J. N., Rothery, P., Roy, D. B., Scott, R. J., Woiwod, I. P.,
494 Birchall, C., Skellern, M. P., Walker, J. H., Baker, P., Browne, E. L., Dewar, A. J. G., Garner, B. H.,
495 Haylock, L. A., Horne, S. L., Mason, N. S., Sands, R. J. N. & Walker, M. J. (2003) Invertebrate responses
496 to the management of genetically modified herbicide-tolerant and conventional spring crops. II. Within-
497 field epigeal and aerial arthropods. *Philosophical transactions of the Royal Society of London. Series B,*
498 *Biological sciences*, **358**, 1847–1862.

499 Heil, M. (2011) Nectar: Generation, regulation and ecological functions. *Trends in Plant Science*, **16**, 191–200.

500 Heinrich, B. (1979a) Resource heterogeneity and patterns of movement in foraging bumblebees. *Oecologia*, **40**,
501 235–245.

502 Heinrich, B. (1979b) “Majoring” and “Minoring” by foraging bumblebees, *Bombus Vagans*: An experimental
503 analysis. *Ecology*, **60**, 245–255.

504 Hendriksma, H.P., Oxman, K.L. & Shafir, S. (2014) Amino acid and carbohydrate tradeoffs by honey bee nectar
505 foragers and their implications for plant-pollinator interactions. *Journal of Insect Physiology*, **69**, 56–64.

506 Howell, A.D. & Alarcón, R. (2007) *Osmia* bees (Hymenoptera: Megachilidae) can detect nectar-rewarding
507 flowers using olfactory cues. *Animal Behaviour*, **74**, 199–205.

508 Hutchings, M.J. & Price, E.A. (1999) *Glechoma hederacea* L. (*Nepeta glechoma* Benth., *N. hederacea* (L.)
509 Trev.). *Ecology*, **87**, 347–364.

510 Jurgens, A. & Dötterl, S. (2004) Chemical composition of anther volatiles in Ranunculaceae: genera-specific
511 profiles in *Anemone*, *Aquilegia*, *Caltha*, *Pulsatilla*, *Ranunculus*, and *Trollius* species. *American Journal of*
512 *Botany*, **91**, 1969–1980.

513 Kessler, A. & Halitschke, R. (2009) Testing the potential for conflicting selection on floral chemical traits by
514 pollinators and herbivores: predictions and case study. *Functional Ecology*, **23**, 901–912.

515 Kim, Y. & Smith, B. (2000) Effect of an amino acid on feeding preferences and learning behavior in the honey
516 bee, *Apis mellifera*. *Journal of Insect Physiology*, **46**, 793–801.

- 517 Kipling, R.P. & Warren, J. (2013) How generalists coexist: the role of floral phenotype and spatial factors in the
518 pollination systems of two *Ranunculus* species. *Journal of Plant Ecology*, **7**, 1–10.
- 519 Kitaoka, T.K. & Nieh, J.C. (2009) Bumble bee pollen foraging regulation: role of pollen quality, storage levels,
520 and odor. *Behavioral Ecology Sociobiology*, **63**, 625–625.
- 521 Klein, A.-M., Vaissière, B.E., Cane, J.H., Steffan-Dewenter, I., Cunningham, S., Kremen, C. & Tscharntke, T.
522 (2007) Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal*
523 *Society B-Biological Sciences*, **274**, 303–313.
- 524 Konzmann, S. & Lunau, K. (2014) Divergent rules for pollen and nectar foraging bumblebees - a laboratory
525 study with artificial flowers offering diluted nectar substitute and pollen surrogate. *PLoS ONE*, **9**, e91900.
- 526 Kremen, C., Williams, N.N. & Thorp, R.W. (2002) Crop pollination from native bees at risk from agricultural
527 intensification. *Proceedings of the National Academy of Sciences of the United States of America*, **99**,
528 16812–16816.
- 529 Lack, A.J. (1982) Competition for pollinations in the ecology of *Centaurea scabiosa* L. and *Centaurea nigra* L.
530 III. Insect visits and the number of successful pollinations. *New Phytologist*, **91**, 321–339.
- 531 Leonhardt, S.D. & Blüthgen, N. (2012) The same, but different: Pollen foraging in honeybee and bumblebee
532 colonies. *Apidologie*, **43**, 449–464.
- 533 Levin, M.D. & Bohart, G.E. (1955) Selection of pollens by honey bees. *American Bee Journal*, **95**, 392–393.
- 534 Mačukanović-Jocić, M., Duletić-Laušević, S. & Jicić, G. (2004) Nectar production in three melliferous species
535 of *Lamiaceae* in natural and experimental conditions. *Acta Veterinaria*, **54**, 475–487.
- 536 Mačukanović-Jocić, M. & Đurđević, L. (2005) Influence of microclimatic conditions on nectar exudation in
537 *Glechoma hirsuta* W.K. *Archives of Biological Sciences*, **57**, 119–126.
- 538 Naimi, B. (2013) *Usdm: Uncertainty analysis for species distribution models* (Version 1.1–12). Available at
539 <http://cran.r-project.org/package=usdm> 01/10/2015
- 540 Nicolson, S.W. (2011) Bee Food: The chemistry and nutritional value of nectar, pollen and mixtures of the two.
541 *African Zoology*, **46**, 197–204.

542 Öckinger, E. & Smith, H.G. (2007) Semi-natural grasslands as population sources for pollinating insects in
543 agricultural landscapes. *Journal of Applied Ecology*, **44**, 50–59.

544 O’Hara, R.B. & Kotze, D.J. (2010) Do not log-transform count data. *Methods in Ecology and Evolution*, **1**, 118–
545 122.

546 Ollerton, J., Erenler, H., Edwards, M. & Crockett, R. (2014) Extinctions of aculeate pollinators in Britain and
547 the role of large-scale agricultural changes. *Science*, **346**, 1360–1362.

548 Pernal, S.F. & Currie, R.W. (2001) The influence of pollen quality on foraging behavior in honeybees (*Apis*
549 *mellifera* L.). *Behavioral Ecology Sociobiology*, **51**, 53–68.

550 Pinheiro, L.A., Torres, L.M., Raimundo, J. & Santos, S.A.P. (2015) Effects of pollen, sugars and honeydew on
551 lifespan and nutrient levels of *Episyrphus balteatus*. *BioControl*, **60**, 47–57.

552 Potts, S.G., Biesmeijer, J.C., Kremen, C., Neumann, P., Schweiger, O. & Kunin W.E. (2010) Global Pollinator
553 Declines; Trends, Impacts and Drivers. *Trends in Ecology and Evolution*, **25**, 345–353.

554 Pyke, G.H. (1978) Optimal foraging: movement patterns of bumblebees between inflorescences. *Theoretical*
555 *Population Biology*, **13**, 72–98.

556 R Development Core Team (2014) R: A language and environment for statistical computing. R Foundation for
557 Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, <http://www.R-project.org>. 01/10/2015

558 Robertson, A.W. Mountjoy, C. Faulkner, B.E. Roberts, V.M. & Macnair M.R. (1999) Bumble bee selection of
559 *Mimulus guttatus* flowers: The effects of pollen quality and reward depletion. *Ecology*, **80**, 2594–2606.

560 Robinson, R.A. & Sutherland, W.J. (2002) Post-war changes in arable farming and biodiversity in Great Britain.
561 *Journal of Applied Ecology*, **39**, 157–176.

562 Roulston, T.H. & Cane, J.H. (2000) Pollen nutritional content and digestibility for animals. *Plant Systematics*
563 *and Evolution*, **222**, 187–209.

564 Roulston, T.H. & Cane, J.H. (2002) The effect of pollen protein concentration on body size in the sweat bee
565 *Lasioglossum zephyrum* (Hymenoptera: Apiformes). *Evolutionary Ecology*, **16**, 49–65.

566 Roulston, T.H., Cane, J.H. & Buchmann, S.L. (2000) What governs protein content of pollen: Pollinator
567 preferences, pollen-pistil interactions, or phylogeny? *Ecological Monographs*, **70**, 617–643.

568 RStudio (2012). *RStudio: Integrated development environment for R* (Version 0.98.1091). Available at
569 <http://www.rstudio.org/>. (Accessed June 11th 2015)

570 Schmidt, J.O. (1982) Pollen foraging preferences of honey bees. *The Southwestern Entomologist*, **7**, 255–259.

571 Sedivy, C., Piskorski, R., Müller, A. & Dorn, S. (2012) Too low to kill: Concentration of the secondary
572 metabolite ranunculin in buttercup pollen does not affect bee larval survival. *Journal of Chemical*
573 *Ecology*, **38**, 996–1002.

574 Silva, E.M. & Dean, B.B. (2004) Patterns of floral nectar production of Onion (*Allium cepa* L.) and the effects
575 of environmental conditions. *Journal of the American Society for Horticultural Science*, **129**, 299–302.

576 Somme, L., Vanderplanck, M., Michez, D., Lombaerde, I., Moerman, R., Wathelet, B., Wattiez, R., Lognay, G.
577 & Jacquemart, A.-L. (2015) Pollen and nectar quality drive the major and minor floral choices of bumble
578 bees. *Apidologie*, **46**, 92–106.

579 Stone, G.N. (1993) Endothermy in the solitary bee *Anthophora plumipes*: Independent measures of
580 thermoregulatory ability, costs of warm-up and the role of body size. *Journal of Experimental Biology*,
581 **174**, 299–320.

582 Strohm, E. & Linsenmair, K.E. (1997) Female size affects provisioning and sex allocation in a digger wasp.
583 *Animal Behaviour*, **54**, 23–34.

584 Tylianakis, J.M., Didham, R.K., Bascompte, J. & Wardle, D. S. (2008). Global change and species interactions
585 in terrestrial ecosystems. *Ecology Letters*, **11**, 1351–1363.

586 Van Damme, E.J. (2008). Plant lectins as part of the plant defense system against insects, in Schaller, A (ed.),
587 *Induced plant resistance to herbivory*, Springer, Netherlands, pp. 285–307.

588 Waser, N.M., Chittka, L., Price, M.V., Williams, N.M. & Ollerton, J. (1996) Generalization in pollination
589 systems, and why it matters. *Ecology*, **77**, 1043–1060.

590 Wickham, H. (2009) *ggplot2: Elegant graphics for data analysis*. Springer-Verlag New York.

- 591 Wille, H., Wille, M., Kilchenmann, V., Imdorf, A. & Bühlmann, G. (1985) Pollenernte und Massenwechsel von
592 drei *Apis mellifera*-Völkern auf demselben Bienenstand in zwei aufeinanderfolgenden Jahren. *Revue*
593 *Suisse de Zoologie*, **92**, 897–914. In: Praz, C., Müller, A. & Dorn, S. (2008) Specialized bees fail to
594 develop on non-host pollen: Do plants chemically protect their pollen? *Ecology*, **89**, 795–804.
- 595 Wolff, D. (2006) Nectar sugar composition and volumes of 47 species of Gentianales from a southern
596 Ecuadorian montane forest. *Annals of Botany*, **97**, 767–77.
- 597 Zuur, A.F., Ieno, E.N. & Elphick, C.S. (2010) A protocol for data exploration to avoid common statistical
598 problems. *Methods in Ecology and Evolution*, **1**, 3–14.
- 599

600 **Table 1** Measurements per inflorescence of nectar (30 replicates) and pollen (10 replicates) quality and quantity (Mean \pm SD) for *Crataegus monogyna*, *Rubus fruticosus*,
601 *Glechoma hederacea*, *Lamium album*, *Ranunculus repens* and *Symphytum officinale* between times and totalled over 24 hours from seven sites in Sussex, UK.

602

Test Plant Species	Nectar (μ l)				Nectar (%)				Pollen (mg)	Pollen (%)	Sugar per Infl. (mg)				Protein per Infl. (mg)
	Morning	Afternoon	Evening	Over 24 Hours	Morning	Afternoon	Evening	Over 24 Hours			Morning	Afternoon	Evening	Over 24 Hours	
<i>Crataegus monogyna</i>	1.1 \pm 1.7	0.4 \pm 0.4	0.4 \pm 0.5	1.9 \pm 2.0	26.13 \pm 2.26	54.95 \pm 3.15	36.54 \pm 1.81	35.03 \pm 9.42	0.11 \pm 0.03	15.39 \pm 2.24	0.6 \pm 0.9	0.6 \pm 0.1	0.3 \pm 0.2	1.2 \pm 1.2	0.02 \pm 0.01
<i>Rubus fruticosus</i>	2.9 \pm 2.8	0.9 \pm 1.2	3.1 \pm 2.1	6.8 \pm 3.7	9.48 \pm 0.52	24.41 \pm 1.87	18.16 \pm 1.26	17.13 \pm 4.72	0.48 \pm 0.23	12.6 \pm 2.28	0.3 \pm 0.2	0.3 \pm 0.3	0.6 \pm 0.5	1.3 \pm 0.7	0.06 \pm 0.02
<i>Glechoma hederacea</i>	1.2 \pm 0.6	0.7 \pm 0.4	0.8 \pm 0.4	2.6 \pm 0.8	29.03 \pm 1.20	30.40 \pm 1.57	29.43 \pm 1.43	29.86 \pm 4.56	0.89 \pm 0.29	19.07 \pm 3.46	0.4 \pm 0.2	0.2 \pm 0.1	0.2 \pm 0.1	0.9 \pm 0.2	0.17 \pm 0.06
<i>Lamium album</i>	1.1 \pm 0.4	0.3 \pm 0.2	0.8 \pm 0.5	2.2 \pm 0.6	19.76 \pm 0.98	32.88 \pm 2.13	27.67 \pm 1.38	25.88 \pm 7.61	1.22 \pm 0.33	4.48 \pm 1.28¹	0.2 \pm 0.1	0.2 \pm 0.1	0.3 \pm 0.2	0.6 \pm 0.3	0.06 \pm 0.02
<i>Ranunculus repens</i>	0.1 \pm 0.1	0.1 \pm 0.1	0.0 \pm 0.0	0.2 \pm 0.2	22.79 \pm 2.25	28.00 \pm NA	29.5 \pm 2.60	24.50 \pm 8.13	0.76 \pm 0.3	12.06 \pm 1.50	0.0 \pm 0.0	0.1 \pm NA	0.0 \pm 0.0	0.1 \pm 0.0	0.09 \pm 0.04
<i>Symphytum officinale</i>	1.8 \pm 0.8	1.3 \pm 0.5	1.2 \pm 0.6	4.3 \pm 1.1	24.07 \pm 1.06	44.59 \pm 0.77	33.25 \pm 1.61	33.88 \pm 4.60	0.95 \pm 0.31	9.71 \pm 1.38	0.5 \pm 0.2	0.7 \pm 0.2	0.5 \pm 0.2	1.7 \pm 0.4	0.09 \pm 0.03

603 ¹This value differed from the previously recorded value of 22.8% in Roulston *et al.* (2000).

604 Bold values indicate the total for nectar measurements over 24 hours, and the total for pollen measurements (for which there was only a single measure per replicate)


605

606

607 **Table 2** Generalized Linear Mixed Models (GLMM) test results investigating the effects of floral abundance, total nectar and pollen quality and interactions between floral
608 abundance and each quality metric on visitation of insect foragers from seven sites in Sussex, UK. Displayed are parameter estimates \pm standard error, z-value and p-value for
609 each explanatory variable in the final model after simplification, and model AIC and log-likelihood tests. P-values are in bold if < 0.05 significance.


Visitation	Explanatory Variables left in model	Parameter Estimate \pm SE	z-value	p-value	AIC	logLik
<i>Apis mellifera</i>	Log flower abundance	-0.066 \pm 0.253	-0.261	0.794	331.2	-159.6
	Total sugar	7.845 \pm 3.288	2.386	0.017		
	Total protein	-2.491 \pm 0.927	-2.687	0.007		
	Log flower abundance \times Total sugar	-1.405 \pm 0.639	-2.201	0.028		
<i>Bombus pratorum</i>	Total sugar	2.838 \pm 0.424	6.687	< 0.001	423.2	-208.6
<i>Bombus pascuorum</i>	Log flower abundance	0.281 \pm 0.249	1.125	0.260	419.5	-203.7
	Total sugar	-1.380 \pm 0.453	-3.048	0.002		
	Total protein	-12.584 \pm 4.158	-2.785	0.005		
	Log flower abundance \times Total protein	2.879 \pm 0.888	3.241	0.001		
All <i>Bombus</i> Abundance	Log flower abundance	0.849 \pm 0.168	5.049	< 0.001	605.5	-297.7
	Total sugar	6.275 \pm 1.889	3.322	< 0.001		
	Log flower abundance \times Total sugar	-1.074 \pm 0.366	-2.932	0.003		
All Insect Abundance	Log flower abundance	0.425 \pm 0.124	3.404	< 0.001	705.7	-347.9
	Total sugar	4.741 \pm 1.522	3.113	0.001		
	Log flower abundance \times Total sugar	-0.776 \pm 0.296	-2.623	0.008		

610




612

613 **Fig. 1** Negative correlation between nectar volume (μl) and nectar sugar concentration (%) averaged between a)
 614 test plant species (Pearson R = -0.49, p = 0.32) and b) test plant species and time period (Pearson R = -0.52, p =
 615 0.02). \blacktriangledown = *Symphytum officinale*, \blacktriangle = *Crataegus monogyna*, \blacksquare = *Glechoma hederacea*, \blacklozenge = *Lamium album*, \bullet =
 616 *Rubus fruticosus* and $+$ = *Ranunculus repens*. Black symbols = Morning (09:00 GMT), dark grey symbols =
 617 Afternoon (15:00) and light grey symbols = Evening (21:00) periods of nectar sampling.




618

619 **Fig. 2** The mean nectar volume (µl) and mean nectar sugar concentration (%) measured from 30 inflorescences
 620 in Morning (09:00 GMT), Afternoon (15:00) and Evening (21:00) time periods averaged from six test plant
 621 species. Significant differences between time periods were identified using post-hoc tests; time periods that do
 622 not share a letter show significant variation ($p < 0.05$), ns = no significance.




623

624 **Fig. 3** Total sugar per inflorescence (mg) of test plant species at Morning (09:00 GMT), Afternoon (15:00) and
 625 Evening (21:00) time periods, with the test statistic ('F' for ANOVA and 'H' for Kruskal-Wallis, respectively)
 626 and significance levels for each analysis of variance between periods of the day for each species. Significant
 627 differences between time periods were identified using post-hoc tests; time periods that do not share a letter
 628 show significant variation ($p < 0.05$), ns = no significance.



629

630 **Fig. 4** Mean insect visits to test plant species in Morning (08:00—10:00 GMT), Afternoon (13:00—15:00) and
 631 Evening (19:00—21:00) time periods, with the test statistic and significance levels of ANOVAs between
 632 periods of the day for each species. Significant differences between time periods were identified using post-hoc
 633 tests; time periods that do not share a letter showed significant variation ($p < 0.05$).



634

635 **Fig. 5** Mean (\pm standard deviation) visits per test plant species to illustrate the positive effect of greater total
 636 sugar on visitation in *Bombus* spp. (above), and total sugar on *Bombus* spp. visits (below) when separated
 637 between sample plots with high or low floral abundance; low floral abundance (10—50 inflorescences per a 4
 638 m², left) shows little or no trend, whilst high floral abundance (51—1000 inflorescences, right) shows a positive
 639 trend. Grey lines and polygons indicate model best fit and 95% Confidence Interval.

640