

Fluctuations and Instability in Sedimentation

Élisabeth Guazzelli¹ and John Hinch²

¹Institut Universitaire des Systèmes Thermiques Industriels, Centre National de la Recherche Scientifique, Aix-Marseille Université, 13453 Marseille Cedex 13, France; email: Elisabeth.Guazzelli@polytech.univ-mrs.fr

²Department of Applied Mathematics and Theoretical Physics, University of Cambridge, Cambridge CB3 0WA, United Kingdom; email: E.J.Hinch@damtp.cam.ac.uk

Annu. Rev. Fluid Mech. 2011. 43:97–116

First published online as a Review in Advance on August 10, 2010

The *Annual Review of Fluid Mechanics* is online at fluid.annualreviews.org

This article's doi:
10.1146/annurev-fluid-122109-160736

Copyright © 2011 by Annual Reviews.
All rights reserved

0066-4189/11/0115-0097\$20.00

Keywords

suspension, low Reynolds number, slender body, diffusion, structural instability

Abstract

This review concentrates on the fluctuations of the velocities of sedimenting spheres, and on the structural instability of a suspension of settling fibers. For many years, theoretical estimates and numerical simulations predicted the fluctuations of the velocities of spheres to increase with the size of the container, whereas experiments found no such variation. Two ideas have increased our understanding. First, the correlation length of the velocity fluctuations was found experimentally to be 20 interparticle separations. Second, in dilute suspensions, a vertical variation in the concentration due to the spreading of the front with the clear fluid can inhibit the velocity fluctuations. In a very dilute regime, a homogeneous suspension of fibers suffers a spontaneous instability in which fast descending fiber-rich columns are separated by rising fiber-sparse columns. In a semidilute regime, the settling is hindered, more so than for spheres.

1. INTRODUCTION

Sedimentation featured in an article in this series by Davis & Acrivos (1985) 25 years ago. That review considered the average settling velocity of spherical particles, which controls the rate of production of sediment and of pure liquid (see **Supplemental Movie 1**; follow the **Supplemental Material link** from the Annual Reviews home page at <http://www.annualreviews.org>). For monodisperse suspensions of spheres, there had been recent theoretical calculations for the dilute limit, a difficult problem because of long-range hydrodynamic interactions. There had also been experimental and theoretical progress on enhanced separation in inclined channels in the Boycott effect. Open questions were identified of whether the interparticle separation played a role in the dilute limit, and the extension to nonspherical particles. The review concluded with a remark that, although the subject had a long history, there were opportunities for future research, a remark well justified by the activity in the past 25 years.

In this review we concentrate on recent progress in understanding fluctuations during the sedimentation of spheres and recent studies of instabilities of suspensions of sedimenting fibers.

2. FLUCTUATIONS IN SETTLING SPHERES

2.1. The Divergence Paradox


Individual particle motions fluctuate about the mean (see **Figure 1**) due to the constantly changing configuration of the suspension microstructure and the resulting long-range multibody hydrodynamic interactions. Consequently, even in a monodisperse suspension with no large differences in Stokes velocities among the particles, there exist distributions of velocities both parallel and perpendicular to gravity. The variance or the standard deviation of the velocity distribution is the simplest measure of the particle velocity fluctuations.

Until recently, there was a paradox in which the variance in the velocities of sedimenting particles was predicted by theory and numerical simulations to depend on the size of the container, whereas experiments saw no such dependency. In this section, we describe the basis of this paradox.

At low Reynolds numbers, the hydrodynamic interactions between particles are long-ranged. The disturbance velocity at a distance r caused by a sphere of radius a sedimenting at V_S is $O(V_S a / r)$,

FALLING CLOUDS

A cloud of particles sedimenting in a viscous fluid evolves into a torus that becomes unstable and breaks up into secondary droplets, which deform into tori themselves in a repeating cascade (Adachi et al. 1978, Machu et al. 2001). This instability occurs even in the complete absence of inertia and without the need to perturb the initial shape (Metzger et al. 2007c). The particles circulate in closed toroidal streamlines as predicted by a continuum approach in which the cloud is modeled as an effective medium of higher density. Fluctuations arising from the multibody character of the hydrodynamic interactions cause particles to depart from these streamlines and to be carried into a downstream tail (Nitsche & Batchelor 1997). Because the lost particles are those located in the circulation rim, this depletes the central region and leads to torus formation. The mechanism responsible for the further expansion of the torus remains unclear, but the breakup can be described as a change in the flow topology that occurs when the torus reaches a critical aspect ratio. Simulations using a point-particle approach containing the minimal physics of the long-range interactions capture this dynamics. Faster breakup is observed for clouds of fibers due to the self-motion of the anisotropic particles (Park et al. 2010) (see **Supplemental Movies 5 and 6**).


Figure 1

Tracking of two marked spheres in the midst of a 30% volume fraction sedimenting suspension of unmarked spheres made optically transparent by matching the index of refraction of the suspending fluid to that of the glass spheres (Nicolai et al. 1995). The particle trajectories are tortuous and exhibit large as well as small loops as the spheres sometimes moved upward against gravity.

where V_S is the Stokes velocity $\frac{2}{9} \frac{(\rho_p - \rho) a^2 g}{\mu}$ (ρ_p is the density of the particle, ρ that of the fluid, μ the viscosity of the fluid, and g the acceleration of gravity). This long-ranged interaction causes problems in theoretical calculations of averaged effects. For the dilute limit, a naive pairwise addition of the disturbance velocities from all the other spheres sedimenting in a spherical domain of radius R would yield a change in the sedimentation velocity of a test sphere of

$$O\left(\int^R V_S \frac{a}{r} n dV\right) = O\left(V_S \phi \left(\frac{R}{a}\right)^2\right),$$

where n is the number density of particles in the suspension, and $\phi = n \frac{4\pi}{3} a^3$ is their volume fraction. This theoretical estimate suggests that the mean settling velocity of the particles is not an intrinsic property of the suspension, but depends on the size, and possibly the shape, of the container. Experimental observations, however, had clearly established the hindered settling result $\langle w \rangle = V_S(1 - k\phi)$, for the mean settling velocity of particles in the dilute limit $\phi \ll 1$, with k a constant ≈ 5 . This is the dilute limit of the Richardson-Zaki law $\langle w \rangle = V_S(1 - \phi)^n$ with $n \approx 5.1$ at low Reynolds numbers.

The paradox for the mean settling velocity was resolved by Batchelor (1972), who noted that there was a multiparticle effect of a backflow of $-5.5 V_S \phi$ at locations where the test sphere could be placed. He split the hydrodynamic interaction into a contribution to the backflow, which gave a divergent integral, and a remainder, which gave a convergent integral. This resulted in his estimate $k = 6.55$ for a uniform distribution of the separation of pairs of particles. An alternative analysis was given by Hinch (1977) in which the leading-order $V_S a/r$ interaction was seen as that of a point force, which could instead be considered as changing the density from that of the suspending liquid to that of the suspension.

Turning now to a calculation of the variance of the fluctuating velocities of the settling particles, a similar naive pairwise addition results in an estimate

$$\langle w^2 \rangle = O\left(V_S^2 \phi \frac{R}{a}\right)$$

(Caffisch & Luke 1985, Tory & Pickard 1986). The question therefore arises again of whether the variance does truly depend on the size of the container or whether it is an intrinsic property that must be calculated by another method.

Hinch (1988) answered this question by considering a blob of size ℓ . This would contain on average $n\ell^3$ particles, but there would be statistical fluctuations of $\sqrt{n\ell^3}$ in the number if the particles were positioned randomly and independently. This fluctuation in the number gives a fluctuation in the weight of the blob of $mg\sqrt{n\ell^3}$, where m is the mass of a particle compensated for buoyancy. Balancing the fluctuation in weight with a Stokes drag on the blob of size ℓ yields a fluctuation in velocity of

$$w' = O\left(\frac{mg\sqrt{n\ell^3}}{6\pi\mu\ell}\right) = O\left(V_S\sqrt{\phi\frac{\ell}{a}}\right). \quad (1)$$

As larger blobs give larger fluctuations in velocity, one would expect to see those corresponding to the largest spherical blob that can be fitted into the container, i.e., limited by the smallest of the height, width, and depth, say L .

The above theoretical estimate is supported by numerical simulations. When particles are positioned randomly and uniformly throughout the container, then the initial value of the variance is found to be proportional to the size of the container, Koch (1994) and Cunha et al. (2002) summing the interaction between point-particles, Ladd (1996, 1997) using a Lattice Boltzmann method, and Bergougnoux et al. (2003) and Mucha et al. (2004) using a truncated Fourier series for point-particles. However, experiments have found that the steady-state value of the variance does not depend on the size of the container (Nicolai & Guazzelli 1995) (see Section 2.2). The paradox for the variance in the velocities of sedimenting particles is therefore that theory and simulations contradict experiments.

This clear disagreement between theory and experiments has become less clear-cut in time with the arrival of new observations that we attempt to explain in Sections 2.2 and 2.3. Segrè et al. (1997) found a dependence on the container size if the container was small, which Brenner (1999) showed followed the theoretical estimate given in Equation 1. The simulations of Bergougnoux et al. (2003), Mucha et al. (2004), and Nguyen & Ladd (2005) found that the velocity fluctuations decay in time from the theoretical estimate. The experiments of Guazzelli (2001), later confirmed by Chehata Gómez et al. (2009), demonstrated that the initial value of the fluctuations does depend on the size of the container and does follow the theoretical estimate. The paradox has now moved to what determines the well-defined steady value of the experimental fluctuations after the decay from the initial value, this steady value normally independent of the size of the container.

Several modifications of the theory have been suggested, aimed at making predictions that do not depend of the size of the container. Koch & Shaqfeh (1991) noted that the long-range hydrodynamic interactions would be screened if each particle was surrounded by sufficient buoyant clear fluid so that the net weight of the volume was precisely equal to that of the suspension in the same volume. However, Ladd (1996) found that there was insufficient volume of clear fluid to make the particle appear neutrally buoyant. Moreover, the volume of clear fluid would have to extend to the interparticle separation $a\phi^{-1/3}$, which would make the suspension look more like a random crystal.

The estimate of the variance in the velocity fluctuations uses the Stokes drag law applied to the blob. Whereas a Reynolds number $Re_p = V_S a / \nu$ based on the radius a is mostly very small in the experiments, a Reynolds number based on the size of the container $Re = Re_p L / a$ may not be small. An estimate of the variance for this case was given by Hinch (1988), $\langle w'^2 \rangle = O(V_S^2 \phi^{2/3} Re_p^{-2/3})$,

which is independent of the size of the container. Koch (1993) and Yin & Koch (2008) have looked at the case of the particle Reynolds number taking moderate values.

A theoretical suggestion by Luke (2000), taken up by Tee et al. (2002), is that a small vertical variation in the concentration of the particles can lead to a density stratification, which can inhibit the velocity fluctuations. We return to this important mechanism in Section 2.3.

Finally we should mention two further ideas. The topic of fluctuations was reviewed from a statistical mechanics perspective by Ramaswamy (2001), using theoretical ideas from Levine et al. (1998). Randomness was characterized in the standard way by a strong diffusion process with white-noise forcing, with sedimentation as a minor linearized perturbation, which it is not. In contrast, Tong & Ackerson (1998) considered particle diffusion small and sedimentation dominant, and drew analogies with turbulent thermal convection. Small structures created next to the bottom heated plate have a size $Ra^{-1/3}$, where Ra is the Rayleigh number, which translates through the analogy into $11a\phi^{-1/3}$; pity there is no analogous heated plate in sedimentation.

2.2. Velocity Fluctuations and Their Correlation Length


The variability of the velocities was recognized in early experiments in the 1970s (see Tory et al. 1992). Later, large velocity fluctuations were measured by tracking marked spheres in an otherwise transparent sedimenting suspension (Ham & Homay 1988, Nicolai et al. 1995) (see **Figure 1**). The measurements were not collected immediately after the initial mixing of the suspension; instead, the suspension was allowed to settle for some time and reach a steady behavior before tracking started. The fluctuations were not seen to be affected by the size of the container in the large containers used (Nicolai & Guazzelli 1995).

Using particle image velocimetry (PIV), Segrè et al. (1997) measured the velocity field of the particles, from which they could construct the spatial correlation function. They found that the horizontal length over which the vertical velocities were correlated was around 20 interparticle separations, $20a\phi^{-1/3}$. This value of the correlation length seems to be a robust result for volume fractions ϕ ranging from 10^{-4} to 0.4 (Guazzelli 2001; Segrè et al. 1997, 2001; Snabre et al. 2009). This correlation length plays a central role in understanding the velocity fluctuations.

When the minimum dimension of the container L was less than $20a\phi^{-1/3}$, Segrè et al. (1997) found that the velocity fluctuations depended on the size of the container and followed the theoretical estimate given in Equation 1 with the size of the blob $\ell = L$. When $L > 20a\phi^{-1/3}$, they found that the steady vertical velocity fluctuations were $2V_S\phi^{1/3}$ independent of L , which corresponds to Equation 1 with $\ell = 20a\phi^{-1/3}$.

These experimental results were confirmed by numerical simulations by Nguyen & Ladd (2004) using a lattice Boltzmann method at $\phi = 0.13$ and $Re_p = 0.1$. With periodic boundary conditions, the velocity fluctuations were constant in time at a value given by Equation 1 with $\ell = L$. However, if impenetrable top and bottom boundaries were used, the velocity fluctuations decayed in time to a steady value, which depended on the size of the container L if $L < 20a\phi^{-1/3}$ and which was independent if $L > 20a\phi^{-1/3}$. The importance of impenetrable top and bottom boundaries had been noted earlier by Ladd (2002) and Cunha et al. (2002), taking up the speculation of Hinch (1988) that convection currents would remove long-wavelength horizontal density fluctuations, leaving the irreducible scale of the interparticle separation.

Whereas the steady-state velocity fluctuations are independent of the size of the container if $L > 20a\phi^{-1/3}$, the early fluctuations do depend on the container. Guazzelli (2001) saw large-scale fluctuations of the size of the width of the container, which dominated the dynamics just after the initial mixing of the suspension (see **Figure 2**). The initial magnitude of the velocities follows Equation 1 with $\ell = L$ (Chehata Gómez et al. 2009). Moreover, the fluctuation magnitudes are


Figure 2

Relaxation of the large-scale fluctuations. The velocity field is from particle-image velocimetry sampling the whole-cell height and width within a laser sheet located in the middle plane, and the concentration profile is from light-attenuation measurements through the suspension (Bergougnoux et al. 2003, Chehata Gómez et al. 2009). The timescale is the Stokes time $t_S = a/V_S$, i.e., the time for an isolated sphere to sediment its radius.

strongly anisotropic with vertical velocities around four times the horizontal velocities. **Figure 2** shows that the initial strong large-scale fluctuations decay in time to weaker small-scale fluctuations. These small-scale fluctuations remain in a steady state until the sedimentation front arrives in the imaging window (see **Figure 3**). This reduction of the initially large fluctuations to a smaller steady value independent of the size of the container, seen initially at low volume fractions ϕ , has been also observed at larger ϕ (Snabre et al. 2009). This behavior is consistent with the above speculation in which heavy blobs fall to the bottom and light blobs rise to the top.

Figure 4 shows a collection of most of the available experimental data for the velocity fluctuations in the steady regime obtained both from particle tracking and from PIV. The velocity fluctuations increase roughly as $\phi^{1/3}$ at low ϕ , although there is some uncertainty in the coefficient. The vertical fluctuations reach a maximum at approximately $\phi = 0.3$, where they are 1.7 times the mean settling speed, and then decrease. The anisotropy between the vertical and horizontal fluctuations, which was four initially, drops to two in the steady state and is even smaller for $\phi > 0.2$.

Similar large velocity fluctuations to those for $0.2 < \phi < 0.4$ have been seen in fluidized beds at low Reynolds number, e.g., by multiple light-scattering (Xue et al. 1992) and by ultrasonic correlation spectroscopy (Cowan et al. 2000). However, although it is often said that a fluidized bed is equivalent to steady sedimentation, the superficial velocity driving the flow in a fluidized




Figure 3

Dominance of the remaining smaller-scale fluctuations until the arrival of the upper sedimentation front. The velocity field is from particle image velocimetry sampling the entire cell cross section for a window covering the lower one-fourth of the cell height (Bergougnoux et al. 2003, Chehata Gómez et al. 2009). The timescale is the Stokes time $t_s = a/V_S$.




Figure 4

Velocity fluctuations normalized by the mean $\langle w \rangle$ in the (a) vertical and (b) horizontal directions versus volume fraction. The dark-blue line is the correlation $3\phi^{1/3}$ and the light-blue line $2\phi^{1/3}$. The red line is the correlation $1.5\phi^{1/3}$ and the orange line $\phi^{1/3}$. Experiments: open diamonds at $Re_p < 10^{-4}$ (Ham & Homsy 1988), filled circles at $Re_p < 10^{-3}$ (Nicolai et al. 1995), light-color filled diamonds at $Re_p \approx 10^{-4}$ (Segrè et al. 1997), dark-color filled diamonds at $Re_p < 10^{-4}$ (Segrè et al. 2001), dark-color filled upside-down triangles at $Re_p < 10^{-4}$ (Chehata Gómez et al. 2009), light-color filled triangles at $Re_p \approx 10^{-3}$ (Snabre et al. 2009). (Insets) Linear plot of relative fluctuations versus volume fraction.

bed vanishes at the walls, and this leads to a convection motion of the particles at low Reynolds numbers.

Although the microstructure of the suspension plays an important role in the behavior, there have been relatively few detailed studies, which often contradict one another, as discussed in Section 2.1 (Koch & Shaqfeh 1991, Ladd 1996). Experiments by Lei et al. (2001) found, developing in time, that large regions had smaller fluctuations in the number of particles than expected for independently randomly positioned particles. Mucha et al. (2004) saw a similar shortfall in numerical simulations. On the other hand, Bergougnoux & Guazzelli (2009) found in experiments that large regions had larger fluctuations than expected, which did not change in time. They also observed small regions devoid of particles and so suggest that the role of the initial mixing on the microstructure should be explored. The structure function $S(\mathbf{k})$ is difficult to access experimentally for non-Brownian particles. In numerical simulations, Nguyen & Ladd (2005) found in the steady state that the horizontal fluctuations in density were suppressed.

2.3. Stratification

The possibility that an unnoticed vertical variation in concentration of the particles might influence experiments was first drawn to our attention by J. Bławdziewicz (private communication, circa 1995). However, the robustness of our experimental results (Nicolai & Guazzelli 1995, Nicolai et al. 1995) to such an uncontrolled variation led us to dismiss the good suggestion. Later Luke (2000) showed how the velocity fluctuations would decay in time in a stably stratified suspension as the density fluctuations moved to their neutral buoyancy level. For that calculation, it was not clear what would create the stratification.

Tee et al. (2002) made two important observations. First, they suggested that the stratification came from the spreading of the interface between the suspension and the clear fluid above. In very dilute suspensions, the interface is diffuse because the self-sharpening caused by hindered settling is negligible. Second, they estimated the small stratification required to influence the velocity fluctuations. For a cube of size L equal to the minimum dimension of the container, they argued that stratification would be important when the variation in the number of particles in the cube due to stratification $L\partial(nL^3)/\partial z$ exceeded the statistical fluctuations $\sqrt{nL^3}$, i.e., when a nondimensional measure of the stratification β exceeded a critical value

$$\beta = -\frac{a}{\phi} \frac{\partial \phi}{\partial z} > \beta_c = \phi^{-1/2} (L/a)^{-5/2}.$$

When the stratification exceeds this very small critical value, the largest blob with statistical fluctuations greater than those from stratification has size

$$\ell = L(\beta_c/\beta)^{2/5},$$

which gives a variance in the velocity fluctuations

$$\langle w'^2 \rangle = O\left(V_S^2 \phi \frac{L}{a} \left(\frac{\beta_c}{\beta}\right)^{2/5}\right).$$

This is independent of the size of the container L . These dominant fluctuations produce a hydrodynamic dispersion with a diffusivity

$$D = w'\ell = kV_S a \phi^{4/5} \left(-a \frac{\partial \phi}{\partial z}\right)^{-3/5},$$

with k a numerical constant.

In a subsequent paper (Mucha & Brenner 2003), numerical simulations were made under different conditions. A local instantaneous diffusivity D was evaluated as the flux of concentration divided by the concentration gradient

$$D(z, t) = - \int_z^\infty \frac{\partial \phi}{\partial t} dz \bigg/ \frac{\partial \phi}{\partial z}.$$

It was found that the diffusivity was not a function of just the concentration ϕ , but collapsed onto a single curve with the above dependence on $\phi^{4/5}(\partial\phi/\partial z)^{-3/5}$, with $k \approx 1$.

The diffusion equation with this nonlinear diffusivity has a similarity solution for the spreading front, with the thickness of the front growing in time as

$$\delta = a\phi^{1/7} (V_{st}/a)^{5/7}.$$

This prediction was tested in experiments by Bergougnoux et al. (2003), who found that the thickness grew linearly in time, mostly because of unavoidable small polydispersity in the size of the spherical particles. Numerical simulations by Bergougnoux et al. (2003) with strictly monodisperse particles could not distinguish between the $t^{5/7}$ power-law growth and a linear growth in time within the range of data available. Numerical simulations in a later paper (Chehata Gómez et al. 2008) favored the linear growth, and experiments presented in the same paper confirmed the linear growth and found how it depended in detail on the polydispersity. It was thought possible that a linear growth might result from heavy blobs falling out the front, leaving it depleted. That said, a universal form was found in both the experiments and the simulations for the concentration profile in the front, reminiscent of the unpublished similarity solution of the diffusion equation with the nonlinear diffusivity.

The velocity fluctuations in the spreading front were studied in numerical simulations by Mucha et al. (2004). Where the stratification was very weak, $\beta < \beta_c$, they found that the variance was proportional to the size of the container. However, where the stratification exceeded the critical value, the variance was reduced by the factor $(\beta_c/\beta)^{2/5}$ given above, although there was considerable scatter in their log-log plot.

To give a cleaner test of the effect of stratification, Tee et al. (2007) set up an experiment and numerical simulations with a controlled uniform stratification in the initial suspension. They found much clearer evidence of the $(\beta_c/\beta)^{2/5}$ reduction of the variance of the velocity fluctuations. The experimental variance was curiously smaller by a factor of 4, corresponding to stratifications 32 times larger. This difference is now thought to result from a gradient in sucrose contributing more to the density stratification than the very dilute concentration of particles.

It seems that the velocity fluctuations are affected by very small vertical gradients in concentration. The small gradients were measured in experiments for the first time by Chehata Gómez et al. (2007) in two illustrative experiments. In one experiment, the weak stratification seemed to control the velocity fluctuations before the arrival of the front. However, in a second experiment, the velocity fluctuations were unaltered while the stratification was negative within the observation window, so if stratification was influencing the velocities, it was not an instantaneous local relation.

The current state of play comes from the more extensive experiments conducted by Chehata Gómez et al. (2009), who also made two sets of numerical simulations, one starting from an initially stratified suspension and the other from a uniform one. Both experiments and simulations found that the initial velocity fluctuations followed the original Cafilisch & Luke scaling (Equation 1) with $\ell = L$. The fluctuations then decayed to a plateau value. In most of the experiments, the plateau value was not related to the local stratification. In the simulations with an initial stratification, the velocity fluctuations decayed by the expected factor of $(\beta_c/\beta)^{1/5}$. In the simulations starting from a uniform suspension, the velocity fluctuations did not decay until the front arrived. Within the

front, the fluctuations were reduced by the stratification factor $(\beta_c/\beta)^{1/5}$ in both experiments and simulations. The data from the simulations were less convincing, possibly due to a delay for the fluctuations to adapt to the changing conditions, and possibly due to the concentration gradient not being constant over the size of the important blobs.

For the level of fluctuations in the plateau before the front arrives, the issue today seems to focus on what is the result of mixing in the experiments. Starting from a concentrated suspension, it is difficult to produce a homogeneous suspension, especially when it is dilute, while avoiding all entrainment of air by the mixer. One would be interested in the initial microstructure, density fluctuations, and possible mean stratification.

2.4. Hydrodynamic Diffusion

We finally turn to the problem of hydrodynamic diffusion, which was the origin of research on the fluctuating motion of the particles because of its importance in mixing, although in recent times the topic has been much neglected. This diffusion process has been coined hydrodynamic diffusion as its origin lies in the hydrodynamic interactions between the particles (Davis 1996). Ham & Homsy (1988) were the first to investigate the nature of the long-time motions of individual spheres in the midst of a sedimenting suspension of like spheres and to relate it to a Fickian diffusion process. Their findings were confirmed and extended by Nicolai et al. (1995) (see **Figure 1**).

After a large number of hydrodynamic interactions, individual particles lose memory of the details of these interactions and undergo a random walk. The long-time diffusive behavior of the fluctuating particle motions was demonstrated by examining an exponential relaxation of the particle velocity autocorrelation functions and the linear growth of the second-order moments of the particle displacements. Interestingly, the measured correlation times were found to be independent of the concentration, with vertical times slightly longer than horizontal times, over a wide range of volume fractions, from 0.05 to 0.4. Self-diffusivities normalized by the measured mean vertical velocity are constant over a substantial range of concentration, from 0.05 to 0.3, and display a strong anisotropy between the vertical and horizontal directions, $D_v^i \approx 8a\langle w \rangle \approx 4D_b^i$, as shown in **Figure 5**. Again hydrodynamic self-diffusivities were seen not to be affected by the size of the container in these experiments collected in the steady state of sedimentation for large containers (Nicolai & Guazzelli 1995). Above a volume fraction of 0.3, there is clearly a different regime with a strong decrease of the self-diffusivities and anisotropy. The basic mechanism for diffusion may be the fluctuating motion of clusters, instead of the fluctuating motion of single particles, a feature that has been observed at high concentrations in numerical simulations (Ladd 1993). This higher-concentration regime deserves more attention.

Only a limited number of numerical simulations have attempted to address hydrodynamic diffusion (Cunha et al. 2002; Koch 1994; Ladd 1993, 1996). This is probably because the predicted divergence of the diffusivity is more severe ($D \sim w^2 \ell \sim \ell^{3/2}$) than that of the velocity fluctuations ($w' \sim \ell^{1/2}$), and the latter has received the most attention; it would be interesting to return to the study of hydrodynamic diffusion. Simulations of hydrodynamic diffusion (and, to a lesser extent, velocity fluctuations) are affected not only by the size of the box but also by its shape. Early dynamic simulations, which include the full hydrodynamic interaction between the particles using a multipole expansion, found an unrealistically large anisotropy in fluctuations, correlation time, and diffusivity (Ladd 1993). This was later attributed to the periodic boundaries imposed in the vertical direction of the cubic box and was found to be remedied by increasing the aspect ratio of the box (Koch 1994). More realistic anisotropy has been found in lattice Boltzmann simulations using an elongated box in the vertical direction (Ladd 1996) or dynamic simulations having an impenetrable lower boundary but still side periodicity (Cunha et al. 2002).




Figure 5

Self-diffusivity [*filled diamonds* at $Re_p < 10^{-4}$ (Ham & Homsey 1988) and *filled circles* at $Re_p < 10^{-3}$ (Nicolai et al. 1995)] in the vertical D_v^S (blue) and horizontal D_h^S (red) directions, normalized by the product of particle radius a and mean velocity $\langle w \rangle$ versus volume fraction.

The self-diffusion discussed above refers to the tracer diffusion of a test particle in the interior of a suspension in which the particle concentration is uniform. But there is another diffusion phenomenon caused by concentration gradients, which acts on a macroscopic scale. This so-called gradient diffusion refers to the drift of particles down a concentration gradient as a result of their fluctuating motion. The broadening of the interface between the sedimenting suspension and the clear fluid above has been attributed partly to this gradient diffusion. The spreading of this front has been measured with different techniques such as light attenuation and magnetic resonance imaging (Davis & Hassen 1988, Lee et al. 1992, Turney et al. 1995a). However, other effects also control the front: polydispersity, which also leads to broadening of the front, but linearly in sedimentation time; hindered settling, which reduces the spreading by a self-sharpening process; and heavy blobs falling out of the front, leaving it depleted. It is thus difficult to deconvolve the simultaneous actions of these effects and to identify the sole effect of gradient diffusion. In particular, even a small polydispersity (causing the front to grow linearly in time) rapidly overcomes the effect of diffusion (causing the front to grow diffusively with the square root of time if the diffusivity is constant). It may make sense then to analyze the early stage of sedimentation if one wants to determine the diffusion coefficient (Lee et al. 1992). When self-sharpening is accounted for (Lee et al. 1992, Martin et al. 1994), the gradient diffusivities are found to be $D^g \approx 20a\langle w \rangle$ for concentration ranging from 0.02 to 0.1 (thus larger than self-diffusivities within the same range), whereas one finds apparent diffusivities $D_{app}^g \approx 8a\langle w \rangle$ if the self-sharpening is ignored (Davis & Hassen 1988, Turney et al. 1995a). However, these calculations should perhaps now be revisited in the light of the ideas emerging in Section 2.3.

3. INSTABILITY IN SETTLING FIBERS

3.1. The Observed Regimes of Sedimentation

The case of nonspherical particles has received much less attention than that of spheres. The fundamental difference is that anisotropic particles orient in flow and drift perpendicular to gravity. Most of the recent focus has been on the sedimentation of suspensions of rod-like particles or

fibers. The sedimentation of disk-like particles or platelets needs further study. The interest in the sedimentation of fibers lies in part in the availability of methods for slender bodies as well as in its importance in industrial applications, such as the fabrication of fiber-reinforced materials and of pulp and paper.

Different regimes can be identified for suspensions of fibers of aspect ratio $A = l/d$ of length l and diameter d . In the dilute regime, $nl^3 < 1$, where n is the number density of particles, each fiber can rotate freely in the large volume β . The semidilute regime is reached for $nl^3 > 1 > nl^2d$. There is a nematic phase transition in which the fibers become aligned at a concentration of $nl^2d \approx 1$, which may correspond to a small volume fraction, $\phi = \pi nld^2/4$, if the aspect ratio of the fiber A is large. Most of the theoretical and numerical studies have been restricted to the dilute regime, whereas both dilute and semidilute regimes have been examined experimentally.

Early experiments on dilute suspensions reported the formation of floc-like inhomogeneities settling at large velocities (Kumar & Ramarao 1991). As the concentration was increased, the number of flocs increased and hindrance occurred. At higher concentrations, a semirigid network of fibers formed, and the settling speed dropped abruptly. Hindered settling was observed in the semidilute regime using magnetic resonance imaging to follow the diffuse interface between the clear fluid and the suspension, but no information was provided concerning the structure of the suspension (Turney et al. 1995b). By measuring the velocity of individual fibers in the dilute regime and the interface speed using optical techniques in the semidilute regime, Anselmet (1989) also reported hindered settling. However, her data are much lower than those of Turney et al. (1995b), with this discrepancy attributed to the small size of her settling vessel. Later, fiber trajectories and orientations were examined in both the dilute and semidilute regimes (Herzhaft et al. 1996, Herzhaft & Guazzelli 1999) by adapting the particle-tracking technique used in the sedimentation of spheres. For all concentrations and aspect ratios studied, the fiber trajectories were found to have large anisotropic fluctuations and a strong alignment in the direction of gravity with occasional flippings. A steady state for the mean settling velocity and the orientation was always achieved.

These experiments identify two distinct regimes. In the dilute regime, visualization of the suspension shows that the fibers clump and form packets separated by sparse regions (see **Figure 6** and **Supplemental Movie 2**). The large velocity gradients between the fast-settling packets and the sparse regions of backflow are considered to be the cause of the flipping motions. In this regime, the mean settling velocity is not hindered and can be larger than the Stokes velocity of





Figure 6

Photograph of an inhomogeneous sedimenting suspension exhibiting packets of fibers for $A = 32$ and $\phi = 0.5\%$ (Herzhaft & Guazzelli 1999).


Figure 7

(a) Mean vertical velocity normalized by the Stokes velocity of a vertical fiber $V_{S\parallel}$ computed using equation 2.1 of Herzhaft & Guazzelli (1999) and (b) velocity fluctuations normalized by the mean in the vertical (blue) and horizontal (red) directions plotted as a function of fiber volume fraction. The solid gray line is the Richardson-Zaki correlation $(1 - \phi)^n$ with $n = 9$. Experiments: $A = 5$ (filled upside-down triangles), $A = 11$ (filled squares), $A = 20$ (filled diamonds), and $A = 32$ (filled circles) at $Re_p \approx 10^{-4}$ (Herzhaft & Guazzelli 1999); $A = 17$ (filled triangles) at $Re_p \approx 10^{-6}$ (Turney et al. 1995b); and $A = 10$ (crosses) at $Re_p \approx 10^{-4}$ (Anselmet 1989). Simulations: $A = 15.6$ (open triangles) (Mackaplow & Shaqfeh 1998), $A = 11$ (open squares), and $A = 32$ (open circles) (Butler & Shaqfeh 2002). (Inset) Close-up of relative velocity fluctuations for low volume fractions.

an isolated vertical fiber $V_{S\parallel}$, as seen in **Figure 7a**. The mean velocity is found to increase at low ϕ , to reach a maximum (more or less pronounced, depending on A) at $\phi \approx 0.005$, and then to decrease with increasing ϕ . No simple scaling has been found to describe this behavior, which is probably a complex function of both ϕ and A . For $\phi > 0.01$, the fibers are still oriented in the direction of gravity, but the mean velocity becomes hindered and depends not on $n\ell^3$ but on the volume fraction ϕ . The hindrance is more severe than in the case of spheres. A Richardson-Zaki law, $(1 - \phi)^n$ with $n \approx 9$, is in good agreement with data having different A 's coming from both fiber-tracking (Herzhaft & Guazzelli 1999) and interface measurements (Turney et al. 1995b) (see **Figure 7a**). In this regime, visualization shows that the fibers form a loosely connected network and the fluid backflow goes up through holes in the network. **Figure 7a** also shows the predictions of a Monte Carlo simulation using a slender-body approximation (Mackaplow & Shaqfeh 1998), which are of the same order of magnitude as the experimental data. But the comparison should be handled with care as this simulation assumes that the suspension is homogeneous and the orientation is isotropic.

The particle velocity fluctuations were found to increase with increasing ϕ across both dilute and semidilute regimes, as shown in **Figure 7b**. These fluctuations are seen to be very strong, reaching values of ≈ 9 times the mean in the vertical direction for $\phi = 0.15$ (see **Figure 7b**). The fluctuation anisotropy is independent of ϕ and A and has a value of ≈ 3 , larger than in the case of spheres. Another important finding is that the orientation distributions, which have a substantial alignment in the direction of gravity, are quite similar whatever ϕ and A .

3.2. The Structural Instability

A linear stability analysis has predicted the instability observed at low concentration (Koch & Shaqfeh 1989). The authors consider alternating tall columns with more and less fibers as the number density of fibers varies horizontally as $n_0 + n_1(t) \cos kx$. At low Reynolds numbers, this

variation in number density drives a vertical flow

$$w_1 = -\frac{n_1 mg}{\mu k^2} \cos kx,$$

where mg is the weight of a fiber adjusted for buoyancy. The shear $\partial w_1/\partial x$ will start to rotate the fibers as

$$\frac{\partial \theta_1}{\partial t} = \frac{\partial w_1}{\partial x}.$$

The small reorientation θ_1 of the anisotropic sedimenting fibers leads to a sideways slip $u_1 = -V_S \theta_1$, with therefore a sideways flux of fibers $n_0 u_1$. A horizontal divergence in this flux changes the initial variation in the number density

$$\frac{\partial}{\partial t} (n_1 \cos kx) + \frac{\partial}{\partial x} (n_0 u_1) = 0.$$

Assembling the above expressions, one finds an instability with a growth rate

$$0 \left(\frac{V_S}{l} (n_0 l^3)^{1/2} \right),$$

where l is the length of the fibers. This explanation simplifies details of the rotation of initially randomly oriented fibers and, as a consequence, has all wavelengths equally unstable. The full analysis of Koch & Shaqfeh (1989) finds that the longest waves are most unstable. The observed instability, however, may depend more on the initial density fluctuations rather than on which grows fastest, particularly because the instability can be expected to become nonlinear by the time the fibers have fallen through their own length.

3.3. Clusters and Streamers

In dilute suspensions at low Reynolds numbers [i.e., having typical particle Reynolds number $Re_p = V_{S\parallel} l/\nu \approx 10^{-4}$ and cell Reynolds number $Re_L = U_{s\parallel} L/\nu = Re_p (L/l) \approx 10^{-2}$], the existence of the instability and enhanced sedimentation has been observed in fiber-tracking experiments (Herzhaft et al. 1996, Herzhaft & Guazzelli 1999). The fibers are observed to orient in the direction of gravity and to form concentrated clusters whose size is of the order of a few fiber lengths and which are evolving objects, constantly dissolving and forming, i.e., capturing and losing fibers. More recent experiments using PIV measurements (Metzger et al. 2005, 2007a) have seen the clusters organize into downward streamers of high density balanced by backflow regions of low density, and the flow structure evolves from long wavelengths to shorter wavelengths (see **Figure 8b** and **Supplemental Movie 3**). Similarly, streamers have been observed when bidisperse suspensions sediment (Davis & Acrivos 1985). Whereas the mean velocity and fiber orientation reach steady states, the flow structure evolves continuously in time without reaching steady state.

The hydrodynamic instability is also seen in zero-Reynolds number numerical simulations of varying degrees of sophistication, ranging from the simplest point-force approximation to the more complex slender-body approximation for the far-field interactions supplemented by a lubrication approximation for the short-range interactions. When using periodic boundary conditions (Butler & Shaqfeh 2002, Gustavsson & Tornberg 2009, Mackaplow & Shaqfeh 1998, Saintillan et al. 2005), the fibers are observed to clump together and to converge into a single dense streamer, which spans the whole height of the periodic box (see **Figure 9**). The dynamics of clusters within the streamer, i.e., their continual dissolution and formation, show strong similarity




Figure 8

Evolving streamers (*a*) in simulation with a bottom bounding wall [time trace of the concentration field over a short interval in the mid-plane of the box, from left to right, $t = 0, 40, 80,$ and 120 (Saintillan et al. 2006b)] and (*b*) in experiments [vertical component of the velocity field extracted from the median plane of the cell by using PIV (Metzger et al. 2007a)]. The timescale is the time for a vertical isolated fiber to sediment half of its length in the experiments and its length in the simulations.




Figure 9

Illustrations of the formation and evolution of clusters in a $4 \times 4 \times 8$ periodic box simulation using slender-body dynamics and comprising 128 fibers (Butler & Shaqfeh 2002). Panel *a* is the initial condition, panel *b* is around the time when the orientation distribution attains steady state, and panels *c* and *d* are snapshots of the clusters at different times. The timescale is the time for an isolated fiber to sediment half of its length.

with experiments. Realistic predictions are obtained by using boxes elongated in the vertical direction. In particular, (a) the mean velocity can reach steady states with values in close agreement with experimental data, and (b) the anisotropy of the fluctuations is ≈ 3 as in experiments, but the predicted values of the fluctuations are lower (see **Figure 7**). It is worth mentioning that all the simulations predict a strong vertical alignment of the fibers although with a degree of anisotropy weaker than in the experiments. A small preferred horizontal alignment seen in the experiments (Herzhaft & Guazzelli 1999, Metzger et al. 2007a) is not captured. A no-flux boundary condition imposed at the bottom of the box has a strong impact on the evolution of the flow structure (Saintillan et al. 2006b). The single large-scale streamer, still seen at initial times, breaks up into smaller streamers in qualitative agreement with the experiments (see **Figure 8** and **Supplemental Movie 4**). The role of the bottom bounding wall may be related to the suppression of the long-wavelength horizontal density fluctuations in suspensions of rigid spheres (see Section 2.2). These simulations capture well the velocity and orientation statistics, but the mean velocity did not achieved a steady state, possibly due to the limited size of the simulation.

Some of the features of the instability are now understood, but the important issue of wavelength selection remains unsolved. The linear stability analysis of Koch & Shaqfeh predicts that the perturbations with the maximum growth rate are those of the largest wavelength. In a bounded system, the container size limits the size of the largest possible wavelength. Indeed, the growth of a single streamer spanning the width of the box is reported in the simulations at short times regardless of the boundary conditions used. However, wavelengths scaling with the container dimension are not observed in experiments at early times (Metzger et al. 2007a). Moreover, the streamer structure evolves slowly from long to short wavelengths without reaching a steady state, and no simple dependence on suspension height, cell cross section, volume fraction, and properties of the fibers and the fluid is observed. The reduction in wavelength is qualitatively reproduced by simulations using a point-particle method and allowing slip but no penetration at the bottom wall (Saintillan et al. 2006b). However, these simulations suggest that wavelength selection is controlled by stratification and hydrodynamic diffusion in a scaling similar to that found for the cutoff length by stratification in the suspension of spheres (Saintillan et al. 2006a), but this mechanism does not account for the wavelength reduction in the experiments (Metzger et al. 2007b).

4. CONCLUDING REMARK AND FUTURE ISSUES

This review has attempted to present the current state of understanding of the sedimentation of non-Brownian particles at low Reynolds numbers. The difficulties that one encounters lie in the long-range nature of the multibody hydrodynamic interactions between particles. The hydrodynamics are determined by the suspension microstructure, i.e., the orientation and relative position of the particles, which is itself determined by the hydrodynamics. This coupling results in a complex collective dynamics, with large fluctuations correlating many thousands of sedimenting spheres and large-scale streamers composed of many sedimenting fibers.

SUMMARY POINTS

1. For a monodisperse suspension of sedimenting spheres, there are large fluctuations in the velocities of the particles, often as large as the mean settling speed, and these fluctuations have a strong anisotropy favoring the vertical direction.

2. The velocity fluctuations have a correlation length of 20 mean interparticle separations, $20a\phi^{1/3}$, a value that seems robust across a wide range of volume fractions. The correlation length is key to understanding whether or not an experimental container is large or not. Few numerical simulations have both sufficient particles and sufficient spatial resolution to be considered large. There is currently no theoretical prediction of the value. Although a few interparticle separations is a natural size, it is difficult to imagine an argument producing the large $20a\phi^{1/3}$, which implies blobs of 8,000 cooperative particles.
3. A weak stratification, i.e., a vertical variation in the concentration of the particles, may be important. Its role in inhibiting velocity fluctuations in the descending front is now well established for very dilute suspensions. In more concentrated suspensions, hindered settling sharpens the front, and so stratification can be expected to be less important.
4. Two distinct regimes of sedimentation have been identified for suspensions of fibers. In the dilute regime, the homogenous suspension is unstable, and the mean sedimentation velocity can exceed the maximum sedimentation velocity of an isolated vertical fiber. In a semidilute regime, the mean velocity is hindered, and the hindrance is more severe than for a suspension of spheres.
5. The hydrodynamic instability that occurs in the dilute regime of sedimentation for suspensions of fibers has been demonstrated by experiments, theory, and simulations. The fibers are observed to orient in the direction of gravity and form clusters that assemble into downward streamers balanced by backflow regions of lower density. This cluster and streamer dynamics are captured well by numerical simulations, but the wavelength selection is not yet resolved.

FUTURE ISSUES

Although the issues discussed in this review are mostly settled, there remain challenges in systems more complex than spheres and fibers in Newtonian fluids.

1. The sedimentation of platelets and deformable particles probably suffers a similar structural instability as fibers (Saintillan et al. 2006b) and should be explored.
2. Sedimentation in a non-Newtonian suspending fluid should be investigated. One of the most spectacular manifestations observed is the formation of particle-rich structures that take the shape of vertical columns in a shear-thinning fluid and that form several conical piles on the bottom (Mora et al. 2005).
3. The influence of finite inertia should be explored. Inertia can induce a qualitative change to the dynamics and structure of a suspension of settling spheres (Hinch 1988; Koch 1993; Yin & Koch 2007, 2008) and to the instability of sedimenting fibers (Kuusela et al. 2003, Shin et al. 2009).

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

We thank L. Bergougnoux, J.E. Butler, B. Metzger, M. Nicolas, D. Saintillan, and P. Snabre for supplying data, figures, and movies. While writing this, E.J.H. was partially supported by award KUK-C1-013-04 made by King Abdullah University of Science and Technology (KAUST).

LITERATURE CITED

- Adachi K, Kiriya S, Koshioka N. 1978. The behaviour of a swarm of particles moving in a viscous fluid. *Chem. Eng. Sci.* 33:115–21
- Anselmet M-C. 1989. *Contribution à l'étude des systèmes fluide particules: suspensions de cylindres, lits fluidisés*. PhD thesis. Univ. Provence
- Batchelor GK. 1972. Sedimentation in a dilute dispersion of spheres. *J. Fluid Mech.* 52:245–68
- Bergougnoux L, Ghicini S, Guazzelli É, Hinch EJ. 2003. Spreading fronts and fluctuations in sedimentation. *Phys. Fluids* 15:1875–87
- Bergougnoux L, Guazzelli É. 2009. Non-Poisson statistics of settling spheres. *Phys. Fluids* 21:091701
- Brenner MP. 1999. Screening mechanisms in sedimentation. *Phys. Fluids* 11:754–72
- Butler JE, Shaqfeh ESG. 2002. Dynamic simulations of the inhomogeneous sedimentation of rigid fibres. *J. Fluid Mech.* 468:205–37
- Caffisch RE, Luke JHC. 1985. Variance in the sedimenting speed of a suspension. *Phys. Fluids* 28:759–60
- Chehata Gómez D, Bergougnoux L, Guazzelli É, Hinch EJ. 2008. Spreading fronts in sedimentation of dilute suspension of spheres. *Phys. Fluids* 20:023302
- Chehata Gómez D, Bergougnoux L, Guazzelli É, Hinch EJ. 2009. Fluctuations and stratification in sedimentation of dilute suspensions of spheres. *Phys. Fluids* 21:093304
- Chehata Gómez D, Bergougnoux L, Hinch EJ, Guazzelli É. 2007. On stratification control of the velocity fluctuations in sedimentation. *Phys. Fluids* 19:098102
- Cowan ML, Page JH, Weitz DA. 2000. Velocity fluctuations in fluidized suspensions probed by ultrasonic correlation spectroscopy. *Phys. Rev. Lett.* 85:453–56
- Cunha FR, Abade GC, Sousa AJ, Hinch EJ. 2002. Modeling and direct simulation of velocity fluctuations and particle-velocity correlations in sedimentation. *J. Fluids Eng.* 124:957–68
- Davis RH. 1996. Hydrodynamic diffusion of suspended particles: a symposium. *J. Fluid Mech.* 310:325–35
- Davis RH, Acrivos A. 1985. Sedimentation of noncolloidal particles at low Reynolds numbers. *Annu. Rev. Fluid Mech.* 17:91–118
- Davis RH, Hassen MA. 1988. Spreading of the interface at the top of a slightly polydisperse sedimenting suspension. *J. Fluid Mech.* 196:107–34; Corrigendum. 1989. *J. Fluid Mech.* 202:598–99
- Guazzelli É. 2001. Evolution of particle-velocity correlations in sedimentation. *Phys. Fluids* 13:1537–40
- Gustavsson K, Tornberg A-K. 2009. Gravity induced sedimentation of slender fibers. *Phys. Fluids* 21:123301
- Ham JM, Homsy GM. 1988. Hindered settling and hydrodynamic dispersion in quiescent sedimenting suspension. *Int. J. Multiphase Flow* 14:533–46
- Herzhaft B, Guazzelli É. 1999. Experimental study of the sedimentation of dilute and semi-dilute suspensions of fibres. *J. Fluid Mech.* 384:133–58
- Herzhaft B, Guazzelli É, Mackaplow MB, Shaqfeh ESG. 1996. Experimental investigation of the sedimentation of a dilute fiber suspension. *Phys. Rev. Lett.* 77:290–93
- Hinch EJ. 1977. An averaged-equation approach to particle interactions in a fluid suspension. *J. Fluid Mech.* 83:695–720
- Hinch EJ. 1988. Sedimentation of small particles. In *Disorder and Mixing*, ed. E. Guyon, J-P Nadal, Y Pomeau, pp. 153–61. Dordrecht: Kluwer Acad.
- Koch DL. 1993. Hydrodynamic diffusion in dilute sedimenting suspensions at moderate Reynolds numbers. *Phys. Fluids A* 5:1141–55
- Koch DL. 1994. Hydrodynamic diffusion in a suspension of sedimenting point particles with periodic boundary conditions. *Phys. Fluids* 6:2894–900
- Koch DL, Shaqfeh ESG. 1989. The instability of a dispersion of sedimenting spheroids. *J. Fluid Mech.* 209:521–42

- Koch DL, Shaqfeh ESG. 1991. Screening mechanisms in sedimenting suspension. *J. Fluid Mech.* 224:275–303
- Kumar P, Ramarao BV. 1991. Enhancement of the sedimentation rates in fibrous suspensions. *Chem. Eng. Commun.* 108:381–401
- Kuusela E, Lahtinen JM, Ala-Nissila T. 2003. Collective effects in settling of spheroids under steady-state sedimentation. *Phys. Rev. Lett.* 90:094502
- Ladd AJC. 1993. Dynamical simulations of sedimenting spheres. *Phys. Fluids A* 5:299–310
- Ladd AJC. 1996. Hydrodynamic screening in sedimenting suspensions of non-Brownian spheres. *Phys. Rev. Lett.* 76:1392–95
- Ladd AJC. 1997. Sedimentation of homogeneous suspensions of non-Brownian spheres. *Phys. Fluids* 9:491–99
- Ladd AJC. 2002. Effect of container walls on the velocity fluctuations of sedimenting spheres. *Phys. Rev. Lett.* 88:048301
- Lee S, Yang Y, Choi C, Lee T. 1992. Combined effect of sedimentation velocity fluctuation and self-sharpening on interface broadening. *Phys. Fluids* 4:2601–606
- Lei X, Ackerson BJ, Tong P. 2001. Settling statistics of hard sphere particles. *Phys. Rev. Lett.* 86:3300–3
- Levine A, Ramaswamy S, Frey E, Bruinsma R. 1998. Screened and unscreened phases in sedimenting suspensions. *Phys. Rev. Lett.* 81:5944–47
- Luke JHC. 2000. Decay of velocity fluctuations in a stably stratified suspension. *Phys. Fluids* 12:1619–21
- Machu G, Meile W, Nitsche LC, Schafinger U. 2001. Coalescence, torus formation and breakup of sedimenting drops: experiments and computer simulations. *J. Fluid Mech.* 447:299–336
- Mackaplow MB, Shaqfeh ESG. 1998. A numerical study of the sedimentation of fibre suspension. *J. Fluid Mech.* 376:149–82
- Martin J, Rakotomalala N, Salin D. 1994. Hydrodynamic dispersion broadening of sedimenting front. *Phys. Fluids* 6:3215–17
- Metzger B, Butler JE, Guazzelli É. 2007a. Experimental investigation of the instability of a sedimenting suspension of fibres. *J. Fluid Mech.* 575:307–32
- Metzger B, Butler JE, Guazzelli É. 2007b. On wavelength selection by stratification in the instability of settling fibers. *Phys. Fluids* 19:098105
- Metzger B, Guazzelli É, Butler JE. 2005. Large-scale streamers in the sedimentation of a dilute fiber suspension. *Phys. Rev. Lett.* 95:164506
- Metzger B, Nicolas M, Guazzelli É. 2007c. Falling clouds of particles in viscous fluids. *J. Fluid Mech.* 580:283–301
- Mora S, Talini L, Allain C. 2005. Structuring sedimentation in a shear-thinning fluid. *Phys. Rev. Lett.* 95:088301
- Mucha PJ, Brenner MP. 2003. Diffusivities and front propagation in sedimentation. *Phys. Fluids* 15:1305–13
- Mucha PJ, Tee S-Y, Weitz DA, Shraiman BI, Brenner MP. 2004. A model for velocity fluctuations in sedimentation. *J. Fluid Mech.* 501:71–104
- Nicolai H, Guazzelli E. 1995. Effect of the vessel size on the hydrodynamic diffusion of sedimenting spheres. *Phys. Fluids* 7:3–5
- Nicolai H, Herzhaft B, Hinch EJ, Oger L, Guazzelli E. 1995. Particle velocity fluctuations and hydrodynamic self-diffusion of sedimenting non-Brownian spheres. *Phys. Fluids* 7:12–23
- Nitsche JM, Batchelor GK. 1997. Break-up of a falling drop containing dispersed particles. *J. Fluid Mech.* 340:161–75
- Nguyen N-Q, Ladd AJC. 2004. Microstructure in a settling suspension of hard spheres. *Phys. Rev. Lett.* E 69:050401
- Nguyen N-Q, Ladd AJC. 2005. Sedimentation of hard-sphere suspensions at low Reynolds number. *J. Fluid Mech.* 525:73–104
- Park J, Metzger B, Guazzelli É, Butler JE. 2010. A cloud of rigid fibres sedimenting in a viscous fluid. *J. Fluid Mech.* 648:351–62
- Ramaswamy S. 2001. Issues in the statistical mechanics of steady sedimentation. *Adv. Phys.* 50:297–341
- Saintillan D, Darve E, Shaqfeh ESG. 2005. A smooth particle-mesh Ewald algorithm for Stokes suspension simulations: the sedimentation of fibers. *Phys. Fluids* 17:033301
- Saintillan D, Shaqfeh ESG, Darve E. 2006a. The effect of stratification on the wavenumber selection in the instability of sedimenting spheroids. *Phys. Fluids* 18:121503

- Saintillan D, Shaqfeh ESG, Darve E. 2006b. The growth of concentration fluctuations in dilute dispersions of orientable and deformable particles under sedimentation. *J. Fluid Mech.* 553:347–88
- Segrè PN, Helbolzheimer E, Chaikin PM. 1997. Long-range correlations in sedimentation. *Phys. Rev. Lett.* 79:2574–77
- Segrè PN, Liu F, Umbanhowar P, Weitz DA. 2001. An effective gravitational temperature for sedimentation. *Nature* 409:594–97
- Shin M, Koch DL, Subramanian G. 2009. Structure and dynamics of dilute suspensions of finite-Reynolds-number settling fibers. *Phys. Fluids* 21:123304
- Snabre P, Pouligny B, Metayer C, Nadal F. 2009. Size segregation and particle velocity fluctuations in settling concentrated suspensions. *Rheol. Acta* 48:855–70
- Tee S-Y, Mucha PJ, Brenner MP, Weitz DA. 2007. Velocity fluctuations of initially stratified sedimenting spheres. *Phys. Fluids* 19:113304
- Tee S-Y, Mucha PJ, Cipelletti L, Manley S, Brenner MP, et al. 2002. Nonuniversal velocity fluctuations of sedimenting particles. *Phys. Rev. Lett.* 89:054501
- Tong P, Ackerson BJ. 1998. Analogies between colloidal sedimentation and turbulent convection at high Prandtl numbers. *Phys. Rev. E* 58:R6931
- Tory EM, Pickard DK. 1986. On the divergence problem in calculating particle velocities in dilute dispersions of identical spheres. *Powder Technol.* 47:39–42
- Tory EM, Kamel MT, Chan Man Fong CF. 1992. Sedimentation is container-size dependent. *Powder Technol.* 73:219–38
- Turney MA, Cheung MK, McCarthy MJ, Powell RL. 1995a. Magnetic resonance imaging study of sedimenting suspensions of noncolloidal spheres. *Phys. Fluids* 7:904–11
- Turney MA, Cheung MK, Powell RL, McCarthy MJ. 1995b. Hindered settling of rod-like particles measured with magnetic resonance imaging. *AIChE J.* 41:251–57
- Xue J-Z, Helbolzheimer E, Rutgers MA, Russel WB, Chaikin PM. 1992. Diffusion, dispersion, and settling of hard spheres. *Phys. Rev. Lett.* 69:1715–18
- Yin X, Koch DL. 2007. Hindered settling velocity and microstructure in suspensions of solid spheres with moderate Reynolds numbers. *Phys. Fluids* 19:093302
- Yin X, Koch DL. 2008. Velocity fluctuations and hydrodynamic diffusion in finite-Reynolds-number sedimenting suspensions. *Phys. Fluids* 20:043305