

Fluorescence and Optical Characteristics of Reduced Flavines and Flavoproteins†

Sandro Ghisla, Vincent Massey,* Jean-Marc Lhoste, and Stephen G. Mayhew‡

ABSTRACT: The fluorescence and absorption properties of a series of reduced flavoproteins have been measured and compared with the properties of suitable model compounds. Contrary to common belief, a number of reduced flavoproteins have been found to exhibit appreciable fluorescence emission with maxima in the range 500–530 nm. In keeping with common observation, the reduced model flavines are devoid of fluorescence in solution at room temperature, but show marked fluorescence emission in the range 476–512 nm at 77°K in rigid glasses. The fluorescence quantum yield of reduced lactate oxidase (emission λ_{max} 507 nm) is increased 4.7 times upon formation of covalent N₅ adducts and the emission maximum is shifted to 476 nm. In the case of the nonfluores-

cent reduced D-amino acid oxidase, formation of covalent adducts leads to appearance of a fluorescence emission maximal at 520 nm. Among the enzymes investigated the absorption spectra of the reduced forms vary markedly. The chromophore of the reduced flavine was found to be very sensitive to variations of the polarity, pH, or viscosity of the solvent and to substitution. An attempt is made to interpret the absorption and fluorescence emission spectra and to correlate the mentioned effects. Thus valid information about the protein environment at the flavine binding site may be obtained. In particular, in the case of fluorescent-reduced enzymes, the vibrational mobility of the reduced flavocoenzyme is restricted by the protein.

The electronic properties of the isoalloxazine chromophore system, which constitutes the redox active moiety of the flavine coenzymes, have been the object of several thorough theoretical and experimental investigations (Sun *et al.*, 1972; Song, 1971, and literature cited therein). These studies have dealt mainly with the oxidized form of the free and protein-bound flavine as well as with model compounds. A distinctive characteristic of oxidized free flavines is their relatively strong fluorescence (quantum yield ~ 0.3 ; Sun *et al.*, 1972), with an emission maximum typically around 520 nm. The energy and intensity of this emission are dependent on solvent polarity and temperature (Koziol, 1969; Sun *et al.*, 1972; Kotaki *et al.*, 1967), on formation of complexes with a variety of molecules (Weber, 1950; Slifkin, 1971), and on the position and properties of substituents (Ghisla and Hemmerich, 1971; Walker *et al.*, 1972; Sun *et al.*, 1972). Similar effects have been observed with riboflavin, FMN, or FAD¹ bound to various apoproteins.

Unfortunately the spectroscopy of reduced flavines has received far less attention than that of the oxidized flavoquinones. A reason for this could be the absence of well-resolved structure in the near-ultraviolet absorption spectrum and of fluorescence at ambient temperature in any solvent. On the other hand, the nonplanarity of the flavine ring in its reduced forms introduces difficulties in theoretical calculations

of the spectroscopic data and leaves some uncertainties in their interpretation. It is commonly assumed that all species of reduced alloxazines and isoalloxazines, both free and protein bound, are nonfluorescent under the usual experimental conditions. One exception previously reported involves isoalloxazines in which the (probably) nonfunctional carbonyl group at position C₄ is reduced by borohydride, but in this case the system retains its redox properties as well as catalytic activity with the enzymes D- and L-amino acid oxidases (Massey *et al.*, 1968), and can therefore be considered to be a modified oxidized flavine. The other reported case is the adduct of lactate oxidase and α -hydroxybutyric acid, the structure of which has not yet been elucidated in detail but which can be assumed to constitute a modified reduced flavine (Walsh *et al.*, 1972).

During the course of our work, we observed that the reduced forms of several flavoproteins exhibit strikingly different absorption spectra. Surprisingly, the reduced form of lactate oxidase was found to have a fairly strong yellow fluorescence in sharp contrast to its oxidized form which is nonfluorescent. These observations prompted a survey of the spectral characteristics of a number of flavoproteins available to us and a more detailed investigation of the absorption and fluorescence spectroscopy of the reduced flavine chromophore. Some of the limitations imposed by the structureless absorption spectra and the absence of fluorescence with free flavines have been overcome in the present work by using appropriately substituted model compounds which were measured in a rigid glassy solvent at 77°K. Under these conditions the absorption spectra appeared far better resolved and the fluorescence could be observed allowing an easier identification of the lowest electronic transition.

Experimental Section

Enzymes. Lactate oxidase was prepared from *Mycobacterium smegmatis* by the method of Lockridge *et al.* (1972). Glucose oxidase was prepared by the method of Swoboda

† From the Department of Biological Chemistry, The University of Michigan Medical School, Ann Arbor, Michigan 48104, and Fondation Curie, Institut du Radium, Section de Biologie, 91405 Orsay, France. Received September 6, 1973. This work was supported by a research grant from the National Institutes of Health (GM11106) to V. M. and a grant from the Schweizerische Naturforschende Gesellschaft to S. G. This work was presented in part at the 50th Anniversary Meeting of the Japanese Biochemical Society, Nagoya, Japan, Sept 1973.

‡ Present address: Laboratorium voor Biochemie, Landbouwhogeschool, Wageningen, The Netherlands.

* Address correspondence to this author at the University of Michigan.

¹ Abbreviations used are: FAD, flavine adenine dinucleotide; FMN, flavine mononucleotide; F_{lox}, oxidized flavine chromophore; F_{red}, reduced flavine chromophore (dihydroflavine).

and Massey (1965). Flavodoxins were isolated from *Peptostreptococcus elsdenii* (Mayhew *et al.*, 1969) and from *Clostridium MP* (Mayhew, 1971). Butyryl coenzyme A dehydrogenase was isolated from *Peptostreptococcus elsdenii* (Engel and Massey, 1971). Shethna flavoprotein was a gift from Dr. Helmut Beinert (Shethna *et al.*, 1966). L-Amino acid oxidase was prepared from *Crotalus adamanteus* venom by the method of Wellner and Meister (1960). D-Amino acid oxidase was prepared from pig kidneys as described by Brumby and Massey (1968). Old yellow enzyme was prepared from brewer's bottom yeast by the method of Matthews and Massey (1969). Oxynitrilase was a gift from Dr. E. Pfeil (Becker and Pfeil, 1966). *p*-Hydroxybenzoate hydroxylase was isolated from *Pseudomonas fluorescens* by the method of Howell *et al.* (1972) and lysine monooxygenase by a modification of the method of Takeda *et al.* (1969). Melilotate hydroxylase was prepared from *Pseudomonas sp* by the method of Strickland and Massey (1973). Thioredoxin reductase from *Escherichia coli* (Zanetti and Williams, 1967) and yeast glutathione reductase (Massey and Williams, 1965) were gifts from Dr. C. H. Williams, Jr. Pig heart lipoyl dehydrogenase was prepared by the method of Massey (1966).

Flavine Models. *N*₃-Carboxymethylumiflavine (lumiflavine 3-acetate) and 3-methyltetraacetylriboflavine were synthesized from lumiflavine and tetraacetylriboflavine according to Hemmerich (1964) and purified by recrystallization.

3-Methyl-4a,5-propano-4a,5-dihydrolumiflavine. To a nitrogen flushed suspension of 0.5 g (1.8 mmol) of 3-methyl-lumiflavine (Hemmerich, 1964) and 2.5 g (25 mmol) of potassium carbonate in a mixture of 20 ml of dimethylformamide and 20 ml of water, 0.9 g (5 mmol) of sodium dithionite was added under stirring. Upon reduction of the lumiflavine (detected by a change in color and partial solubilization of the suspended material), 3.6 g (18 mmol) of 1,3-dibromopropane was added and the mixture stirred at room temperature. Completion of the reaction is indicated by disappearance of the fluorescence and 450-nm absorption of the oxidized lumiflavine in an acidified and aerated sample of the reaction mixture. The reaction mixture was then poured into 100 ml of cold water and the yellow precipitate was washed thoroughly with dilute acetic acid and water and recrystallized from methanol-water: yield, 180 mg (30%) of pure product; mp 188–190°; nuclear magnetic resonance (nmr) (CDCl₃) 6.8 and 6.7 (2 H, C₆ and C₉ protons), 3.65 and 3.25 (6 H, N₃CH₃ and N₁₀CH₃), 2.22 (6 H, C₇H₃ + C₈CH₃), and 4.40–1.60 ppm (6 H, multiplet, protons of propano bridge). The compound is very sensitive to alkaline hydrolysis. *Anal.* Calcd for C₁₇H₂₀N₄O₂ (mol wt 312.37): C, 65.37; H, 6.45; N, 17.94. Found: C, 65.31; H, 6.30; N, 17.55.

3-Methyl-5-ethyl-1,5-dihydrolumiflavine was prepared by reduction of 3-methyl-5-ethylumiflavoquinonium perchlorate (Ghisla *et al.*, 1973) with H₂-Pt in acetonitrile and then diluted with ethanol for spectroscopic measurements.

3-Methyl-4a-ethoxy-5-ethyl-4a,5-dihydrolumiflavine is formed directly upon solution of 3-methyl-5-ethylumiflavoquinonium perchlorate in absolute ethanol. 1,3-Dimethyl-1,5-dihydrolumiflavine was a gift from Dr. F. Müller. FMN and FAD were obtained from Sigma and purified by chromatography on DEAE-cellulose (Massey and Swoboda, 1963). All other chemicals were of reagent grade from commercially available sources.

Methods. Absorption spectra were recorded with a Cary Model 17 or 15 spectrophotometer. Fluorescence emission and excitation spectra were recorded at room temperature with the ratio-recording fluorimeter described by Casola *et*

al. (1966). The temperature was kept constant at 25 ± 0.3° during the measurements. The lifetime of the emission from reduced lactate oxidase was measured with a TRW fluorescence lifetime fluorimeter, using 0.01 M imidazole-HCl buffer (pH 7.0) at ambient temperature. The reduction of the flavine chromophore was performed with the agents listed in Table I under anaerobic conditions in "Thunberg"-type cuvetts. Catalytic reduction of the model flavines was carried out with H₂-Pd or H₂-Pt on charcoal in a vessel separated from the cuvet by a sintered glass filter. Upon completion of reduction, which can be monitored by the disappearance of the fluorescence, the reduced flavine was sucked anaerobically into the cuvet by application of vacuum. The cuvet cells had dimensions of 10 mm in the emission path and of 3 mm in the excitation path for most measurements of flavoproteins.

The measurements at 77°K were carried out essentially as described by Eley *et al.* (1970). The flavine models (~10⁻⁵ M), dissolved in absolute ethanol, were measured in 4-mm o.d. quartz tubes.

The reductions were carried out with Pt- or Pd-H₂ and monitored by disappearance of the absorption and fluorescence of the oxidized species at 295°K. The tubes were then sealed under vacuum. As the extinction coefficients refer to the room temperature spectra they must be corrected for solvent contraction at 77°K.

The flavoenzymes were dissolved in the most suitable buffer systems as given in the literature to yield concentrations of 10⁻⁵–3 × 10⁻⁵ M. Fluorescence intensities relative to *N*₃-carboxymethylumiflavine (5 × 10⁻⁷–4 × 10⁻⁶ M) were measured in the same cell under the same experimental conditions. This reference compound was used because of its stability against photodecomposition. At λ_{exc} 450 nm and λ_{emiss} 520 nm it has a quantum yield of 134% compared to FMN (both compounds are 10⁻⁶ M in 0.05 M phosphate buffer at pH 6 at 26°). The emission intensity of the flavoproteins was measured at the λ_{max} of the emission. The excitation wavelength was chosen at the absorption λ_{max} in the 450-nm region for the oxidized flavoproteins and at the absorption λ_{max} in the 350-nm region for the reduced flavoproteins (*cf.* Table I). The relative intensities of the protein emission spectra are not corrected for possible self-absorption and for quantum yields and are reported for comparative purposes only.

Results and Discussion

Spectroscopy of Reduced Flavines in Solution. The 2e⁻ reduction of F1_{ox} can lead to several forms of F1_{red}, which can be stabilized by appropriate substitution. The chemistry of these different isomeric classes has been reviewed elsewhere (Ghisla *et al.*, 1973). In the present work only the 1,5- and 4a,5-dihydroflavines, which have been recognized as biologically relevant, will be discussed.

Unsubstituted 1,5-Dihydroflavines. The absorption spectrum of nonsubstituted 1,5-dihydroflavines is characterized by a poorly structured line shape in the near-ultraviolet (uv) and visible range which is rather sensitive to the nature of the solvent. In general, in nonpolar and aprotic solvents (at room temperature) a better resolution is observed (Figures 1–3). In ethanol at 77°K, which forms a clear glass, this weakly absorbing region clearly appears to consist of three different electronic transitions having their maximum probability at 404, 340, and 296 nm, respectively (Figure 3). In this rigid solvent a weak fluorescence, the excitation spectrum of which is identical with the absorption spectrum, is observed with a maximum intensity at 495 nm. This fluorescence is the mirror

TABLE 1: Absorption and Fluorescence Characteristics of Flavoproteins.^a

Enzyme	Coenzyme	Oxidized State (F1 _{ox})				Reduced State (1,5-H ₂ F1 _{red})					Reduction Method
		Absorption		Fluorescence		Absorption			Fluorescence		
				λ _{max} (nm)	ε (10 ³ M ⁻¹ cm ⁻¹)				λ _{max} (nm)	Rel Inten (% of FMN)	
		λ _{max} (nm)	ε (10 ³ M ⁻¹ cm ⁻¹)			Spectral Type ^b					
Lactate oxidase	FMN	460 373	11.3	—	—	360	4.8	4	507	7.0	NaBH ₄ , substrate, Na ₂ S ₂ O ₄
Lactate oxidase (red) N ₅ CH ₂ COOH adduct ^c	FMN					365	5.0	4	476	34	CH ₃ CH(OH)COOH- BrCH ₂ COOH
Glucose oxidase	FAD	452 383	14.1 13.3	—	—	355	6.2	4	—	—	NaBH ₄
D-Amino acid oxidase	FAD	455 380	11.3 9.7	530	1.5	415 s 355	2.9 4.1	2n	—	—	Substrate, Na ₂ S ₂ O ₄ , EDTA- <i>hν</i>
D-Amino acid oxidase (red) C _{4a} CH ₂ C ₆ H ₅ adduct ^c	FAD					365 320 s	5.8 7.4	5	520	3.9	C ₆ H ₅ CH ₂ COOH- <i>hν</i>
L-Amino acid oxidase	FAD	460 383	11.3	—	—	450 s 400 s	2.5 4.5	2n	520	3.2	Na ₂ S ₂ O ₄ , substrate
L-Lysine monooxygenase	FAD	462 385	11.2 11.2	—	—	410 s 350	2.8 5.5	2a	512	2.1	Lysine, Na ₂ S ₂ O ₄
Melilotate hydroxylase	FAD	454 384	11.3 10.3	530	1.2	410 350	2.5 5.1	2n	—	—	EDTA- <i>hν</i>
<i>p</i> -Hydroxybenzoate hydroxylase	FAD	450 374	11.3	525	1.2	420 350	2.1 5.1	2n	—	—	EDTA- <i>hν</i>
Oxynitrilase	FAD	460 390	11.2 12.4	—	—	345	5.0	4	500	2.4	Na ₂ S ₂ O ₄
Lipoic acid dehydrogenase	FAD	455 365 355	11.3 9.2 9.1	—	45	360	3.5	4	—	—	Na ₂ S ₂ O ₄
Glutathione reductase	FAD	460 370	11.3 10.4	—	—	<i>d</i>	<i>d</i>		—	—	Na ₂ S ₂ O ₄
Thioredoxin reductase	FAD	480s 457 380	9.3 11.3 11.7	520	15.5	425 320	2.4 9.0	3n,lt	—	—	Na ₂ S ₂ O ₄ , NaBH ₄
Butyryl-CoA dehydrogenase	FAD	448 376	10.6 8.2	530	5.5	370 s	3.4	1a	520	0.8	Na ₂ S ₂ O ₄
Flavodoxin (<i>P. elsdentii</i>)	FMN	447	10.6	—	—	450 s 365	1.7 5.5	3a,lt	530	0.9	Na ₂ S ₂ O ₄
Flavodoxin (<i>Clostridium MP</i>)	FMN	378 447	9.1 10.6	—	—	313 450 s 367	9.5 1.5 5.25	3a, lt	530	0.75	Na ₂ S ₂ O ₄
“Shetna flavoprotein”	FMN	378 372	9.1 9.6	—	—	315 <i>d</i>	8.3 <i>d</i>		—	—	Na ₂ S ₂ O ₄
“Old yellow enzyme”	FMN	462 380	10.6 10.6	530	3.6	340	4.7	4	—	—	Na ₂ S ₂ O ₄

^a For general references concerning the preparation and properties of the listed enzymes and compounds *cf.* the Experimental Section. A dash (—) indicates that no detectable fluorescence was observed; spaces left blank indicate that no information is available or that the form does not exist in the oxidized state. ^b For the classification see the text. ^c These adducts were prepared as will be reported elsewhere (S. Ghisla and V. Massey, manuscript in preparation; *cf.* also legend to Figure 9). The structures were attributed based on the characteristic reactivity of the modified coenzymes obtained after release from the protein (Ghisla *et al.*, 1973). In the case of D-amino acid oxidase the photoreaction of C₆H₅CH₂COO⁻ with the oxidized enzyme appears to yield predominantly C_{4a} substitution with some formation of the N₅ isomer. ^d Full reduction of the enzyme was not achieved experimentally.

image, in a wavenumber scale, of the first transition, which is poorly resolved in polar solvents like water, dimethyl-

formamide, or ethanol at room temperature, but easily identified in the glassy ethanol spectrum. In toluene (Figure 2) a




FIGURE 1: Absorption spectra of FMN in aqueous solution in the oxidized and reduced state and in the neutral (n) and anionic state (a): (---) FMN_{ox} in 0.033 N NaOH; (···) FMN_{ox} at pH 5 in 0.09 M citrate buffer; (—) FMN_{red} at pH 8.5 in 0.1 M pyrophosphate buffer; (- - -) FMN_{red} at pH 5 in 0.09 M citrate buffer. The reductions were achieved by illumination in the presence of EDTA (Frisell *et al.*, 1959).

similar but less pronounced resolution is observed at room temperature, although no fluorescence is detectable.

The flavohydroquinone anion (Figures 1–3), which is formed upon deprotonation at N₁, exhibits similar spectra to the neutral species, but with well-defined different characteristics. The relative extent of the long-wavelength absorption of the neutral and anion species appears to vary with the nature of the solvent. In water or toluene a blue shift is apparent for the anion whereas in ethanol or dimethylformamide the spectrum is insensitive to ionization at room temperature. However, in glassy ethanol the better defined first absorption band appears shifted to the red in the anion, as compared to the second and




FIGURE 2: Absorption spectra of N₃-methyltetraacetylriboflavine in toluene in the oxidized and reduced state: (···) the neutral oxidized flavine for comparison; (—) reduced flavine in the neutral state (n); (- - -) reduced flavine in the presence of 5% tri-*n*-butylamine (a). The reduction was carried out with H₂-Pd as explained in the Experimental Section. (In the presence of a 10 molar excess of benzoic acid the spectrum (—, n) is practically unchanged. Ionization is possibly incomplete even in the presence of 5% tributylamine.)




FIGURE 3: Absorption and fluorescence spectra of reduced tetraacetylriboflavine ($R = \text{CH}_2(\text{CHOAc})_2\text{CH}_2\text{OAc}$) in ethanol at room (295°K) and liquid nitrogen (77°K) temperature. The anion was obtained in the presence of 5% triethylamine. The extinction coefficients refer to the room temperature spectra. They must be corrected for solvent contraction at 77°K. The reduction was carried out with H₂-Pt and the samples were sealed under vacuum in 4-mm o.d. quartz tubes (*cf.* Experimental Section for further details).

third absorption bands which remain constant (Figure 3). This is supported by the 15-nm shift to the red of the low-temperature fluorescence relative to that of the neutral species.

The F1_{red} cation is obtained by protonation of the neutral species at N₅ at low pH and does not absorb above 330 nm (Ghisla *et al.*, 1973). Clearly in this case the conjugation through the central ring is inhibited.

Substituted 1,5-Dihydrolumiflavines. In comparison to the three-banded absorption spectrum of unsubstituted 1,5-dihydroflavines alkyl substitution at N₅ appears to result in a single and structureless absorption band above 300 nm centered around 325–350 nm, depending on the nature of the substituent (Figure 4; Ghisla *et al.*, 1973). N₁ mono- (Figure 4) or N₁,N₅ disubstitution (not shown) results in a similar effect. However, both the low-temperature absorption spectra




FIGURE 4: Absorption and fluorescence spectra of N₅-substituted F1_{red} in ethanol at room (295°K) and liquid nitrogen temperature (77°K) and of N₁-substituted F1_{red} in dioxane at room temperature (*cf.* Figure 3 and Experimental Section for further details).




FIGURE 5: Absorption and fluorescence spectra of 4a,5-disubstituted Fl_{red} in ethanol at room (295°K) and liquid nitrogen temperature (77°K) (cf. Figure 3 and Experimental Section for further details).

and the relatively strong fluorescence observed in ethanol at 77°K for the N_5 substituted compounds indicate that this absorption band consists of two unresolved transitions which must correspond to the electronic transitions of the unsubstituted species, with the first transition being blue shifted upon alkylation. (A similar blue shift of the third transition is also observed.) Furthermore, N_5 but not N_1 alkylation appears to intensify the first transition. Thus, this transition contributes to most of the absorption band of N_5 -substituted compounds, with a maximum around 350 nm (cf. Figure 4).

Substituted 4a,5-Dihydrolumiflavines. This isomeric form of Fl_{red} is not the "normal" product of reduction of Fl_{ox} , but its derivatives are possible intermediates during flavoprotein catalysis (Hemmerich *et al.*, 1967; Hamilton, 1971; Massey *et al.*, 1969; Walsh *et al.*, 1971; Main *et al.*, 1972). The line shape of its absorption spectrum is similar to that of the N_1 , N_5 -dihydroflavines, but is even more sensitive to the nature of the substitution (Figures 5 and 6). Very characteristic for this class of compounds is the presence of three absorption bands in the 280–360-nm region, the intensity of which varies considerably depending on the nature of the substituents (Walker *et al.*, 1970; Ghisla *et al.*, 1973). In the case of C_{4a} -ethoxy- N_5 -ethylflavines these bands are well resolved and can clearly be attributed to three different electronic transitions. In ethanol the lowest energy absorption band is very sensitive to temperature. Although its long-wavelength edge does not shift upon cooling, it rises more steeply and its maximum appears red shifted in the glass at 77°K. Furthermore, the nonfluorescing compound (at room temperature) exhibits an intense fluorescence at 77°K, the emission spectrum of which is a better image of the low temperature than of the room temperature absorption spectrum. This suggests that a large change in geometry occurs during the lifetime of the fluorescing state at room temperature, which is prevented in the rigid solvent at low temperature thus resulting in a larger overlap of the absorption and fluorescence spectra. In order to better understand this phenomenon, a C_{4a} , N_5 -propano Fl_{red} was synthesized (Figure 6) in which the "bridge" blocks the pyramidal N_5 inversion and is expected to induce a much more rigid structure even in fluid solvents. In fact, this flavine was found to be slightly fluorescent at room temperature in ethanol and very strongly fluorescent at 77°K. On the other hand, it ex-




FIGURE 6: Absorption and fluorescence spectra of 4a,5-bridged Fl_{red} under the same conditions as described in Figures 3 and 5.

hibits a less-resolved absorption spectrum in the 300-nm region and a shift of the first band as compared to C_{4a} -ethoxy- N_5 -ethyl Fl_{red} (Figure 5) at ambient temperature, but the same maximum at ~ 380 nm as the latter model at 77°K. However, with similar compounds in which the steric and electronic properties of the C_{4a} and/or N_5 substituent are varied systematically a continuity in the absorption spectra of this series can be observed (Walker *et al.*, 1970; Ghisla *et al.*, 1973). Probably an interplay of steric interactions from the vicinal C_{4a} and N_5 substituents (which would affect the planarity of the chromophore) and of electronic effects is responsible for the variations of the intensity and the shifts of these bands.

A peculiarity of reduced flavines is their ability to flex along their N_{10} – N_5 axis to form a butterfly-shaped molecule (Kierkegaard *et al.*, 1971). In the normal reduced flavine the angle so formed between the two resulting planes in the molecule is probably small; with N_1 - and N_5 -substituted reduced flavine models this angle has been shown by X-ray crystallography to be considerably larger (20–35°) (Kierkegaard *et al.*, 1971). In connection with this, it has been pointed out that the degree of bending in these molecules is typically reflected by their absorption spectra (Hemmerich *et al.*, 1971), where decrease of the 400–450- and the 290–300-nm absorption together with the appearance of a pronounced maximum in the 320–355-nm region reflects a larger angle between the two planes of the molecule.

From the present work it appears that the low-energy absorption spectrum of reduced flavines consists of at least three transitions which may be assumed to correspond to $\pi \rightarrow \pi^*$ electronic excitations. These transitions, in particular the first one, are moderately forbidden, either because they are polarized in plane and along the short axis of the molecule or, alternatively, if they are long-axis polarized, because the folding of the flavine ring diminishes their probability. The planarity of the system may also affect the energy of the transitions. In the more planar compounds this should result in a red shift of the whole spectrum including the fluorescence. However, the substitution effects described above suggest that the first transition, also observed in the fluorescence, should be short-axis polarized, corresponding mostly to charge transfer upon excitation from the electron-rich C_{4a} – N_5 region to the rest of the molecule. Then, as indicated by the results described




FIGURE 7: Absorption, fluorescence emission, and fluorescence excitation spectra of flavodoxin from *P. elsdenii*, 2×10^{-5} M, in 0.1 M pyrophosphate buffer (pH 8.5) at 25°: (---) absorption of the oxidized enzyme; (—) absorption spectrum of the enzyme reduced with 1 equiv of dithionite in the apparatus described by Foust *et al.* (1969); (---) fluorescence emission (λ_{exc} 368 nm); (-·-) fluorescence excitation spectrum (λ_{emiss} 530 nm) of the reduced enzyme. The oxidized enzyme is devoid of fluorescence.

above, the substitution by electron donating groups at one of these atoms should increase the intensity of the transition.

On the other hand, it appears that the absence of fluorescence for most reduced flavines in fluid solvents and its relative intensity in glassy solvents at 77°K, $N_1-H, N_5-H < N_1-H, N_5-R \ll C_{4a}-R, N_5-R$, is also related to steric factors with the most efficient radiationless decay processes in the less rigid structures. Thus, both the rigidity of the molecule and that of its environment appear to govern the fluorescence yield. The mechanism of fluorescence quenching in the reduced flavines is probably quite different from that observed with $F1_{ox}$ and its anion and cation, the latter being also fluorescent only in rigid media, where the dynamic proton quenching is minimized (Eley *et al.*, 1970).

As a consequence of its peculiar configuration the free reduced flavine undergoes facile (vibrational) inversion processes of which the kinetic and thermodynamic parameters have been investigated (Tauscher *et al.*, 1973). These vibration processes thus probably play an important role in the radiationless decay of the reduced flavine excited states. However, some of the inversion frequencies can be rather slow even at room temperature (Tauscher *et al.*, 1973), as compared to the lifetime of the first fluorescent excited state. It is probable that a "full inversion" of the molecule is not required for radiationless dissipation of energy.

Absorption Spectra of Reduced Flavoproteins. Table I represents a survey of the spectral and fluorescence properties of reduced flavoproteins and similar data for the oxidized flavoproteins for comparative purposes. The spectra of the different reduced flavoenzymes can all reasonably well be simulated by the spectra of the models discussed above, under appropriate conditions. Therefore the spectra of the reduced enzymes in Table I have been classified according to their closest similarity to one of the model systems displayed in Figures 1–5 in their neutral (n) or anionic (a) form at ambient temperature or at 77°K (lt) (low temperature).

As a general trend it is found that the absorption maxima of the protein-bound reduced coenzymes are slightly shifted to longer wavelengths (5–25 nm, *cf.* Table I) indicating a more planar structure as compared to the corresponding free reduced flavines, and that the resolution of the bands is better. Likewise, a better resolution has been documented with oxidized flavoproteins (Palmer and Massey, 1968).

A major effect on the spectra of the flavine molecule is induced by changes in the ionization state as shown in Figures 1, 2, and 3. Free flavines such as FMN and FAD exhibit pK values between their neutral and anionic forms at ~ 10 (Walaas and Walaas, 1956) for the oxidized state, 8.5 for the semiquinone state (Ehrenberg *et al.*, 1967), and ~ 6.5 for the fully reduced state (Lowe and Clark, 1956), which are reflected in their electronic spectra (Figure 1). For flavoproteins on the other hand, in most cases where information is available, no ionization has been observed over the limited pH region in which such enzymes are stable. Some exceptions are known; for example, in the oxidized state D-amino acid oxidase has a pK of 9.5 (Massey and Ganther, 1965) and glycolate oxidase a still lower pK of 8.0 (Schuman and Massey, 1971). In the semiquinoid form, glucose oxidase (Massey and Palmer, 1966) and lysine monooxygenase (M. Flashner and V. Massey, unpublished results) are the only known cases to exhibit a pK close to that of the free coenzymes.

In the fully reduced state, only the flavodoxins have been examined in any detail, and in three such proteins a pK in the region 5.8–6.7 has been determined from the effects of pH on the redox potentials (Mayhew, 1971). This pK corresponds closely with a pK of the fully reduced FMN; however, changes of pH in the range 8–5.4 have very little effect on the spectra. The reduced form of the flavodoxin from *P. elsdenii*, a typical example of this class, has an absorption with a distinct shoulder centered at 450 nm and two peaks in the near-ultraviolet, at 368 and 313 nm (Figure 7). Comparison with the models in Figures 2 and 3 and the fact that the fluorescence is shifted on these proteins (~ 500 –510 \rightarrow 530 nm) (Table I) by a similar amount as upon formation of the anion in the models (495 \rightarrow 510 nm) (Figure 3) suggest that in reduced flavodoxins the flavine is in its anionic form and has a rather planar conformation. A similar conclusion has been reached for the Shetna flavoprotein by Edmonson and Tollin (1971) on the basis of the pH dependence of reduction.

Within the reduced flavoproteins studied, many have absorption spectra with features similar to those of the neutral 1,5-dihydroflavine chromophore in an aprotic, unpolar solvent (*cf.* Figure 2 and Table I). It should be pointed out that with most of these, *e.g.*, *p*-hydroxybenzoate hydroxylase, melilotate hydroxylase, and D-amino acid oxidase, the shortest wavelength band is not well resolved from the uv absorption of the protein itself as it is in the case of flavodoxins. Of the other reduced flavoproteins examined, thioredoxin reductase is the only one which appears to have in its reduced form a spectrum which corresponds to that of the neutral 1,5-dihydroflavine chromophore in rigid medium, *i.e.*, it has a two-banded spectrum with absorption maxima at 425 and 320 nm, both peaks shifted about 20 nm to longer wavelengths than those seen with $F1_{red}$ at 77°K (*cf.* Figure 3). In this case, the ~ 350 -nm transition is probably hidden under the more intense third transition.

A distinctly different type of spectrum is exhibited by reduced lactate oxidase (Figure 8). Comparison with Figure 4 shows that this enzyme has spectral characteristics similar to the substituted models, *i.e.*, it has very little absorbance in the 400–450-nm region in contrast to normal reduced flavine

(Figures 1–3) and has but a single absorbance peak maximal at 360 nm. Furthermore, it differs significantly from the anion form of $F1_{red}$ by lacking apparently completely the near-ultraviolet band in the region of 300 nm. These properties indicate that in this case the first and third transitions are probably blue shifted, as observed with the nonplanar models (*cf.* Figure 4). Several reduced flavoproteins which have been characterized as 4 in Table I show essentially the same spectral features as lactate oxidase, *viz.* old yellow enzyme, glucose oxidase, and lipoyl dehydrogenase, but with the latter two enzymes the blue shift of the first transition is not so pronounced.

Fluorescence Properties of Reduced Flavoproteins. The emission of reduced lactate oxidase has been shown to be fluorescence by the observation that the emission lifetime is <20 nsec. In view of the similarity of this emission with that observed with other reduced flavoproteins (*cf.* below) and with model compounds (*cf.* above), it is reasonable to assume that the emission has the same origin in all cases. A systematic study of the emission lifetimes of these reduced flavoproteins and compounds is planned.

Figure 8 shows the absorption and fluorescence spectra of lactate oxidase in its oxidized and reduced states as a typical example. While the oxidized enzyme is devoid of visible fluorescence, the reduced enzyme displays a relatively strong fluorescence emission, maximal at 507 nm, with an intensity at its maximum 7% that of free FMN (*cf.* Table I). That the emission observed from reduced lactate oxidase originates from the reduced flavine chromophore is shown by the good correspondence of the fluorescence excitation spectrum with that of the absorption spectrum (Figure 8). Addition of pyruvate under anaerobic conditions to reduced lactate oxidase results in the appearance of a long-wavelength absorption band, which has been attributed to formation of a charge-transfer complex between reduced enzyme and pyruvate (Lockridge *et al.*, 1972). The dissociation constant for this complex has been estimated as 2×10^{-3} M from spectral titration studies (Lockridge *et al.*, 1972). It is found that the addition of pyruvate results in a quenching of the fluorescence of the reduced enzyme, with almost complete quenching at very high pyruvate concentrations. From such fluorescence quenching experiments a K_d of 1.5×10^{-3} M was obtained, in reasonable agreement with the value obtained from absorption spectrum changes. A similar quenching was observed on addition of lysine or ornithine to reduced lysine monooxygenase (M. S. Flashner and V. Massey, manuscript in preparation). These results support the interpretation of the long-wavelength species being due to a molecular complex of the charge-transfer type; quenching of fluorescence in (charge-transfer) complexes in general and in complexes formed by oxidized flavine with a variety of molecules has been reported previously (Weber, 1950; Gibson *et al.*, 1962).

While the majority of the flavoproteins tested did not show detectable fluorescence in their (unmodified) reduced state, in accord with the previous literature, a number did show significant emission. The maximum as well as the intensity of this fluorescence emission were found to vary considerably from flavoprotein to flavoprotein. The two most typical examples are represented by the reduced lactate oxidase (Figure 8) and by the reduced flavodoxins, *e.g.*, from *P. elsdenii* (Figure 7) where the fluorescence emission maximum, while of low intensity, is shifted to 530 nm. That the observed fluorescence was not due to incomplete reduction (*i.e.*, that it was not due to oxidized flavine) was shown in all cases by the close correspondence of the fluorescence excitation spectrum with the




FIGURE 8: Absorption, fluorescence emission, and fluorescence excitation spectra of lactate oxidase, 3×10^{-5} M in 0.01 M imidazole-HCl buffer (pH 7.0), at 25° : (---) absorption spectrum of the oxidized enzyme; (—) absorption of the enzyme reduced with L-lactate, after the decay of the initially formed, nonfluorescent complex (*cf.* text for explanations); (---) fluorescence emission (λ_{exc} 360 nm); (· · ·) fluorescence excitation spectrum (λ_{emiss} 507 nm) of the reduced enzyme. The oxidized enzyme does not show detectable fluorescence.

absorption spectrum of the *reduced* enzyme. Furthermore, the spectra obtained were generally independent of the reducing agent used, *e.g.*, substrate, dithionite, EDTA-light, or borohydride. An interesting relation is seen wherein those enzymes which exhibit fluorescence in the reduced state are for the most part nonfluorescent in the oxidized state and *vice versa*.

As the occurrence of fluorescence in many reduced flavoproteins cannot be correlated with known properties of the enzymes it is probable that it is affected by several factors which play an interdependent role. The fluorescence yield of $F1_{ox}$ has been shown to be quenched in model systems upon formation of complexes with various heteroaromatic, mostly electron-rich molecules as, *e.g.*, tyrosine or tryptophan (Weber, 1950; Slifkin, 1971; Tollin, 1968). An analogous quenching is reported in this work for the complex formed by the reduced flavine of lactate oxidase and pyruvate. A strong quenching of the fluorescence has also been observed when sulfur-containing residues are covalently linked to the flavine nucleus (Ghisla and Hemmerich, 1971; Müller and Hemmerich, 1966). It is therefore reasonable to assume that the occurrence of fluorescence emission in (oxidized as well as reduced) flavoproteins is conditioned by the presence of specific amino acids close to the flavine binding site and also to the presence of specific (quenching) hydrogen bridges between protein and coenzyme. In agreement with this assumption a tryptophan and a methionine group in flavodoxin from *Clostridium MP* (Anderson *et al.*, 1972; R. B. Burnett, G. D. Darling, D. L. Kendall, M. E. Lequesne, S. G. Mayhew, and M. L. Ludwig, private communication) and a tyrosine in flavodoxin from *Desulfovibrio vulgaris* (Watenpaugh *et al.*, 1972) have been shown to be located in Van der Waals contact with the coenzyme by X-ray crystallography. It should be pointed out, however, that, due to the "opposite" chemical and electronic properties of $F1_{ox}$ (electron deficient, acceptor) and $F1_{red}$ (electron rich, donor), different interactions should be expected at the two oxidation levels. Thus, compounds which quench the $F1_{ox}$ fluorescence by formation of complexes possibly do not affect the quantum yield of $F1_{red}$ and *vice versa*.

On the other hand the results obtained in the model system and in particular at 77°K clearly suggest that the main factor leading to appearance of fluorescence in reduced flavoproteins




FIGURE 9: Absorption and fluorescence spectra of lactate oxidase and of its reduced N_3CH_2COOH derivative; 8×10^{-5} M in 0.01 M imidazole-HCl buffer (pH 7), at 25° : (---) absorption spectrum of the oxidized enzyme; (···) absorption spectrum of the enzyme reduced with lactate (*cf.* Figure 8) after addition of 10^{-2} M $BrCH_2COOH$; (—) absorption and (---) fluorescence emission of the covalent N_3CH_2COOH adduct. This adduct was formed by irradiation of the reduced enzyme in the presence of bromoacetate with light in a way similar to that described by Walker *et al.* (1970).

is the rigidity introduced by the protein frame. Thus, it is reasonable to conclude that such a "freezing" of the vibrational inversion processes mentioned above occurs at least in those flavoproteins which show a fluorescence in the reduced state (*cf.* Table I). In many flavoproteins, as, *e.g.*, thioredoxin reductase (Table I; Zanetti and Williams, 1967) which have spectra indicating extensive restriction of mobility, a fluorescence emission should be expected. Probably in such cases of potentially fluorescent reduced flavoproteins the emission is quenched by other factors, *e.g.*, interactions with specific protein residues as mentioned above or by dynamic quenching through molecules of the solvent (H^+) which may retain free access to the coenzyme binding site.

Another line of evidence which provides support for the interpretation made above is the fact that substitution in the reduced flavoproteins leads to a strong increase in the fluorescence emission intensity in a similar way as observed with the models. The most illustrative example is provided by reduced lactate oxidase. When bromoacetate is added, the absorption maximum of the normal reduced enzyme (*cf.* Figure 4) is blue shifted to 353 nm (Figure 9) and the fluorescence quenched by 65%. Upon formation of the covalent N_3CH_2COOH adduct (as described in the legend to Figure 9) the fluorescence emission maximum is blue shifted to 476 nm and increased 4.7-fold in intensity compared to the free reduced enzyme, and thus has a quantum yield of $\sim 85\%$ that of free FMN. Furthermore, the spectral effects induced by this substitution are entirely consistent with the changes observed upon immobilization of the N_3 -substituted model at $77^\circ K$ (Figure 4); they consist of a red shift and intensification of the ~ 350 -nm band and a marked decrease of the absorption in the 390–450-nm region. As with the model this would be attributed to a blue shift and intensification of the first transition also apparent in the blue shift of the fluorescence.

Similar fluorescence emissions have been obtained with other reduced covalent derivatives of lactate oxidase. Thus, the inactivated enzyme obtained in the "suicide" reaction of

this enzyme with an acetylenic substrate, 2-hydroxy-3-butyric acid (Walsh *et al.*, 1972), exhibits a fluorescence with an intensity 26% that of free FMN. An equally significant situation applies with D-amino acid oxidase. In the reduced form this enzyme does not display detectable fluorescence. However, on introduction of the bulky benzyl residue at position 4a or N_3 by photoalkylation with phenyl acetate by the method of Walker *et al.* (1970), fluorescence is displayed (*cf.* Table I). Preliminary results with photoalkylation of other flavoproteins indicate that this may be a fairly general phenomenon.

Conclusions

From the comparisons and studies described above it may be concluded that the polarity of the protein environment at the flavine binding site and the interactions between prosthetic group and protein vary considerably within different classes of flavoproteins. Specific hydrogen bridges and the constraint induced by the protein structure affecting in particular the planarity of the reduced coenzyme could explain the observed differences in the absorption spectra. In some cases, in particular with reduced lactate oxidase, the spectral and fluorescence characteristics discussed above indicate that the reduced coenzyme is probably immobilized in a nonplanar form. In other cases, *e.g.*, with thioredoxin reductase, it appears that the reduced coenzyme shows similar characteristics as more planar models in a rigid environment.

Finally we wish to emphasize, that contrary to commonly held beliefs, the reduced flavine chromophore is intrinsically fluorescent. In the case of free reduced flavines in solution, this fluorescence is not displayed, presumably because of energy dissipation to the environment, *e.g.*, through the vibration processes described by Tauscher *et al.* (1973) and discussed above. This concept is in full accord with the reappearance of fluorescence emission from reduced flavines in rigid media at low temperature. In the case of reduced fluorescent flavoproteins the constraint introduced by the protein may hinder the same vibration processes thus reducing the radiationless decay to the extent that fluorescence emission reappears at room temperature.

Of particular interest is the apparent effect of ring substitution on the quantum yield of fluorescence. In view of the emphasis in the current literature on the possibility of covalent intermediates in flavoprotein catalysis, this work suggests that observation of fluorescence characteristics in partial reactions of flavoenzymes with their substrates may yield valuable information on the validity of these hypotheses.

Acknowledgment

We wish to thank Dr. M. S. Flashner for the data on lysine monooxygenase and for many helpful discussions and to Dr. J. D. Shore for help in the measurement of the fluorescence lifetime.

References

- Anderson, R. D., Apgar, P. A., Burnett, R. M., Darling, G. C., LeQuesne, M. E., Mayhew, S. G., and Ludwig, M. L. (1972), *Proc. Nat. Acad. Sci. U. S.* 69, 3189.
- Becker, W., and Pfeil, E. (1966), *Biochem. Z.* 346, 301.
- Brumby, P. E., and Massey, V. (1968), *Biochem. Prep.* 12, 29.
- Casola, L., Brumby, P. E., and Massey, V. (1966), *J. Biol. Chem.* 241, 4977.
- Edmonson, D., and Tollin, G. (1971), *Biochemistry* 10, 133.

- Ehrenberg, A., Müller, F., and Hemmerich, P. (1967), *Eur. J. Biochem.* 21, 286.
- Eley, M., Lhoste, J. M., Lee, C. Y., Cormier, M. J., and Hemmerich, P. (1970), *Biochemistry* 9, 2902.
- Engel, P. C., and Massey, V. (1971), *Biochem. J.* 125, 879.
- Foust, G. P., Burleigh, B. D., Mayhew, S. G., Williams, C. H., Jr., and Massey, V. (1969), *Anal. Biochem.* 27, 530.
- Frisell, W. R., Chung, C. W., and Mackenzie, C. G. (1959), *J. Biol. Chem.* 234, 1297.
- Ghisla, S., Hartmann, U., Hemmerich, P., and Müller, F. (1973), *Justus Leibigs Ann. Chem.*, 1388.
- Ghisla, S., and Hemmerich, P. (1971), *FEBS (Fed. Eur. Biochem. Soc.) Lett.* 16, 229.
- Gibson, Q. H., Massey, V., and Atherton, N. M. (1962), *Biochem. J.* 85, 369.
- Hamilton, G. A. (1971), *Progr. Bioorg. Chem.* 1, 83.
- Hemmerich, P. (1964), *Helv. Chim. Acta* 47, 464.
- Hemmerich, P., Ghisla, S., Hartmann, U., and Müller, F. (1971), in *Flavins and Flavoproteins*, Kamin, H., Ed., Baltimore, Md., University Park Press, pp 83–105.
- Hemmerich, P., Massey, V., and Weber, G. (1967), *Nature (London)* 213, 728.
- Howell, L. G., Spector, T., and Massey, V. (1972), *J. Biol. Chem.* 247, 4340.
- Kierkegaard, P., Norrestam, R., Werner, P., Csoeregh, I., Glehn, M., Karlsson, R., Leijonmark, M., Roennquist, O., Stensland, B., Tillberg, O., and Torbjørsson, L. (1971), in *Flavins and Flavoproteins*, Kamin, H., Ed., Baltimore, Md., University Park Press, pp 1–22.
- Kotaki, A., Maoi, M., Okuda, J., and Yagi, K. (1967), *J. Biochem. (Tokyo)* 61, 404.
- Koziol, J. (1969), *Photochem. Photobiol.* 9, 45.
- Lockridge, O., Massey, V., and Sullivan, P. A. (1972), *J. Biol. Chem.* 247, 8097.
- Lowe, H. J., and Clark, W. M. (1956), *J. Biol. Chem.* 221, 983.
- Main, L., Kasperek, G. J., and Bruice, T. C. (1972), *Biochemistry* 11, 3991.
- Massey, V. (1966), *Methods Enzymol.* 9, 272.
- Massey, V., Curti, B., Müller, F., and Mayhew, S. G. (1968), *J. Biol. Chem.* 243, 1329.
- Massey, V., and Ganther, H. (1965), *Biochemistry* 4, 1161.
- Massey, V., Müller, F., Feldberg, R., Schuman, M., Sullivan, P., Howell, L. G., Mayhew, S. G., Matthews, R. G., and Foust, G. P. (1969), *J. Biol. Chem.* 244, 3999.
- Massey, V., and Palmer, G. (1966), *Biochemistry* 5, 3181.
- Massey, V., and Swoboda, B. E. P. (1963), *Biochem. Z.* 338, 474.
- Massey, V., and Williams, C. H., Jr. (1965), *J. Biol. Chem.* 240, 4470.
- Matthews, R. G., and Massey, V. (1969), *J. Biol. Chem.* 244, 1779.
- Mayhew, S. G. (1971), *Biochim. Biophys. Acta* 235, 276.
- Mayhew, S. G., Foust, G. P., and Massey, V. (1969), *J. Biol. Chem.* 244, 803.
- Müller, F., and Hemmerich, P. (1966), *Helv. Chim. Acta* 49, 2352.
- Palmer, G., and Massey, V. (1968), in *Biological Oxidations*, Singer, T. P., Ed., New York, N. Y., Interscience, pp 263–300.
- Schuman, M., and Massey, V. (1971), *Biochim. Biophys. Acta* 227, 500.
- Shethna, Y., Wilson, P. W., and Beinert, H. (1966), *Biochim. Biophys. Acta* 113, 225.
- Slifkin, M. A. (1971), *Charge Transfer Interactions of Biomolecules*, New York, N. Y., Academic Press, pp 132–172.
- Song, P. S. (1971), in *Flavins and Flavoproteins*, Kamin, H., Ed., Baltimore, Md., University Park Press, pp 37–61, and literature cited therein.
- Strickland, S., and Massey, V. (1973), *J. Biol. Chem.* 248, 2944.
- Sun, M., Moore, T. A., and Song, P. S. (1972), *J. Amer. Chem. Soc.* 94, 1730.
- Swoboda, B. E. P., and Massey, V. (1965), *J. Biol. Chem.* 240, 2209.
- Takeda, H., Yamamoto, S., Kojima, Y., and Hayaishi, O. (1969), *J. Biol. Chem.* 244, 2935.
- Tauscher, L., Ghisla, S., and Hemmerich, P. (1973), *Helv. Chim. Acta* 56, 630.
- Tollin, G. (1968), in *Molecular Associations in Biology*, Pullman, B., Ed., New York, N. Y., Academic Press, pp 393–409.
- Walaas, E., and Walaas, O. (1956), *Acta Chem. Scand.* 10, 122.
- Walker, W. H., Hemmerich, P., and Massey, V. (1970), *Eur. J. Biochem.* 13, 258.
- Walker, W. H., Singer, T. P., Ghisla, S., and Hemmerich, P. (1972), *Eur. J. Biochem.* 26, 279.
- Walsh, C. T., Schonbrunn, A., and Abeles, R. H. (1971), *J. Biol. Chem.* 246, 6855.
- Walsh, C. T., Schonbrunn, A., Babson, J., Lockridge, O., Massey, V., and Abeles, R. H. (1972), *J. Biol. Chem.* 247, 6004.
- Watenpaugh, K. D., Sieker, L. C., Jensen, L. H., LeGall, J., and Dubourdieu, M. (1972), *Proc. Nat. Acad. Sci. U. S. A.* 69, 3185.
- Weber, G. (1950), *Biochem. J.* 47, 114.
- Wellner, O., and Meister, A. (1960), *J. Biol. Chem.* 235, 2013.
- Zanetti, G., and Williams, C. H., Jr. (1967), *J. Biol. Chem.* 242, 5232.