



Title	Fluorescence characteristics of dissolved organic matter in the deep waters of the Okhotsk Sea and the northwestern North Pacific Ocean
Author(s)	Yamashita, Youhei; Cory, Rose M.; Nishioka, Jun; Kuma, Kenshi; Tanoue, Eiichiro; Jaffé, Rudolf
Citation	Deep Sea Research Part II: Topical Studies in Oceanography, 57(16), 1478-1485 <a href="https://doi.org/10.1016/j.dsr2.2010.02.016">https://doi.org/10.1016/j.dsr2.2010.02.016</a>
Issue Date	2010-08-15
Doc URL	<a href="http://hdl.handle.net/2115/47865">http://hdl.handle.net/2115/47865</a>
Type	article (author version)
File Information	DSR2-57-16_1478-1485.pdf



[Instructions for use](#)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

**Fluorescence characteristics of dissolved organic matter in the deep waters of the  
Okhotsk Sea and the northwestern North Pacific Ocean**

Youhei Yamashita<sup>1\*</sup>, Rose M. Cory<sup>2</sup>, Jun Nishioka<sup>3</sup>, Kenshi Kuma<sup>4</sup>, Eiichiro Tanoue<sup>5</sup>,  
Rudolf Jaffé<sup>1</sup>

<sup>1</sup>Southeast Environmental Research Center, and Department of Chemistry and  
Biochemistry, Florida International University, Miami, Florida, 33199, USA

<sup>2</sup>Department of Environmental Sciences and Engineering, University of North Carolina,  
Chapel Hill, Chapel Hill, North Carolina, 27599, USA

<sup>3</sup>Pan-Okhotsk Research Center, Institute of Low Temperature Science, Hokkaido  
University, Sapporo, 060-0819, Japan

<sup>4</sup>Graduate School of Environmental Sciences, Hokkaido University, Sapporo, 060-0810,  
Japan

<sup>5</sup>Graduate School of Environmental Studies, Nagoya University, Nagoya, 464-8601,  
Japan

\*Corresponding author at: Present address: Faculty of Environmental Earth Science,  
Hokkaido University, N10, W5, Kita-ku, Sapporo, Hokkaido 060-0810, Japan, E-mail  
address: yamashiy@ees.hokudai.ac.jp

1 **Abstract**

2 Fluorescent dissolved organic matter (DOM), a fraction of chromophoric DOM, has  
3 been known to be produced in the deep ocean and has been considered to be bio-  
4 refractory. However, the factors controlling fluorescence properties of DOM in the deep  
5 ocean are still not well understood. In this study, we determined the fluorescence  
6 properties of DOM in the deep waters of the Okhotsk Sea and the northwestern North  
7 Pacific Ocean using excitation-emission matrix (EEM) fluorescence and parallel factor  
8 analysis (PARAFAC). One protein-like, two humic-like components, and one uncertain  
9 component, that might be derived from a fluorometer artifact, were identified by EEM-  
10 PARAFAC. Fluorescence intensity levels of the protein-like component were highest in  
11 the surface waters, decreased with depth, but did not change systematically in the  
12 bathypelagic layer (1000 m - bottom). Fluorescence characteristics of the two humic-like  
13 components were similar to those traditionally defined as marine and terrestrial humic-  
14 like fluorophores, respectively. The fluorescence intensity levels of the two humic-like  
15 components were lowest in the surface waters, increased with depth in the mesopelagic  
16 layer (200 - 1000 m), and then slightly decreased with depth in the bathypelagic layer.  
17 The ratio of the two humic-like components remained in a relatively narrow range in the  
18 bathypelagic layer compared to that in the surface layer, suggesting a similar composition  
19 of humic-like fluorophores in this layer. In addition, the fluorescence intensities of the  
20 two humic-like components were linearly correlated to apparent oxygen utilization  
21 (AOU) in the bathypelagic layer, suggesting that both humic-like components are  
22 produced *in situ* as organic matter is oxidized biologically. These findings imply that  
23 optical characteristics of humic-like fluorophores once formed might not be altered

1 further biologically or geochemically in the deep ocean. On the other hand, relationships  
2 of fluorescence intensities with AOU and Fe(III) solubility were different between the  
3 two humic-like components in the mesopelagic layer, suggesting different environmental  
4 dynamics and biogeochemical roles for the two humic-like components.

5

6 **Keywords;** Dissolved organic matter (DOM), Fluorescence, Excitation-emission matrix  
7 (EEM) fluorescence, parallel factor analysis (PARAFAC), Biological production, Deep  
8 water, Okhotsk Sea, northwestern North Pacific

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

1 **1. Introduction**

2 Dissolved organic matter (DOM) in the ocean constitutes one of the largest reduced  
3 carbon pools in the global carbon cycle (700 PgC; Hedges and Keil, 1995).  
4 Concentrations of dissolved organic carbon (DOC) are usually highest in the surface  
5 layer, decrease sharply with depth, and then do not change significantly in the  
6 mesopelagic (200 - 1000 m), bathypelagic (1000 – 4000 m), and abyssopelagic (4000 m -  
7 bottom) layers (Hansell, 2002). In terms of volume, however, the deep ocean occupies a  
8 major part of the marine system, and thus, the majority of DOM (~650 PgC) occurs in the  
9 deep ocean (Ogawa and Tanoue, 2003). It is therefore clearly important to better  
10 understand the composition and environmental dynamics of DOM in the deep sea.

11 In the deep ocean, DOC concentrations have been reported to slightly decrease along  
12 with deep ocean circulation, i.e., with increasing water mass age, indicating the removal  
13 of DOM in this layer (Hansell and Carlson, 1998). Radiocarbon measurements in bulk  
14 DOC showed “old” average apparent age (4000 - 6000 yrs; Williams and Druffel, 1987),  
15 however, “young” radiocarbon ages of neutral sugars in high molecular weight (HMW)-  
16 DOM were also reported, suggesting that an important fraction of deep sea DOM may be  
17 introduced by dissolution from large, rapidly sinking particles (Repeta and Aluwihare,  
18 2006). In terms of quality, nuclear magnetic resonance analysis of HMW-DOM clarified  
19 that major chemical forms of HMW-DOM are similar to those of freshly produced  
20 organic matter (Benner et al., 1992; McCarthy et al., 1997; Aluwihare et al., 2002).  
21 However, contributions of humic substances, e.g., aromaticity in HMW-DOM, were  
22 greater in the deep HMW-DOM compared to the surface HMW-DOM (Benner et al.,  
23 1992; McCarthy et al., 1997; Aluwihare et al., 2002).

1 Chromophoric DOM (CDOM), the optically active fraction of DOM, occurs  
2 ubiquitously in the ocean and aquatic environments in general, and can be separated into  
3 two major classes, i.e., humic-like components and biogenic components such as protein-  
4 like fluorophores (Coble, 1996; Yamashita and Tanoue, 2003; 2009; Maie et al., 2006,  
5 2007, 2008). Vertical profiles in the ocean of biogenic components in CDOM showed a  
6 pattern similar to those of DOC (Mopper and Schultz, 1993; Yamashita and Tanoue,  
7 2009). Humic-like components in CDOM have been considered as a fraction of  
8 molecularly-uncharacterized components or humic-like substances (Hedges et al., 2000).  
9 Levels of humic-like components in CDOM, determined by absorption coefficients and  
10 fluorescence intensity, have been found to be lowest in surface waters, a trend commonly  
11 attributed to photodegradation. The components increase with depth in the mesopelagic,  
12 but show little or no gradient at greater depths (Hayase et al., 1988; Chen and Bada,  
13 1992; Mopper and Schultz, 1993; Hayase and Shinozuka, 1995; Nelson et al., 2007;  
14 Yamashita et al., 2007; Yamashita and Tanoue, 2008; 2009). Thus, the contribution of  
15 humic-like components in CDOM to bulk DOM is larger in deep waters compared to  
16 surface waters. This view is also consistent with the differences in chemical forms of  
17 HMW-DOM between surface and deep waters.

18 Such vertical distribution profiles of humic-like components in CDOM suggest that  
19 the biogeochemical roles of CDOM in deep waters can be more important than those in  
20 surface waters. For example, vertical profiles of the humic-like fluorescence intensity  
21 were related to those of Fe(III) solubility in the mesopelagic and bathypelagic layers but  
22 not in the surface layer, suggesting that humic-like components in CDOM are major  
23 factor in controlling the Fe(III) solubility and the dissolved Fe concentration in the deep

1 ocean through their complexation with Fe(III) as natural organic ligands (Tani et al.,  
2 2003; Takata et al., 2004; 2005; Kitayama et al., 2009).

3 Recently, the basin scale distribution of the humic-like fluorescence intensity was  
4 examined for the Pacific Ocean (Yamashita and Tanoue, 2008), finding a continuous  
5 increase in fluorescence intensity in the bathypelagic and abyssopelagic layers from the  
6 Southern Ocean to the northern North Pacific, while a decrease in DOC concentration  
7 was evident (Hansell and Carlson, 1998). A strong linear relationship between the  
8 fluorescence intensity and apparent oxygen utilization (AOU) was also reported in these  
9 layers for the Pacific (Yamashita and Tanoue, 2008). Since AOU increases with oxygen  
10 consumption during organic matter oxidation processes, it is apparent that bio-refractory  
11 humic-like components in CDOM are produced *in situ* as organic matter is oxidized  
12 biologically. In addition, Yamashita and Tanoue (2008) reported variations of AOU-  
13 fluorescence intensity relationships between the bathypelagic (+ abyssopelagic) and  
14 mesopelagic layers and suggested that the cause leading to these different relationships  
15 was the distribution of different water masses having different pre-formed CDOM levels  
16 between these layers.

17 Previous studies assumed that fluorescence properties were not significantly different  
18 across water depths and oceanic regions, and therefore, the levels of humic-like  
19 components in CDOM were determined by the fluorescence intensity of single excitation-  
20 emission pairs (e.g., 320 nm excitation and 420 nm emission) (Hayase et al., 1988; Chen  
21 and Bada, 1992; Hayase and Shinozuka, 1995; Yamashita et al., 2007; Yamashita and  
22 Tanoue, 2008). Thus, it is still unclear whether optically similar DOM is produced *in situ*

1 and accumulated in the deep ocean, or if fluorescence characteristics change with  
2 increasing AOU.

3 In the present study, CDOM in surface and deep waters of the Okhotsk Sea and the  
4 northwestern North Pacific Ocean were characterized by excitation-emission matrix  
5 (EEM) fluorescence with parallel factor analysis (PARAFAC). The PARAFAC  
6 methodology was successfully applied to decompose EEMs into a number of  
7 fluorescence components (Stedmon et al., 2003; Stedmon and Markager, 2005a; Cory  
8 and McKnight, 2005; Murphy et al., 2008; Jaffé et al., 2008; Yamashita et al., 2008). The  
9 main objective of the present study was to examine the dynamics of the CDOM  
10 components obtained through EEM-PARAFAC. Special attention was paid to clarify the  
11 similarity/dissimilarity of fluorescence characteristics for the deep ocean, an area ideally  
12 suited to study the effects on DOM composition by microbial activity in the absence of  
13 photochemical interferences.

14

## 15 **2. Materials and methods**

16 Seawater samples were collected from surface to bottom layers at 4 stations in the  
17 Okhotsk Sea and northwestern North Pacific Ocean (Fig. 1) during May and June 2000.  
18 In this paper, we defined depths from 200 to 1000 m and from 1000 m to the bottom as  
19 the mesopelagic and bathypelagic layers, respectively. Sampling and filtration procedure  
20 were described in detail elsewhere (Tani et al., 2003). Samples were filtered through 0.22  
21  $\mu\text{m}$  filter and kept frozen ( $< -20\text{ }^{\circ}\text{C}$ ) on board. In the laboratory, frozen samples were  
22 thawed and used for measurements of Fe(III) solubility and fluorescence intensity at 320  
23 nm excitation and 420 nm emission (Tani et al., 2003). Aliquots for EEM measurements



1 were taken in low density polyethylene tubes and then re-frozen. These tubes were pre-  
2 cleaned using the same procedure as used for trace metal sampling (Bruland et al., 1979).  
3 The effects of freezing on EEMs of oceanic DOM are not clear. However, using samples  
4 from central North Pacific surface and deep layers, we determined differences in humic-  
5 like fluorescence intensity at 320 nm excitation and 420 nm emission before and after  
6 freezing, finding that differences were  $-0.1 \pm 1.7\%$  ( $n = 9$ ) relative to fresh samples  
7 analyzed prior to any freezing or thawing cycles. The freezing effects on protein-like  
8 fluorophores found in marine DOM samples are also unknown. However, a recent study  
9 on the effects of freezing and thawing on DOM fluorescence documented up to  $\pm 50\%$   
10 change in fluorescence intensity for the humic and protein peaks in EEMs of riverine  
11 DOM that had been filtered prior to analysis with  $1.2 \mu\text{m}$  GF/C filters (Spencer et al.  
12 2007). It is likely that in the latter study, some of the changes in fluorescence upon  
13 freezing may be attributed to rupture of microbial cells or particulate origin, which would  
14 not be expected in our study where  $0.22 \mu\text{m}$  filters were used for sample preparation.

15 The results of bulk fluorescence intensity at 320 nm excitation and 420 nm emission,  
16 Fe(III) solubility, temperature, salinity, sigma-theta, nutrients, and AOU at 4 stations can  
17 be found elsewhere (Tani et al., 2003).

18 Just before the EEM measurement, samples were thawed and allowed to stand until  
19 reaching near room temperature. The measurements of EEM were carried out using a  
20 fluorescence spectrometer (Hitachi F-4500) according to Yamashita and Tanoue (2003).  
21 The specific instrumental components were corrected for excitation and emission  
22 according to protocols recommended by the manufacturer. The EEMs were generated by  
23 scanning emission spectra from 225 to 500 nm at 1 nm intervals with 5 nm increments of

1 the excitation wavelength from 225 to 400 nm. Several post acquisition steps were  
2 involved in the correction of EEMs. First, the EEM of Milli-Q water, which was freshly  
3 produced and measured every day, was subtracted from the EEM of each sample. Second,  
4 fluorescence intensities were corrected to the area under the water Raman peak  
5 (excitation = 350 nm), analyzed daily, and then were converted to quinine sulfate unit  
6 (QSU) using a calibration with quinine sulfate monohydrate in a solution of  $0.05 \text{ mol L}^{-1}$   
7  $\text{H}_2\text{SO}_4$ . Inner filter corrections were not applied for these samples because of the  
8 extremely low absorbance for open ocean samples (Nelson et al., 2007; Yamashita and  
9 Tanoue, 2009).

10 PARAFAC modeling for EEM data has been described in detail elsewhere (Stedmon  
11 et al., 2003; Stedmon and Bro, 2008). The modeling was carried out in MATLAB with  
12 the DOMFluor toolbox (Stedmon and Bro, 2008) using 80 EEMs obtained from surface  
13 to deep waters. The EEM wavelength ranges used were 260-400 nm and 300-500 nm for  
14 excitation and emission, respectively. The determination of the correct number of  
15 components was primarily achieved by the split half analysis and random initialization  
16 (Stedmon and Bro, 2008). In this study, the fluorescence intensities of each PARAFAC  
17 component were reported as QSU.

18

### 19 **3. Results and discussion**

#### 20 *3.1. EEM fluorescence of DOM in the deep ocean*

21 Typical EEM spectra obtained from surface and deep waters are presented in Fig. 2.  
22 Both EEMs showed high levels of fluorescence intensity at around 275 nm excitation and  
23 305 nm emission. This peak corresponds to the tyrosine-like fluorophore or peak B

1 (Coble, 1996; Yamashita and Tanoue, 2003) and showed higher fluorescence intensity in  
2 surface waters compared to deep waters. While a detectable fluorescence signal was  
3 found for tyrosine-like fluorophores, tryptophan-like fluorophore were not observed in  
4 samples from this study region. Such fluorescence characteristics of protein-like  
5 fluorophores have been previously identified in the oceanic surface and deep waters  
6 (Mopper and Schultz, 1993; Yamashita and Tanoue, 2003; 2004a).

7 Both, surface and deep water EEMs also showed a high fluorescence intensity at <  
8 260 nm excitation and 400 - 500 nm emission, and the fluorescence intensity of this peak  
9 was higher in deep waters compared to surface waters (Fig. 2). This peak corresponds to  
10 the humic-like peak A (Coble, 1996) and has been reported to be present from surface to  
11 deep waters in the open ocean (Mopper and Schultz, 1993; Coble, 1996). Low levels of  
12 this peak in the surface water compared to deep water were also found in the Sargasso  
13 Sea (Mopper and Schutlz, 1993). In addition to peak A, the region at 290-350 nm  
14 excitation and 380-400 nm emission presented another peak in both surface and deep  
15 waters (Fig. 2). According to Coble (1996), this peak corresponds to the marine humic-  
16 like peak M and/or humic-like peak C. This peak was red-shifted in the deep water  
17 compared with that in the surface water (Fig. 2). Even though only a few studies applied  
18 EEMs to deep ocean DOM (Mopper and Schultz, 1993; Coble, 1996), similar red-shift,  
19 i.e., dominance of marine humic-like fluorophore (peak M) in the surface water and the  
20 replacement to humic-like fluorophore (peak C) with depth, have been suggested (Coble,  
21 1996).

22

23 *3.2. PARAFAC components*

1 EEMs showed differences in fluorescence characteristics of DOM between the surface  
2 and deep waters (Fig. 2), however, it is hard to evaluate such differences applying the  
3 peak picking technique. PARAFAC was, therefore, applied for quantitative evaluation of  
4 differences in EEMs. A four component PARAFAC model was validated and selected as  
5 most suitable for this dataset (Fig. 3). Spectral characteristics of components identified in  
6 the region studied were similar to those previously found in other aquatic environments  
7 (Table 1). The selection of the four component model does not mean that the EEMs only  
8 contained four different fluorophores, but that these selected four components were the  
9 most representative fluorescent groups in this dataset.

10 Component 1 was comprised of two excitation maxima (< 260 nm and 370 nm) with  
11 an emission maximum at 466 nm and was categorized as mixture of the traditional  
12 humic-like peaks A and C which were usually representative fluorophores in terrestrial  
13 environments and low salinity waters in coastal environments (Coble, 1996; Conmy et al.,  
14 2004). The spectral features were also similar to a terrestrial humic-like component found  
15 in the surface water from coastal to oceanic regions (C3/P3; Murphy et al., 2008). On the  
16 other hand, this component is also similar to a component reported as ubiquitous in  
17 coastal regions (4; Stedmon and Markager, 2005a). A similar component was also  
18 identified in an Antarctic lake, which lacks a higher plant contribution from its watershed  
19 (Cory and McKnight, 2005). Thus, component 1 showed similar spectral characteristics  
20 with terrestrial (higher plant derived) fluorophores and fluorophores related to microbial  
21 reworking of organic matter.

22 There was an intensive peak at 325 nm excitation and 385 nm emission for component  
23 2. This peak was similar to the marine humic-like peak M traditionally defined (Coble,

1 1996) and was similar to marine humic-like components found in the PARAFAC model  
2 from coastal environments (C6; Yamashita et al., 2008) as well as surface waters from  
3 coastal to oceanic regions (Murphy et al., 2008). Again, a similar component was also  
4 identified in an Antarctic lake (C3; Cory and McKnight, 2005), and was found to be  
5 prominent in wastewater samples (6; Stedmon and Markager, 2005a), thus providing  
6 evidence that this component is associated with recent biological activity. Therefore, this  
7 component could be characterized as representing a group of humic-like fluorophores  
8 strongly associated with recent biological activity.

9 Component 3 did not appear as a peak in EEM and has not been commonly reported in  
10 previous PARAFAC studies (Cory and McKnight, 2005; Stedmon and Markager, 2005a;  
11 Stedmon et al., 2007; Yamashita et al., 2008; Yamashita and Jaffé, 2008). Only one study,  
12 using three fluorometers (i.e., SPEX FluoroLog-2, SPEX FluoroMax-2, and Varian Cary  
13 Eclipse), identified similar component by PARAFAC modeling for samples from surface  
14 waters in coastal to oceanic transects (P4; Murphy et al., 2008). This component has an  
15 uncertain origin, and is likely an analytical artifact of the fluorometer (Murphy et al.,  
16 2008). These results suggest that the peak A region can be affected by fluorometer  
17 artifacts and that caution is needed in the interpretation of EEMs for especially low  
18 CDOM level samples such as oceanic waters. Thus, based on the above, this component  
19 was not discussed further in this study.

20 Only one component was found in the region of protein-like fluorophores, in contrast  
21 to many previous studies that have identified up to four protein-like components (Cory  
22 and McKnight, 2005; Stedmon and Markager, 2005a; Murphy et al., 2008; Yamashita  
23 and Jaffé, 2008; Yamashita et al., 2008). Fluorescence characteristics of component 4

1 were almost identical to those of free tyrosine (Yamashita and Tanoue, 2003) and were  
2 similar to those of a tyrosine-like component found in previous PARAFAC studies (Cory  
3 and McKnight, 2005; Stedmon and Markager, 2005a; Murphy et al., 2008; Yamashita et  
4 al., 2008). The other common protein-like component, i.e., the tryptophan-like  
5 component, was not observed in this study. The dominance of tyrosine-like fluorophore  
6 in oceanic waters was also observed previously (Mopper and Schultz, 1993; Yamashita  
7 and Tanoue, 2003).

8

### 9 *3.3. Vertical profiles of PARAFAC components*

10 Levels of tyrosine-like component 4 were highest in the surface waters, decreased  
11 with depth, and then did not change systematically towards the bottom (Fig. 4). Similar  
12 vertical characteristics were also observed for protein-like fluorophores in EEMs at the  
13 Sargasso Sea (Mopper and Schultz, 1993) and the Sagami Bay, Japan (Yamashita and  
14 Tanoue, 2004a). Yamashita and Tanoue (2003) found the linear relationships between  
15 tyrosine-like fluorescence intensities and tyrosine concentrations as well as  
16 concentrations of total hydrolyzable amino acids from coastal to oceanic environments.  
17 Even though such relationships were not studied for the deep ocean, the vertical profiles  
18 of dissolved amino acids in the open ocean (Kaiser and Benner, 2008; 2009) were similar  
19 to those found for protein-like component 4 in this study. Such similarity of vertical  
20 characteristics between dissolved amino acids and tyrosine-like component 4 suggest that  
21 tyrosine-like component 4 is likely derived from tyrosine-containing molecules in the  
22 open ocean.

1 Vertical profiles of humic-like components 1 (corresponding to a mixture of the  
2 traditionally-defined peaks A and C, Table 1) and 2 (corresponding to the traditionally-  
3 defined peak M, Table 1) were similar to each other (Fig. 4). The lowest fluorescence  
4 intensity of humic-like components were found in the surface waters, followed by an  
5 increase with depth, maximizing at 600~1000 meters, and then showing a slight decrease  
6 with depth. In the mesopelagic layer, however, fluorescence intensity gradients for  
7 component 1 were greater than those for component 2. Vertical profiles of both humic-  
8 like components were almost identical to those of fluorescence intensity at 320 nm  
9 excitation and 420 nm emission at the same stations (Tani et al., 2003). The lowest levels  
10 of both humic-like components in the surface layer may have resulted from the  
11 degradation of humic-like fluorophores in the photic zone. This is most likely due to  
12 photodegradation since it is well known that, while humic-like fluorescence intensity  
13 remarkably decreases during photo-irradiation (Mopper et al., 1991; Chen and Bada,  
14 1992; Nieto-Cid et al., 2006) it tends to increase during microbial degradation of organic  
15 matter (Yamashita and Tanoue, 2004; Nieto-Cid et al., 2006).

16 Ratios of humic-like component 1 to component 2, corresponding to the ratio of the  
17 traditionally-defined peak C/peak M (Table 1), were lowest (1.0~1.2) in the surface  
18 waters, increased with depth, and then remained in a relatively narrow range (1.4~1.5) in  
19 the bathypelagic layer (Fig. 4). Vertical characteristics indicated that humic-like  
20 components 1 and 2 were present at similar levels in the surface water, but humic-like  
21 component 1 was enriched compared to component 2 in the bathypelagic layer. The red-  
22 shift of the humic-like fluorophores in the deep water compared to the surface water (Fig.  
23 2) could be explained by the abovementioned change in ratio, i.e., the change in humic-

1 like composition, with depth (Fig. 4). The shift of peak position from peak C  
2 (corresponding to component 1) to peak M (corresponding to component 2) was also  
3 observed during photodegradation but not during biodegradation experiments using  
4 coastal waters (Moran et al., 2000). Thus, the observation of a low ratio for the surface  
5 waters in conjunction with the lowest fluorescence intensities of humic-like components  
6 suggest that component 1 may be more photo-reactive compared to component 2.

#### 8 *3.4. Fluorescence characteristics of in situ-produced CDOM*

9 Strong evidence for the microbial production of humic-like CDOM during organic  
10 matter degradation processes has been provided by several authors (Chen and Bada,  
11 1992; Hayase and Shinozuka, 1995; Nelson et al., 1998; 2007; Nieto-Cid, et al., 2005;  
12 Yamashita et al., 2007; Yamashita and Tanoue, 2008) and through *in vitro* experiments  
13 (Parlanti et al., 2000; Rochelle-Newall and Fisher, 2002; Nelson et al., 2004; Yamashtia  
14 and Tanoue, 2004b; Stedmon and Markager, 2005b; Nieto-Cid et al., 2006). Figure 5  
15 shows relationships between AOU and fluorescence intensities of humic-like components  
16 1 and 2 in the mesopelagic and bathypelagic layers. In the bathypelagic layer, the  
17 relatively simple water mass distribution, primarily supplied from Circumpolar Deep  
18 Water (Matsumoto, 2007), allowed for a better examination of relationship between AOU  
19 and fluorescence intensities (Yamashita and Tanoue, 2008). Therefore, in this study, *in*  
20 *situ*-produced humic-like fluorophores are discussed for the bathypelagic layer.

21 Fluorescence intensities of both humic-like components were linearly related to AOU  
22 in the bathypelagic layer (Fig. 5; [Comp. 1] =  $0.0029 \times [\text{AOU}] + 0.45$ ,  $R^2 = 0.82$ ,  $n = 16$ ,  
23  $p < 0.001$ ; [Comp. 2] =  $0.0018 \times [\text{AOU}] + 0.36$ ,  $R^2 = 0.69$ ,  $n = 16$ ,  $p < 0.001$ ). The *in situ*



1 production of bio-refractory humic-like fluorophores were indicated from the linear  
2 relationship between AOU and fluorescence intensity at 320 nm excitation and 420 nm  
3 emission in the bathypelagic layer from the Southern Ocean to northern North Pacific  
4 where AOU ranged from 130 to 310  $\mu\text{mol kg}^{-1}$  (Yamashita and Tanoue, 2008). The AOU  
5 range in the bathypelagic layer observed here (175 to 301  $\mu\text{mol kg}^{-1}$ ) was slightly  
6 narrower than that described in the Pacific Ocean (Yamashita and Tanoue, 2008), but still  
7 similar, suggesting that the relationships between AOU and fluorophores for the  
8 bathypelagic layers in the Okhotsk Sea and the northwestern North Pacific Ocean are  
9 comparable to that for the Pacific bathypelagic layer.

10 The red-shifted humic-like fluorophore (peak C) has been well known to be a  
11 terrestrial humic-like fluorophore, i.e., the dominant fluorophore in terrestrial  
12 environments (Coble, 1996). However, the linear relationships between AOU and humic-  
13 like components 1 and 2 in the bathypelagic layer suggest that both component 1  
14 (corresponding to traditional peak C) and component 2 (corresponding to traditional peak  
15 M) are produced *in situ* in the deep ocean. PARAFAC components with fluorescence  
16 features similar to the traditional peak C were found from an Antarctic lake where no  
17 higher plant sources are evident (Table 1; Cory and McKnight, 2005). The microbial  
18 production of all humic-like PARAFAC components including components  
19 corresponding to the traditionally-defined peak C, was reported for mesocosm  
20 experiments (Stedmon and Markager, 2005b). Results from such PARAFAC-based  
21 studies suggest that fluorophores showing similar fluorescence characteristics to  
22 component 1 can originate from either terrestrial or microbial sources of organic matter.  
23 It should be noted that the traditionally-defined peak C is often used as a tracer for

1 terrestrial CDOM in coastal environments (e.g., Chen et al., 2004; Conmy et al., 2004).  
2 Thus, further studies are necessary to clarify the dynamics of fluorophores corresponding  
3 to component 1 for the wide range aquatic environments.

4 In addition, ratios of humic-like component 1 to component 2 were in a relatively  
5 narrow range (Fig. 4) and were not linearly related to AOU in the bathypelagic layer  
6 ( $[C1/C2] = 0.0004 \times [AOU] + 1.35, R^2 = 0.07, n = 16, p > 0.01$ ). Such results suggest that  
7 the relative proportion of *in situ*-produced fluorophores did not change with increase in  
8 AOU in the deep ocean. Similar optical properties of *in situ*-produced humic like CDOM  
9 were also suggested from the uniform distribution of the ratio of fluorescence intensity to  
10 absorption coefficient in the bathypelagic layer along a transect from the equatorial to  
11 northern North Pacific (Yamashita and Tanoue, 2009). Thus, similarities of optical  
12 characteristics of humic-like CDOM in the bathypelagic layer suggest that the  
13 composition of *in situ* produced humic-like CDOM may not be significantly altered  
14 biologically and geochemically.

15

### 16 3.5. Fluorescence characteristics and its environmental implications in the mesopelagic 17 layer

18 Plots between AOU and component 1 in the mesopelagic layer had a similar linear  
19 relationship with that in the bathypelagic layer (Fig. 5). On the other hand, relationships  
20 between AOU and component 2 in the mesopelagic layer largely deviated from those in  
21 the bathypelagic layer (Fig. 5). Yamashita and Tanoue (2008) found different AOU-  
22 fluorescence relationships between the mesopelagic and bathypelagic layers in the Pacific  
23 Ocean and indicated that differences in levels of pre-formed fluorophores yield different

1 relationships. Thus, relationships between AOU and component 1 and 2 suggested that  
2 pre-formed levels of component 1 were similar between the mesopelagic and  
3 bathypelagic layers, but those of component 2 in the mesopelagic layer were high  
4 compared to the bathypelagic layer.

5 The water mass in the bathypelagic layer is supplied primarily from Circumpolar Deep  
6 Water (Matsumoto, 2007), but major water mass in the mesopelagic layer are derived  
7 from North Pacific Intermediate Water (NPIW) in the Oyashio region and the Okhotsk  
8 Sea Intermediate Water (OSIW) which contribute to the formation of the NPIW (Yasuda,  
9 2004). The Amur River discharges into the Okhotsk Sea, and as a consequence, high  
10 levels of terrestrial biomarker as well as terrestrial CDOM were found in the NPIW for  
11 the central North Pacific (Hernes and Benner, 2002; Yamashita and Tanoue, 2009).  
12 Therefore, high levels of pre-formed component 2 in the mesopelagic layer might be the  
13 result from terrestrial environments, discharged through the Amur River, and thus adding  
14 to the fraction formed during OSIW and NPIW.

15 In general, however, the dominance of component 1 (corresponding to peak C; Table  
16 1) compared to component 2 (corresponding to peak M; Table 1) would be expected for  
17 terrestrial environments, as mentioned above, but not for deep ocean waters. Such  
18 apparent contradiction might be explained by photodegradation of humic-like  
19 fluorophores. The higher photo-labile property of component 1 compared to component 2  
20 was suggested by vertical profiles of their ratio (Fig. 4). The blue-shift of peak position  
21 was also found for humic-like fluorophores in estuarine DOMs during photodegradation  
22 (Moran et al., 2000), implying that peak C is more photo-reactive compared to peak M.  
23 Cory et al. (2007) also found that a PARAFAC component similar to component 1 was

1 more photodegradable than other PARAFAC components. Thus, the component 1  
2 produced in the surface waters might be photodegraded readily, and thus,  
3 photodegradation processes might keep component 1 at low levels in the surface layer,  
4 irrespective of oceanic region differences. A fraction of component 1 might also be  
5 riverine derived and could be selectively photo-degraded compared to component 2 in the  
6 coastal margin of the Okhotsk Sea.

7 Different distributional patterns between components 1 and 2 in the mesopelagic layer  
8 (Fig. 5) suggested differences in the biogeochemical roles between the two humic-like  
9 components. Figure 6 shows the relationships between Fe(III) solubility and fluorescence  
10 intensity of components 1 and 2 in the mesopelagic layer. Linear relationships between  
11 Fe(III) solubility and humic-like fluorescence intensity were found in the deep ocean,  
12 suggesting that humic-like components in CDOM may be a major factor controlling the  
13 Fe(III) solubility and the dissolved Fe concentration in the deep ocean through the  
14 complexation of Fe(III) as natural organic ligands (Tani et al., 2003; Takata et al., 2004;  
15 2005; Kitayama et al., 2009). Interestingly, Fe(III) solubility linearly correlated with  
16 levels of component 1 ( $R^2 = 0.81$ ,  $n = 20$ ,  $p < 0.001$ ), but not with those of component 2  
17 ( $R^2 = 0.29$ ,  $n = 20$ ,  $p > 0.01$ ). Such differences in relationships implies that component 1  
18 is likely to be important fluorophores as ligand for Fe(III) compared to component 2.  
19 Such differences in interactivity of trace metals with DOM fluorophores have previously  
20 been reported (Yamashita and Jaffé, 2008). In addition, extremely high concentrations of  
21 dissolved and particulate Fe in the OSIW and the NPIW compared to other water masses  
22 were found in the same region (Nishioka et al., 2007). These results suggest that  
23 component 1 may be an important contributor to Fe(III) solubility and, subsequently, may

1 regulate the high concentration of dissolved Fe in the mesopelagic layer of the North  
2 Pacific Ocean.

3

#### 4 **4. Conclusions**

5 The *in situ* production of bio-refractory humic-like fluorophores was demonstrated  
6 based on linear relationship between humic-like fluorescence intensity and AOU as well  
7 as nutrient concentrations in the mesopelagic and bathypelagic layers (Hayase et al.,  
8 1988; Chen and Bada, 1992; Hayase and Shinozuka, 1995; Yamashita et al., 2007;  
9 Yamashita and Tanoue, 2008). To clarify the similarity/dissimilarity of humic-like  
10 fluorophores in the deep ocean, EEM-PARAFAC was for the first time applied to DOM  
11 in the deep ocean. PARAFAC statistically decomposed EEMs into one protein-like and  
12 two humic-like components at the Okhotsk Sea and the northwestern North Pacific Ocean.

13 The spectral characteristics of one of the humic-like components were similar to the  
14 microbial/marine humic-like fluorophore (peak M), while those of the other were similar  
15 to the terrestrial humic-like fluorophore (peak C) as traditionally assigned. The vertical  
16 profiles of the two humic-like components were similar and showed lowest levels in the  
17 surface waters, peaked in the mesopelagic layer, and decreased slightly with depth in the  
18 bathypelagic layer. Levels of both humic-like components were linearly correlated to  
19 AOU in the bathypelagic layer, implying that both humic-like components are *in situ*  
20 produced during microbial degradation processes of organic matter. In addition, the ratio  
21 of the two humic-like components remained in a relatively narrow range and was not  
22 related to AOU variations in the bathypelagic layer, suggesting that *in situ* produced

1 humic-like fluorophores once formed might not be altered biologically or geochemically  
2 along this layer.

3

#### 4 **Acknowledgements**

5 We thank S. Nakabayashi (JAMSTEC), M. Kusakabe (NIRS) and Captain M.  
6 Akimoto and the crew of the RV Mirai for assistance during the sampling cruise, D.  
7 Hansell for editorial assistance, and two anonymous reviewers for helpful comments and  
8 suggestions that helped improve the quality of this manuscript. Y.Y. thanks the College  
9 of Arts and Science at FIU for financial support. This study was partially supported by  
10 Grant-in-Aid for Scientific Research (No. 20241005) from the JSPS. This is SERC  
11 contribution #477.

12

1 **References**

- 2 Aluwihare, L.I., Repeta, D.J., Chen, R.F., 2002. Chemical composition and cycling of  
3 dissolved organic matter in the Mid-Atlantic Bight. *Deep-Sea Research II* 49, 4421-  
4 4437.
- 5 Benner, R., Pakulski, J.D. McCarthy, M., Hedges, J.I., Hatcher, P.G., 1992. Bulk  
6 chemical characteristics of dissolved organic matter in the ocean. *Science* 255, 1561-  
7 1564.
- 8 Bruland, K.W., Franks, R.P., Knauer, G.A., Martin, J.H., 1979. Sampling and analytical  
9 methods for the determination of copper, cadmium, zinc, and nickel at the nanogram  
10 per liter level in seawater. *Analytica Chimica Acta* 105, 233–245.
- 11 Chen, R.F., Bada, J.L., 1992. The fluorescence of dissolved organic matter in seawater.  
12 *Marine Chemistry* 37, 191-221.
- 13 Chen, R.F., Bissett, P., Coble, P., Conmy, R., Gardner, G.B., Moran, M.A., Wang, X.,  
14 Wells, M.L., Whelan, P., Zepp, R.G., 2004. Chromophoric dissolved organic mater  
15 (CDOM) source characterization in the Louisiana Bight. *Marine Chemistry* 89, 257-  
16 272.
- 17 Coble, P.G., 1996. Characterization of marine and terrestrial DOM in seawater using  
18 excitation-emission matrix spectroscopy. *Marine Chemistry* 51, 325-46.
- 19 Conmy, R.N., Coble, P.G., Chen, R.F., Gardner, G.B., 2004. Optical properties of  
20 dissolved organic matter in the Northern Gulf of Mexico. *Marine Chemistry* 89, 127-  
21 144.

1 Cory, R.M., McKnight, D.M., 2005. Fluorescence spectroscopy reveals ubiquitous  
2 presence of oxidized and reduced quinines in dissolved organic matter.  
3 *Environmental Science and Technology* 39, 5142-8149.

4 Cory, R. M., McKnight, D. M., Chin, Y.-P., Miller, P., Jaros C. L., 2007. Chemical  
5 characteristics of fulvic acids from Arctic surface waters: Microbial contributions and  
6 photochemical transformations. *Journal of Geophysical Research* 112, G04S51.

7 Hansell, D.A., 2002. DOC in the global ocean carbon cycle. In: Hansell, D.A., Carlson,  
8 C.A. (Eds.), *Biogeochemistry of Marine Dissolved Organic Matter*. Academic Press,  
9 San Diego, pp. 685-715.

10 Hansell, D.A., Carlson, C.A., 1998. Deep-ocean gradients in the concentration of  
11 dissolved organic carbon. *Nature* 395, 263-266.

12 Hayase, K., Shinozuka, N., 1995. Vertical distribution of fluorescent organic matter along  
13 with AOU and nutrients in the equatorial Central Pacific. *Marine Chemistry* 48, 283-  
14 290.

15 Hayase, K., Tsubota, H., Sunada, I., Goda, S., Yamazaki, H., 1988. Vertical distribution  
16 of fluorescent organic matter in the North Pacific. *Marine Chemistry* 25, 373-381.

17 Hedges, J.I., Keil, R.G., 1995. Sedimentary organic matter preservation: an assessment  
18 and speculative synthesis. *Marine Chemistry* 49, 81-115.

19 Hedges, J.I., Eglinton, G., Hatcher, P.G., Kirchman, D.L., Arnosti, C. Derenne, S.,  
20 Evershed, R.P., Kögel-Knabner, I., de Leeuw, J.W., Littke, R., Michaelis, W.,  
21 Rulkötter, J., 2000. The molecularly-uncharacterized component of nonliving organic  
22 matter in natural environments. *Organic Geochemistry* 31, 945-958.



1 Hernes, P.J., Benner, R., 2002. Transport and diagenesis of dissolved and particulate  
2 terrigenous organic matter in the North Pacific Ocean. *Deep-Sea Research I* 49, 2119-  
3 2132.

4 Jaffé, R., McKnight, D., Maie, N., Cory, R., McDowell, W.H., Cambell, J.L., 2008.  
5 Spatial and temporal variations in DOM composition in ecosystems: The importance  
6 of long-term monitoring of optical properties. *Journal of Geophysical Research* 113,  
7 G04032.

8 Kaiser, K., Benner, R., 2008. Major bacterial contribution to the ocean reservoir of  
9 detrital organic carbon and nitrogen. *Limnology and Oceanography* 53, 99-112.

10 Kaiser, K., Benner, R., 2009. Biochemical composition and size distribution of organic  
11 matter at the Pacific and Atlantic Time-Series Stations. *Marine Chemistry* 113, 63-77.

12 Kitayama, S., Kuma, K., Manabe, E., Sugie, K., Takata, H., Isoda, Y., Toya, K., Saitoh,  
13 S., Takagi, S., Kamei, Y., Sakaoka, K., 2009. Controls on iron distribution in the  
14 deep-water column of the North Pacific Ocean: Iron(III) hydroxide solubility and  
15 marine humic-type dissolved organic matter. *Journal of Geophysical Research*, in  
16 press.

17 Maie, N., Parish, K.J., Watanabe, A., Knicker, H. Benner, R., Abe, T., Kaiser, K., Jaffé,  
18 R., 2006. Chemical characteristics of dissolved organic nitrogen in an oligotrophic  
19 subtropical coastal ecosystems. *Geochimica Cosmochimica Acta* 70, 4491-4506.

20 Maie, N., Pisani, O., Jaffé, R., 2008. Mangrove tannins in aquatic ecosystems: Their fate  
21 and possible influence of dissolved organic carbon and nitrogen cycling. *Limnology  
22 and Oceanography* 53, 160-171.

- 1 Maie, N., Scully, N.M., Pisani, O., Jaffé, R., 2007. Composition of a protein-like  
2 fluorophore of dissolved organic matter in coastal wetlands and estuarine ecosystems.  
3 Water Research 41, 563-570.
- 4 Matsumoto, K., 2007. Radiocarbon-based circulation age of the world oceans. Journal of  
5 Geophysical Research 112, C09004.
- 6 McCarthy, M., Pratum, T., Hedges, J., Benner, R., 1997. Chemical composition of  
7 dissolved organic nitrogen in the ocean. Nature 390, 150-153.
- 8 Mopper, K., Schultz, C.A., 1993. Fluorescence as a possible tool for studying the nature  
9 and water column distribution of DOC components. Marine Chemistry 41, 229-238.
- 10 Mopper, K., Zhou, X., Kieber, R.J., Kieber, D.J., Sikorski, R.J., Jones, R.D., 1991.  
11 Photochemical degradation of dissolved organic carbon and its impact on the oceanic  
12 carbon cycle. Nature 353, 60-62.
- 13 Moran, M.A., Sheldon Jr., W.M., Zepp, R.G., 2000. Carbon loss and optical property  
14 changes during long-term photochemical and biological degradation of estuarine  
15 dissolved organic matter. Limnology and Oceanography 45, 1254-1264.
- 16 Murphy K.R., Stedmon, C.A., Waite, T.D., Ruiz, G.M., 2008. Distinguishing between  
17 terrestrial and autochthonous organic matter sources in marine environments using  
18 fluorescence spectroscopy. Marine Chemistry 108, 40-58.
- 19 Nelson, N.B., Carlson, C.A., Steinberg, D.K., 2004. Production of chromophoric  
20 dissolved organic matter by Sargasso Sea microbes. Marine Chemistry 89, 273-287.
- 21 Nelson, N.B., Siegel, D.A., Carlson, C.A., Swan, C., Smethie Jr., W.M., Khatiwala, S.,  
22 2007. Hydrography of chromophoric dissolved organic matter in the North Atlantic.  
23 Deep-Sea Research I 54, 710-731.

- 1 Nelson, N.B., Siegel, D.A., Michaels, A.F., 1998. Seasonal dynamics of colored  
2 dissolved material in the Sargasso Sea. *Deep-Sea Research I* 45, 931-957.
- 3 Nieto-Cid, M., Álvarez-Salgado, X.A., Gago, J., Pérez, F.F., 2005. DOM fluorescence, a  
4 tracer for biogeochemical processes in a coastal upwelling system (NW Iberian  
5 Peninsula). *Marine Ecology Progress Series* 297, 33-50.
- 6 Nieto-Cid, M., Álvarez-Salgado, X.A., Pérez, F.F., 2006. Microbial and photochemical  
7 reactivity of fluorescent dissolved organic matter in a coastal upwelling system.  
8 *Limnology and Oceanography* 51, 1391-1400.
- 9 Nishioka, J., Ono, T., Saito, H., Nakatsuka, T., Takeda, S., Yoshimura, T., Suzuki, K.,  
10 Kuma, K., Nakabayashi, S., Tsumune, D., Mitsudera, H., Jhonson, W.K., Tsuda, A.,  
11 2007. Iron supply to the western subarctic Pacific: Importance of iron export from the  
12 Sea of Okhotsk. *Journal of Geophysical Research* 112, C10012.
- 13 Ogawa, H., Tanoue, E., 2003. Dissolved organic matter in oceanic waters. *Journal of*  
14 *Oceanography* 59, 129-147.
- 15 Parlanti, E., Wörz, K., Geoffroy, L., Lamotte, M., 2000. Dissolved organic matter  
16 fluorescence spectroscopy as a tool to estimate biological activity in a coastal zone  
17 submitted to anthropogenic inputs. *Organic Geochemistry* 31, 1765-1781.
- 18 Repeta, D.J., Aluwihare, L. I., 2006. Radiocarbon analysis of neutral sugars in high-  
19 molecular-weight dissolved organic carbon: Implications for organic carbon cycling.  
20 *Limnology and Oceanography* 51, 1045-1053.
- 21 Rochelle-Newall, E.J., Fisher, T.R., 2002. Production of chromophoric dissolved organic  
22 matter fluorescence in marine and estuarine environments: an investigation into the  
23 role of phytoplankton. *Marine Chemistry* 77, 7-21.

- 1 Stedmon, C.A., Bro, R., 2008. Characterizing dissolved organic matter fluorescence with  
2 parallel factor analysis: a tutorial. *Limnology and Oceanography: Methods* 6, 572-579.
- 3 Stedmon, C.A., Markager, S., 2005a. Resolving the variability in dissolved organic  
4 matter fluorescence in a temperate estuary and its catchment using PARAFAC  
5 analysis. *Limnology and Oceanography* 50, 686-697.
- 6 Stedmon, C.A., Markager, S., 2005b. Tracing the production and degradation of  
7 autochthonous fractions of dissolved organic matter by fluorescence analysis.  
8 *Limnology and Oceanography* 50, 1415-1426.
- 9 Stedmon, C.A., Markager, S., Bro, R., 2003. Tracing dissolved organic matter in aquatic  
10 environments using a new approach to fluorescence spectroscopy. *Marine Chemistry*  
11 82, 239-254.
- 12 Stedmon, C.A., Thomas, D.N., Granskog, M., Kaartokallio, H., Papadimitriou, S., Kuosa,  
13 H., 2007. Characteristics of dissolved organic matter in Baltic coastal sea ice:  
14 Allochthonous or autochthonous origins? *Environmental Science and Technology* 41,  
15 7273-7279.
- 16 Spencer, R.G.M., Bolton, L., Baker, A., 2007. Freeze/thaw and pH effects on freshwater  
17 dissolved organic matter fluorescence and absorbance properties from a number of  
18 UK locations. *Water Research* 41, 2941-2950.
- 19 Takata, H., Kuma, K., Iwade, S., Isoda, Y., Kuroda, H., Senjyu, T., 2005. Comparative  
20 vertical distribution of iron in the Japan Sea, the Bering Sea, and the western North  
21 Pacific Ocean. *Journal of Geophysical Research* 110, C07004.
- 22 Takata, H., Kuma, K., Iwade, S., Yamajoh, Y., Yamaguchi, A., Takagi, S., Sakaoka, K.,  
23 Yamashita, Y., Tanoue, E., Midorikawa, T., Kimura, K., Nishioka, J., 2004. Spatial

1        variability of iron in the surface water of the northwestern North Pacific Ocean.  
2        *Marine Chemistry* 86, 139-157.

3        Tani, H., Nishioka, J., Kuma, K., Takata, H., Yamashita, Y., Tanoue, E., Midorikawa, T.,  
4        2003. Iron(III) hydroxide solubility and humic-type fluorescent organic matter in the  
5        deep water column of the Okhotsk Sea and the northwestern North Pacific Ocean.  
6        *Deep-Sea Research I* 50, 1063-1078.

7        Williams, P.M., Druffel, E.R.M., 1987. Radiocarbon in dissolved organic matter in the  
8        central North Pacific Ocean. *Nature* 330, 246-248.

9        Yamashita, Y., Jaffé, R., 2008. Characterizing the interactions between trace metals and  
10        dissolved organic matter using excitation-emission matrix and parallel factor  
11        analysis. *Environmental Science and Technology* 42, 7374-7379.

12        Yamashita, Y., Tanoue, E., 2003. Chemical characterization of protein-like fluorophores  
13        in DOM in relation to aromatic amino acids. *Marine Chemistry* 82, 255-271.

14        Yamashita, Y., Tanoue, E., 2004a. Chemical characteristics of amino acid-containing  
15        dissolved organic matter in seawater. *Organic Geochemistry* 35, 679-692.

16        Yamashita and Tanoue, 2004b. In situ production of chromophoric dissolved organic  
17        matter in coastal environments. *Geophysical Research Letter* 31, L14302.

18        Yamashita, Y., Tanoue, E., 2008. Production of bio-refractory fluorescent dissolved  
19        organic matter in the ocean interior. *Nature Geoscience* 1, 579-582.

20        Yamashita, Y., Tanoue, E., 2009. Basin scale distribution of chromophoric dissolved  
21        organic matter in the Pacific Ocean. *Limnology and Oceanography* 54, 598-609.

22        Yamashita, Y., Jaffé, R., Maie, N., Tanoue, E., 2008. Assessing the dynamics of  
23        dissolved organic matter (DOM) in coastal environments by excitation and emission

1 matrix fluorescence and parallel factor analysis (EEM-PARAFAC). *Limnology and*  
2 *Oceanography* 53, 1900-1908.

3 Yamashita, Y., Tsukasaki, A., Nishida, T., Tanoue, E., 2007. Vertical and horizontal  
4 distribution of fluorescent dissolved organic matter in the Southern Ocean. *Marine*  
5 *Chemistry* 106, 498-509.

6 Yasuda, I., 2004. North Pacific Intermediate Water: Progress in SAGE (SubArctic Gyre  
7 Project) and related projects. *Journal of Oceanography* 60, 385-395.

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

1 **Figure captions**

2 Figure 1. Sampling locations in the Okhotsk Sea (Stns A and B) and the northwestern  
3 North Pacific Ocean (Stns C and D).

4

5 Figure 2. Examples of EEMs from the surface water (20 m at Stn A) and the deep water  
6 (1500 m at Stn C).

7

8 Figure 3. Spectral characteristics and validation of four component model by  
9 PARAFAC. The grey lines in right columns show the results of split half validation.

10

11 Figure 4. Vertical profiles of two humic-like components (components 1 and 2), one  
12 protein-like component (component 4), and ratio of two humic-like components in the  
13 Okhotsk Sea (Stns A and B) and the northwestern North Pacific Ocean (Stns C and D).  
14 The mesopelagic layer (200 -1000 m) was shaded.

15

16 Figure 5. Relationships between AOU and humic-like components 1 and 2 in the  
17 mesopelagic and bathypelagic layers. Solid lines indicate linear relationships between  
18 AOU and humic-like components 1 and 2 in the bathypelagic layer, respectively.

19

20 Figure 6. Relationships between Fe(III) solubility and humic-like components 1 and 2 in  
21 the mesopelagic layer.

Table 1. Characteristics of four components derived from PARAFAC model compared with those previous studies<sup>a</sup>

Component	Excitation maximum (nm)	Emission maximum (nm)	Coble (1996)	Stedmon and Markager (2005a) <sup>b</sup>	Cory and McKnight (2005)	Yamashita et al. (2008) <sup>d</sup>	Murphy et al. (2008) <sup>d</sup>
1	<260 (370)	466	A/C	4 (Ter/Aut)	SQ2 <sup>c</sup>	-	C3/P3 (Ter)
2	325 (<260)	385	M	6 (Ant)	C3 <sup>c</sup>	C6 (Mar)	C2/P1 (Mar)
3	<260	-	-	-	-	-	P4
4	275	306	B	8 (Tyr-like)	Tyr-like <sup>c</sup>	C7 (Tyr-like)	C1/P5 (Tyr-like)

<sup>a</sup> symbols and acronyms used for PARAFAC components previously reported could be found in cited literature.

<sup>b</sup> Ter, Aut and Ant means origin of terrestrial, autochthonous and anthropogenic, respectively.

<sup>c</sup> were identified in the Antarctic data set only (Cory and McKnight, 2005), indicating microbial origin.

<sup>d</sup> Ter and Mar means terrestrial and marine humic-like components, respectively.






Figure 1. Yamashita et al




Figure 2. Yamashita et al




Figure 3. Yamashita et al




Figure 4. Yamashita et al




Figure 5. Yamashita et al




Figure 6. Yamashita et al