


## Fluoride anion binding by cyclic boronic esters: influence of backbone chelate on receptor integrity

Christopher Bresner, Joanna K. Day, Natalie D. Coombs, Ian A. Fallis, Simon Aldridge,  
Simon J. Coles, and Michael B. Hursthouse.

### Supporting Information

1.	WinEQNMR fit of $^1\text{H}$ NMR titration data and determination of K for <b>4</b>	2
2.	ES(-) mass spectra for $[\mathbf{4}\cdot\text{F}]^-$	4
3.	Cyclic voltametry traces for $(\text{FcB})_2\text{calix}$ ( <b>6</b> ) with/without fluoride	6
4.	$^1\text{H}$ NMR spectra for the reaction of $(\text{PhB})_2\text{calix}$ ( <b>8</b> ) with $[\text{nBu}_4\text{N}]\text{F}\cdot x\text{H}_2\text{O}$ and with water	8
5.	Crystallographic data for $\text{FcBO}_2\text{C}_2\text{H}_4$ ( <b>3a</b> )	10
6.	Crystallographic data for $\text{fc}(\text{BO}_2\text{C}_2\text{H}_4)_2$ ( <b>3b</b> )	11
7.	Crystallographic data for $\text{fcB}_2\text{calix}$ ( <b>7</b> )	12

1. WinEQNMR fit of  $^1\text{H}$  NMR titration data and determination of K for  $\text{FcBO}_2\text{C}_2\text{H}_2\text{Ph}_2$  (4)



Calculations by WinEQNMR Version 1.20 by Michael J. Hynes  
 Program run at 13:44:34 on 03/09/2006

DATA FOR 1:1 COMPLEX USING CHEMICAL SHIFT (Titration.FIT)

Reaction: FcB(OR)2 + F- = [FcB(OR)2F]-

FILE: titration.FIT

IDEAL DATA: K1 = 30; DELTA FcB(OR)2 = 5.32; DELTA complex = 4.35

File prepared by S. Aldridge, February 01 2006

NO.	A	PARAMETER	DELTA	ERROR	CONDITION	DESCRIPTION
1	1	3.57930E+01	1.000E+00	9.817E+00	4.399E+01	K1
2	1	5.37827E+00	5.000E-02	1.044E-02	1.761E+00	SHIFT FcB(OR)2
3	1	4.36904E+00	5.000E-02	5.889E-02	4.049E+01	SHIFT complex

ORMS ERROR = 1.59E-02 MAX ERROR = 2.69E-02 AT OBS.NO. 9

RESIDUALS SQUARED = 2.02E-03

RFACTOR = 0.2746 PERCENT


NO.	A	EXPT. DEL	CALC. DEL	RESIDUAL	% DEV	WEIGHT	Fluoride
FcB(OR)2							
pH							
1	1	5.3041E+00	5.3158E+00	-1.1693E-02	-2.2046E-01	1.0000E+00	7.8700E-03
9.7664E-02		0.0000E+00					
2	1	5.2514E+00	5.2524E+00	-1.0300E-03	-1.9613E-02	1.0000E+00	1.6130E-02
9.7664E-02		0.0000E+00					
3	1	5.2013E+00	5.1906E+00	1.0710E-02	2.0591E-01	1.0000E+00	2.4510E-02
9.7664E-02		0.0000E+00					
4	1	5.1283E+00	5.1314E+00	-3.1276E-03	-6.0987E-02	1.0000E+00	3.2890E-02
9.7664E-02		0.0000E+00					
5	1	5.0326E+00	5.0236E+00	9.0141E-03	1.7911E-01	1.0000E+00	4.9400E-02
9.7664E-02		0.0000E+00					
6	1	4.9249E+00	4.9271E+00	-2.2011E-03	-4.4693E-02	1.0000E+00	6.6170E-02
9.7664E-02		0.0000E+00					
7	1	4.8019E+00	4.7782E+00	2.3716E-02	4.9389E-01	1.0000E+00	9.9190E-02
9.7664E-02		0.0000E+00					
8	1	4.7221E+00	4.7222E+00	-6.1989E-05	-1.3127E-03	1.0000E+00	1.1557E-01
9.7664E-02		0.0000E+00					
9	1	4.6503E+00	4.6772E+00	-2.6946E-02	-5.7945E-01	1.0000E+00	1.3157E-01
9.7664E-02		0.0000E+00					
10	1	4.5966E+00	4.6096E+00	-1.2972E-02	-2.8222E-01	1.0000E+00	1.6396E-01
9.7664E-02		0.0000E+00					
11	1	4.5683E+00	4.5537E+00	1.4612E-02	3.1985E-01	1.0000E+00	2.0498E-01
9.7664E-02		0.0000E+00					


TOLERANCE ON SUM OF SQUARES 0.0300

TOLERANCE ON EIGEN VALUES 0.0001


CONVERGANCE AFTER 2 ITERATIONS

## 2. ES(-) mass spectra for [4F]<sup>-</sup>






3. Cyclic voltametry traces for (FcB)<sub>2</sub>calix (6)




Black: CV for 6; grey: CV for 6 plus trace fluoride.



Black: CV for [FcBF<sub>3</sub>]; grey: CV for **6** plus excess fluoride.

4.  $^1\text{H}$  NMR spectra for the reaction of  $(\text{PhB})_2\text{calix}$  (**8**) with  $[\text{nBu}_4\text{N}]\text{F}\cdot x\text{H}_2\text{O}$



Top:  $(\text{PhB})_2\text{calix}$  (**8**) in  $\text{CDCl}_3$


Middle:  $(\text{PhB})_2\text{calix}$  (**8**) plus excess  $[\text{nBu}_4\text{N}]\text{F}\cdot x\text{H}_2\text{O}$  in  $\text{CDCl}_3$

Bottom:  $\text{H}_4(\text{calix})$  plus excess  $[\text{nBu}_4\text{N}]\text{F}\cdot x\text{H}_2\text{O}$  in  $\text{CDCl}_3$

The resonance indicated with the asterisk is due to the  $\text{NCH}_2$  protons of the  $[\text{nBu}_4\text{N}]^+$  counter-ion and is strongly dependent on concentration and ionic strength.



**$^1\text{H}$  NMR spectra for the reaction of  $(\text{PhB})_2\text{calix}$  (**8**) with water**




Top:  $(\text{PhB})_2\text{calix}$  (**8**) in  $\text{CDCl}_3$

Top Middle:  $(\text{PhB})_2\text{calix}$  (**8**) plus excess water in  $\text{CDCl}_3$

Bottom Middle:  $\text{PhB}(\text{OH})_2$  in  $\text{CDCl}_3$

Bottom:  $\text{H}_4(\text{calix})$  in  $\text{CDCl}_3$


## 5. Crystallographic data for $\text{FcBO}_2\text{C}_2\text{H}_4$ (**3a**)



Crystal data and structure refinement for **3a**:

Empirical formula	C <sub>12</sub> H <sub>13</sub> B Fe O <sub>2</sub>	
Formula weight	255.88	
Temperature	150(2) K	
Wavelength	0.71073 Å	
Crystal system	Orthorhombic	
Space group	<i>p</i> 21 21 21	
Unit cell dimensions	<i>a</i> = 5.81790(10) Å	$\alpha = 90^\circ$
	<i>b</i> = 9.9233(2) Å	$\beta = 90^\circ$
	<i>c</i> = 18.4741(5) Å	$\gamma = 90^\circ$
Volume	1066.56(4) Å <sup>3</sup>	
<i>Z</i>	4	
Density (calculated)	1.594 Mg/m <sup>3</sup>	
Absorption coefficient	1.391 mm <sup>-1</sup>	
<i>F</i> (000)	528	
Crystal size	0.60 x 0.25 x 0.20 mm <sup>3</sup>	
Theta range for data collection	3.67 to 26.36°	
Index ranges	-5 ≤ <i>h</i> ≤ 7, -12 ≤ <i>k</i> ≤ 12, -23 ≤ <i>l</i> ≤ 23	
Reflections collected	8431	
Independent reflections	2138 [ <i>R</i> (int) = 0.0670]	
Completeness to theta = 26.36°	99.7 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.7684 and 0.4892	
Refinement method	Full-matrix least-squares on <i>F</i> <sup>2</sup>	
Data / restraints / parameters	2138 / 0 / 145	
Goodness-of-fit on <i>F</i> <sup>2</sup>	1.029	
Final <i>R</i> indices [ <i>I</i> > 2σ( <i>I</i> )]	<i>R</i> 1 = 0.0328, <i>wR</i> 2 = 0.0795	
<i>R</i> indices (all data)	<i>R</i> 1 = 0.0347, <i>wR</i> 2 = 0.0812	
Absolute structure parameter	0.03(2)	
Largest diff. peak and hole	0.332 and -0.681 e.Å <sup>-3</sup>	

## 6. Crystallographic data for $\text{fc}(\text{BO}_2\text{C}_2\text{H}_4)_2$ (**3b**)



Crystal data and structure refinement for **3b**:

Empirical formula	C <sub>14</sub> H <sub>16</sub> B <sub>2</sub> Fe O <sub>4</sub>	
Formula weight	325.74	
Temperature	150(2) K	
Wavelength	0.71073 Å	
Crystal system	Monoclinic	
Space group	C 2/m	
Unit cell dimensions	a = 7.0116(4) Å b = 10.0523(7) Å c = 9.8912(7) Å	α = 90° β = 104.211(3)° γ = 90°
Volume	675.82(8) Å <sup>3</sup>	
Z	2	
Density (calculated)	1.601 Mg/m <sup>3</sup>	
Absorption coefficient	1.126 mm <sup>-1</sup>	
F(000)	336	
Crystal size	0.25 x 0.20 x 0.08 mm <sup>3</sup>	
Theta range for data collection	3.62 to 26.33°	
Index ranges	-8 ≤ h ≤ 8, -10 ≤ k ≤ 12, -12 ≤ l ≤ 12	
Reflections collected	1901	
Independent reflections	740 [R(int) = 0.0634]	
Completeness to theta = 26.33°	99.7 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.9153 and 0.7661	
Refinement method	Full-matrix least-squares on F <sup>2</sup>	
Data / restraints / parameters	740 / 0 / 53	
Goodness-of-fit on F <sup>2</sup>	1.097	
Final R indices [I > 2σ(I)]	R1 = 0.0449, wR2 = 0.0934	
R indices (all data)	R1 = 0.0535, wR2 = 0.0966	
Largest diff. peak and hole	0.606 and -0.547 e.Å <sup>-3</sup>	

## 7. Crystallographic data for fcB<sub>2</sub>calix (7)




Table 1. Crystal data and structure refinement for 7:

Empirical formula	C <sub>57</sub> H <sub>67</sub> B <sub>2</sub> Fe O <sub>4</sub>	
Formula weight	893.58	
Temperature	150(2) K	
Wavelength	0.71073 Å	
Crystal system	Triclinic	
Space group	P -1	
Unit cell dimensions	a = 10.7685(3) Å	α = 89.514(2)°.
	b = 12.7505(4) Å	β = 73.968(2)°.
	c = 19.0095(6) Å	γ = 76.0250(10)°.
Volume	2429.57(13) Å <sup>3</sup>	
Z	2	
Density (calculated)	1.221 Mg/m <sup>3</sup>	
Absorption coefficient	0.356 mm <sup>-1</sup>	
F(000)	954	
Crystal size	0.38 x 0.20 x 0.13 mm <sup>3</sup>	
Theta range for data collection	2.74 to 27.21°.	
Index ranges	-13 ≤ h ≤ 13, -16 ≤ k ≤ 16, -23 ≤ l ≤ 24	
Reflections collected	37381	
Independent reflections	10716 [R(int) = 0.3243]	
Completeness to theta = 27.21°	98.7 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.9551 and 0.8764	
Refinement method	Full-matrix least-squares on F <sup>2</sup>	
Data / restraints / parameters	10716 / 6 / 587	
Goodness-of-fit on F <sup>2</sup>	1.025	
Final R indices [I > 2σ(I)]	R1 = 0.1237, wR2 = 0.3054	
R indices (all data)	R1 = 0.1874, wR2 = 0.3554	
Extinction coefficient	0.017(4)	
Largest diff. peak and hole	2.075 and -2.014 e.Å <sup>-3</sup>	