

Focus-formation of replication protein A, activation of checkpoint system and DNA repair synthesis induced by DNA double-strand breaks in *Xenopus* egg extract

Takayuki Kobayashi¹, Shusuke Tada^{1,*}, Takashi Tsuyama¹, Hiromu Murofushi², Masayuki Seki¹ and Takemi Enomoto¹

¹Molecular Cell Biology Laboratory, Graduate School of Pharmaceutical Sciences, Tohoku University, Aoba-ku, Sendai, Miyagi 980-8578, Japan

²Department of Biophysics and Biochemistry, Graduate School of Science, University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan

*Author for correspondence (e-mail: tada@mail.pharm.tohoku.ac.jp)

Accepted 12 May 2002

Journal of Cell Science 115, 3159-3169 (2002) © The Company of Biologists Ltd

Summary

The response to DNA damage was analyzed using a cell-free system consisting of *Xenopus* egg extract and demembrated sperm nuclei. In the absence of DNA-damaging agents, detergent-resistant accumulation of replication protein A appeared in nuclei after a 30 minute incubation, and a considerable portion of the replication protein A signals disappeared during a further 30 minute incubation. Similar replication protein A accumulation was observed in the nuclei after a 30 minute incubation in the extract containing camptothecin, whereas a further 30 minute incubation generated discrete replication protein A foci. The addition of camptothecin also induced formation of γ -H2AX foci, which have been previously shown to localize at sites of DSBs. Analysis of the time course of DNA replication and results obtained using geminin, an inhibitor of licensing for DNA replication, suggest that the discrete replication protein A foci formed in response to camptothecin-induced DNA damage occur in a DNA-replication-dependent manner. When the nuclei were incubated in the extract containing *EcoRI*, discrete

replication protein A foci were observed at 30 minutes as well as at 60 and 90 minutes after incubation, and the focus-formation of replication protein A was not sensitive to geminin. DNA replication was almost completely inhibited in the presence of *EcoRI* and the inhibition was sensitive to caffeine, an inhibitor of ataxia telangiectasia mutated protein (ATM) and ATM- and Rad3-related protein (ATR). However, the focus-formation of replication protein A in the presence of *EcoRI* was not influenced by caffeine treatment. *EcoRI*-induced incorporation of biotin-dUTP into chromatin was observed following geminin-mediated inhibition of DNA replication, suggesting that the incorporation was the result of DNA repair. The biotin-dUTP signal co-localized with replication protein A foci and was not significantly suppressed or stimulated by the addition of caffeine.

Key words: Double-strand DNA break, DNA repair, Checkpoint, *Xenopus* egg extract, Replication protein A, Camptothecin, *EcoRI*

Introduction

DNA double-strand breaks (DSBs) are a serious threat to living cells, and are thought to be caused by ionizing radiation, certain chemotherapeutic drugs and mechanical stress as well as stalled replication forks (Haber, 2000; Karran, 2000; Khanna and Jackson, 2001). Once cells recognize lesions on the DNA, they activate a series of pathways to repair the damage and to arrest the cell cycle in order to gain time for the repair. Although tremendous progress has been made in recent years in elucidating the molecular basis of these processes, there is still a need to clarify the connection between recognition of the DNA lesions, repair and the cellular responses such as checkpoint controls or apoptosis.

Replication protein A (RPA) is the eukaryotic single-stranded DNA binding protein complex, composed of three subunits (Wold, 1997). The complex was originally identified as a factor participating in DNA replication of a cell-free simian virus 40 (SV40) replication system (Fairman and Stillman, 1988; Wobbe et al., 1987; Wold and Kelly, 1988). Observations have since indicated that it functions in many aspects of DNA metabolism

such as DNA repair, recombination and transcription as well as DNA replication (Wold, 1997). Functions of RPA in response to DNA damage have also been suggested. Ataxia telangiectasia-mutated protein kinase (ATM) and DNA-dependent protein kinase (DNA-PK), which are involved in the primary response to DNA damage (Durocher and Jackson, 2001; Shiloh, 2001), phosphorylate RPA in a DNA-damage-dependent manner (Boubnov and Weaver, 1995; Oakley et al., 2001; Shao et al., 1999).

Moreover, RPA has been suggested to associate with the damaged locus on the chromatin. In addition to this association with UV-damaged DNA (Burns et al., 1996; Lao et al., 2000), a number of reports have suggested a link between RPA and DSB repair. Rad51 protein, the eukaryotic homologue of *Escherichia coli* RecA recombinase, formed foci that increased in number in response to several genotoxic treatments, including ionizing radiation (Haaf et al., 1995). In the cells in which Rad51 foci formed after ionizing radiation, RPA also formed foci, most of which co-localized with the Rad51 foci and sites of DNA repair (Golub et al., 1998; Haaf et al., 1999).

In addition, localization of BLM and WRN in the nuclear foci containing Rad51 and RPA was also reported (Bischof et al., 2001; Sakamoto et al., 2001). BLM and WRN are eukaryotic DNA helicases homologous to *Escherichia coli* recombination-associated DNA helicase, RecQ, and mutations in these genes result in Bloom and Werner syndromes, respectively, which are genetic disorders characterized by genomic instability (Enomoto, 2001; Mohaghegh and Hickson, 2001). These findings suggest that the RPA-Rad51 foci are sites of DNA repair by homologous recombination.

The cell-free system established using extracts derived from *Xenopus* eggs is useful for biochemical and cytological analysis of the dynamic processes involved in the cell cycle (Blow, 2001; Lohka and Masui, 1983). Because the extract is made from freshly laid eggs that are synchronized at metaphase of meiosis II, upon addition of sperm chromatin, subsequent cell cycle events, such as DNA replication, progress highly synchronously in the extract. Therefore, this system has been widely used in analyses of the mitotic process, the control of chromosomal DNA replication, DNA replication checkpoints, etc. Recently, several attempts have been made using this cell-free system to examine the checkpoint system in response to DNA damage (Costanzo et al., 2000; Guo and Dunphy, 2000; Guo et al., 2000) and the repair of DSBs by nonhomologous end joining (Labhart, 1999). Hence, the cell-free system offers a useful means to analyse the dynamic processes in response to and repair of DSBs.

In studies using the *Xenopus* cell-free system, focus-formation of RPA was implicated in the initiation of DNA replication (Adachi and Laemmli, 1992; Adachi and Laemmli, 1994; Chen et al., 2001). The RPA foci in the nuclei that formed in *Xenopus* egg extracts were originally identified as components of DNA pre-replication centers (Adachi and Laemmli, 1992; Adachi and Laemmli, 1994). Focus forming activity 1 (FFA-1), a *Xenopus* homologue of WRN (Yan et al., 1998), was identified as a protein supporting the formation of RPA foci (Yan and Newport, 1995), and an analysis using the dominant-negative form of FFA-1 suggested a function for FFA-1 in DNA replication (Chen et al., 2001). Thus, focus-formation of RPA in *Xenopus* egg extracts has been mainly analyzed in terms of initiation of DNA replication, although its occurrence in response to DNA damage was also reported (Tchang and Mechali, 1999).

The cellular response to DSBs is a complicated linkage of cellular processes, such as recognition of DSBs, activation of checkpoint pathways, selection of an appropriate DNA repair system, repair of lesions and restoration of normal cell cycle events. To understand the molecular basis behind these dynamic cellular responses, we have employed a cell-free system consisting of extract derived from *Xenopus* eggs. In the present study, we were able to reproduce several important steps in the cellular response to DSBs. As a result, we demonstrated focus-formation of RPA in response to DSBs, inhibition of DNA replication by checkpoint pathways, and DNA lesion-induced DNA synthesis that was co-localized with RPA foci.

Materials and Methods

Preparation of *Xenopus* egg extract and demembrated sperm

Metaphase arrested cell-free extracts derived from *Xenopus* eggs were

prepared as described (Chong et al., 1997). In brief, freshly laid *Xenopus* eggs were dejellied in a 2% cysteine solution, pH 7.6, containing 1 mM EGTA, and rinsed with 15 mM Tris-HCl, pH 7.6, 110 mM NaCl, 2 mM KCl, 1 mM MgSO₄, 2 mM NaHCO₃, 0.5 mM Na₂HPO₄ and 2 mM EGTA, and then with 50 mM Hepes-KOH, pH 7.6, 50 mM KCl, 5 mM MgCl₂, 2 mM β-mercaptoethanol and 5 mM EGTA. Then, the eggs were crushed by centrifugation (10,000 g, 10 minutes). The cytoplasmic fraction was cleared by centrifugation (84,000 g, 20 minutes) after an addition of cytochalasin B (10 μg/ml). The resultant extract was supplemented with 1% glycerol and stored in liquid nitrogen.

When required, the extracts were released into interphase by addition of 0.3 mM CaCl₂ and incubated for 15 minutes at 23°C. The released extract was then supplemented with 250 μg/ml of cycloheximide, 25 mM phosphocreatin and 15 μg/ml of creatin phosphokinase.

Xenopus sperm nuclei were prepared after demembration by lysolecithin as described (Chong et al., 1997) and stored at -80°C until use.

Preparation of anti-RPA antiserum

Xenopus egg extract was diluted with a buffer consisting of 50 mM Hepes-KOH, pH 7.4, 5 mM MgCl₂, 2 mM β-mercaptoethanol and 0.5 M NaCl. After an addition of ammonium sulfate (0.21 g/ml), precipitates were collected by centrifugation and dissolved in buffer-1 (20 mM Tris-HCl, pH 7.5, 1 mM EDTA, 0.2 M NaCl, 10% glycerol, 0.2 mM DTT, 0.1 mM phenylmethylsulfonyl fluoride and 0.1 mg/ml each of pepstatin, leupeptin and chymostatin) followed by dialysis against the same buffer. The dialysate was loaded onto a DEAE-cellulose column. After washing the column with buffer-1 containing 0.05 M NaCl, RPA was eluted with a linear gradient of 0.05-1 M NaCl in the same buffer. RPA was eluted at 0.35 M NaCl. The RPA fractions were pooled and loaded onto a single-stranded DNA column. After washing the column with buffer-1 containing 0.5 M NaCl, RPA was eluted from the column with the same buffer containing 2 M NaCl.

The purified *Xenopus* RPA was electrophoresed on SDS-polyacrylamide gels. After brief staining of the gels with Coomassie Brilliant Blue, the protein band of the 32 kDa subunit was dissected and homogenized in phosphate-buffered saline (PBS; 137 mM NaCl, 2.68 mM KCl, 8.04 mM Na₂HPO₄, 1.47 mM KH₂PO₄). Gel homogenate was mixed with complete or incomplete adjuvant and injected into rabbits to prepare antiserum against *Xenopus* RPA 32 kDa subunit.

Immunofluorescence microscopy

Preparation of coverslips

Demembrated sperm nuclei (10,000 sperm heads) were incubated in 10 μl of *Xenopus* egg extract at 23°C for given periods. During the incubation, 0.05 units/μl of *Eco*RI (New England BioLabs), 25 μM camptothecin (Sigma), 5 mM caffeine (Wako, Osaka, Japan), and 3 μg/ml of geminin were added to the reaction mixture as indicated. For experiments to detect DNA synthesis, 10 μM biotin-16-dUTP (Roche) was also added to the extract prior to the incubation. Then, the nuclei that formed around the sperm chromatin were fixed by addition of 90 μl of ice-cold extraction buffer (EB; 50 mM KCl, 50 mM Hepes-KOH, pH 7.6, 5 mM MgCl₂) containing 3.7% formaldehyde and the fixed mixture was diluted with 90 μl of ice-cold EB containing 0.5% Triton X-100. After incubation on ice for 2 minutes, the sample was supplemented with 10 μl of 37% formaldehyde and further incubated for 10 minutes on ice. The mixture was then overlaid onto EB containing 30% sucrose and fixed on glass coverslips by centrifugation (400 g for 20 minutes).

Observation of RPA and phosphorylated-H2AX (γ-H2AX) accumulation in nuclei

Each coverslip mounted with sample was rinsed three times with

0.05% Tween-20 in PBS, and reacted with rabbit anti-*Xenopus* RPA antisera or antibody against phosphorylated peptide derived from γ -H2AX (Trevigen Inc) diluted in 0.05% Tween-20-PBS supplemented with 10% nonfat milk for 2 hours at room temperature or overnight at 4°C. Next, the coverslip was rinsed three times with 0.05% Tween-20-PBS and incubated with FITC-labelled donkey anti-rabbit IgG (Jackson Immuno Research Laboratories). It was then rinsed once with 0.05% Tween 20-PBS, incubated for 5 minutes in 0.05% Tween 20-PBS containing 20 μ g/ml of propidium iodide (Wako), and rinsed three times with 0.05% Tween 20-PBS and once with PBS. The coverslip was mounted on a glass slide with PermaFluor Aqueous Mountant (Thermo Shandon, Pittsburgh, PA).

Observation of DNA synthesis

The preparation used for experiments to detect DNA synthesis with biotin-dUTP was similar to that for the detection of RPA signals with a few changes. After fixation onto a coverslip and incubation with anti-RPA antisera, the coverslip was incubated with Texas Red-labelled anti-rabbit IgG (H+L) (Vector Laboratories, Burlingame CA) and Fluorescein-labelled streptavidin (Vector Laboratories). It was then rinsed twice with 0.05% Tween-20-PBS and once with PBS. The coverslip was mounted on a glass slide with PermaFluor Aqueous Mountant (Thermo Shandon).

Microscopic observations

The prepared samples were observed using a Laser Scanning Confocal Imaging System, MRC-1024 (Bio-Rad). A fluorescence microscope (Leica) was also used for experiments to count numbers of discrete RPA foci or focus-positive nuclei. We confirmed that negative controls using non-immune serum generate no or very little signal upon these observations (data not shown).

Isolation of detergent-insoluble nuclear fraction

Demembrated sperm nuclei (8000 sperm heads) were incubated in 8 μ l of *Xenopus* egg extract at 23°C for indicated periods. The extract was diluted in 1 ml of nuclear isolation buffer (NIB; 50 mM Hepes-KOH, pH 7.6, 50 mM KCl, 2 mM DTT, 0.5 mM spermidine, 0.15 mM spermine, 1 μ g/ml leupeptine and 1 μ g/ml pepstatin) supplemented with 0.1% Triton X-100 and 2.5 mM ATP (TNIBA), and then underlayered with the same buffer containing 15% sucrose. The nuclei were precipitated at 9000 *g* in a swinging bucket centrifuge for 5 minutes at 4°C. After washing with 1 ml TNIBA, the nuclear precipitate was subjected to 15% SDS-polyacrylamide gel electrophoresis for immunoblotting or micrococcal nuclease treatment.

Micrococcal nuclease (MNase) treatment of detergent-insoluble nuclear fraction

Isolated nuclear fraction was suspended in 12 μ l of TNIBA supplemented with 1 mM CaCl₂ and 0.2 units MNase, and incubated for 10 minutes at 37°C. EGTA (1 mM final concentration) was added to terminate the reaction and the mixture was separated into supernatant and precipitated fractions by centrifugation at 1300 *g* for 4 minutes at 4°C. The resultant fractions were subjected to 15% SDS-polyacrylamide gel electrophoresis for immunoblotting analysis.

Assay for DNA synthesis

DNA synthesizing activity was measured by the method as described (Chong et al., 1997), with slight modification. In brief, *Xenopus* egg extract containing 1.85 kBq/ μ l of [α -³²P]dCTP was supplemented with 0.05 units/ μ l *Eco*RI (New England BioLabs), 25 μ M camptothecin (Sigma), 5 mM caffeine (Wako), and 3 μ g/ml of geminin as indicated. The reaction was initiated by the addition of




Fig. 1. Characterization of anti-RPA p32 antibody by immunoblotting. 1 μ l of interphase *Xenopus* egg extract was subjected to SDS-electrophoresis on a 10% polyacrylamide gel. The electrophoresed proteins were transferred onto PVDF-membrane and the protein bands that reacted with the antibody were visualized as described in Materials and Methods.

demembrated sperm nuclei (1000 sperm heads/ μ l extract). After incubation at 23°C for a given period, the reaction was terminated and the mixture was digested by 0.2 mg/ml proteinase K in a buffer containing 20 mM Tris-HCl, pH 7.5, 5 mM EDTA, and 0.5% SDS. The digested mixture was then precipitated with cold 10% TCA, 2% Na₄P₂O₇-10H₂O and filtered through a 25 mm GF/C disc (Whatman). After washing with 5% TCA, 0.5% Na₄P₂O₇-10H₂O and then with ethanol, the filter was dried and immersed in water. The radioactivity on the filter was measured by detecting emission caused by the Cerenkov effect.

Immunoblotting

The samples for immunoblotting were electrophoresed on an SDS-polyacrylamide gel and electrically transferred onto Hybond-P PVDF-membrane (Amersham Pharmacia Biotech). The membrane was blocked with PBS containing 3% bovine serum albumin (BSA) and incubated with anti-RPA p32 antiserum diluted 3000-fold with 0.05% Tween-20 and 3% BSA in PBS. After washing with 0.05% Tween-20-PBS and incubating with horseradish peroxidase-conjugated anti-rabbit IgG, the protein bands that reacted with the anti-RPA antiserum were visualized with ECL western blotting detection reagents (Amersham Pharmacia Biotech).

Preparation of recombinant geminin

Recombinant geminin was expressed from geminin H cDNA lacking a destruction box (geminin^{DEL}) and affinity-purified as described (McGarry and Kirschner, 1998).

Results

Production of antibody against the 32 kDa subunit of *Xenopus* RPA

To observe the behavior of RPA in response to DNA-lesions in *Xenopus* egg extract, we raised an antibody against the middle-sized subunit of *Xenopus* RPA heterotrimer. *Xenopus* RPA was purified from *Xenopus* egg extract as described in Materials and Methods and the purified fraction was subjected to SDS-polyacrylamide gel electrophoresis. The protein band corresponding to the 32 kDa subunit was excised and rabbits were immunized. The obtained antiserum showed good reaction specificity towards 32 kDa protein in crude *Xenopus* egg extracts (Fig. 1).

Accumulation of RPA in the nuclei formed in *Xenopus* egg extract

To address the accumulation of RPA in the nuclei formed in




Fig. 2. Focus-formation of RPA in the nuclei incubated with *Xenopus* egg extract. (A) Demembrated *Xenopus* sperm nuclei were incubated at 23°C for 30 minutes (a-c), 60 minutes (d-f) or 90 minutes (g-i). RPA resistant to detergent treatment is shown in panels a, c, d, f, g and i. DNA was visualized using propidium iodide (b,e,h). Panels c, f and i are enlarged images of the nuclei indicated by arrowheads in panels a, d and g, respectively. The same nuclei in the 'DNA' panels are also indicated by arrowheads. Bars, 50 μ m (a,b,d,e,g,h); 20 μ m (c,f,i). (B) A detergent-insoluble nuclear fraction was prepared from the nuclei incubated in *Xenopus* egg extract for 30 minutes, 60 minutes or 90 minutes. The p32 subunit of RPA in the nuclear fraction was visualized by immunoblotting after SDS-polyacrylamide gel electrophoresis. (C) The detergent-insoluble nuclear fraction prepared from 30 minute incubated nuclei was incubated for 10 minutes at 37°C with (lanes 1,2) or without (lanes 3,4) MNase, and the reaction mixture was centrifuged. The resultant supernatant (lanes 1,3) and precipitated (lanes 2,4) fractions were subjected to SDS-polyacrylamide gel electrophoresis and immunoblotting.

Xenopus egg extracts, we first examined the behavior of RPA in the absence of DNA-damage-inducing agents by indirect immunofluorescence microscopy (Fig. 2A). The time course of nuclear formation indicated that rounded nuclei appeared around 30 minutes after addition of demembrated *Xenopus* sperm (data not shown), when DNA replication was about to start in the nuclei (Fig. 3C, closed circle). At this time, the accumulation of RPA occurred in most of the newly formed nuclei (Fig. 2Aa,b). An enlarged image of a representative nucleus indicated that RPA accumulated evenly in the nucleus (Fig. 2Ac). More sensitive observations revealed that the RPA signal appeared to form a large number of minute dots distributed in the whole nucleus (data not shown). However, a considerable portion of the RPA signals disappeared during a further 30 minute incubation (Fig. 2Ad,f), and did not reappear upon further incubation (Fig. 2Ag,i). When geminin, an

endogenous inhibitor of the licensing for DNA replication (McGarry and Kirschner, 1998; Tada et al., 2001), was added to the extract, the RPA accumulation observed in the damage non-inducing condition was suppressed almost completely (data not shown).

We then examined chromatin association of RPA under the same conditions by immunoblotting analysis. The 32 kDa band that reacted with the anti-RPA antibody was observed in the detergent-insoluble nuclear fraction isolated after a 30 minute incubation, and the intensity of the band was gradually reduced during further 30 and 60 minute incubations (Fig. 2B). This band was hardly observed in the chromatin fraction isolated before nuclear formation or in the nuclear fraction isolated after incubation in geminin-treated extract (data not shown). To confirm that the detected RPA was associated with chromatin, the nuclear fraction was isolated after a 30 minute incubation, briefly incubated with or without micrococcal nuclease (MNase), and separated into supernatant and precipitated fractions. After MNase treatment, RPA was detected in the supernatant fraction (Fig. 2C), indicating that RPA associates with chromatin.

These results suggest that the minute RPA dots observed here reflect the accumulation of RPA on pre-replication centers reported previously (Adachi and Laemmli, 1992), which form prior to DNA replication and are removed from the sperm chromatin during the progression of DNA replication.

Induction of RPA focus-formation by camptothecin

Next, we observed the behavior of RPA in the nuclei that formed in the extract containing a DNA damaging agent, camptothecin (CPT). CPT is an inhibitor of DNA topoisomerase I and is known to induce DSBs under certain conditions. The nuclei formed after a 30 minute incubation in the extract containing CPT were smaller than those formed in the absence of CPT (Fig. 3Ab; compare with Fig. 2Ab), possibly reflecting some influence of CPT on the remodeling of chromatin structure during the transition from tightly condensed sperm to nuclei. Although the enlargement of nuclei was slightly delayed in the presence of CPT, it occurred to almost the same degree as in the absence of CPT during a longer incubation (Fig. 3Ae,h).

In spite of the difference in nuclear size, RPA had accumulated in 97% of nuclei after 30 minute incubation (Fig. 3Aa). Enlarged images of representative nuclei indicated that the accumulation took place evenly in the nucleus (Fig. 3Ac). In contrast to the results obtained in the damage non-inducing condition, RPA-signals were observed in 98% and 94% of the nuclei after 60 and 90 minute incubation, respectively (Fig. 3Ad,g). In addition, the CPT-induced RPA signal observed after 60 and 90 minutes formed discrete foci in contrast to the RPA accumulation in the nuclei after 30 minutes (Fig. 3Af,i). Both the number of RPA-positive nuclei and the number of foci in the nucleus after 60 minute incubation were increased being dependent on the concentration of CPT in the extract (data not shown).

To examine whether the CPT-induced discrete RPA-foci were associated with chromatin, we prepared a detergent-insoluble nuclear fraction from nuclei incubated for 60 minutes in CPT-containing extract (Fig. 3B). The amount of RPA in the nuclear fraction was increased by CPT treatment, and RPA was




Fig. 3. Focus-formation of RPA and DNA replication in the nuclei incubated with *Xenopus* egg extract containing CPT. (A) Demembrated *Xenopus* sperm nuclei were incubated at 23°C, for the periods indicated, in the presence of CPT. RPA localization (a,c,d,f,g,i) and DNA (b,e,h) were visualized. Panels c, f and i are enlarged images of the nuclei indicated by arrowheads in panels a, d and g, respectively. The same nuclei in the 'DNA' panels are also indicated by arrowheads. Bars, 50 μ m (a,b,d,e,g,h); 20 μ m (c,f,i). Numbers of RPA-accumulated nuclei/observed nuclei from two independent experiments are shown below the panels. The percentages of RPA-accumulated nuclei are also presented in parentheses. (B) A detergent-insoluble nuclear fraction was prepared from the nuclei incubated at 23°C for 60 minutes in the absence (lane 1) or the presence (lanes 2-6) of 25 μ M CPT. The nuclear fraction was subjected to SDS-polyacrylamide gel electrophoresis directly (lanes 1,2), or after incubation with (lanes 3,4) or without (lanes 5,6) MNase followed by centrifugal separation of supernatant (lanes 3,5) and precipitated (lanes 4,6) fractions. The p32 subunit of RPA was visualized by immunoblotting. (C) The sperm nuclei were incubated in the extract at 23°C for the periods indicated in the presence of [α - 32 P]dCTP. Circles, squares and triangles represent DNA synthesis in the absence, and presence of CPT and CPT+caffeine, respectively. The extent of DNA synthesis is represented as a percentage of accumulative DNA synthesis after a 90 minute incubation under damage-noninducing conditions. 100% DNA synthesis is estimated to be an amount equivalent to the duplication of about 90% of input DNA. Open symbols indicate accumulative DNA synthesis during a 60 minute incubation in the presence of geminin. (D) The sperm nuclei were incubated in the extract containing CPT for 60 minutes at 23°C in the presence of geminin. Localization of RPA (a,c) and DNA (b) is shown. Panel c is an enlarged image of the nucleus indicated by an arrowhead in panel a. The same nucleus in panel b is also indicated by an arrowhead. Bars, 50 μ m (a,b); 20 μ m (c).

liberated from the nuclear fraction by brief treatment of MNase, indicating the association of RPA with chromatin.

Fig. 3C shows the time-course of DNA synthesis in the presence and absence of CPT. The level of DNA synthesis was lower in the presence of CPT and reached a plateau after 50 minute incubation, remaining at about 40% of the level attained in the absence of CPT. DNA synthesis in the presence or absence of CPT was almost completely inhibited by the presence of geminin, indicating that the observed DNA synthesis was due to DNA replication.

CPT is thought to induce DSBs in a DNA replication-dependent manner (Avemann et al., 1988; D'Arpa et al., 1990; Holm et al., 1989; Hsiang et al., 1989; Ryan et al., 1991). Thus, it is likely that the RPA foci that appeared after a 60 minute incubation in the presence of CPT are formed by generation of

DSBs and the formation of RPA foci is prevented by inhibition of DNA replication. This notion is further supported by the fact that geminin inhibited the focus-formation of RPA in the presence of CPT (Fig. 3D).

DSBs reportedly induce rapid phosphorylation of a serine residue within the C-terminus region of histone H2AX, one of three types of H2A histones (Rogakou et al., 1998; Rogakou et al., 1999). This phosphorylation is thought to be a reliable marker for the formation of DSBs because of the specificity of this reaction (Rogakou et al., 1999). To obtain evidence for DSB formation by the presence of CPT in our experimental conditions, we performed indirect immunofluorescence staining by using an antibody that specifically recognizes the phosphorylated form of H2AX (γ -H2AX). Formation of γ -H2AX foci was not observed in the absence of DNA damaging




Fig. 4. Formation of γ -H2AX foci in the nuclei incubated in *Xenopus* egg extract containing CPT. Demembrated sperm nuclei were incubated in *Xenopus* egg extract with (B) or without (A) CPT. γ -H2AX foci formation (a,c,e,g) and DNA (b,d,f,h) in the absence (a,b,c,d) or the presence (e,f,g,h) of geminin were visualized. Panels c, d, g and h are enlarged images of the nuclei indicated by arrowheads in panels a, b, e and f, respectively. Bars, 50 μ m (a,b,e,f); 20 μ m (c,d,g,h).

agents (Fig. 4A). By contrast, a large number of γ -H2AX foci were induced by CPT-treatment and the induction was suppressed almost completely by inhibiting DNA replication with geminin (Fig. 4B).

Induction of RPA focus-formation by *EcoRI*

Although our observations and previous reports strongly suggested that CPT caused DSBs in a DNA-replication-dependent manner, we have not succeeded in direct detection of the CPT-induced DSBs in our system. To eliminate the uncertainty of DSB formation by CPT and the complexity caused by the influence of DNA replication, we devised a simple and more direct way to introduce DSBs using restriction enzymes.

Fig. 5A shows the results obtained with the *Xenopus* egg extract containing 0.05 units/ μ l of *EcoRI*, which produces a double-strand DNA cleavage with a four-nucleotide-overhang at the 3'-end of the break. This concentration of *EcoRI* did not affect the nuclear formation in the *Xenopus* egg extract (Fig.




Fig. 5. Focus-formation of RPA in nuclei incubated with *Xenopus* egg extract containing *EcoRI*. (A) Demembrated *Xenopus* sperm nuclei were incubated at 23°C for the periods indicated in the presence of *EcoRI*. Localization of RPA (a,c,d,f,g,i) and DNA (b,e,h) is shown. Panels c, f and i are enlarged images of the nuclei indicated by arrowheads in panels a, d and g, respectively. The same nuclei in the 'DNA' panels are also indicated by arrowheads. Bars, 50 μ m (a,b,d,e,g,h) or 20 μ m (c,f,i). Numbers of RPA-accumulated nuclei/observed nuclei from two independent experiments are shown below the panels. The percentages of RPA-accumulated nuclei are also presented in parentheses. (B) A detergent-insoluble nuclear fraction was prepared from the nuclei incubated at 23°C for 60 minutes in the absence (lane 1) or the presence (lanes 2-6) of 0.05 units/ μ l *EcoRI*. The nuclear fraction was subjected to SDS-polyacrylamide gel electrophoresis directly (lanes 1,2), or after incubation with (lanes 3,4) or without (lanes 5,6) MNase followed by centrifugal separation of supernatant (lanes 3,5) and precipitated (lanes 4,6) fractions. The p32 subunit of RPA was visualized by immunoblotting. (C) The sperm nuclei were incubated in the extract containing *EcoRI* for 60 minutes at 23°C in the presence of geminin. Localization of RPA (a,c) and DNA (b) is shown. Panel c is an enlarged image of the nucleus indicated by an arrowhead in panel a. The same nucleus is also indicated by an arrowhead in panel b. Bars, 50 μ m (a,b) or 20 μ m (c).

5A,b,e,h), which took place about 30 minutes after the addition of the demembrated sperm, and the nuclei gradually enlarged during the incubation. Under these conditions, RPA foci appeared in 100%, 99% and 98% of the observed nuclei at 30 minutes, 60 minutes and 90 minutes, respectively (Fig. 5A,a,d,g). After the *EcoRI*-treatment, smeared DNA bands, centered at about 50 kbp, were detected by agarose gel electrophoresis, indicating that the restriction enzyme actually digested sperm DNA under these conditions (data not shown). The formation of RPA foci was also observed using *EcoRV* and *PstI*, which form blunt and 5' overhung DNA ends, respectively, or by the addition of *EcoRI* after nuclear formation (data not shown). Both the number of RPA foci-positive nuclei and the number of foci in the nucleus increased in response to the amount of *EcoRI* (data not shown). Immunoblot analysis indicated that RPA in the detergent-insoluble nuclear fraction was increased by the addition of *EcoRI* and released from the nuclear fraction by the treatment of MNase (Fig. 5B).

The enlarged images of representative nuclei at 60 minutes and 90 minutes were almost identical to those formed in the presence of CPT (Fig. 5Af,i). By contrast, the RPA signals after a 30 minute incubation in the presence of *EcoRI* were more similar to those at 60 minutes than after 30 minutes in the presence of CPT (Fig. 5Ac). The difference at 30 minutes may reflect the difference between *EcoRI* and CPT, which are thought to induce DSBs independently of and dependent on DNA replication, respectively. Indeed, the focus-formation of RPA in the nuclei that formed in the presence of *EcoRI* was insensitive to the inhibition of DNA replication by the addition of geminin (Fig. 5C).

Suppression of DNA replication induced by DNA damage was released by caffeine

Next, we assessed DNA replicating activity in the extract containing *EcoRI* (Fig. 6). While the CPT-treatment allowed nuclei to replicate DNA at approximately 40% of the level attained in the absence of DNA-damaging agent (Fig. 3C), the *EcoRI*-treatment prevented DNA replication almost completely. The reduction of DNA replication in the presence of *EcoRI* or CPT is likely to be due to the activation of a checkpoint system by DSB formation.

To confirm the involvement of a DNA-damage-induced checkpoint system in the reduction of DNA replication, we used caffeine as an inhibitor of ATM and ATR. Although caffeine alone had little or no effect on DNA synthesis in the absence of DNA damaging agents, the ability to synthesize DNA in the presence of *EcoRI* was restored to control levels by addition of caffeine. This restored DNA synthesis was sensitive to geminin treatment, indicating that DNA replication activity was suppressed by a DNA-damage-induced checkpoint system containing caffeine-sensitive components. Replication of DNA was also restored by caffeine in the nuclei that formed in the presence of CPT (Fig. 3C, closed triangle).

RPA focus-formation induced by DNA damage was not influenced by the caffeine-sensitive checkpoint pathway
Next, we examined whether caffeine also affects the formation




Fig. 6. Suppression of DNA replication by *EcoRI*-treatment. Demembrated *Xenopus* sperm nuclei were incubated for 90 minutes at 23°C in the extract with or without *EcoRI*, caffeine and geminin in the presence of [α - 32 P]dCTP. The extent of DNA synthesis is represented as a percentage of accumulative DNA synthesis after a 90 minute incubation in the absence of DNA damaging agents. 100% DNA synthesis is estimated to be an amount equivalent to the duplication of about 90% of input DNA.

of RPA foci in the presence of *EcoRI*. The *EcoRI*-induced RPA focus-formation proceeded in the presence of caffeine (Fig. 7A). This result indicates that the RPA focus-formation does not occur downstream of the activation of ATM or ATR kinase and the DNA lesion-induced DNA replication arrest. A similar result was obtained with the nuclei formed in the presence of CPT (Fig. 7B).

Detection of DNA repair synthesis that is co-localized with RPA foci and is insensitive to caffeine treatment

We next attempted to detect DSB-induced DNA repair synthesis. Because geminin efficiently inhibited DNA replication, it seemed likely that DNA repair synthesis in geminin-treated nuclei would be easily detected because of the low background of DNA replication. However, as shown in Fig. 3C and Fig. 6, DNA synthesis in the geminin-treated nuclei was hardly detectable by the standard assay. Then, we tried microscopic observation of DNA synthesis after the incorporation of biotin-conjugated dUTP.

Although DNA replication gave quite intense signals of DNA synthesis (data not shown), geminin efficiently suppressed the signal due to DNA replication as expected (Fig. 8a). When *EcoRI* was added to the extract containing geminin, DNA synthesis was detected in the nuclei (Fig. 8d,g). *EcoRI*-induced RPA foci in the nuclei were also detected under these conditions, similarly to the previous experiments (Fig. 8e,h), and the location of the loci used to synthesize DNA showed good correlation with that of the RPA foci (Fig. 8f,i). Similar to the focus-formation of RPA, caffeine treatment did not have any apparent effect on DNA repair synthesis observed in this system (Fig. 8j,m) or on the relation of loci of the DNA synthesis and the RPA foci (Fig. 8l,o), suggesting that at least part of the DSB repair process is not dependent on ATM or ATR activation.




Fig. 7. Caffeine did not affect focus-formation of RPA. Demembrated *Xenopus* sperm nuclei were incubated in the extract containing 5 mM caffeine for 60 minutes at 23°C in the presence of *Eco*RI (A) or CPT (B). Localization of RPA (a,c) and DNA (b) is shown. Panel c is an enlarged image of the nucleus indicated by an arrowhead in panel a. The same nucleus is also indicated by an arrowhead in panel b. Bars, 50 μ m (a,b) or 20 μ m (c).

Discussion

To analyze responses to the formation of DSBs, we developed a cell-free system consisting of extracts derived from *Xenopus* eggs. We examined the detergent-insensitive localization of RPA in the nuclei that formed in *Xenopus* egg extract. The antibody used in this study was raised against *Xenopus* RPA-p32, the middle-sized subunit of the RPA-heterotrimer. This antibody recognized a 32 kDa protein in crude *Xenopus* egg extract and in a chromatin-associated fraction in immunoblot analysis. It also recognized phosphorylated forms of the protein (data not shown).

In the extract containing no DNA damaging agents, RPA-accumulation occurred evenly in the nuclei after a 30 minute incubation, and the accumulation was completely inhibited by geminin-treatment. Geminin is an endogenous protein that selectively binds to and inhibits Cdt1/RLF-B (Tada et al., 2001; Wohlschlegel et al., 2000). Cdt1/RLF-B possesses an activity essential for chromatin loading of MCM/P1 protein complex (Tada et al., 1999; Maiorano et al., 2000; Nishitani et al., 2000), a component of the pre-replicative complex necessary for the initiation of DNA replication (Blow, 2001; Takisawa et al., 2000; Tye, 1999). Therefore, geminin inhibits the initiation of DNA replication and is likely to have no or very little effect on other processes in living cells, including progression of the cell cycle. Thus, geminin is a good tool for eliminating the influence of DNA replication. The sensitivity of the RPA signals to geminin suggests that the accumulation of RPA is dependent on the initiation of DNA replication as described previously (Adachi and Laemmli, 1992; Adachi and Laemmli, 1994; Chen et al., 2001).

The RPA-signal observed by indirect immunofluorescence was significantly reduced at 60 minutes and almost absent at 90 minutes, when DNA synthesis is almost complete (Fig. 3C, closed circle) and most of the MCM proteins were removed from the chromatin (Thömmes et al., 1997) in *Xenopus* egg extracts. Thus, the disappearance of the RPA-

signal during further incubation seems to reflect the removal of RPA from the chromatin during the progression of DNA replication. This notion was supported by the result of immunoblot analysis detecting RPA in the detergent-insoluble nuclear fraction. The sequential alteration in the behavior of RPA seemed to differ between the present study and the study reported previously (Chen et al., 2001). They observed discrete RPA signal even after completion of DNA replication under conditions similar to our damage non-inducing conditions. The discrepancy may be caused by the difference in the method used for fixation before immunostaining. A wash of fixed nuclei with 0.1% Triton X-100 followed immobilization of nuclei on coverslips by centrifugation in a protocol used by Chen et al., whereas, in our experiments, fixed nuclei were washed with 0.24% Triton X-100 before immobilization. Another possibility for this discrepancy is the difference in antibodies used for the immunostaining. While Chen et al., used antibodies against the largest subunit or all three subunits of RPA, the antibody we used for this study was specific to the p32 subunit of RPA.

CPT is an inhibitor of DNA topoisomerase I and is known to trap DNA topoisomerase I cleavage complexes resulting in enzyme-linked breaks in the substrate DNA (Hsiang et al., 1985; Hsiang and Liu, 1988; Pommier et al., 1998). The formation of DSBs by CPT is thought to depend on DNA replication on the basis of several lines of evidence (Avemann et al., 1988; D'Arpa et al., 1990; Holm et al., 1989; Hsiang et al., 1989; Ryan et al., 1991). We analyzed RPA and γ -H2AX foci formation in the nuclei incubated in CPT-containing *Xenopus* egg extract and the results further support this notion. Our result also suggested that the nuclear formation was slightly retarded in the extract containing CPT, although the influence was not so effective since no retardation of nuclear enlargement was observed after a 60 minute incubation. Thus, it is possible that CPT disturbed the decondensation of extremely condensed sperm chromatin and the effect of CPT on decondensation induced the formation of the RPA-foci observed in the present study. However, when CPT was added after chromatin decondensation or nuclear formation, it still caused the formation of the RPA foci (data not shown), suggesting that the retardation of the nuclear formation has no or very little effect on the CPT-induced formation of the nuclear RPA foci.

DNA replication in CPT-treated nuclei started after a 30 minute incubation, similar to that observed in the absence of DNA-damage-inducing agents, but ceased at 50 minutes even though the amount of replicated DNA was only about 40% of that in the control nuclei. This result suggests that DNA lesions arose after the start of DNA replication and activated a DNA-damage checkpoint pathway that inhibited progression of DNA replication. Indeed, the reduction of DNA replicating activity was completely recovered by treatment with caffeine, an inhibitor of ATM and ATR, which play key roles in checkpoint pathways.

Moreover, detailed observations revealed that the RPA accumulation after a 30 minute incubation in CPT-containing extract appeared to be composed of a large number of minute RPA dots evenly distributed in the nuclei (data not shown). The pattern of RPA accumulation was similar to that after a 30 minute incubation in the control extract rather than in *Eco*RI-containing extract, suggesting that the presence of CPT in the




Fig. 8. *EcoRI*-induced DNA synthesis was co-localized with RPA foci and insensitive to caffeine treatment. Demembrated *Xenopus* sperm nuclei were incubated in the extract containing geminin and biotin-dUTP for 30 minutes at 23°C in the absence (a-c), or presence of *EcoRI* (d-i) or *EcoRI*+caffeine (j-o). Localization of biotin-dUTP (a,d,g,j,m) and RPA (b,e,h,k,n) is shown in green and red, respectively. Merged images are also shown in panels c, f, i, l and o. Panels g, h, i, m, n and o are enlarged images of the nuclei indicated by arrowheads in panels d, e, f, j, k and l, respectively. Bars, 50 μm (a-f,j-l); 20 μm (g-i,m-o).

extract did not affect the formation of fine RPA dots in the nuclei at 30 minutes, when DNA replication was about to start. In contrast, discrete RPA foci were observed after prolonged incubation, whereas very few discrete RPA foci were observed in the nuclei in the absence of DNA damaging agents. These results suggest that the fine RPA dots at 30 minutes formed at sites where DNA replication was about to start and the discrete foci that appeared after prolonged incubation were located where DNA lesions had been induced during DNA replication by CPT-treatment.

The notion that the RPA foci formed where DNA lesions were induced, was further supported by the experiment using *EcoRI* in which discrete RPA foci were observed at 30 minutes as well as after prolonged incubation in the absence of DNA replication. Geminin-insensitive, DNA replication-independent DNA synthesis was induced by *EcoRI* under these experimental conditions, and was co-localized with *EcoRI*-

induced RPA foci. Therefore, the DNA synthesis is likely to be due to the repair of DSBs, further confirming that RPA accumulated at the loci of DNA lesions.

It is possible that the RPA foci observed in the *EcoRI*-treated or CPT-treated nuclei after a 60 and 90 minute incubation resulted from the failure to extinguish the RPA foci that formed in the nuclei after a 30 minute incubation by inhibiting progression of DNA replication. However, the results obtained in the present study provide evidence against this idea. First, the formation of RPA foci in *EcoRI*-treated nuclei was insensitive to geminin, which eliminates the formation of fine RPA foci after a 30 minute incubation in the absence of DNA damaging agents. Second, although caffeine-treatment overcame the inhibition of DNA replication by the checkpoint mechanism, it had no effect on the formation of RPA foci in the *EcoRI*- or CPT-treated nuclei.

The process inhibiting the initiation of DNA replication by a caffeine-sensitive checkpoint system has been discussed in a previous report (Costanzo et al., 2000). The authors showed that the association of Cdc45 with chromatin was suppressed by ATM, activated by linearized plasmids in *Xenopus* egg extracts. They also showed that in this ATM-containing fraction, the activity of Cdk2 and its ability to associate with cyclin E was inhibited. They explained these results by suggesting that Cdc45 loading on chromatin is suppressed by the inhibition of Cdk2/Cyclin E protein kinase. The inhibition probably occurs via the activation of ATM, followed by the activation of Chk2/Cds1 protein kinase, and the suppression of Cdc25A protein phosphatase, which activates Cdk2/Cyclin E (Costanzo et al., 2000; Falck et al., 2001). All of the above processes should be suppressed after caffeine-treatment via inhibition of the ATM activation.

The result that caffeine did not inhibit the formation of DNA-lesion-induced RPA foci or DNA synthesis suggests that the focus-formation and at least a part of the repair process occurs independently of downstream events of ATM or ATR activation. However, it does not necessarily exclude the involvement of DNA-PK. The concentration of caffeine needed to inhibit DNA-PK activity by 50% is 10 mM (Sarkaria et al., 1999), higher than levels used in the present study. In addition, DNA-PK has been reported to phosphorylate RPA in response to DNA damage (Niu et al., 1997; Shao et al., 1999; Zernik-Kobak et al., 1997), which suggests that the behavior of RPA is controlled by the phosphorylation catalyzed by DNA-PK.

We have reported here the establishment of a simple cell-free system using *Xenopus* egg extract, which may permit further study into the recognition and repair of DSBs. Using restriction enzymes and recombinant geminin, the use of which in intact cells is very difficult, we have obtained clear results that indicate accumulation of RPA on chromatin containing DSBs, suppression of DNA replication by DNA-damage-induced checkpoint mechanisms, and co-localization of the DSB-induced DNA synthesis loci and the RPA foci. We have also shown that the recognition of DNA lesions by RPA and DNA synthesis induced by DSB-formation occurs independently of the downstream events of ATM or ATR activation, including the arrest of DNA replication. We think that this system will open the way to detailed analyses regarding the recognition and the repair processes of DSBs.

We thank Ben Hodgson and Dana Branzei for revision of English in the manuscript. This work was supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan, and a Health Sciences Research Grant from the Ministry of Health, Labour and Welfare of Japan.

References

- Adachi, Y. and Laemmli, U. K.** (1992). Identification of nuclear pre-replication centers poised for DNA synthesis in *Xenopus* egg extracts: immunolocalization study of replication protein A. *J. Cell Biol.* **119**, 1-15.
- Adachi, Y. and Laemmli, U. K.** (1994). Study of the cell cycle-dependent assembly of the DNA pre-replication centres in *Xenopus* egg extracts. *EMBO J.* **13**, 4153-4164.
- Avemann, K., Knippers, R., Koller, T. and Sogo, J. M.** (1988). Camptothecin, a specific inhibitor of type I DNA topoisomerase, induces DNA breakage at replication forks. *Mol. Cell Biol.* **8**, 3026-3034.
- Bischof, O., Kim, S. H., Irving, J., Beresten, S., Ellis, N. A. and Campisi, J.** (2001). Regulation and localization of the Bloom syndrome protein in response to DNA damage. *J. Cell Biol.* **153**, 367-380.
- Blow, J. J.** (2001). Control of chromosomal DNA replication in the early *Xenopus* embryo. *EMBO J.* **20**, 3293-3297.
- Boubnov, N. V. and Weaver, D. T.** (1995). *scid* cells are deficient in Ku and replication protein A phosphorylation by the DNA-dependent protein kinase. *Mol. Cell Biol.* **15**, 5700-5706.
- Burns, J. L., Guzder, S. N., Sung, P., Prakash, S. and Prakash, L.** (1996). An affinity of human replication protein A for ultraviolet-damaged DNA. Implications for damage recognition in nucleotide excision repair. *J. Biol. Chem.* **271**, 11607-11610.
- Chen, C. Y., Graham, J. and Yan, H.** (2001). Evidence for a replication function of FFA-1, the *Xenopus* orthologue of Werner syndrome protein. *J. Cell Biol.* **152**, 985-996.
- Chong, J. P., Thommes, P., Rowles, A., Mahbubani, H. M. and Blow, J. J.** (1997). Characterization of the *Xenopus* replication licensing system. *Methods Enzymol.* **283**, 549-564.
- Costanzo, V., Robertson, K., Ying, C. Y., Kim, E., Avvedimento, E., Gottesman, M., Grieco, D. and Gautier, J.** (2000). Reconstitution of an ATM-dependent checkpoint that inhibits chromosomal DNA replication following DNA damage. *Mol. Cell* **6**, 649-659.
- D'Arpa, P., Beardmore, C. and Liu, L. F.** (1990). Involvement of nucleic acid synthesis in cell killing mechanisms of topoisomerase poisons. *Cancer Res.* **50**, 6919-6924.
- Durocher, D. and Jackson, S. P.** (2001). DNA-PK, ATM and ATR as sensors of DNA damage: variations on a theme? *Curr. Opin. Cell Biol.* **13**, 225-231.
- Enomoto, T.** (2001). Functions of recQ family helicases: possible involvement of Bloom's and Werner's syndrome gene products in guarding genome integrity during DNA replication. *J. Biochem. (Tokyo)* **129**, 501-507.
- Fairman, M. P. and Stillman, B.** (1988). Cellular factors required for multiple stages of SV40 DNA replication in vitro. *EMBO J.* **7**, 1211-1218.
- Falck, J., Mailand, N., Syljuasen, R. G., Bartek, J. and Lukas, J.** (2001). The ATM-Chk2-Cdc25A checkpoint pathway guards against radioresistant DNA synthesis. *Nature* **410**, 842-847.
- Golub, E. I., Gupta, R. C., Haaf, T., Wold, M. S. and Radding, C. M.** (1998). Interaction of human rad51 recombination protein with single-stranded DNA binding protein, RPA. *Nucleic Acids Res.* **26**, 5388-5393.
- Guo, Z. and Dunphy, W. G.** (2000). Response of *Xenopus* Cds1 in cell-free extracts to DNA templates with double-stranded ends. *Mol. Biol. Cell* **11**, 1535-1546.
- Guo, Z., Kumagai, A., Wang, S. X. and Dunphy, W. G.** (2000). Requirement for Atr in phosphorylation of Chk1 and cell cycle regulation in response to DNA replication blocks and UV-damaged DNA in *Xenopus* egg extracts. *Genes Dev.* **14**, 2745-2756.
- Haaf, T., Golub, E. I., Reddy, G., Radding, C. M. and Ward, D. C.** (1995). Nuclear foci of mammalian Rad51 recombination protein in somatic cells after DNA damage and its localization in synaptonemal complexes. *Proc. Natl. Acad. Sci. USA* **92**, 2298-2302.
- Haaf, T., Raderschall, E., Reddy, G., Ward, D. C., Radding, C. M. and Golub, E. I.** (1999). Sequestration of mammalian Rad51-recombination protein into micronuclei. *J. Cell Biol.* **144**, 11-20.
- Haber, J. E.** (2000). Partners and pathways repairing a double-strand break. *Trends Genet.* **16**, 259-264.
- Holm, C., Covey, J. M., Kerrigan, D. and Pommier, Y.** (1989). Differential requirement of DNA replication for the cytotoxicity of DNA topoisomerase I and II inhibitors in Chinese hamster DC3F cells. *Cancer Res.* **49**, 6365-6368.
- Hsiang, Y. H. and Liu, L. F.** (1988). Identification of mammalian DNA topoisomerase I as an intracellular target of the anticancer drug camptothecin. *Cancer Res.* **48**, 1722-1726.
- Hsiang, Y. H., Hertzberg, R., Hecht, S. and Liu, L. F.** (1985). Camptothecin induces protein-linked DNA breaks via mammalian DNA topoisomerase I. *J. Biol. Chem.* **260**, 14873-14878.
- Hsiang, Y. H., Lihou, M. G. and Liu, L. F.** (1989). Arrest of replication forks by drug-stabilized topoisomerase I-DNA cleavable complexes as a mechanism of cell killing by camptothecin. *Cancer Res.* **49**, 5077-5082.
- Karran, P.** (2000). DNA double strand break repair in mammalian cells. *Curr. Opin. Genet. Dev.* **10**, 144-150.
- Khanna, K. K. and Jackson, S. P.** (2001). DNA double-strand breaks: signaling, repair and the cancer connection. *Nat. Genet.* **27**, 247-254.
- Labhart, P.** (1999). Nonhomologous DNA end joining in cell-free systems. *Eur. J. Biochem.* **265**, 849-861.
- Lao, Y., Gomes, X. V., Ren, Y., Taylor, J. S. and Wold, M. S.** (2000). Replication protein A interactions with DNA. III. Molecular basis of recognition of damaged DNA. *Biochemistry* **39**, 850-859.
- Lohka, M. J. and Masui, Y.** (1983). Formation in vitro of sperm pronuclei and mitotic chromosomes induced by amphibian ooplasmic components. *Science* **220**, 719-721.
- Maiorano, D., Moreau, J. and Mechali, M.** (2000). XCDT1 is required for the assembly of pre-replicative complexes in *Xenopus laevis*. *Nature* **404**, 622-625.
- McGarry, T. J. and Kirschner, M. W.** (1998). Geminin, an inhibitor of DNA replication, is degraded during mitosis. *Cell* **93**, 1043-1053.
- Mohaghegh, P. and Hickson, I. D.** (2001). DNA helicase deficiencies associated with cancer predisposition and premature ageing disorders. *Hum. Mol. Genet.* **10**, 741-746.
- Nishitani, H., Lygerou, Z., Nishimoto, T. and Nurse, P.** (2000). The Cdt1 protein is required to license DNA for replication in fission yeast. *Nature* **404**, 625-628.
- Niu, H., Erdjument-Bromage, H., Pan, Z. Q., Lee, S. H., Tempst, P. and Hurwitz, J.** (1997). Mapping of amino acid residues in the p34 subunit of human single-stranded DNA-binding protein phosphorylated by DNA-dependent protein kinase and Cdc2 kinase in vitro. *J. Biol. Chem.* **272**, 12634-12641.
- Oakley, G. G., Loberg, L. I., Yao, J., Risinger, M. A., Yunker, R. L., Zernik-Kobak, M., Khanna, K. K., Lavin, M. F., Carty, M. P. and Dixon, K.** (2001). UV-induced hyperphosphorylation of replication protein A depends on DNA replication and expression of ATM protein. *Mol. Biol. Cell* **12**, 1199-1213.
- Pommier, Y., Pourquier, P., Fan, Y. and Strumberg, D.** (1998). Mechanism of action of eukaryotic DNA topoisomerase I and drugs targeted to the enzyme. *Biochim. Biophys. Acta* **1400**, 83-105.
- Rogakou, E. P., Pilch, D. R., Orr, A. H., Ivanova, V. S. and Bonner, W. M.** (1998). DNA double-stranded breaks induce histone H2AX phosphorylation on serine 139. *J. Biol. Chem.* **273**, 5858-5868.
- Rogakou, E. P., Boon, C., Redon, C. and Bonner, W. M.** (1999). Megabase chromatin domains involved in DNA double-strand breaks in vivo. *J. Cell Biol.* **146**, 905-916.
- Ryan, A. J., Squires, S., Strutt, H. L. and Johnson, R. T.** (1991). Camptothecin cytotoxicity in mammalian cells is associated with the induction of persistent double strand breaks in replicating DNA. *Nucleic Acids Res.* **19**, 3295-3300.
- Sakamoto, S., Nishikawa, K., Heo, S. J., Goto, M., Furuichi, Y. and Shimamoto, A.** (2001). Werner helicase relocates into nuclear foci in response to DNA damaging agents and co-localizes with RPA and Rad51. *Genes Cells* **6**, 421-430.
- Sarkaria, J. N., Busby, E. C., Tibbetts, R. S., Roos, P., Taya, Y., Karnitz, L. M. and Abraham, R. T.** (1999). Inhibition of ATM and ATR kinase activities by the radiosensitizing agent, caffeine. *Cancer Res.* **59**, 4375-4382.
- Shao, R. G., Cao, C. X., Zhang, H., Kohn, K. W., Wold, M. S. and Pommier, Y.** (1999). Replication-mediated DNA damage by camptothecin induces phosphorylation of RPA by DNA-dependent protein kinase and dissociates RPA:DNA-PK complexes. *EMBO J.* **18**, 1397-1406.
- Shiloh, Y.** (2001). ATM and ATR: networking cellular responses to DNA damage. *Curr. Opin. Genet. Dev.* **11**, 71-77.
- Tada, S., Chong, J. P. J., Mahbubani, H. M. and Blow, J. J.** (1999). The RLF-B component of the replication licensing system is distinct from Cdc6 and functions after Cdc6 binds to chromatin. *Curr. Biol.* **9**, 211-214.
- Tada, S., Li, A., Maiorano, D., Mechali, M. and Blow, J. J.** (2001).

- Repression of origin assembly in metaphase depends on inhibition of RLF-B/Cdt1 by geminin. *Nat. Cell Biol.* **3**, 107-113.
- Takisawa, H., Mimura, S. and Kubota, Y.** (2000). Eukaryotic DNA replication: from pre-replication complex to initiation complex. *Curr. Opin. Cell Biol.* **12**, 690-696.
- Tchang, F. and Mechali, M.** (1999). Nuclear import of p53 during *Xenopus laevis* early development in relation to DNA replication and DNA repair. *Exp. Cell Res.* **251**, 46-56.
- Thömmes, P., Kubota, Y., Takisawa, H. and Blow, J. J.** (1997). The RLF-M component of the replication licensing system forms complexes containing all six MCM/P1 polypeptides. *EMBO J.* **16**, 3312-3319.
- Tye, B. K.** (1999). MCM proteins in DNA replication. *Annu. Rev. Biochem.* **68**, 649-686.
- Wobbe, C. R., Weissbach, L., Borowiec, J. A., Dean, F. B., Murakami, Y., Bullock, P. and Hurwitz, J.** (1987). Replication of simian virus 40 origin-containing DNA in vitro with purified proteins. *Proc. Natl. Acad. Sci. USA* **84**, 1834-1838.
- Wohlschlegel, J. A., Dwyer, B. T., Dhar, S. K., Cvetic, C., Walter, J. C. and Dutta, A.** (2000). Inhibition of eukaryotic DNA replication by geminin binding to Cdt1. *Science* **290**, 2309-2312.
- Wold, M. S.** (1997). Replication protein A: a heterotrimeric, single-stranded DNA-binding protein required for eukaryotic DNA metabolism. *Annu. Rev. Biochem.* **66**, 61-92.
- Wold, M. S. and Kelly, T.** (1988). Purification and characterization of replication protein A, a cellular protein required for in vitro replication of simian virus 40 DNA. *Proc. Natl. Acad. Sci. USA* **85**, 2523-2527.
- Yan, H. and Newport, J.** (1995). FFA-1, a protein that promotes the formation of replication centers within nuclei. *Science* **269**, 1883-1885.
- Yan, H., Chen, C. Y., Kobayashi, R. and Newport, J.** (1998). Replication focus-forming activity 1 and the Werner syndrome gene product. *Nat. Genet.* **19**, 375-378.
- Zernik-Kobak, M., Vasunia, K., Connelly, M., Anderson, C. W. and Dixon, K.** (1997). Sites of UV-induced phosphorylation of the p34 subunit of replication protein A from HeLa cells. *J. Biol. Chem.* **272**, 23896-23904.