

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Roman L. Hruska U.S. Meat Animal Research
Center

U.S. Department of Agriculture: Agricultural
Research Service, Lincoln, Nebraska

1987

Follicle Stimulating Hormone, Luteinizing Hormone and Testicular Leydig Cell Responses to Estradiol Immunization in Ile-de-France Rams

B. D. Schanbacher
USDA-ARS

J. Pelletier
INRA

M. T. Hochereau-de Reviers
INRA

Follow this and additional works at: <http://digitalcommons.unl.edu/hruskareports>

Schanbacher, B. D.; Pelletier, J.; and Hochereau-de Reviers, M. T., "Follicle Stimulating Hormone, Luteinizing Hormone and Testicular Leydig Cell Responses to Estradiol Immunization in Ile-de-France Rams" (1987). *Roman L. Hruska U.S. Meat Animal Research Center*. 301.

<http://digitalcommons.unl.edu/hruskareports/301>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Roman L. Hruska U.S. Meat Animal Research Center by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Follicle Stimulating Hormone, Luteinizing Hormone and Testicular Leydig Cell Responses to Estradiol Immunization in Ile-de-France Rams

B. D. SCHANBACHER,* J. PELLETIER,† AND M. T. HOCHEREAU-DE REVIERS†

Active immunization of Ile-de-France rams against estradiol (E_2) resulted in the production of E_2 -neutralizing antibodies and an elevation in the plasma concentrations of FSH, LH, and testosterone. The presence of E_2 antibodies did not affect the testosterone metabolic clearance rate, indicating that the immunization-mediated 10-fold increase in plasma testosterone was the result of a 10-fold increase in testicular testosterone production. Testis weights, as well as nuclear and cytoplasmic volumes of individual peritubular and perivascular Leydig cells, were greater in E_2 -immunized rams than in albumin-immunized controls. Leydig cell numbers were not affected by treatment. The E_2 antibodies were capable not only of neutralizing the inhibitory effects of endogenous E_2 on gonadotropin levels in intact rams, but were able to block the effects of exogenously administered E_2 on their FSH and LH secretory response to castration. It is concluded that circulating E_2 in the ram is involved in pituitary-testicular endocrine homeostasis and that E_2 immunoneutralization can be employed to enhance testosterone secretion in this species.

Key words: estradiol immunoneutralization, gonadotropins, testosterone secretion, ram testes.

J Androl 1987; 8:97-102.

Estradiol-17 β (E_2) in peripheral blood results from gonadal (Payne et al, 1976; Dorrington et al, 1978) and extragonadal (Longcope et al, 1978; Worgul et al, 1981) production. Whereas intratesticular estrogen acts locally to regulate steroid biosynthesis in Leydig cells (Hsueh et al, 1978; Melner and Abney, 1980; Moger, 1980), peripheral estrogens can affect Leydig cell function indirectly via their feedback actions on the hypothalamic-pituitary axis (Swerdlhoff and Odell, 1968; deJong et al, 1975).

Reprint requests: B.D. Schanbacher, Ph.D., R. L. Hruska U.S. Meat Animal Research Center, Clay Center, Nebraska 68933.

Submitted for publication February 12, 1986; revised version received July 9, 1986; accepted for publication August 26, 1986.

*From the I.N.R.A., Station de Physiologie de la Reproduction, France† and U.S.D.A., R. L. Hruska U.S. Meat Animal Research Center, Clay Center, Nebraska**

In male sheep, E_2 is a potent regulator of gonadotropin secretion (Riggs and Malven, 1974; D'Occhio et al, 1983), presumably by acting through specific receptors at the levels of the hypothalamus (Pelletier and Caraty, 1981) and/or the pituitary (Thieulant and Pelletier, 1979; Glass et al, 1984). The physiologic importance of picomolar concentrations of circulating E_2 in the feedback regulation of LH secretion in intact rams has been demonstrated only recently (Schanbacher, 1984). Active immunization of Suffolk rams against E_2 elicited a two- to three-fold increase in mean serum LH concentrations and a parallel increase in mean serum testosterone concentrations. These data suggest that circulating E_2 plays a physiologic role in the regulation of pituitary LH secretion and that E_2 immunoneutralization can be employed to stimulate the reproductive-endocrine axis of rams. The following study was undertaken in the Ile-de-France breed to extend our previous findings regarding estradiol immunization effects in the domestic ram. The parameters assessed included: 1) FSH secretion; 2) testosterone production and metabolic clearance rates; 3) Leydig cell morphology; and 4) antibody effects on exogenous as well as endogenous E_2 .

Materials and Methods

Animals and Immunization

Four-year-old Ile-de-France rams weighing approximately 80 kg were assigned to the following study. All

animals were kept at the INRA research facility near Nouzilly and were fed alfalfa hay, wheat straw, and grain supplement as required for body weight maintenance. Water, salt, and mineral supplement were provided *ad libitum*. Experimentation began in October at the time of primary immunization and extended into the nonbreeding season, a time when testosterone levels generally are suppressed.

Rams were randomly assigned to one of two groups. Five rams were actively immunized against the carrier protein bovine serum albumin (BSA; 1 mg) while each of a second group of five rams was immunized with 1 mg of an E₂-6-CMO-BSA conjugate (E₂:BSA; Steraloids, Inc., Wilton, NH). Antigens were dissolved in 1 ml saline and then emulsified with 1 ml Freund's complete adjuvant as described previously (Schanbacher, 1984). Injections were made in 20 or more subcutaneous sites over the ribs. BSA and E₂:BSA rams were given the respective antigens as booster injections (1 mg each) in December and February.

Experiment 1

BSA- and E₂:BSA-immunized rams were sampled in April (ovine nonbreeding season) via jugular venipuncture at 20-minute intervals for 8 hours. Plasma from these samples were stored frozen at -20 C for subsequent determination of E₂ antibody binding titer and concentrations of FSH, LH, and testosterone. All ten rams were subsequently castrated under pentothal anesthesia. Pieces of testes were fixed in Bouin Holland's solution (right testis) or in glutaraldehyde:collidine buffer followed by osmium (left testis). Ten-micron sections were stained by the Feulgen reaction and counterstained with Alcian blue for histologic assessment of peritubular and perivascular Leydig cells (Lunstra and Schanbacher 1987). The area of individual cells, total number of cells, and total volume of these cells were determined by morphometric means (Hochereau-de Reviers et al, 1979). Fifteen days postcastration, when circulating testosterone levels were not detectable, estimates of testosterone metabolic clearance rates for BSA- and E₂:BSA-immunized animals were ascertained. Each animal was injected intravenously with 1 mg of testosterone dissolved in 2 ml of 10% ethanolic saline at 1000 hours. Plasma samples were collected at 5-minute intervals starting 10 minutes after injection through 60 minutes and again at 90 and 120 minutes. The natural log transformation of plasma testosterone concentrations was used to calculate the decay constant ($-\alpha$) used in derivation of the testosterone metabolic clearance rate (Terqui, 1978). The equation, $r = Ae^{-\alpha t}$, simplifies to $MCR_T = \alpha/A$, where r is testosterone in fraction of the injected dose per liter of plasma, t is time expressed in minutes, and A the fractional dose intercept. Testosterone daily production rates of immunized rams were computed from the product of mean plasma testosterone concentrations and the testosterone metabolic clearance rate (Baird et al, 1969).

Experiment 2

The BSA- and E₂:BSA-immunized rams were implanted with a 6-cm Silastic capsule containing estradiol (Schanbacher, 1984) at the time of castration to assess their

susceptibility to E₂ negative feedback. This dose of E₂ is known to suppress LH secretion, especially during the nonbreeding season. The postcastration gonadotropin response, including FSH and LH, was determined in plasma samples collected on days 4, 7, and 11. Blood samples were taken again via jugular venipuncture at 20-minute intervals, but for only 3 hours (0900-1200 hours) on each of the respective days. Plasma LH concentrations were determined in all samples, whereas FSH and E₂ antibody binding titers were determined in plasma pools prepared from samples of each animal. E₂ implants were removed from all animals after blood sampling on day 11.

Hormone Assays and E₂ Antibody Titers

Plasma FSH (Blanc and Poirier, 1979) and LH (Pelletier et al, 1982) were determined by double antibody radioimmunoassay procedures described previously. Determinations were made in duplicate using 100 μ l of plasma. Assay sensitivities and intraassay coefficients of variation were: FSH, 1 ng/ml HG-FSH 225 (35 \times NIH-FSH-S3) and 7.5%; LH, 300 pg/ml LH-M-3 (1.8 \times NIH-LH-S1) and 8.9%.

Plasma testosterone concentrations were also determined by double antibody radioimmunoassay (Garnier et al, 1978) but in duplicate 50- μ l ethylacetate:cyclohexane (1:1, v:v) extracts. Extraction efficiency using 10 volumes of extracting solvent averaged $92 \pm 6\%$. The minimal detectable amount of testosterone was 200 pg/ml and the intraassay coefficient of variation was 11.8% for a plasma pool containing 2.5 ng/ml.

E₂ antibody titers were determined by a charcoal precipitation assay (Schanbacher, 1984) and the results expressed as the percentage of radiolabeled E₂ specifically bound in 1000-fold plasma dilutions. In brief, diluted plasma was incubated overnight at 4 C with 20,000 dpm (18 pg) tritium-labeled E₂ (Amersham Radiochemical Centre, England) and free and bound tracer were separated the next day by dextran-treated charcoal. The supernatant (bound fraction) was counted by liquid scintillation spectroscopy.

Morphometric Analyses

Intertubular tissue relative volume was assessed on 20 fields per left testis with an ocular reticule integrator (25 points) at a $\times 256$ magnification. Leydig cell relative volume was assessed similarly at a $\times 1,600$ magnification. Total intertubular and Leydig cell volumes were then calculated. Mean cross sectional areas of Leydig cells or Leydig cell nuclei were measured with an automatic planimeter on at least 20 peritubular cells per testis adjacent to stage 8 tubules of the seminiferous epithelium cycle (Roosen-Runge and Giesel, 1950). Perivascular Leydig cells were measured independently of the stages of the seminiferous epithelium.

Assuming there were similar numbers of peritubular and perivascular Leydig cells, mean Leydig cell volume and total number of Leydig cells per testis were calculated. Perivascular Leydig cell cross sectional areas were also measured on four glutaraldehyde-fixed testes per treatment group. Two- μ m semithin sections, stained with Toluidine Blue, were prepared and Leydig cells were measured only when their nuclei contained an equatorial nucleolus.

TABLE 1. (Experiment 1). Mean plasma FSH, LH and Testosterone Concentrations and Testosterone Metabolic Clearance Rate and Production Rate Estimates after Immunization Against Bovine Serum Albumin or an Estradiol:BSA Conjugate in Mature Ile-de-France Rams*

	BSA Rams	E ₂ :BSA Rams
E ₂ binding titer (B/Bo at 1:1000)	< 0.5	39.8 ± 5.2
FSH		
Mean conc. (ng/ml)	1.9 ± 0.2	4.3 ± 0.9†
LH		
Mean conc. (ng/ml)	1.4 ± 0.5	3.3 ± 0.8†
Peak interval (min)	336 ± 59	110 ± 14†
Peak amplitude (ng/ml)	3.4 ± 1.2	8.2 ± 1.3†
Testosterone		
Mean conc. (ng/ml)	2.5 ± 0.3	23.7 ± 4.9†
MCR (l/d)‡	1959 ± 51	1886 ± 47
PR (mg/d)‡	6.3 ± 0.7	64.2 ± 3.7†

*Mean plasma hormone concentrations (± SEM, n = 5) were calculated from an 8-hour intensive bleed in April (primary immunizations were given in October with booster injections in December and February).

†Significantly different from BSA control value, $P < 0.01$.

‡MCR = metabolic clearance rate.

PCR = daily production rate.

Statistical Analyses

All treatment means for BSA- and E₂:BSA-immunized rams were compared utilizing student's *t*-test (Freund et al, 1960).

Results

Experiment 1

Immunization of mature Ile-de-France rams against E₂-6-CMO-BSA resulted in the production of antibodies capable of binding tritiated E₂ (Table 1). Whereas approximately 40% of the tracer was bound in E₂:BSA ram plasma diluted 1000-fold, plasma from BSA-immunized rams bound less than 0.5% of the E₂ tracer. Competitive inhibition curves yielded relative cross reactivities of 0.8% for testosterone and less than 0.1% for dihydrotestosterone.

E₂:BSA-immunized rams were found to have elevated plasma FSH, LH, and testosterone when compared with BSA-immunized controls (Table 1). Plasma LH and testosterone were secreted episodically in all rams with the frequency and amplitude of the LH secretory episodes being greater in E₂:BSA-immunized rams. The linear decay curves of exogenously administered testosterone between 20 and 60 minutes were nearly identical (Fig. 1), suggesting that




Fig. 1. Disappearance of immunoreactive testosterone in plasma of acutely castrated, albumin- (BSA) and estradiol: albumin-(E₂:BSA) immunized rams after a 1-mg intravenous bolus injection at time zero. Metabolic clearance rates were determined from the linear decay curves observed between 20 and 60 minutes (inset) to be similar for BSA- and E₂:BSA-immunized rams.

disappearance of testosterone from the plasma compartment of BSA- and E₂:BSA-immunized rams was the same. Indeed, the calculated metabolic clearance rate for testosterone of E₂:BSA rams was not significantly different from that of BSA-immunized controls. Consequently, the observed 10-fold increase in plasma testosterone of E₂:BSA-immunized rams reflected a 10-fold increase in testosterone daily production rate (Table 1).

Testis weights were significantly heavier in E₂:BSA-immunized rams. Light microscopic inspection revealed an increase in nuclear and cytoplasmic volume occupied by individual Leydig cells of these rams (Table 2). This increase in cell size was observed for both peritubular and perivascular Leydig cells. An increase in perivascular Leydig cells was also observed in semithin sections from E₂-immunized rams (64.5 ± 4.5 vs. 85.4 ± 12.5 μm²). The total volume of Leydig cells per testis was significantly increased in

TABLE 2. (Experiment 1). Testis Weight and Leydig Cell Size and Number in Mature Ile-de-France Rams after Immunization Against Bovine Serum Albumin or an Estradiol:BSA Conjugate*

	BSA Rams	E ₂ :BSA Rams
Testis weight (gm)	173.4 ± 11.4	219.3 ± 12.4†
Cross sectional area (μm ²)		
Peritubular Leydig cells	37.9 ± 2.2	49.0 ± 1.7†
Perivascular Leydig cells	45.9 ± 2.1	57.9 ± 3.6†
Perivascular Leydig cell nuclei	21.6 ± 0.5	25.7 ± 0.9†
Leydig cell number/testis (10 ⁶)	13.1 ± 0.34	12.3 ± 0.6
Leydig cell total volume/testis (cm ³)	2.62 ± 0.13	3.66 ± 0.31†

*Mean ± SEM, n = 5) determined from morphometric analyses of testis tissue collected in April.

†Significantly different from BSA control value, $P < 0.01$.

E₂-immunized animals although the calculated number of Leydig cells per testis was not significantly different between groups.

Experiment 2

While a small but significant ($P < 0.05$) FSH rise was observed in BSA-immunized, E₂-implanted animals following castration, a pronounced postcastration rise in both plasma FSH and LH was observed in E₂:BSA-immunized, E₂-implanted animals possessing circulating E₂ antibodies (Table 3). The acute gonadotropin response to castration in the former group and the high plasma levels of FSH and LH achieved by day 11 suggest that the biologic activity of exogenous E₂ was neutralized by the circulating antibody. Excess binding (E₂ antibody) was observed

in the plasma of E₂:BSA rams during the three sampling periods despite their continuous exposure to E₂.

Discussion

Active immunization of mature Ile-de-France rams against E₂-6-CMO-BSA resulted in the production of circulating antibodies to E₂. This method of neutralizing the biologic activity of a steroid (Nieschlag et al, 1975), and of E₂ in particular (Nishihara and Takahashi, 1983; Schanbacher, 1984), has proven itself useful in determining the biologic actions of endogenous circulating E₂. Similarly, the action of exogenous E₂ can also be blocked as shown in Experiment 2 of the present study. When compared with passive immunization, active immunization has the advantage of requiring small amounts of foreign protein (i.e., antigen vs. gamma globulin) and prolonged *in vivo* antibody production can produce a condition of chronic hormone deficiency.

Immunoneutralization of circulating E₂ in Ile-de-France rams resulted in elevated plasma FSH, LH, and testosterone concentrations. The increase in LH and testosterone secretion in E₂-immunized Ile-de-France rams confirms similar observations in E₂-immunized male rats (Nishihara and Takahashi, 1983), E₂-immunized Suffolk rams (Schanbacher, 1984) and adult crossbred rams passively immunized with an antisera raised against E₂ (Sanford, 1985). Curiously, passive immunization of Merino ram lambs against E₂ stimulated growth of the testis but did not affect mean plasma levels of FSH, LH, and testosterone (Land et al, 1981). Increased plasma testosterone appears to be the result of gonadotropin-stimulated synthesis and secretion as reflected in the 10-fold increase of the testosterone daily production

TABLE 3. (Experiment 2). Mean Plasma FSH and LH Concentrations after Castration and Estradiol Implant Administration in Bovine Serum Albumin- and Estradiol:BSA-immunized Ile-de-France Rams*

	Days Postcastration:Implantation			
	-	4	7	11
E ₂ binding titer (B/Bo at 1:1000)				
BSA	< 0.5	< 0.5	< 0.5	< 0.5
E ₂ :BSA	36.0 ± 5.8†	33.5 ± 5.9†	34.0 ± 6.3†	32.9 ± 7.8†
FSH (ng/ml)				
BSA	1.7 ± 0.2	4.0 ± 0.2	4.2 ± 0.1	6.3 ± 1.7
E ₂ :BSA	4.5 ± 0.4†	17.4 ± 1.7†	24.4 ± 0.5†	26.8 ± 3.2†
LH (ng/ml)				
BSA	1.4 ± 0.3	1.2 ± 0.2	1.1 ± 0.1	1.0 ± 0.2
E ₂ :BSA	3.1 ± 0.7†	5.8 ± 0.3†	7.1 ± 0.5†	7.9 ± 0.5†

*Mean ± SEM, n = 5.

†Significantly different from the corresponding BSA control value, $P < 0.01$.

rate. Although the E₂ antibody cross-reacted slightly with testosterone and dihydrotestosterone, the testosterone metabolic clearance rate was unaffected by immunization. Nishihara and Takahashi (1983) drew the same conclusion in E₂-immunized male rats. In their study, active immunization employed the same antigen as that used in the present study, but the testosterone metabolic clearance rate and the testosterone daily production rate were determined utilizing the method of constant infusion of tritiated testosterone. From these results, it must be concluded that interactions between circulating E₂ antibody and testosterone are negligible and that testosterone production is increased.

Observations that concurrent increases in testosterone and plasma LH levels occur in rams immunized against E₂ lead us to conclude that E₂ action on the brain, rather than testosterone and its aromatization by the brain, accounts for a major part of steroid feedback control of LH secretion. Therefore, it appears that testicular or peripheral E₂ production is of greater physiologic significance to the ram than E₂ production derived from central aromatization. Results of this E₂ immunization experiment also suggest chronic regulation of FSH secretion by E₂. It has been reported that E₂ circulates in intact rams (Schanbacher and Ford, 1976; Sanford et al, 1982) and that E₂ implants can produce dose-dependent suppression of serum FSH in castrate rams (Schanbacher, 1979). However, the biologic significance of E₂-regulated FSH secretion remains uncertain. Perhaps low levels of E₂ synergize with testicular inhibin to regulate FSH. Interestingly, Barenton and Pelletier (1983) and Barenton et al (1983) showed that E₂ production by the ram testis parallels the onset of spermatogenesis and testosterone secretion.

The susceptibility of gonadotropin secretion (both LH and FSH) to E₂ feedback and the efficacy by which circulating E₂ antibody neutralizes the biologic actions of E₂ were demonstrated in Experiment 2. Utilizing the acutely-castrated, E₂-treated ram model (Schanbacher, 1984), we observed a maintenance of plasma LH levels in BSA-immunized rams comparable to that of precastration or normal intact ram values for 11 days (until the E₂ implants were removed). Plasma FSH levels during the same postcastration:implant sampling period were only moderately increased over intact levels, suggesting that E₂ may also participate in the feedback regulation of FSH secretion. In contrast, both plasma LH and FSH were increased markedly in E₂:BSA rams (Table 3). We conclude from these data that the circulating E₂ antibodies are

capable of neutralizing (inactivating) exogenous as well as endogenous sources of estrogen.

Testosterone secretion in E₂:BSA-immunized rams was markedly enhanced and suggests that E₂ immunization may provide a means for elevating circulating androgen levels, particularly during the nonbreeding season (Schanbacher, 1984). Elevated peripheral testosterone levels were associated with hypertrophy of both peritubular and perivascular Leydig cells. This response to E₂ immunization may have been mediated peripherally (indirectly) as a result of increased gonadotropin release or perhaps, via circulating E₂ antibody, directly upon the testis. For example, the high affinity E₂ antibodies within the testicular vascular system may quickly and efficiently neutralize E₂ of testicular origin to remove its possible inhibitory influence on steroidogenic enzymes within the testis. The presence of estrogen receptors in Leydig cells (Brinkman et al, 1972; van Beurden-Lamers et al, 1974) and the observed inhibitory effects of E₂ on testicular steroid biosynthesis *in vivo* (Hsueh et al, 1978; Kalla et al, 1980; Nozu et al, 1981) and *in vitro* (Brinkman et al, 1980) support an existing hypothesis for possible direct effects of E₂ on the testis.

In conclusion, plasma FSH, LH, and testosterone levels are increased in mature Ile-de-France rams during the spring, or nonbreeding season, following E₂ immunization. These responses could prove beneficial for several facets of male reproductive function, including spermatogenesis, sperm cell viability, and libido.

Acknowledgment

The authors wish to thank Christine Perreau, Didier Chesneau, Odile Moulin, and Dr. Michel Terqui for their technical assistance during this study, and Debbie Brown for typing this manuscript.

References

- Baird DT, Horton R, Longcope C, Tait JF. Steroid dynamics under steady-state conditions. *Recent Prog Horm Res* 1969; 25: 611-664.
- Barenton B, Pelletier J. Seasonal changes in testicular gonadotropin receptors and steroid content in the ram. *Endocrinology* 1983; 112:1441-1446.
- Barenton B, Hochereau-de Reviers MT, Perreau C, Saumande J. Changes in testicular gonadotropin receptors and steroid content through postnatal development until puberty in the lamb. *Endocrinology* 1983; 112:1447-1453.
- Blanc MR, Poirier JC. A new homologous radioimmunoassay for ovine follicle stimulating hormone: development and characterization. *Ann Biol Anim Biochem Biophys* 1979; 19:1011-1026.
- Brinkman AO, Leemborg FG, Roodnat EM, DeJong FH, van der Molen HJ. A specific action of estradiol on enzymes involved in testicular steroidogenesis. *Biol Reprod* 1980; 23:801-809.
- Brinkman AO, Mulder E, Lamars-Stahlhofen GJM, van der Molen HJ. An estradiol receptor in rat testis interstitial

- tissue. *FEBS Lett* 1972; 26:301-305.
- deJong FH, Ulienbroek J Th J, van der Molen HJ. Oestradiol-17 β , testosterone, and gonadotropins in oestradiol-17 β -treated intact adult male rats. *J Endocrinol* 1975; 65:281-282.
- D'Occhio MJ, Schanbacher BD, Kinder JE. Androgenic and oestrogenic steroid participation in feedback control of luteinizing hormone secretion in male sheep. *Acta Endocrinol, Copenh* 1983; 102:499-504.
- Dorrington JT, Fritz IB, Armstrong DT. Control of testicular estrogen synthesis. *Biol Reprod* 1978; 18:55-64.
- Freund JE, Livermore PE, Miller I. *Experimental statistics*. Englewood Cliffs, NJ: Prentice-Hall, Inc., 1960; 16-18.
- Garnier DH, Cotla Y, Terqui M. Androgen radioimmunoassay in the ram: results of direct plasma testosterone and dehydroepiandrosterone measurement and physiological evaluation. *Ann Biol Anim Biochem Biophys* 1978; 18:265-281.
- Glass JD, Amann RP, Nett TM. Effects of season and sex on the distribution of cytosolic estrogen receptors within the brain and the anterior pituitary gland of sheep. *Biol Reprod* 1984; 30:894-902.
- Hochereau-de Reviere MT, Blanc MR, Cahoreau C, Courrot M, Dacheux JL, Pisselet C. Histological testicular parameters in bilateral cryptorchid adult rams. *Ann Biol Anim Biochem Biophys* 1979; 19:1141-1146.
- Hsueh AJW, Dufau ML, Catt KJ. Direct inhibitory effect of estrogen on Leydig cell function of hypophysectomized rats. *Endocrinology* 1978; 103:1096-1102.
- Kalla NR, Nisula BC, Menard R, Loriaux D. The effect of estradiol on testicular testosterone biosynthesis. *Endocrinology* 1980; 106:35-39.
- Land RB, Baird DT, Carr WR. Increased testicular growth of Tasmanian Merino ram lambs treated with antisera to oestrogens. *J Reprod Fertil* 1981; 62:151-158.
- Longcope C, Pratt JH, Schneider SH, Fineberg SE. Aromatization of androgens by muscle and adipose tissue in vivo. *J Clin Endocrinol Metab* 1978; 46:146-151.
- Lunstra DD, Schanbacher BD. Leydig cell morphology and testosterone secretion in cryptorchid Suffolk rams. *Biol Reprod* 1987 (In press).
- Melner MH, Abney TO. The direct effect of 17 β -estradiol on LH-stimulated testosterone production in hypophysectomized rats. *J Steroid Biochem* 1980; 13:203-210.
- Moger WH. Direct effect of estrogens on the endocrine function of the mammalian testis. *Can J Physiol Pharmacol* 1980; 58:1011-1022.
- Nieschlag E, Usadel KH, Wickings EJ, Kley HK, Wuttke W. Effects of active immunization with steroids on endocrine and reproductive functions in male animals. In: Nieschlag E, ed. *Immunization with hormones in reproduction research*. Amsterdam: North-Holland Publishing Co., 1975; 155-172.
- Nishihara M, Takahashi M. Effects of active immunization against oestradiol-17 β on luteinizing hormone and testosterone in male rats. *Biol Reprod* 1983; 29:1092-1097.
- Nozu K, Matsuura S, Catt KJ, Dufau ML. Modulation of Leydig cell androgen biosynthesis and cytochrome P-450 levels during estrogen treatment and human chorionic gonadotropin-induced desensitization. *J Biol Chem* 1981; 256:10012-10017.
- Payne AH, Kelch RP, Musich SS, Halpern ME. Intratesticular site of aromatization in the human. *J Clin Endocrinol Metab* 1976; 42:1081-1087.
- Pelletier J, Caraty A. Characterization of cytosolic 5 α -DHT and 17 β -estradiol receptors in the ram hypothalamus. *J Steroid Biochem* 1981; 14:603-611.
- Pelletier J, Garnier DH, de Reviere MM, Terqui M, Ortavant R. Seasonal variation in LH and testosterone release in rams of two breeds. *J Reprod Fertil* 1982; 64:341-346.
- Riggs BL, Malven PV. Spontaneous patterns of LH release in castrate male sheep and the effects of exogenous estradiol. *J Anim Sci* 1974; 38:1239-1244.
- Roosen-Runge EC, Giesel LO. Quantitative studies on spermatogenesis in the albino rat. *Am J Anat* 1950; 1:1-30.
- Sanford LM. Evidence that estrogen regulation of testosterone secretion in adult rams is mediated by both indirect (gonadotropin dependent) and direct (gonadotropin independent) means. *J Androl* 1985; 6:306-314.
- Sanford LM, Simaraks S, Palmer WM, Howland BE. Circulating estrogen levels in the ram: influence of season and mating and their relationship to testosterone levels and mating frequency. *Can J Anim Sci* 1982; 62:85-93.
- Schanbacher BD. The regulation of FSH secretion in rams. *J Reprod Fertil [Suppl]* 1979; 26:15-16.
- Schanbacher BD. Regulation of luteinizing hormone secretion in male sheep by endogenous estrogen. *Endocrinology* 1984; 115:944-950.
- Schanbacher BD, Ford JJ. Seasonal profiles of plasma luteinizing hormone, testosterone and estradiol in the ram. *Endocrinology* 1976; 99:752-757.
- Swerdlow RS, Odell WD. Feedback control of male gonadotropin secretion. *Lancet* 1968; 2:683.
- Terqui M. Contribution a l'etude des oestrogenes chez la brebis et la truie. These D'Etat, L'Universite PARIS, 1978.
- Thieulant ML, Pelletier J. Evidence for androgen and estrogen receptors in castrated ram pituitary cytosol: influence of time after castration. *J Steroid Biochem* 1979; 10:677-687.
- Van Beurden-Lamers WMD, Brinkman AO, Mulder E, van der Molen HJ. High affinity binding of oestradiol-17 β by cytosols from testis interstitial tissue, pituitary, adrenal, liver and accessory sex glands of the male rat. *Biochem J* 1974; 140:495-502.
- Worgul TJ, Santen RJ, Samojlik E, Irwin G, Falvo RE. Evidence that brain aromatization regulates LH secretion in the male dog. *Am J Physiol* 1981; 241:E246-E249.