



Foraging at Solid Urban Waste Disposal Sites as Risk Factor for Cephalosporin and Colistin Resistant *Escherichia coli* Carriage in White Storks (*Ciconia ciconia*)

OPEN ACCESS

Edited by:

Sebastian Guenther,
University of Greifswald, Germany

Reviewed by:

Jian-Hua Liu,
South China Agricultural University,
China
Dorina Timofte,
The University of Liverpool,
United Kingdom

***Correspondence:**

Ursula Höfle
ursula.hofle@uclm.es
Lourdes Migura-Garcia
lourdes.migura@irta.cat

† These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Antimicrobials, Resistance
and Chemotherapy,
a section of the journal
Frontiers in Microbiology

Received: 30 December 2019

Accepted: 29 May 2020

Published: 28 July 2020

Citation:

Höfle U, Jose Gonzalez-Lopez J,
Camacho MC, Solà-Ginés M,
Moreno-Mingorance A,
Manuel Hernández J, De La Puente J,
Pineda-Pampliega J, Aguirre JI,
Torres-Medina F, Ramis A, Majó N,
Blas J and Migura-Garcia L (2020)
Foraging at Solid Urban Waste
Disposal Sites as Risk Factor
for Cephalosporin and Colistin
Resistant *Escherichia coli* Carriage
in White Storks (*Ciconia ciconia*).
Front. Microbiol. 11:1397.
doi: 10.3389/fmicb.2020.01397

Ursula Höfle^{1*}, Juan Jose Gonzalez-Lopez², Maria Cruz Camacho¹, Marc Solà-Ginés³,
Albert Moreno-Mingorance², Jose Manuel Hernández⁴, Javier De la Puente⁵,
Javier Pineda-Pampliega⁶, José Ignacio Aguirre⁶, Fernando Torres-Medina^{7,8},
Antoni Ramis^{3,9}, Natalia Majó^{3,9}, Julio Blas⁷ and Lourdes Migura-Garcia^{3*†}

¹ SaBio (Health and Biotechnology) Working Group, Instituto de Investigación en Recursos Cinéticos IREC (CSIC-UCLM-JCCM), Ciudad Real, Spain, ² Servei de Microbiologia, Hospital Vall d'Hebron, Universitat Autònoma de Barcelona, Barcelona, Spain, ³ IRTA, Centre de Recerca en Sanitat Animal (CRESA, IRTA-UAB), Universitat Autònoma de Barcelona (UAB), Bellaterra, Spain, ⁴ Independent Researcher, Cañada de Calatrava, Spain, ⁵ Bird Monitoring Unit, SEO/BirdLife, Madrid, Spain, ⁶ Department of Biodiversity, Ecology and Evolution, Faculty of Biology, Complutense University of Madrid, Madrid, Spain, ⁷ Departamento de Biología de la Conservación, Estación Biológica de Doñana, Consejo Superior de Investigaciones Científicas, Seville, Spain, ⁸ Department of Biology, University of Saskatchewan, Saskatoon, SK, Canada, ⁹ Departament de Sanitat i Anatomia Animals, Universitat Autònoma de Barcelona (UAB), Bellaterra, Spain

White stork (*Ciconia ciconia*) may act as a reservoir and vehicle of cephalosporin resistant (CR) *Escherichia coli*. Between 2011 and 2014, we sampled white storks from colonies exposed to different degrees of anthropic pressure across the major areas of natural distribution of white storks in Spain. Cloacal swab samples ($n = 467$) were obtained from individuals belonging to 12 different colonies from six different regions. Additionally, 70 samples were collected from recently deposited droppings at the base of nesting platforms. We phenotypically characterized *E. coli* isolates, confirmed presence of CR genes and classified plasmids. Risk factors for acquiring these genes were assessed. Overall, 8.8% (41 out of 467) storks carried CR *E. coli* in their cloaca and five (7.1%) were identified from recently deposited droppings; therefore, 46 isolates were further characterized. Of them, 20 contained *bla*_{CTX-M-1}, nine *bla*_{CMY-2}, six *bla*_{CTX-M-14}, four *bla*_{SHV-12}, three *bla*_{CTX-M-15}, two *bla*_{CTX-M-32}, one *bla*_{CTX-M-1} together with *bla*_{CMY-2}, and one *bla*_{CTX-M-1} together with *bla*_{SHV-12}. All were multidrug-resistant, and four harbored the plasmid-mediated colistin resistance *mcr-1* gene. CR genes were associated with the presence of IncI1, IncFIB, and IncN replicon families. *Xba*I-macrorestriction analysis revealed a great diversity among most of the *Xba*I-PFGE types, but indistinguishable types were also seen with isolates obtained from different locations. Clonal complex 10 was the most common among CR *E. coli* and two *bla*_{CTX-M-15} positive isolates were identified as B2-ST131. Carriage of CR *E. coli* was significantly higher in colonies located close to solid urban waste disposal sites in which foraging on human waste was more likely and in one case to cattle grazing.

The co-occurrence of *bla*_{CMY-2} and *mcr-1* on plasmids of *E. coli* isolated from wild birds as early as 2011 is of note, as the earliest previous report of *mcr-1* in wild birds is from 2016. Our study shows that foraging at landfills and in association with cattle grazing are important risk factors for the acquisition of CR *E. coli* in white storks.

Keywords: *Escherichia coli*, virulence factors, white stork *Ciconia*, solid urban waste landfills, cephalosporin resistance

INTRODUCTION

Over the past years, the presence of antimicrobial resistant bacteria in wildlife, particularly cephalosporin resistant (CR) *Escherichia coli* has become a problem of increasing concern in public health (Guenther et al., 2011). The genes encoding antimicrobial resistance are frequently located on plasmids, which may mediate the horizontal transfer of such genes to other bacteria (Carattoli, 2013; Dierikx et al., 2013). The overuse of antimicrobials in both human and veterinary medicine has been considered one of the main factors contributing to the dissemination of antimicrobial resistant bacteria, increasing their detection in humans, food-producing animals, and food (Jouini et al., 2007; Smet et al., 2010). Furthermore, antimicrobial residues from urban and livestock sources may persist over time in soil and aquatic environments (Kemper, 2008). As a consequence, the emergence of antimicrobial resistant bacteria in the environment may be facilitated (Martínez, 2009), representing a global threat of major concern for human, animal, and environmental health.

Some studies have demonstrated the presence of biologically active antibiotic residues in animal and human waste, such as sewage and manure (Kummerer, 2009; Cycoń et al., 2019; Menz et al., 2019). In the case of wild birds, most of the studies have associated the influence of human activities such as farming, presence of dumpsites, or even tourism with the detection of antibiotic resistant bacteria (Allen et al., 2010; Ahlstrom et al., 2019). In addition, several reports have suggested the importance of wildlife in the dissemination of CR *E. coli* (Pinto et al., 2010; Veldman et al., 2013; Báez et al., 2015). Since wild birds in their natural environment are not treated with antibiotics (Santos et al., 2013), they are potential sentinels of multidrug resistant bacteria discharged into the environment.

The role of wild birds as disseminators of antibiotic-resistant bacteria between distant ecosystems is difficult to estimate (Báez et al., 2015; Ahlstrom et al., 2019). White storks (*Ciconia ciconia*), like many other free-living migratory birds, can become long-distance vectors of CR *E. coli*. They feed nearby pastures and plowed fields, marshy wetlands, rice fields, and more recently on solid human waste disposal sites, and display long-distance movement patterns within and between continents (Flack et al., 2016; Bécares et al., 2019). Due to their mobility and contact with fecal contamination in pastures and surface waters, they may effectively acquire and spread disease and antimicrobial-resistant bacteria (Szczepanska et al., 2015) and become a potential source of CR *E. coli* for humans and farm animals (Tryjanowski et al., 2006; Keller et al., 2011). White storks may breed in open farmland with access to marshy wetlands but also, very often,

they cohabit with humans, making use of manmade facilities such as roofs of buildings, telephone or electric power line poles, or other constructions for their nests. If such individuals acquire CR *E. coli* during feeding, they can act as potential reservoirs of resistant bacteria (Liakopoulos et al., 2016a,b).

There have been several studies reporting detection of CR *E. coli* in wild birds (Veldman et al., 2013; Zurfluh et al., 2013; Báez et al., 2015) and Alcalá et al. (2016) described for the first time the presence of CR *E. coli* in a white stork in Spain. However, to our knowledge this is the first study to analyze carriage of CR *E. coli* in white storks, in detail, in a large number of colonies subjected to different degrees of anthropic pressure with the aim of determining the epidemiology of CR *E. coli* in this species and their role as potential spreaders of CR *E. coli*. For this purpose, the isolates have been extensively characterized by antimicrobial susceptibility testing (AST), the identification of extended-spectrum β -lactamase (ESBL) and plasmid-mediated AmpC β -lactamase (pAmpC) genes, and the plasmids harboring these genes, phylotyping, genotypic relationships and the detection of virulence genes associated to avian pathogenic *E. coli* (APEC).


MATERIALS AND METHODS

Ethics Statement

Sampling of white storks was in all cases associated to ringing or radio-tagging activities. None of the storks were specifically captured or handled for the purpose of this study. All ringing and radio-tagging activities were carried out under the pertinent permits from the local authorities (regional governments of Castilla – La Mancha, Madrid, Extremadura, Castilla y Leon, and Andalucía) and by registered specialized personnel. Handling and sampling of the storks was carried out following all applicable international, national, and/or institutional guidelines for the care and ethical use of animals, specifically directive 2010/63/EU and Spanish laws 9/2003 and 32/2007, and Royal decrees 178/2004 and 1201/2005.

Field Sampling

The study area covered most of the natural distribution of white storks in Spain. We collected cloacal swab samples ($n = 467$) from storks from 12 different colonies located in five different regions. Sampling occurred more intensively and in consecutive years in colonies situated in South-central Spain. Hence, six colonies in South-central Spain were sampled in 2013 and 2014. From two colonies, samples were obtained only in 2013 and four additional colonies were sampled only in 2014 (**Figure 1** and **Table 1**).



In addition, recently deposited droppings were recovered at the base of the nesting platforms in three of the six colonies in South-central Spain in 2011.

To take specific aspects of the feeding and spatial ecology of white storks into account, samples were obtained during the reproductive season (approximately February–July), when white storks are spatially bound to their colony/nest. During this period, foraging occurs mostly in a spatially defined area and can be linked fairly clearly to a specific type of habitat/food source, allowing for the comparison of the effect of feeding on natural and/or human origin (waste) resources (Alonso et al., 1991; Tortosa et al., 2002). In contrast, during the migration and wintering periods, the spatial distribution of foraging increases, as well as the intensity of use of human waste disposal sites as a predictable food source, making association of storks to a specific habitat more difficult (Ciach and Kruszyk, 2010; Bécares et al., 2019). The continuous predictable availability of food on landfills has a profound effect on the migratory patterns of several avian species including white storks (Plaza and Lambertucci, 2017). While juvenile white storks and some adult birds still migrate to Africa, the majority have shortened their migratory distance considerably. As a result, many Spanish white storks winter in Spain or close to, landfills and are joined by birds from central and northern Europe, that either are on stopover on their route toward Africa, or remain in Spain for the winter (Gilbert et al., 2016; Bécares et al., 2019).

Samples included, on one hand recently deposited droppings collected below nesting platforms in three colonies in South-central Spain ($n = 70$) in March 2011 (just prior to egg-laying), thus presumably belonging to the adults breeding on each particular platform. On the other hand, sampling comprised cloacal swab samples from adult and nestling white storks collected during the breeding period (May–July) in 2013 ($n = 226$) and 2014 ($n = 241$).

The largest part of the cloacal swab samples (one per bird $n = 441$) was collected from chicks during the ringing activities performed at their nests (nestlings). This allowed for the collection of a considerable number of samples without the need of stressful, costly and time-consuming capture. In addition, cloacal swab samples were obtained from 26 breeding adults captured for satellite/gps data logger transmitter fitting, in their nests (the presence of small chicks at their nest was certified), meaning that the samples were obtained in a period where foraging was also concentrated in a defined type of habitat. Tarsus length and body weight were recorded in all white storks in order to enable calculation of a scaled mass index (SMI), subsequently used as a proxy for body condition.

Recently deposited droppings were collected using sterile cotton tip swabs and small zip-lock bags. The samples were kept at 4°C until arrival at the laboratory and divided into a subsample for storage at –80°C and a sample for immediate processing. Cloacal swab samples were obtained using sterile

TABLE 1 | Location, distance to closest human waste disposal site, presence of livestock, and type, and number of samples and presence of CR *E. coli* obtained from the different white stork colonies.

Colony designation	Province	Distance to human waste disposal site (km)	Live-stock	Year	Age	Number of samples	CR <i>E. coli</i>
Abenójar	Ciudad Real	60.95	Extensive sheep	2011	Droppings	23	0 (0%)
				2013	Nestling	28	0 (0%)
					Adult	2	0 (0%)
					Nestling	29	0 (0%)
Almodóvar	Ciudad Real	50.58 (formerly 1 km)	Extensive sheep	2011	Droppings	10	0 (0%)
				2013	Nestling	30	2 (6.7%)
					Adult	2	0 (0%)
					Nestling	30	3 (10%)
Alcázar de San Juan	Ciudad Real	0.77	Extensive sheep	2011	Droppings	37	5 (13.5%)
				2013	Nestling	8	0 (0%)
					Adult	3	0 (0%)
					Nestling	13	1 (7.7%)
Cabañeros	Ciudad Real	64.91	No	2013	Nestling	15	0 (0%)
					Adult	2	0 (0%)
				2014	Nestling	19	0 (0%)
Alcalá de Henares	Madrid	2.609	No	2014	Nestling	14	0 (0%)
Pinilla del Valle	Madrid	30.417	Extensive cattle	2013	Nestling	31	0 (0%)
					Adult	2	0 (0%)
				2014	Nestling	29	0 (0%)
Prado Herrero	Madrid	11.425	Cattle (breeder)	2013	Nestling	30	6 (20%)
				2014	Nestling	57	8 (14%)
Perales del Río	Madrid	6.343	No	2014	Nestling	20	2 (10%)
Malpartida	Cáceres	6.547	Extensive sheep	2013	Nestling	43	16 (37.2%)
					Adult	1	0 (0%)
Doñana, Dehesa de Abajo	Sevilla	33.203	Extensive cattle	2014	Nestling	18	1 (5.6%)
					Adult	7	2 (28.6%)
Doñana, Matagordas	Huelva	40.370	Extensive cattle	2014	Adult	6	0 (0%)
León	León	38.279	Extensive cattle	2013	Nestling	28	0 (0%)
					Adult	1	0 (0%)

cotton swabs in AMIES transport medium (Deltalab, Barcelona, Spain). They were kept at 4°C until arrival to the laboratory, where they were processed in less than 12 h after sampling. Samples were plated onto MacConkey agar (Oxoid, Basingstoke, United Kingdom) supplemented with 4 mg/L cefotaxime (Sigma-Aldrich Chemical, Madrid, Spain), and incubated overnight at 37°C. Lactose positive colonies morphologically compatible with *E. coli* were considered indicative of growth above breakpoint concentrations of cefotaxime based on the Clinical and Laboratory Standards Institute (CLSI) document, describing clinical resistance for cefotaxime ≥ 4 (CLSI, 2020). From these plates, three lactose-positive colonies were stored at -80°C in 30% glycerol BHI (Brain-Heart Infusion broth, Scharlau Microbiología, Barcelona, Spain). Subsequently, one representative colony was selected for further

characterization in the present study and confirmed as *E. coli* by PCR (Heininger et al., 1999).

Antimicrobial Susceptibility Testing

The combined disc method was used for the confirmatory tests for ESBL-producing *E. coli* (CLSI, 2020). Therefore, the following discs were used: cefotaxime, 30 μg ; cefotaxime + clavulanic acid, 30 + 10 μg ; ceftazidime, 30 μg ; ceftazidime + clavulanic acid, 30 + 10 μg (Oxoid, Basingstoke, United Kingdom). For the identification of pAmpC-producing *E. coli*, the cefoxitin disc (30 μg) was used. Additionally, the cefepime disc (30 μg) was also used. Moreover, all isolates were subjected to susceptibility testing by broth microdilution (VetMIC GN-mo, National Veterinary Institute, Uppsala, Sweden) to the following antimicrobial agents: ampicillin (1–128 mg/L), cefotaxime

(0.016–2 mg/L), ceftazidime (0.25–16 mg/L), nalidixic acid (1–128 mg/L), ciprofloxacin (0.008–1 mg/L), gentamicin (0.12–16 mg/L), streptomycin (2–256 mg/L), kanamycin (8–16 mg/L), chloramphenicol (2–64 mg/L), florfenicol (4–32 mg/L), trimethoprim (1–128 mg/L), sulfamethoxazole (8–1,024 mg/L), tetracycline (1–128 mg/L), and colistin (0.5–4 mg/L). The *E. coli* ATCC 25922 was used as a control strain. Isolates were considered to be wild-type or non-wild-type based on epidemiological cut-off (ECOFF) values defined by EUCAST¹. ECOFF separates wild-type populations from isolates that have developed reduced susceptibility for the antimicrobial and may differ from clinical breakpoints used for clinical settings (EFSA, 2017).

Antimicrobial Resistance Genes

All cefotaxime resistant (CR) isolates were tested by PCR methods for the presence of the *bla*_{CTX-M}, *bla*_{SHV}, *bla*_{CMY-1}, and *bla*_{CMY-2} genes as previously described by Hasman et al. (2005). The isolates showing resistance to colistin were also tested for the presence of *mcr* genes (Rebelo et al., 2018). Sequencing of both strands of amplicons was performed.

Phylogeny, Macrorestriction Analysis and Multi Locus Sequence Typing

The isolates were classified in phylogenetic groups (A, B1, B2, C, D, E, or F) by PCR, according to a method previously described by Clermont et al. (2000, 2013).

Genomic DNA macrorestriction analysis by pulsed-field gel electrophoresis (PFGE) was performed to determine the genomic relatedness among the isolates. The experiments were carried out as described in the PulseNet protocol (Ribot et al., 2006). *Salmonella enterica* serovar Branderup H9812 was used as a size marker. The results were analyzed by Fingerprinting II Informatix software (Applied Maths, Sint-Martens-Latem, Belgium). Isolates were considered to have a different PFGE-type when changes in at least one band were detected after digestion with *Xba*I. The analysis of the PFGE-types was carried out using the Dice similarity coefficient and the unweighted-pair group method with arithmetic averages (optimization of 1.25% and position tolerance of 1.25%).

Multi locus sequence typing (MLST) was performed according to the protocol and primers specified on the *E. coli* MLST website² (Wirth et al., 2006). Sequences were analyzed using Vector NTI advance 11 (InforMax, Inc., Bethesda, MD, United States). The allelic profile and the sequence types (STs) were obtained through the electronic database at the *E. coli* MLST website.

Whole-genome sequencing (WGS) was performed with *E. coli* isolates which revealed a new allele sequence or new allele combination after the Sanger sequencing. WGS was performed on a MiSeq Instrument (Illumina, San Diego, CA, United States) using the Nextera DNA Flex Library Prep Kit (Illumina) and paired-end sequenced with a MiSeq Reagent kit v3 (Illumina) with an average insertion size of 300 bp.

The obtained sequences were submitted to Enterobase³ to obtain new allele/STs designation.

Detection of Virulence-Associated Genes

All 46 isolates were tested by multiplex PCR for the genes described previously by Johnson et al. (2008) as the minimal predictors of APEC virulence; *iroN*, *ompT*, *hlyF*, *iutA*, and *iss* (Figure 2).

Mating Experiments and Plasmid Characterization

Filter mating experiments were carried out to assess the transferability of ESBL-encoding plasmids by conjugation. For the analysis, one representative isolate from each PFGE-type was selected. Mating assays were performed as previously described (Bielak et al., 2011), using the isolates as donors and rifampicin-resistant *E. coli* HB101 as a recipient. Transconjugants were selected on LB agar plates supplemented with rifampicin (150 mg/L) and ceftriaxone (1 mg/L). The identity of the transconjugants was confirmed by PFGE and the presence of the corresponding CR gene was confirmed by PCR.

Plasmid DNA was purified from the subset of 42 isolates (one isolate as representative of PFGE-type) using a Qiagen Plasmid Midi kit (Qiagen, Hilden, Germany) according to the manufacturer's recommendations. Plasmids were introduced to electrocompetent plasmid free *E. coli* BL21 cells by electroporation. Transformants were selected in LB agar plates containing ceftriaxone (1 mg/L), and PCR assays were performed to confirm the presence of ESBL and/or pAmpC genes. The presence of a unique plasmid in the transformants and their corresponding sizes were analyzed using S1 nuclease digestion followed by PFGE (S1-PFGE) (Barton et al., 1995). Additionally, plasmids were classified by PCR-based replicon typing (Carattoli et al., 2005).

Statistical Analysis

As a proxy to the degree of exposure of storks to human waste foraged food, we used the distance of each colony to the nearest human waste disposal site. This is based on the reduced home range of storks during chick raising and incubation and the assumption that storks are less likely to undertake long foraging flights during this period as compared to other stages. This hypothesis is supported by satellite transmitter data from radio-tagged adult storks from colonies in South-central Spain that show distinct foraging either in natural habitat or at human waste disposal sites (Bécares et al., 2019). In addition, the distance between a stork's nest and solid human waste landfills has been shown to correlate with the frequency and intensity with which the storks forage on waste (Djerdali et al., 2016; Gilbert et al., 2016).

For this, the coordinates for each colony/nest sampled, in parallel to the coordinates of all known human waste disposal sites, were projected onto a scaled flat reference system in the geographic information system QGIS (QGIS desktop 2.8.1),

¹<http://www.eucast.org/>

²<http://mlst.warwick.ac.uk/mlst/dbs/Ecoli>

³<http://enterobase.warwick.ac.uk/>

ECOFFs, to at least three antimicrobial families). Furthermore, 54% (25/46) exhibited a non-wild-type phenotype to more than seven antimicrobial classes. MIC values determined that all isolates exhibited a non-wild-type phenotype to cephalosporins (100% resistance to cefotaxime and 96% to ceftazidime), with 20 isolates yielding amplicons for *bla*_{CTX-M-1}, nine *bla*_{CMY-2}, six *bla*_{CTX-M-14}, four *bla*_{SHV-12}, three *bla*_{CTX-M-15}, two *bla*_{CTX-M-32}, one *bla*_{CTX-M-1} together with *bla*_{CMY-2}, and one *bla*_{CTX-M-1} together with *bla*_{SHV-12}. Additionally, 100% of the isolates were non-wild-type to ampicillin, 85% to sulfamethoxazole, 80% to ciprofloxacin and nalidixic acid, 67% to tetracycline, 65% to trimethoprim, 54% to streptomycin, 33% to gentamicin, 22% to kanamycin, 9% to chloramphenicol, and 7% to florfenicol (Figure 2).

In addition, four isolates from recently deposited droppings collected at nests located at the colony closest to a landfill (Alcázar de San Juan, Table 1) were phenotypically resistant to colistin (MIC ≥ 4) and were confirmed to carry the *mcr-1* gene (id. 173, 176, 178, and 184).

Phylogeny, PFGE, and MLST

The phylotyping analysis identified six different phylogenetic groups among the 46 isolates (Figure 2). Of them, 14 belonged to phylogroup B1 (30%), 12 to group A (26%), eight to group E (17%), four to group B2 (9%), four to group F (9%), and one to group C (2%). No phylogroup could be determined for three of the isolates (7%).

Among the 46 CR *E. coli* isolates, 42 different macro-restriction profiles were detected. However, during 2013, a cluster of isolates sharing a similar PFGE profile (100% similarity) and identical MLST type (ST2309) was identified. These isolates (506, 626, and 635) were obtained from two different colonies located within a radius of about 250 km, in Madrid and Cáceres.

The MLST analysis also revealed a high degree of diversity. Overall, 30 different STs were found among the 46 isolates (mean of number of isolates per ST, 1.53; range, 1–6) (Figure 2).

ST10 was the most prevalent type (6 isolates; 13%), followed by ST155, ST602, and ST2309 (3 isolates; 6.5%, each) and by ST48, ST93, ST131, ST457, and ST1844 (2 isolates; 4.3%, each). ST8877 was firstly identified in this study. Grouping STs by clonal complexes (CC) showed that CC10 was the most prevalent including 22% of the isolates, followed by CC38, CC155, CC168, and CC446 in the 3% of the population respectively (Figure 2).

Detection of Virulence Genes Associated to APEC

Out of 46 isolates, 70, 70, 70, 57, and 52% yielded amplicons for *ompT*, *hlyF*, *iutA*, *iss*, and *iroN*, respectively. Accordingly, 25 isolates (54%) were considered APEC, since they had four to five of the genes mentioned above (Figure 2).

Conjugation and Transformation Experiments

Among the 46 CR isolates, a subset of 42 representing each *Xba*I-macrorestriction profile were selected for conjugation and transformation experiments. Of them, 15 were able to transfer the CR genes by conjugation. Additionally, 33 transferred the CR genes to the BL21 electrocompetent strain. S1-nuclease PFGE confirmed the presence of a unique plasmid in all selected isolates except in two of them (518 and 598), which contained more than one plasmid. Sizes of plasmids varied between 40 and 150 kb approximately (Table 2). PCR-based replicon typing showed that IncII was the most common replicon and was associated to *bla*_{CTX-M-1}, *bla*_{CTX-M-14}, *bla*_{CMY-2}, and *bla*_{SHV-12} genes. IncII was present in 25 of the isolates, followed by IncFIB and IncN. The replicons IncFIA, IncFIC, IncA/C, IncK, and IncL/M were also detected (Table 2). Four isolates presented between two and five replicons on the same plasmid (517, 631, 176, and 184), and none of the tested replicons were detected in seven of the transformants or transconjugants (434, 514, 554, 508, 173, 626, and 634).

TABLE 2 | Characteristics of plasmids harboring ESBL and/or pAmpC genes from 42 cephalosporin-resistant *E. coli* isolates.

Gene and isolate	Conjugation result	Transformation result	Inc type(s)							Plasmid size (kb)
			II	FIB	N	K	FIA	FIC	A/C	
<i>bla</i> _{CTX-M-1}										
516	TC1b		+							100
524	TC3a		+							90
530.13		TF4a	+							60
586	TC5a	TF5a	+							100
598	–	–								
597	TC7a	TF7a	+							100
610		TF9a	+							100
615		TF10b	+							90
619		TF11a	+							100
627		TF12a	+							100

(Continued)

TABLE 2 | Continued

Gene and isolate	Conjugation result	Transformation result	Inc type(s)							Plasmid size (kb)	
			I1	FIB	N	K	FIA	FIC	A/C		L/M
706		TF14a	+								100
433		TF15a	+								110
434	TC16b	TF16a									110
495	TC17a	TF17a		+							50
514		TF18a									60
520		TF19a	+								100
522		TF38a	+								110
530.14		TF20a			+						40
554	TC21a	TF21a									50
<i>bla</i> _{CTX-M-32}											
517		TF2a		+				+	+		130
593		TF6a	+								100
<i>bla</i> _{CTX-M-14}											
626		TF30a									110
547		TF31a	+								110
582	TC32a	TF32a				+					80
584	TC33a	TF33a	+								90
<i>bla</i> _{CTX-M-15}											
631		TF13a		+			+				150
634	TC39b										60
518	-	-									
<i>bla</i> _{CMY-2}											
599		TF22a							+		120
508		TF23a									100
565	TC24a	TF24a	+								80
173		TF25a									90
176		TF26a	+	+	+			+		+	90
178		TF27a	+								90
184	TC28b		+		+	+					40
215		TF29b			+						40
<i>bla</i> _{SHV-12}											
587	TC34a		+								90
600	TC35c		+								90
605		TF36a	+								100
606		TF8b	+								100
<i>*bla</i> _{CTX-M-1} + <i>bla</i> _{CMY-2}											
519		TF37a	+								100
<i>*bla</i> _{CTX-M-1} + <i>bla</i> _{SHV-12}											
465	TC40a			+							110

Transconjugants (TC) and transformants (TF) used for PCR-based replicon typing method are in boldface. Two isolates (518 and 598) presented more than one plasmid in the TC and/or TF isolates, therefore they are not included in the table. *Only *bla*_{CTX-M-1} was transferred to the transformant and transconjugant isolate, respectively.

Risk Factors for the Acquisition of CR *E. coli* by White Storks

Cephalosporin resistant *E. coli* prevalence did not vary significantly between nestling (8.8%, 39 out of 441) and adult storks (7.7%, 2 out of 26). Sampling year [$F(1,359) = 2.538$, $p = 0.112$], Sex [$F(1,359) = 0.517$, $p = 0.473$], number of siblings in the nest [$F(1,359) = 1.645$, $p = 0.179$], and livestock grazing close to the colony [$F(2,359) = 1.78$, $p = 0.17$] did not affect CR *E. coli* carriage in white stork nestlings. However, distance of the colony to human waste disposal sites had a significant effect on

the prevalence of CR *E. coli* [GLMM $F(1,359) = 8.641$, $p = 0.004$], with nestlings from colonies close to landfills being significantly more likely to carry CR *E. coli* (Figure 1 and Table 3). Carriage of CR *E. coli* [Student's t -test $F(1,359) = 2.946$, $p = 0.258$] or APEC [Student's t -test $F(1,359) = 1.921$, $p = 0.164$] had no significant effect on the body condition of white stork nestlings. However, white stork nestlings from colonies close to human waste disposal sites were in significantly better body condition than nestlings from colonies far from human waste disposal sites [ANOVA $F(7,341) = 8.469$, $p < 0.001$].

TABLE 3 | Results of generalized mixed linear model (GLMM) for cephalosporin resistant (CR) *E. coli* carriage in white stork nestlings in Spain.

Model	Predictor	Test model effects	Description	F	Standard Error	t	p
CR <i>E. coli</i>	Livestock presence	0.304	None	0.468	1.121	0.418	0.677
			Ext sheep	-1.08	0.758	-1.424	0.155
			Ext cattle	0	-	-	-
	Sex	0.170	F	0.319	0.444	0.719	0.473
			M	0	-	-	-
	Sampling year	0.112	2013	-0.921	0.578	-1.593	0.112
			2014	0	-	-	-
	Distance landfill (m)	0.004	Continuous	0.000	0.000	2.909	0.004
	Number of siblings in nest	0.179	Single chick	1.729	0.921	1.877	0.061
			Two siblings	1.54	0.775	1.988	0.048
			Three siblings	0.816	0.661	1.234	0.218
			Four siblings	0	-	-	-

Statistically significant results are highlighted in bold.

DISCUSSION

White storks are free-living colonial birds that frequently use humanized habitats. In the recent past, they have adapted to foraging at solid urban waste disposal sites and introduced American crayfish (*Procambarus clarkii*) in rice fields and small water bodies as predictable food sources during winter. This has led to a significant modification of migratory behavior of a large part of the adult stork population of Western Europe as well as, in the case of solid urban waste disposal sites, to the exposure to numerous contaminants and pathogens (Flack et al., 2016; Gilbert et al., 2016).

As storks cohabit in the proximities of humans and food producing animals, they are able to acquire bacteria from these and transport them to other locations in their annual or regular movements (Tryjanowski et al., 2006). The present study takes advantage of the fact that even though during winter most storks forage on human waste; during breeding the nesting site and colony determine the foraging ecology and allow for the characterization of risk factors for the exposure of white storks to CR *E. coli* (Alonso et al., 1991; Tortosa et al., 2002; Bécares et al., 2019). Our results suggest that the stork nestlings more likely fed human waste by their parents (using distance to rubbish dump as a proxy) were more likely to carry CR *E. coli*. CR *E. coli* carriage did not affect body condition in “infected” nestlings. Hence, the present study has demonstrated the presence of CR *E. coli* in white storks feeding on human waste and their capacity to disseminate and transport the resistant bacteria. In particular, similar clones with the same resistance genes have been detected in nestlings from colonies in Madrid and Cáceres, located within a 250 km distance. These results suggest a common source of contamination, together with the capacity of the birds to spread the same clones during their movements. In this context, a colony that is now at a more than 50 km flight distance from the closest human waste disposal site that was sealed in 2005, but satellite transmitter data from two adults evidenced occasional visits to this landfill even during the breeding season

(Table 1 and Figure 1). According to our results, extensive livestock farming in natural habitats (present in the proximity of at least five of six natural habitat colonies) appears to be less of a risk factor for the acquisition of CR *E. coli* by white storks. However, nestlings from one colony located on a farm that raises cattle and where storks and cattle share the pastures, also situated at a flight distance of 11 km to the nearest landfill (Prado Herrero, Table 1), most commonly carried CR *E. coli* with the *bla*_{CTX-M-1} gene, which is commonly described in cattle in Europe (Schmid et al., 2013). Storks frequently collect cattle dung and place it in the nest just prior to, and a few weeks after hatching, presumably to help chicks maintain body temperature (Tortosa and Villafuerte, 1999). This behavior as well as foraging of the adults on contaminated pastures (especially on dung beetles) is a likely route of exposure of the nestlings to CR *E. coli* excreted by cattle.

Some studies suggest that wildlife also play an important role in the global dissemination of antimicrobial resistance genes, transferring mobile genetic elements between bacterial communities (Allen et al., 2010). In our study, the presence of multidrug-resistant *E. coli* demonstrated the potential capacity of white stork to disseminate and transport ESBL and/or AmpC encoding genes located in conjugative plasmids. These isolates were all resistant to third generation cephalosporins and most of them were also resistant to fluoroquinolones. These two families of antimicrobials are nominated of critical importance in human health since they are widely used for the treatment of human infections by Gram-negative bacteria (European Centre for Disease Prevention and Control [ECDC], 2018). In addition, four of the isolates, obtained in 2011 from recently deposited droppings, co-harbored *bla*_{CMY-2} and *mcr-1*, although on different plasmids. They were also resistant to multiple antibiotics including third-generation cephalosporins, fluoroquinolones, colistin, and aminoglycosides, among others, making these isolates resistant to almost all last-line antimicrobials available. Interestingly, all *mcr-1* carrying isolates presented different PFGE profiles and belonged to the CC10, two were ST10, one was ST48 and one was ST1286. This high risk CC has previously been

reported as the most commonly associated with *mcr-1* globally, not only in humans but also in animals and in environmental samples (Caltagirone et al., 2017; Matamoros et al., 2017; García-Meniño et al., 2018; Lalaoui et al., 2019). All STs found within this study, except for ST8877, have previously been identified in third-generation CR isolates of animal origin (Dierikx et al., 2013; Solà-Ginés et al., 2015; Alcalá et al., 2016; Batalha de Jesus et al., 2019). Of note, CR *E. coli* CC10 has been reported as one of the most prevalent lineages on farms, food, and waste water plants, and the second most abundant in humans in Spain (Ojer-Usoz et al., 2017). Additionally, ST131, ST69, and ST10 are the most prevalent ExPEC lineages found in humans (Manges et al., 2019), and all of them have also been identified in this study. This fact also highlights the importance of wildlife as potential reservoirs, not only of antimicrobial resistance genes, but also of human pathogenic bacteria belonging to high-risk clones.

Several studies have suggested a relationship between the different *E. coli* phylogroups and the virulence of the isolates (Picard et al., 1999; Clermont et al., 2011). Accordingly, non-virulent isolates mainly belong to A, B1, and C phylogenetic groups, whereas the most virulent phylogroups are associated to B2 followed by D (Russo and Johnson, 2000). In this study, most of the isolates belonged to phylogroups A and B1, and only four isolates were B2. On the contrary, analyzing phylogroups and virulence genes in this strain collection, we have observed that almost all B1 isolates had the APEC virulence-associated genes, whereas only one B2 strain had the APEC genes. This may be due to the presence of virulence factors associated to mobile genetic elements such as in the case of the previously described ColV plasmid (Johnson and Nolan, 2009), which contained several virulence genes within the same plasmid (*hlyF*, *ompT*, *iss*, and *cvaC*). Therefore, the acquisition of such type of plasmid may confer B1 isolates with more virulence factors than expected. Carriage of *E. coli* isolates harboring APEC-associated genes in the cloaca of white stork nestlings was apparently not related to solid urban waste disposal site origin of food or any of the other factors; neither did it affect body condition.

The present study showed a high variability of CR genes among the isolates, *bla*_{CTX-M-1} being the most prevalent (48%), followed by *bla*_{CMY-2} and *bla*_{CTX-M-14} (20 and 13%, respectively). These *bla*_{CTX-M} types are the most frequent variants in ESBL producers in animals and food of animal origin, while *bla*_{CTX-M-15} has been mostly described in human medicine (Valverde et al., 2015) and occasionally on farms (Cameron-Veas et al., 2015). This study has demonstrated the presence of the *bla*_{CTX-M-15} gene in three isolates of white stork harbored in plasmids of different sizes and in one occasion contained in an IncFIB replicon, the most common incompatibility group for *bla*_{CTX-M-15} isolated from humans in Spain (Carattoli, 2009). Additionally, the remaining two *bla*_{CTX-M-15} isolates were B2-ST131. They were obtained from two different colonies, one of which is closely associated to a village with part of the nests on urban structures, and confirmed consumption of human residues. The other colony is in a natural environment with no confirmed consumption of human residues during the breeding season, nevertheless this colony is situated at 30.4 km flight distance from Madrid, and thus exposure to human

waste cannot be excluded completely. In fact satellite transmitter data from an adult stork from this colony shows extensive use of solid urban waste disposal sites outside the breeding period (Bécares et al., 2019).

Significant differences were found between location of the colonies and presence of CR genes. For instance, in the colony located in Cáceres, the most prevalent gene was *bla*_{CTX-M-1} associated to 90 and 100 kb plasmids, whereas in the colony located in Alcázar de San Juan (Ciudad Real) the most prevalent was *bla*_{CMY-2} harbored in two types of plasmids (40 and 90 kb). The former was also the colony in which nestlings carried the greatest variety of CR genes (5) as compared to the other colonies.

The most common replicons found in the study were IncII (60%), followed by IncFIB (12%), and IncN (10%). To date, these three replicons are the most commonly found in plasmids carrying CR genes in *E. coli* isolated from poultry (Wang et al., 2013; Solà-Ginés et al., 2015). Almost all of the CR genes containing IncII replicons were located in plasmids of the same sizes, between 90 and 110 kb; specifically, 100 kb in the case of *bla*_{CTX-M-1}. However, in the case of two isolates, isolation of the plasmid carrying the CR gene could not be achieved. During transformation and conjugation experiments, co-transfer of plasmids of different molecular weights was observed. The entry of two or more plasmids into a single transformation procedure is not uncommon; Weston et al. (1979) demonstrated that 20% of the transformants could be double transformants (Weston et al., 1979). This type of transformation can be frequent if it is performed with high amounts of plasmid DNA (Goldsmith et al., 2007).

CONCLUSION

In conclusion, results from this study confirm that exposure of white storks to human waste constitutes a risk factor for the acquisition of CR genes as the carriage of CR *E. coli* was significantly higher in colonies located close to solid urban waste disposal sites. As all individuals included in this study are free-living birds, they have not been treated with antimicrobials. Since they are not subjected to any direct antibiotic pressure and they cohabit in the proximities of humans, the presence of these resistance traits is a reflection of the anthropic pressure of human activities and particularly exposure to human waste and farming practices. In fact, outside the breeding season adult and juvenile storks have been observed to use human waste disposal sites indistinctive of the habitat of the colony of origin (Bécares et al., 2019). The distinctive pattern of prevalence of CR genes in white stork nestlings in association to foraging behavior of their parents during the breeding season, in addition to recent findings that show that a large proportion of storks are exposed to human waste from solid urban waste landfills during wintering, suggests that persistence of CR genes in bacteria of the digestive tract of white storks is relatively short. This makes white storks excellent sentinels for local antibiotic pressure in their foraging habitat. Nevertheless, due to their great mobility and migratory habits, the white stork can also act as carriers of CR

E. coli, contributing to the global dissemination of antimicrobial resistance genes and plasmids.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation, to any qualified researcher.

ETHICS STATEMENT

Ethical review and approval was not required for the animal study because handling of the storks included in the study had been approved for the specific ongoing studies in each colony, and our study was based on fecal droppings or cloacal swabs only.

AUTHOR CONTRIBUTIONS

MC, AR, NM, and UH designed the study. UH, MC, JM, JD, JP-P, JA, FT-M, and JB sampled and collected relevant biological data from the stork colonies and isolated the isolates for further characterization. MS-G and LM-G carried out the identification, phenotypic, and molecular characterization of isolates and plasmids. JJ and AM-M performed MLST and WGS. UH and MC performed the statistical analysis. UH, MS-G, and LM-G drafted the manuscript. All authors revised the draft manuscript and contributed to writing the final version of the manuscript.

FUNDING

This work was partially supported by the grants AGL2013-47852-R and RTI2018-095586-B-C22 from the Ministerio de Economía y Competitividad (MINECO), project RTA2011-00111-C03 funded by the Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria (INIA) and by the CERCA program from Generalitat de Catalunya.

REFERENCES

- Ahlstrom, C. A., Bonnedahl, J., Woksepp, H., Hernandez, J., Reed, J. A., Tibbitts, L., et al. (2019). Satellite tracking of gulls and genomic characterization of faecal bacteria reveals environmentally mediated acquisition and dispersal of antimicrobial-resistant *Escherichia coli* on the Kenai Peninsula. *Alaska. Mol. Ecol.* 28, 2531–2545. doi: 10.1111/mec.15101
- Alcalá, L., Alonso, C. A., Simon, C., González-Esteban, C., Oros, J., Rezusta, A., et al. (2016). Wild Birds, frequent carriers of extended-spectrum beta-lactamase (ESBL) producing *Escherichia coli* of CTX-M and SHV-12 Types. *Microb. Ecol.* 72, 861–869. doi: 10.1007/s00248-015-0718-0
- Allen, H. K., Donato, J., Wang, H. H., Cloud-Hansen, K. A., Davies, J., and Handelsman, J. (2010). Call of the wild: antibiotic resistance genes in natural environments. *Nat. Rev. Microbiol.* 8, 251–259. doi: 10.1038/nrmicro.2312
- Alonso, J. C., Alonso, J. A., and Carrascal, L. M. (1991). Habitat selection by foraging White Storks, *Ciconia*, during the breeding season. *Can. J. Zool.* 69, 1957–1962. doi: 10.1139/z91-270

The VHIR-HUVH was supported by Plan Nacional de I+D+i 2013–2016, Instituto de Salud Carlos III and the Spanish Ministry of Health (FIS PI15/00604) and the Subdirección General de Redes y Centros de Investigación Cooperativa, MINECO, Spanish Network for Research in Infectious Diseases (REIPI RD16/0016/0003) – co-financed by European Development Regional Fund “A way to achieve Europe. AM-M is a Ph.D. student registered with the Universidad Autònoma de Barcelona. The contract of LM-G was supported by INIA and the European Social Fund. JP-P was funded by a grant from the Complutense University of Madrid (CT45/15-CT46/15). FT-M was supported by the Isabel Maria Lopez Martinez Memorial Scholarship at the University of Saskatchewan (Canada). Trapping activities of adult birds were partially supported in the framework of SEO/BirdLife Migra program and financed by Fundación Iberdrola España and Storch Schweiz, by grant CGL2012-32544 from the MINECO, and by grant 511/2012 from the Organismo Autónomo de Parques Nacionales.

ACKNOWLEDGMENTS

We are indebted to Cabañeros National Park and the urban residues treatment plant of Alcázar de San Juan for permission of access to white stork colonies. We gratefully acknowledge the help of A. Herrera-Dueñas; the owners and workers of Prado Herrero Farm; J. Prieto in Alcalá de Henares; the forest officers in Pinilla and the ringing group Monticola, in particular R. Moréno-Opo and V. de la Torre in “La Torrecilla” for help with sample collection in the Community of Madrid, and S. Palacios and J. Ayala for field work in Seville and Doñana, as well as the contribution of the following bird ringers in sample collection in colonies in Malpartida de Cáceres: J. M. Abad, T. Álvarez, M. Giraldo, C. Domínguez, M. Iglesias y M. Cardalliaguet, and León: I. Rodríguez, B. Fuertes, D. Miguélez, I. Roa, F. de la Calzada, J. García, F. Ortega, D. Prada, and M. Cendoya (GIA-León). An earlier version of this article constitutes one of the chapters of the Ph.D. thesis of one of the authors (MS-G), published by Solà-Ginés (2018).

- Báez, J., Hernández-García, M., Guamparito, C., Díaz, S., Olave, A., Guerrero, K., et al. (2015). Molecular characterization and genetic diversity of ESBL-producing *Escherichia coli* colonizing the migratory Franklin's gulls (*Leucophaeus pipixcan*) in Antofagasta. North of Chile. *Microb. Drug Resist.* 21, 111–116. doi: 10.1089/mdr.2014.0158
- Barton, B. M., Harding, G. P., and Zuccarelli, A. J. (1995). A general method for detecting and sizing large plasmids. *Anal. Biochem.* 226, 235–240. doi: 10.1006/abio.1995.1220
- Batalha de Jesus, A. A., Freitas, A. A. R., de Souza, J. C., Martin, S. N., Botelho, L. A. B., Girão, V. B. C., et al. (2019). High-level multidrug-resistant *Escherichia coli* isolates from wild birds in a large urban environment. *Microb. Drug Resist.* 25, 167–172. doi: 10.1089/mdr.2018.0180
- Bécares, J., Blas, J., López-López, P., Schulz, H., Torres-Medina, F., Flack, A., et al. (2019). *Migration and Spatial Ecology of the White Stork in Spain. Migración y Ecología espacial de la Cigüeña Blanca en España. Monografía n.º 5 del Programa Migra.* Madrid: SEO/BirdLife.
- Bielak, E., Bergenholtz, R. D., Jorgensen, M. S., Sorensen, S. J., Hansen, L. H., and Hasman, H. (2011). Investigation of diversity of plasmids carrying the

- bla*TEM-52 gene. *J. Antimicrob. Chemother.* 66, 2465–2474. doi: 10.1093/jac/dkr331
- Caltagirone, M., Nucleo, E., Spalla, M., Zara, F., Novazzi, F., Marchetti, V. M., et al. (2017). Occurrence of extended spectrum β -Lactamases, KPC-Type, and MCR-1.2-producing *Enterobacteriaceae* from wells, river water, and wastewater treatment plants in oltrepò pavese area, Northern Italy. *Front. Microbiol.* 8:2232. doi: 10.3389/fmicb.2017.02232
- Cameron-Veas, K., Solà-Ginés, M., Moreno, M. A., Fraile, L., and Migura-García, L. (2015). Impact of the use of beta-lactam antimicrobials on the emergence of *Escherichia coli* isolates resistant to cephalosporins under standard pig-rearing conditions. *Appl. Environ. Microbiol.* 81, 1782–1787. doi: 10.1128/aem.03916-14
- Carattoli, A. (2009). Resistance plasmid families in *Enterobacteriaceae*. *Antimicrob. Agents Chemother.* 53, 2227–2238. doi: 10.1128/aac.01707-08
- Carattoli, A. (2013). Plasmids and the spread of resistance. *Int. J. Med. Microbiol.* 303, 298–304. doi: 10.1016/j.ijmm.2013.02.001
- Carattoli, A., Bertini, A., Villa, L., Falbo, V., Hopkins, K. L., and Threlfall, E. J. (2005). Identification of plasmids by PCR-based replicon typing. *J. Microbiol. Methods* 63, 219–228. doi: 10.1016/j.mimet.2005.03.018
- Ciach, M., and Kruszyk, R. (2010). Foraging of White Storks *Ciconia* on rubbish dumps on non-breeding grounds. *Waterbirds* 33, 101–104. doi: 10.1675/063.033.0112
- Clermont, O., Bonacorsi, S., and Bingen, E. (2000). Rapid and simple determination of the *Escherichia coli* phylogenetic group. *Appl. Environ. Microbiol.* 66, 4555–4558. doi: 10.1128/aem.66.10.4555-4558.2000
- Clermont, O., Christenson, J. K., Denamur, E., and Gordon, D. M. (2013). The Clermont *Escherichia coli* phylo-typing method revisited: improvement of specificity and detection of new phylo-groups. *Environ. Microbiol. Rep.* 5, 58–65. doi: 10.1111/1758-2229.12019
- Clermont, O., Olier, M., Hoede, C., Diancourt, L., Brisse, S., Keroudean, M., et al. (2011). Animal and human pathogenic *Escherichia coli* isolates share common genetic backgrounds. *Infect. Genet. Evol.* 11, 654–662. doi: 10.1016/j.meegid.2011.02.005
- CLSI (2020). *Clinical and Laboratory Standards Institute Performance Standards for Antimicrobial Susceptibility Testing: 30th Edition, Informational Supplement M100-ED30:2020*. Wayne, PA: CLSI.
- Cycoń, M., Mrozik, A., and Piotrowska-Seget, Z. (2019). Antibiotics in the soil environment—degradation and their impact on microbial activity and diversity. *Front. Microbiol.* 10:338. doi: 10.3389/fmicb.2019.00338
- Dierix, C., Van Der Goot, J., Fabri, T., Van Essen-Zandbergen, A., Smith, H., and Mevius, D. (2013). Extended-spectrum-beta-lactamase- and AmpC-beta-lactamase-producing *Escherichia coli* in Dutch broilers and broiler farmers. *J. Antimicrob. Chemother.* 68, 60–67. doi: 10.1093/jac/dks349
- Djerdali, S., Guerrero-Casado, J., and Tortosa, F. S. (2016). The effects of colony size interacting with extra food supply on the breeding success of the White Stork (*Ciconia ciconia*). *J. Ornithol.* 157, 941–947. doi: 10.1007/s10336-016-1343-5
- EFSA (2017). The European Union summary report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food in 2015. European Food Safety Authority (EFSA) and European Centre for Disease Prevention and Control (ECDC). *EFSA J.* 15:5812.
- European Centre for Disease Prevention and Control [ECDC] (2018). *Antimicrobial Consumption. In: ECDC. Annual Epidemiological Report for 2017*. Stockholm: ECDC.
- Flack, A., Fiedler, W., Blas, J., Pokrovsky, I., Kaatz, M., Mitropolsky, M., et al. (2016). Costs of migratory decisions: a comparison across eight white stork populations. *Sci. Adv.* 2:e1500931. doi: 10.1126/sciadv.1500931
- García-Meniño, I., García, V., Mora, A., Díaz-Jiménez, D., Flament-Simon, S. C., Alonso, M. P., et al. (2018). Swine enteric colibacillosis in Spain: pathogenic potential of mcr-1 ST10 and ST131 *E. coli* isolates. *Front. Microbiol.* 9:2659. doi: 10.3389/fmicb.2018.02659
- Gilbert, N. I., Correia, R. A., Silva, J. P., Pacheco, C., Catry, I., Atkinson, P. W., et al. (2016). Are white storks addicted to junk food? Impacts of human waste disposal site use on the movement and behaviour of resident white storks (*Ciconia ciconia*) from a partially migratory population. *Mov. Ecol.* 4:7.
- Goldsmith, M., Kiss, C., Bradbury, A. R., and Tawfik, D. S. (2007). Avoiding and controlling double transformation artifacts. *Protein Eng Des Sel* 20, 315–318. doi: 10.1093/protein/gzm026
- Guenther, S., Ewers, C., and Wieler, L. H. (2011). Extended-spectrum beta-lactamases producing *E. coli* in wildlife, yet another form of environmental pollution? *Front. Microbiol.* 2:246. doi: 10.3389/fmicb.2011.00246
- Hasman, H., Mevius, D., Veldman, K., Olesen, I., and Aarestrup, F. M. (2005). beta-Lactamases among extended-spectrum beta-lactamase (ESBL)-resistant *Salmonella* from poultry, poultry products and human patients in The Netherlands. *J. Antimicrob. Chemother.* 56, 115–121.
- Heininger, A., Binder, M., Schmidt, S., Unertl, K., Botzenhart, K., and Doring, G. (1999). PCR and blood culture for detection of *Escherichia coli* bacteremia in rats. *J. Clin. Microbiol.* 37, 2479–2482. doi: 10.1128/jcm.37.8.2479-2482.1999
- Johnson, T. J., and Nolan, L. K. (2009). Pathogenomics of the virulence plasmids of *Escherichia coli*. *Microbiol. Mol. Biol. Rev.* 73, 750–774. doi: 10.1128/mmb.00015-09
- Johnson, T. J., Wannemuehler, Y., Doetkott, C., Johnson, S. J., Rosenberger, S. C., and Nolan, L. K. (2008). Identification of minimal predictors of avian pathogenic *Escherichia coli* virulence for use as a rapid diagnostic tool. *J. Clin. Microbiol.* 46, 3987–3996. doi: 10.1128/jcm.00816-08
- Jouini, A., Vinue, L., Slama, K. B., Sáenz, Y., Klibi, N., Hammami, S., et al. (2007). Characterization of CTX-M and SHV extended-spectrum beta-lactamases and associated resistance genes in *Escherichia coli* isolates of food samples in Tunisia. *J. Antimicrob. Chemother.* 60, 1137–1141. doi: 10.1093/jac/dkm316
- Keller, J. I., Shriver, W. G., Waldenstrom, J., Griekspoor, P., and Olsen, B. (2011). Prevalence of *Campylobacter* in wild birds of the mid-Atlantic region. *USA. J. Wildl. Dis.* 47, 750–754. doi: 10.7589/0090-3558-47.3.750
- Kemper, N. (2008). Veterinary antibiotics in the aquatic and terrestrial environment. *Ecol. Ind.* 8, 1–13. doi: 10.1016/j.ecolind.2007.06.002
- Kummerer, K. (2009). Antibiotics in the aquatic environment—a review—part I. *Chemosphere* 75, 417–434. doi: 10.1016/j.chemosphere.2008.11.086
- Laloui, R., Djukovic, A., Bakour, S., Sanz, J., Gonzalez-Barbera, E. M., Salavert, M., et al. (2019). Detection of plasmid-mediated colistin resistance, *mcr-1* gene, in *Escherichia coli* isolated from high-risk patients with acute leukemia in Spain. *J. Infect. Chemother.* 25, 605–609. doi: 10.1016/j.jiac.2019.03.007
- Liakopoulos, A., Mevius, D. J., Olsen, B., and Bonnedahl, J. (2016a). The colistin resistance *mcr-1* gene is going wild. *J. Antimicrob. Chemother.* 71, 2335–2336. doi: 10.1093/jac/dkw262
- Liakopoulos, A., Olsen, B., Geurts, Y., Artursson, K., Berg, C., Mevius, D. J., et al. (2016b). Molecular characterization of extended-spectrum-cephalosporin-resistant *Enterobacteriaceae* from wild kelp gulls in south America. *Antimicrob. Agents Chemother.* 60, 6924–6927. doi: 10.1128/aac.01120-16
- Manges, A. R., Geum, H. M., Guo, A., Edens, T. J., Fibke, C. D., and Pitout, J. D. D. (2019*). Global extraintestinal pathogenic *Escherichia coli* (ExPEC) lineages. *Clin. Microbiol. Rev.* 32:e00135-18.
- Martínez, J. L. (2009). The role of natural environments in the evolution of resistance traits in pathogenic bacteria. *Proc. Biol. Sci.* 276, 2521–2530. doi: 10.1098/rspb.2009.0320
- Matamoros, S., van Hattem, J. M., Arcilla, M. S., Willemsse, N., Melles, D. C., Penders, J., et al. (2017). Global phylogenetic analysis of *Escherichia coli* and plasmids carrying the *mcr-1* gene indicates bacterial diversity but plasmid restriction. *Sci. Rep.* 7:15364.
- Menz, J., Olsson, O., and Kummerer, K. (2019). Antibiotic residues in livestock manure: does the EU risk assessment sufficiently protect against microbial toxicity and selection of resistant bacteria in the environment? *J. Hazard Mater.* 5:120807. doi: 10.1016/j.jhazmat.2019.120807
- Ojer-Usoz, E., González, D., and Vitas, A. I. (2017). Clonal diversity of ESBL-producing *Escherichia coli* isolated from environmental, human and food samples. *Int. J. Environ. Res. Public Health* 14:E676.
- Peig, J., and Green, A. J. (2009). New perspectives for estimating body condition from mass/length data: the scaled mass index as an alternative method. *Oikos* 118, 1883–1891. doi: 10.1111/j.1600-0706.2009.17643.x
- Picard, B., García, J. S., Gouriou, S., Duriez, P., Brahimi, N., Bingen, E., et al. (1999). The link between phylogeny and virulence in *Escherichia coli* extraintestinal infection. *Infect. Immun.* 67, 546–553. doi: 10.1128/iai.67.2.546-553.1999
- Pinto, L., Radhouani, H., Coelho, C., Martins Da Costa, P., Simoes, R., Brandao, R. M., et al. (2010). Genetic detection of extended-spectrum beta-lactamase-containing *Escherichia coli* isolates from birds of prey from Serra da Estrela Natural Reserve in Portugal. *Appl. Environ. Microbiol.* 76, 4118–4120. doi: 10.1128/aem.02761-09

- Plaza, P. I., and Lambertucci, S. A. (2017). How are garbage dumps impacting vertebrate demography, health, and conservation? *Global Ecol. Conserv.* 12, 9–20.
- Rebelo, A. R., Bortolaia, V., Kjeldgaard, J. S., Pedersen, S. K., Leekitcharoenphon, P., Hansen, I. M., et al. (2018). Multiplex PCR for detection of plasmid-mediated colistin resistance determinants, *mcr-1*, *mcr-2*, *mcr-3*, *mcr-4* and *mcr-5* for surveillance purposes. *Eur. Surveill.* 23, 17–00672.
- Ribot, E. M., Fair, M. A., Gautom, R., Cameron, D. N., Hunter, S. B., Swaminathan, B., et al. (2006). Standardization of pulsed-field gel electrophoresis protocols for the subtyping of *Escherichia coli* O157:H7, *Salmonella*, and *Shigella* for PulseNet. *Foodborne Pathog. Dis.* 3, 59–67.
- Russo, T. A., and Johnson, J. R. (2000). Proposal for a new inclusive designation for extraintestinal pathogenic isolates of *Escherichia coli*: ExPEC. *J. Infect. Dis.* 181, 1753–1754.
- Santos, T., Silva, N., Igrejas, G., Rodrigues, P., Micael, J., Rodrigues, T., et al. (2013). Dissemination of antibiotic resistant *Enterococcus* spp. and *Escherichia coli* from wild birds of Azores Archipelago. *Anaerobe* 24, 25–31.
- Schmid, A., Hormansdorfer, S., Messelhauser, U., Kasbohrer, A., Sauter-Louis, C., and Mansfeld, R. (2013). Prevalence of extended-spectrum beta-lactamase-producing *Escherichia coli* on Bavarian dairy and beef cattle farms. *Appl. Environ. Microbiol.* 79, 3027–3032.
- Smet, A., Martel, A., Persoons, D., Dewulf, J., Heyndrickx, M., Herman, L., et al. (2010). Broad-spectrum beta-lactamases among *Enterobacteriaceae* of animal origin: molecular aspects, mobility and impact on public health. *FEMS Microbiol. Rev.* 34, 295–316.
- Solà-Ginés, M. (2018). *Molecular Epidemiology and Mechanisms of Cephalosporin Resistance in Escherichia coli of Different Origins: Broilers, Flies and White Storks*. Dissertation, Universidad Autónoma de Barcelona, Bellaterra.
- Solà-Ginés, M., Cameron-Veas, K., Badiola, I., Dolz, R., Majó, N., Dahbi, G., et al. (2015). Diversity of multi-drug resistant avian pathogenic *Escherichia coli* (APEC) causing outbreaks of colibacillosis in broilers during 2012 in Spain. *PLoS One* 10:e0143191. doi: 10.1371/journal.pone.0143191
- Szczepanska, B., Kaminski, P., Andrzejewska, M., Spica, D., Kartanas, E., Ulrich, W., et al. (2015). Prevalence, virulence, and antimicrobial resistance of *Campylobacter jejuni* and *Campylobacter coli* in white stork *Ciconia* in Poland. *Foodborne Pathog. Dis.* 12, 24–31.
- Tortosa, F. S., Caballero, J. M., and Reyes-López, J. (2002). Effect of rubbish dumps on breeding success in the *White Stork* in Southern Spain. *Waterbirds* 25, 39–43.
- Tortosa, F. S., and Villafuerte, R. (1999). Effect of nest microclimate on effective endothermy in White Stork *Ciconia* nestlings. *Bird Study* 46, 336–341.
- Tryjanowski, P., Sparks, T. H., and Jerzak, L. (2006). *The White Stork in Poland: Studies in Biology, Ecology and Conservation*. Poznan: Bogucki WN.
- Valverde, A., Turrientes, M. C., Norman, F., San Martín, E., Moreno, L., Pérez-Molina, J. A., et al. (2015). CTX-M-15-non-ST131 *Escherichia coli* isolates are mainly responsible of faecal carriage with ESBL-producing *Enterobacteriaceae* in travellers, immigrants and those visiting friends and relatives. *Clin. Microbiol. Infect.* 21, e251–e254.
- Veldman, K., Van Tulden, P., Kant, A., Testerink, J., and Mevius, D. (2013). Characteristics of cefotaxime-resistant *Escherichia coli* from wild birds in the Netherlands. *Appl. Environ. Microbiol.* 79, 7556–7561.
- Wang, J., Stephan, R., Karczmarczyk, M., Yan, Q., Hachler, H., and Fanning, S. (2013). Molecular characterization of bla ESBL-harboring conjugative plasmids identified in multi-drug resistant *Escherichia coli* isolated from food-producing animals and healthy humans. *Front. Microbiol.* 4:188. doi: 10.3389/fmicb.2013.00188
- Weston, A., Humphreys, G. O., Brown, M. G., and Saunders, J. R. (1979). Simultaneous transformation of *Escherichia coli* by pairs of compatible and incompatible plasmid DNA molecules. *Mol. Gen. Genet.* 172, 113–118.
- Wirth, T., Falush, D., Lan, R., Colles, F., Mensa, P., Wieler, L. H., et al. (2006). Sex and virulence in *Escherichia coli*: an evolutionary perspective. *Mol. Microbiol.* 60, 1136–1151.
- Zurfluh, K., Nuesch-Inderbinen, M., Stephan, R., and Hachler, H. (2013). Higher-generation cephalosporin-resistant *Escherichia coli* in feral birds in Switzerland. *Int J. Antimicrob. Agents* 41, 296–297.

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Höfle, Jose Gonzalez-Lopez, Camacho, Solà-Ginés, Moreno-Mingorance, Manuel Hernández, De La Puente, Pineda-Pampliega, Aguirre, Torres-Medina, Ramis, Majó, Blas and Migura-García. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.