

Forcing k -repetitions in degree sequences

Yair Caro

Department of Exact Sciences
University of Haifa at Oranim
Tivon 36006, Israel

yacaro@kvgeva.org.il

Asaf Shapira *

School of Mathematics
Tel-Aviv University
Tel-Aviv 69978, Israel

asafico@tau.ac.il

Raphael Yuster

Department of Mathematics
University of Haifa
Haifa 31905, Israel

raphy@math.haifa.ac.il

Submitted: Jun 24, 2013; Accepted: Jan 22, 2014; Published: Feb 7, 2014

Mathematics Subject Classifications: 05C07

Abstract

One of the most basic results in graph theory states that every graph with at least two vertices has two vertices with the same degree. Since there are graphs without 3 vertices of the same degree, it is natural to ask if for any fixed k , every graph G is “close” to a graph G' with k vertices of the same degree. Our main result in this paper is that this is indeed the case. Specifically, we show that for any positive integer k , there is a constant $C = C(k)$, so that given any graph G , one can remove from G at most C vertices and thus obtain a new graph G' that contains at least $\min\{k, |G| - C\}$ vertices of the same degree.

Our main tool is a multidimensional zero-sum theorem for integer sequences, which we prove using an old geometric approach of Alon and Berman.

Keywords: degree sequence; repetition

1 Introduction

All graphs considered in this paper are finite, simple, and undirected. Graph theory notation follows [3].

Perhaps one of the most obvious properties shared by all graphs (with at least two vertices) is that they have at least two vertices with the same degree. In general, the *repetition number* of a graph G , denoted by $\text{rep}(G)$, is the maximum multiplicity of a vertex degree in G . So, any graph with at least two vertices has $\text{rep}(G) \geq 2$, and there are simple constructions showing that equality holds for infinitely many graphs. Repetition numbers of graphs (and hypergraphs) have been studied by several researchers. See [2, 4, 5, 6, 7, 8, 9] for some representing works.

*Supported in part by ISF Grant 224/11 and a Marie-Curie CIG Grant 303320.

Since infinitely many graphs have $\text{rep}(G) = 2$, it seems interesting to determine how many vertices one must delete from a graph in order to increase its repetition number to 3 or higher. We state the problem more formally as follows. For a given positive integer k , let $C = C(k)$ denote the least integer such that any graph with n vertices has an induced subgraph with at least $n - C$ vertices whose repetition number is at least $\min\{k, n - C\}$. Stated otherwise, we can delete at most $C(k)$ vertices from any graph and obtain a subgraph with k repeated degrees (or a regular subgraph in case the subgraph has less than k vertices). Trivially $C(1) = C(2) = 0$. A simple lower bound of $C(k) \geq k - 2$ is easily obtained from the fact that there are infinitely many graphs with $\text{rep}(G) = 2$ and precisely one pair of vertices with the same degree. For general k , it is not obvious, however, that $C(k)$ is finite. The main result of this paper shows that it is.

Before we state our main result, let us consider a more general setting, where the graph in question is the complete graph with integer edge weights in $\{0, \dots, r\}$. The (weighted) degree of a vertex v , denoted by $\text{deg}(v)$, is the sum of the weights of the edges incident with it. Likewise, the repetition number of an edge-weighted graph is the maximum multiplicity of a vertex degree. Observe that, unlike the unweighted case, already for $r = 2$ we do not necessarily have two vertices with the same degree, as can be seen by assigning the weights 0, 1, 2 to the edges of a triangle. The related well-studied graph parameter of *irregularity strength* asks for the smallest r for which the edges of a graph can be weighted with $\{1, \dots, r\}$ (weight 0 represents non-edges) such that all vertex degrees are distinct (see [11, 13]).

For given positive integers k, r , let $C = C(k, r)$ denote the least integer such that any complete graph with n vertices and edge weights in $\{0, \dots, r\}$ has a complete subgraph with at least $n - C$ vertices whose repetition number is at least $\min\{k, n - C\}$. Notice that $C(k) = C(k, 1)$. Our main result is then the following.

Theorem 1. *For positive integers k, r there exists a constant $C = C(k, r)$ such that the following holds. Any complete graph with n vertices whose edges have weights in $\{0, \dots, r\}$ contains a complete subgraph with at least $n - C$ vertices whose repetition number is at least $\min\{k, n - C\}$.*

The bound we obtain for $C(k, r)$ via the proof of Theorem 1 is exponential in terms of k already for $r = 1$. In the case $r = 1$, it satisfies $C(k) = C(k, 1) \leq (8k)^k$. Together with a nontrivial lower bound of $C(k) \geq \Omega(k \log k)$ that we prove in Section 5 we obtain the following corollary.

Corollary 2. *We have*

$$\Omega(k \log k) \leq C(k) \leq (8k)^k .$$

It would be of course interesting to close the exponential gap between the upper bound and lower bound in the above corollary. Already for the first nontrivial case, $C(3)$, the proof of Theorem 1 gives the explicit bound $C(3) \leq 203$. An alternative ad hoc argument shows that $3 \leq C(3) \leq 6$ (see Section 4). Even the exact value of $C(3)$ is an open problem. When trying to adapt these ad hoc arguments to obtain an upper bound for $C(4)$ one quickly sees that the case analysis becomes significantly involved. Instead, our proof

of Theorem 1 is obtained by reducing the problem (after some additional combinatorial arguments are applied) to another problem in additive combinatorics, which may be interesting in its own right.

Let $[-r, r]^d$ denote the set of d -dimensional vectors over $\{-r, \dots, r\}$. The main tool used in the proof of Theorem 1 is the following theorem.

Theorem 3. *For positive integers r, d, q the following holds. Any sequence of $n \geq (\lceil q/r \rceil + 2)(2rd + 1)^d$ elements of $[-r, r]^d$ whose sum is in $[-q, q]^d$ contains a nonempty proper subsequence whose sum is zero.*

It is noteworthy to mention here a result of Erdős et al. [10] who proved that for every positive integer k , there is a number $N(k)$ such that for all $n > N(k)$, if the edges of the complete graph K_n are red-blue colored, then there is a monochromatic complete subgraph K_k (say blue) the degree of whose vertices in the blue-colored graph differ by at most $R(k) - 2$ where $R(k)$ is the diagonal Ramsey number. Our proof of Theorem 1 shows, in fact, that for $n > N(k)$ and any red-blue coloring, we can delete some $M(k)$ vertices such that there is a monochromatic K_k (say blue) with equal degrees in the blue-colored graph.

We finally mention that our result here shows that any graph G is “close” to another graph G' with k vertices of the same degree, where by close we mean that few vertices need to be removed. It is of course natural to ask what happens if we are allowed to remove (or also add) edges. As it turns out this variant of the problem is not hard. Indeed, it is easy to see that in this case it is enough to remove $O(k^2)$ edges in order to force k vertices of the same degree. It is also easy to see that $\Omega(k^2)$ edge removals are required. We omit the details.

The rest of this paper is organized as follows. Section 2 contains the proof of Theorem 3. The proof of Theorem 1 is given in Section 3. Section 4 considers the special case $C(3)$. A non-linear lower bound for $C(k)$ is established in Section 5.

2 Proof of Theorem 3

Our proof of Theorem 3 uses a geometric approach similar to the one used by Alon and Berman in [1]. To this end, we need the following result of Sevast'yanov [14].

Lemma 4 (Sevast'yanov [14]). *Let V be any normed d -dimensional space. Suppose $v_1, \dots, v_n \in V$ where $\|v_i\| \leq 1$ and $\sum_{i=1}^n v_i = 0$. Then there is a permutation π on $\{1, \dots, n\}$ such that for all $j = 1, \dots, n$,*

$$\left\| \sum_{i=1}^j v_{\pi(i)} \right\| \leq d.$$

Proof (Theorem 3): Consider a sequence $X = [x_1, \dots, x_n]$ of $n \geq (\lceil q/r \rceil + 2)(2rd + 1)^d$ elements of $[-r, r]^d$ whose sum is $w \in [-q, q]^d$. It is easy to see that since $w \in [-r, r]^d$, we can

always find $p = \lceil q/r \rceil$ vectors $x_{n+1}, \dots, x_{n+p} \in [-r, r]^d$ whose sum is $-w$. Adding these “artificial” vectors to X we obtain a zero-sum collection $X' = [x_1, \dots, x_n, x_{n+1}, \dots, x_{n+p}]$. For every $1 \leq i \leq n + p$ set $v_i = x_i/r$, and consider $X'' = [v_1, \dots, v_{n+p}]$. Then X'' is a zero-sum sequence with $\|v_i\|_\infty \leq 1$ for every $v_i \in X''$. By Lemma 4 (with the ℓ_∞ norm), there is a permutation π on $\{1, \dots, n + p\}$ such that for all $j = 1, \dots, n + p$, we have $\left\| \sum_{i=1}^j v_{\pi(i)} \right\|_\infty \leq d$. Observe now that each coordinate of any v_i is a rational of the form t/r where $t \in \{-r, \dots, r\}$. Hence, any possible sum of a subset of vectors of X'' whose ℓ_∞ norm is at most d is a vector with rational coordinates where each coordinate is of the form t/r where $t \in \{-rd, \dots, rd\}$. Hence, there are at most $(2rd + 1)^d$ such possible sums. As any prefix sum of the elements of X'' ordered by the permutation π has ℓ_∞ norm at most d , we have, by the pigeonhole principle, that some prefix sum value repeats at least $(n + p)/(2rd + 1)^d$ times. Since $n \geq (p + 2)(2rd + 1)^d$, some prefix sum value, call it z , repeats at least $p + 2$ times.

Consider therefore $p + 2$ locations j_1, \dots, j_{p+2} for which $\sum_{i=1}^{j_\ell} v_{\pi(i)} = z$ for $\ell = 1, \dots, p + 2$. This means that for $\ell = 1, \dots, p + 1$, we have

$$\sum_{i=j_\ell+1}^{j_{\ell+1}} v_{\pi(i)} = 0.$$

As we added only p artificial vectors to X' (and thus also to X''), one of these $p + 1$ collections must give us the collection we want. Specifically, there exists some ℓ for which $\pi(j_\ell + 1), \dots, \pi(j_{\ell+1}) \in \{1, \dots, n\}$, implying that we have a non-empty and proper collection of vectors from X (namely $x_{\pi(j_\ell+1)}, \dots, x_{\pi(j_{\ell+1})}$) whose sum is zero. \square

The following is an immediate corollary of Theorem 3.

Corollary 5. *For positive integers r, d, q the following holds. Any sequence of $n \geq (\lceil q/r \rceil + 2)(2rd + 1)^d$ elements of $[-r, r]^d$ whose sum, denoted by z , is in $[-q, q]^d$ contains a subsequence of length at most $(\lceil q/r \rceil + 2)(2rd + 1)^d$ whose sum is z .*

3 Proof of Theorem 1

We start with the following simple lemma.

Lemma 6. *Suppose that G is a complete graph with n vertices whose edges have weights in $\{0, \dots, r\}$. If $n \geq s^2$ where s is a positive integer, then there is a set S of s vertices with the property that $|\deg(x) - \deg(y)| \leq sr$ for any $x, y \in S$.*

Proof. Clearly, the difference between the minimum degree of G and the maximum degree of G is at most $(n - 2)r$. We need to prove that for any positive integer s , if $n \geq s^2$, then there is a set S of s vertices with the property that $|\deg(x) - \deg(y)| \leq sr$ for any $x, y \in s$. To see this, assume that the vertices are $\{v_1, \dots, v_n\}$ where $\deg(v_i) \leq \deg(v_{i+1})$. Let $S_j = \{v_{(s-1)j+1}, \dots, v_{(s-1)(j+1)+1}\}$ for $j = 0, \dots, \lfloor (n - 1)/(s - 1) \rfloor - 1$. Observe that

the last vertex of S_j is also the first vertex of S_{j+1} . Now, if some S_j has the desired property of S , we are done. Otherwise the difference between $\deg(v_1)$ and the last vertex of $S_{\lfloor (n-1)/(s-1) \rfloor - 1}$ is at least

$$(sr + 1) \cdot \lfloor \frac{n-1}{s-1} \rfloor \geq (sr + 1) \frac{n-s+1}{s-1} > (n-2)r$$

where the last inequality follows from the assumption $n \geq s^2$. As this contradicts the maximum gap between the minimum and maximum degree, the lemma follows. \square

We note that for the case $r = 1$, a slightly stronger result follows from a result of Erdős et al [10] using the Erdős-Gallai criterion for degree sequences. It is proved that for any $n \geq s \geq 2$ there is a set S of s vertices with the property that $|\deg(x) - \deg(y)| \leq s - 2$ for any $x, y \in S$.

Recall that $R_r(k)$ is the *multicolored Ramsey number*, which is the least integer s such that in any coloring of the edges of the complete graph with s vertices using r colors, there is a monochromatic K_k (see [12]).

Proof (Theorem 1): Let $s = R_{r+1}(k)$ be the multicolored Ramsey number, and set

$$C = C(k, r) = \max\{s^2, k + (s + 2)(2r(k - 1) + 1)^{k-1}\}.$$

Consider a complete graph with n vertices whose edges have weights in $\{0, \dots, r\}$. We may assume that $n \geq C$ as otherwise we can just delete, say, $n - 2 \leq C$ vertices and obtain a regular subgraph with two vertices. Since $n \geq C \geq s^2$, we have by Lemma 6 that there is a set S of s vertices with the property that $|\deg(x) - \deg(y)| \leq sr$ for any $x, y \in S$. Thus, for some $p \in \{0, \dots, r\}$ the induced complete subgraph $G[S]$ has an induced complete subgraph on k vertices, denoted by K , such that the edge (x, y) for $x, y \in K$ has weight p .

Without loss of generality, $V(G) \setminus K = \{v_1, \dots, v_{n-k}\}$ and $K = \{v_{n-k+1}, \dots, v_n\}$. Let $w(v_i, v_j)$ denote the weight of the edge connecting v_i and v_j . We construct a sequence of $n-k$ vectors x_1, \dots, x_{n-k} of $[-r, r]^{k-1}$ as follows. Coordinate j of x_i is $w(v_{n-k+j}, v_i) - w(v_n, v_i)$ for $i = 1, \dots, n-k$ and $j = 1, \dots, k-1$. Observe that indeed $w(v_{n-k+j}, v_i) - w(v_n, v_i) \in \{-r, \dots, r\}$ as required. What can be said about the sum of all the j 'th coordinates?

$$\begin{aligned} \sum_{i=1}^{n-k} (w(v_{n-k+j}, v_i) - w(v_n, v_i)) &= \sum_{i=1}^{n-k} w(v_{n-k+j}, v_i) - \sum_{i=1}^{n-k} w(v_n, v_i) \\ &= (\deg(v_{n-k+j}) - p(k-1)) - (\deg(v_n) - p(k-1)) \\ &= \deg(v_{n-k+j}) - \deg(v_n) \leq sr. \end{aligned}$$

Hence,

$$z = \sum_{i=1}^{n-k} x_i \in [-sr, sr]^{k-1}.$$

Since $n - k \geq C - k \geq (s + 2)(2r(k - 1) + 1)^{k-1}$, we have by Corollary 5 with $d = k - 1$ and $q = sr$ that there is a subsequence of X of size at most $(s + 2)(2r(k - 1) + 1)^{k-1}$ whose

sum is z . Deleting the vertices of G corresponding to the elements of this subsequence results in a subgraph with at least $n - (s + 2)(2r(k - 1) + 1)^{k-1} \geq n - C$ vertices in which all the k vertices of K have the same degree. \square

The following corollary is immediate from the bound $C(k, 1) \leq \max\{s^2, k + (s + 2)(2(k - 1) + 1)^{k-1}\}$ given in the proof of Theorem 1, together with the well known facts that $s = R_2(k) < 4^k$ and $R_2(3) = 6$.

Corollary 7. *We have $C(k) \leq (8k)^k$ and $C(3) \leq 203$.*

4 Equating three vertices

In this section we prove that $3 \leq C(3) \leq 6$.

Proposition 8. *For any graph G with at least 5 vertices, one can delete at most 6 vertices such that the subgraph obtained has at least three vertices with the same degree. Consequently, $C(3) \leq 6$.*

Proof. By a result of [10], G has a set X of 5 vertices such that $|\deg(u) - \deg(v)| \leq 3$ for $u, v \in X$. Now, either X has a triangle, or else X has an independent set of size 3, or else X induces a C_5 . In any case, this implies that X has three vertices x, y, z with $\deg(x) \leq \deg(y) \leq \deg(z)$ such that the following holds: (x, y) is an edge if and only if (x, z) is an edge. Furthermore, if (y, z) is an edge, then $\{x, y, z\}$ induce a triangle. We use this property implicitly throughout the remainder of the proof. We may also assume $\deg(z) > \deg(x)$ otherwise x, y, z already have the same degree. Throughout the proof we denote by $N(\cdot)$ the set of neighbors of a vertex in the current G (that is, in the graph G after some vertices have possibly been deleted). Similarly, we denote by $\deg(\cdot)$ the degree of a vertex in the current G .

Consider first the case $\deg(x) < \deg(y) = \deg(z)$ in the original G . If $(N(z) \setminus N(x)) \cap (N(y) \setminus N(x)) \neq \emptyset$ we can delete a vertex of this intersection and decrease the degrees of y and z by 1 without affecting the degree of x . Otherwise, if $(N(z) \setminus N(x)) \cap (N(y) \setminus N(x)) = \emptyset$ we can delete a vertex of $N(z) \setminus N(x)$ and a vertex of $N(y) \setminus N(x)$ and decrease the degrees of y and z by 1 without affecting the degree of x . Observe that in any case we delete at most two vertices. Repeating this process at most three times we eventually obtain $\deg(x) = \deg(y) = \deg(z)$. Overall, we have deleted at most 6 vertices.

Consider next the case $\deg(x) \leq \deg(y) < \deg(z)$ in the original G . Let $p = \deg(z) - \deg(y)$ and let $q = \deg(y) - \deg(x)$, and observe that $p + q \leq 3$. Let us first equate $\deg(z)$ and $\deg(y)$ by deleting some $u \in N(z) \setminus N(y)$. Observe that $u \neq x$. We always prefer to delete a vertex u that is non-adjacent to x , as long as there is such a vertex u . Overall, we have deleted p vertices. The problem is that in the current graph we may have that $\deg(x)$ also decreased by some amount $r \leq p$. Suppose first that $r = 0$. As in the previous case, we may need to delete $2q$ additional vertices to equate the degrees of y and z to that of x . Overall, we have deleted at most $p + 2q \leq 6$ vertices. If $r > 0$ then this means that at some point, when we deleted a vertex u , that vertex also had to be adjacent to x .

Hence, in the current graph, $(N(z) \setminus N(x)) \subset (N(y) \setminus N(x))$. So, we may simply delete r additional vertices (all of them from $N(z) \setminus N(x)$) to equate the degrees of y and z to that of x . Overall, we deleted $p + r \leq 2p \leq 6$ vertices. \square

The graph in Figure 1 proves that $C(3) \geq 3$. If we delete vertices 1, 2, 8, then in the resulting graph, vertices 4, 5, 7 have the same degree. It is easy to check that deleting any two vertices does not yield a graph with repetition number 3. In fact, a computer verification asserts that this graph is the smallest graph (in terms of the number of vertices) for which we need to delete more than two vertices to obtain repetition number 3.




Figure 1: A graph showing that $C(3) \geq 3$.

5 A lower bound for $C(k)$

In this section we prove that $C(k) = \Omega(k \log k)$. Our construction is based upon an additional building block: graphs that have the property that all of their induced subgraphs have repetition number which is not too large.

Lemma 9. *For any integer n , there exists a graph D_n with n vertices such that no induced subgraph of D_n has more than $3n/\ln n$ vertices with the same degree.*

Proof. Consider the non-increasing sequence of integers $a_i = \lfloor 2n/(i \ln n) \rfloor$. Let $S_i = \sum_{j=1}^i a_j$ and let s be the largest integer such that $S_s \leq n$. Observe that $s \leq \sqrt{n}$ since

$$S_{\lceil \sqrt{n} \rceil} = \sum_{i=1}^{\lceil \sqrt{n} \rceil} a_i \geq \frac{2n}{\ln n} \left(\sum_{i=1}^{\lceil \sqrt{n} \rceil} \frac{1}{i} \right) > n.$$

Note that we also have $n - S_s \leq a_1 = \lceil 2n/(\ln n) \rceil$.

Construct a graph D_n with n vertices as follows. Take s vertex-disjoint cliques B_1, \dots, B_s where $B_i = K_{a_i}$ is a clique on a_i vertices. The remaining set of $n - S_s$ vertices of D_n is a set denoted by B_{s+1} consisting of $n - S_s$ isolated vertices.

Let H be any induced subgraph of D_n , and suppose that K is a set of k vertices of H with the same degree in H . Let j be the largest index such that K contains a vertex from B_j . Observe that if $j = s + 1$, then all vertices of K have degree 0 in H and thus contain at most one vertex from each B_i for $i \geq 1$. In this case we have

$$k \leq |B_0| + s = n - S_s + s \leq \lceil 2n/(\ln n) \rceil + \sqrt{n} \leq \frac{3n}{\ln n}.$$

If $j \leq s$, then all vertices of K have the same degree in H which is at most $a_j - 1$ and thus K contains at most a_j vertices from each B_i for $i = 1, \dots, j$. Hence

$$k \leq ja_j = j \lceil 2n/(j \ln n) \rceil \leq \frac{2n}{\ln n} + \sqrt{n} \leq \frac{3n}{\ln n}.$$

□

Theorem 10. $C(k) = \Omega(k \log k)$.

Proof. We will prove the slightly stronger assertion that for k sufficiently large, there are infinitely many graphs such that one cannot equate the degrees of k vertices by removing fewer than $(k \ln k)/10$ vertices.

Let H_n be any graph with n vertices and with $\text{rep}(H_n) = 2$. Denote its vertices by $\{h_1, \dots, h_n\}$ where $\text{deg}(h_i) \leq \text{deg}(h_{i+1})$. Let $H_n(q)$ be the graph obtained from H_n by the following procedure. First, take a q -blowup of H_n , namely, each h_i is replaced by an independent set of q vertices, denoted by $B_i = \{h_{i,1}, \dots, h_{i,q}\}$. We connect each vertex of B_i to each vertex of B_j if and only if $(h_i, h_j) \in E(H_n)$. Otherwise, $B_i \cup B_j$ is an independent set. Next, for each $i = 1, \dots, n$ we place the graph D_q of Lemma 9 inside B_i . Observe that $H_n(q)$ has nq vertices, and, furthermore, since $\text{rep}(H_n) = 2$, we have that for $i > j$, $\text{deg}(h_i) - \text{deg}(h_j) \geq \lfloor (i - j)/2 \rfloor$. Since $\text{deg}(h_i)q \leq \text{deg}(h_{i,x}) < (\text{deg}(h_i) + 1)q$ for all $x = 1, \dots, q$, we have that for $i > j$,

$$\text{deg}(h_{i,x}) - \text{deg}(h_{j,y}) \geq \frac{q(i - j)}{2} - \frac{3q}{2}. \tag{1}$$

We claim that for, say, $q = 5k$, any induced graph obtained from $H_n(q)$ by deleting fewer than $(k \ln k)/10$ vertices, does not have k vertices with the same degree. Indeed, assume otherwise and suppose we can delete a set of vertices X of size at most $(k \ln k)/10$ from $H_n(q)$ such that the resulting subgraph H' has a set K of k vertices with the same degree. Let $B'_i = B_i \setminus X$ for $i = 1, \dots, n$ and notice that if two vertices of K belong to B_i , then they have the same degree in the subgraph induced by B'_i . Notice also that in $H_n(q)$, the degrees of any two vertices of K differ by at most $(k \ln k)/10$. Hence, if $v \in B_i$

and $u \in B_j$ are two vertices of K , then $|i - j| \leq (\ln k)/20 - 1$, as otherwise, by (1), their degrees in $H_n(q)$ would differ by at least

$$\frac{q(\ln k/20 - 1)}{2} - \frac{3q}{2} > \frac{k \ln k}{10}$$

where in the last inequality we used $q = 5k$ and that k is sufficiently large. It follows that some B'_i contains at least $20k/\ln k$ vertices of K . However, as the subgraph induced by B'_i was obtained from D_q by deleting vertices, we have by Lemma 9 that it cannot have more than $3q/\ln q$ vertices of the same degree. But we now arrive at a contradiction since $3q/\ln q < 20k/\ln k$. \square

References

- [1] N. Alon and K.A. Berman. Regular hypergraphs, Gordon's Lemma, Steinitz' Lemma and invariant theory. *Journal of Combinatorial Theory, Series A*, 43(1):91–97, 1986.
- [2] P. Balister, B. Bollobás, J. Lehel, and M. Morayne. Repeated degrees in random uniform hypergraphs. *SIAM Journal on Discrete Mathematics*, 27(1):145–154, 2013.
- [3] B. Bollobás. *Extremal Graph Theory*. Academic Press, 1978.
- [4] B. Bollobás. Degree multiplicities and independent sets in K_4 -free graphs. *Discrete Mathematics*, 158(1):27–35, 1996.
- [5] B. Bollobás and A. Scott. Independent sets and repeated degrees. *Discrete Mathematics*, 170(1):41–49, 1997.
- [6] Y. Caro and D. West. Repetition number of graphs. *Electronic Journal of Combinatorics*, 16(1):#R7, 2009.
- [7] Y. Caro and R. Yuster. Large induced subgraphs with equated maximum degree. *Discrete Mathematics*, 310(4):742–747, 2010.
- [8] G. Chen, J. Hutchinson, W. Piotrowski, W. Shreve, and B. Wei. Degree sequences with repeated values. *Ars Combinatoria*, 59:33–44, 2001.
- [9] G. Chen and R. Schelp. Ramsey problems with bounded degree spread. *Combinatorics, Probability, and Computing*, 2:263–269, 1993.
- [10] P. Erdős, G. Chen, C. Rousseau, and R. Schelp. Ramsey problems involving degrees in edge-colored complete graphs of vertices belonging to monochromatic subgraphs. *European Journal of Combinatorics*, 14(3):183–189, 1993.
- [11] M. Kalkowski, M. Karonski, and F. Pfender. A new upper bound for the irregularity strength of graphs. *SIAM Journal on Discrete Mathematics*, 25(3):1319–1321, 2011.
- [12] S. Radziszowski. Small Ramsey numbers. *Electronic Journal of Combinatorics*, DS1:94 pages, dynamic survey, 2014.
- [13] B. Seamone. The 1-2-3 conjecture and related problems: a survey. [arXiv:1211.5122](https://arxiv.org/abs/1211.5122).
- [14] S.V. Sevast'yanov. On approximate solutions of scheduling problems. *Metody Diskretnogo Analiza*, 32:66–75, 1978 (in Russian).