

FOXL2: a central transcription factor of the ovary

Adrien Georges^{1,2}, Aurelie Auguste^{1,2}, Laurianne Bessi re^{1,2}, Anne Vanet^{1,2}, Anne-Laure Todeschini^{1,2} and Reiner A Veitia^{1,2}

¹CNRS UMR 7592, Institut Jacques Monod, 15 Rue H el ne Brion, 75013 Paris, France

²Universit  Paris Diderot, Paris VII, Paris, France

Correspondence should be addressed to R A Veitia
Email
veitia.reiner@ijm.univ-paris-diderot.fr

Abstract

Forkhead box L2 (FOXL2) is a gene encoding a forkhead transcription factor preferentially expressed in the ovary, the eyelids and the pituitary gland. Its germline mutations are responsible for the blepharophimosis ptosis epicanthus inversus syndrome, which includes eyelid and mild craniofacial defects associated with primary ovarian insufficiency. Recent studies have shown the involvement of FOXL2 in virtually all stages of ovarian development and function, as well as in granulosa cell (GC)-related pathologies. A central role of FOXL2 is the lifetime maintenance of GC identity through the repression of testis-specific genes. Recently, a highly recurrent somatic FOXL2 mutation leading to the p.C134W substitution has been linked to the development of GC tumours in the adult, which account for up to 5% of ovarian malignancies. In this review, we summarise data on FOXL2 modulators, targets, partners and post-translational modifications. Despite the progresses made thus far, a better understanding of the impact of FOXL2 mutations and of the molecular aspects of its function is required to rationalise its implication in various pathophysiological processes.

Key Words

- ▶ cancer
- ▶ oestrogen receptors
- ▶ ovary
- ▶ ovarian function
- ▶ FOXL2

Journal of Molecular Endocrinology
(2014) 52, R17–R33

Introduction

The mammalian ovary is a complex organ, ensuring various functions essential to the reproductive process. Its most obvious role is the production and release of functional female gametes, the oocytes. In addition, the ovary also has a critical endocrine function, including notably the production of oestrogens, which are essential to the development of female genitalia, and progesterone, which is necessary for the establishment of pregnancy. These functions are tightly coordinated throughout folliculogenesis, a dynamic process involving a continuous differentiation of three cell types: the oocyte and the surrounding granulosa and theca cells. A simplified summary of this process is outlined in Fig. 1.

The coordinated differentiation of the oocyte and the surrounding somatic cells is under the control of a large

number of signaling factors including gonadotropins (follicle-stimulating hormone (FSH) and luteinizing hormone (LH)), members of the transforming growth factor β (TGF β) family (GDF-9, BMP15, BMP4, BMP7, AMH, FST, etc.), growth factors (IGF1, FGF2, KITL, etc.) and sex hormones (estradiol, testosterone, progesterone, etc.), which mediate cellular and tissue-level communication (Edson *et al.* 2010). Granulosa cells (GCs) play a central role in the coordination of folliculogenesis through their direct communication with the oocyte and theca cells, their sensitivity to pituitary hormones and their ability to produce oestrogens, anti-M llerian hormone (AMH), activins etc. Although many signals targeting GCs are known, the way these cells integrate this information at the transcriptional level is still poorly understood. For example,




Figure 1

Outline of the main steps of folliculogenesis. Folliculogenesis starts with primordial follicles that are subsequently recruited to become primary follicles. Proliferating GCs progressively form several layers around the oocyte (secondary to late preantral follicles). From this stage, a layer of theca cells surrounds the follicle, and the follicles start to produce oestrogens. Theca cells produce androgens, which are converted into oestrogens in granulosa cells. Whereas follicular growth to the secondary/preantral stage is independent of gonadotrophins, progression beyond this stage strictly depends on FSH stimulation. As GC continue to

proliferate, the antrum is formed and separates *de facto* granulosa cells into two general populations: cumulus and mural GCs. At this stage, selection occurs between growing follicles, and only one or a limited number of follicles continue growing to the preovulatory stage while others undergo atresia. Two layers of theca (theca interna and externa) can be distinguished around selected follicles. After ovulation, which is triggered by a peak of FSH and LH, theca cells and mural granulosa cells luteinise to produce progesterone.

the reasons why GCs become dependent on FSH for their proliferation at the antral stage or the precise mechanisms by which the FSH/LH surge triggers ovulation remain unclear. Similarly, the precise signal(s) triggering follicle recruitment for maturation are yet to be determined.

The dysregulation of GC differentiation is involved in major ovarian pathologies, such as primary ovarian insufficiency (POI), polycystic ovary syndrome (PCOS) or GC tumours (GCTs). POI is defined by the absence of menses during 4 consecutive months, low oestrogen levels and an abnormally high level of FSH occurring before the age of 40 years. Although these symptoms are close to those of menopause, it happens occasionally that ovarian

function can reappear and allow pregnancy (De Vos *et al.* 2011). Understanding the molecular mechanisms underlying GC differentiation may therefore provide insights into the pathogenesis and treatment of these diseases. Conversely, the identification of genetic determinants for these conditions may be highly informative in the study of ovarian function.

Although POI is a rather multifactorial condition, heterogeneous genetic causes have been identified in familial and sporadic cases. Some of the altered genes are known for their role in folliculogenesis such as *FSHB*, *FSHR*, *FOXO3A*, *ESR1*, *LHB*, *NR5A1*, *NOBOX* and others (Qin *et al.* 2007, De Vos *et al.* 2011, Lakhali *et al.* 2012).

An example of a gene associated with familial POI is the one encoding the transcription factor (TF) forkhead box L2 (FOXL2), whose mutations are responsible for the blepharophimosis ptosis epicanthus inversus syndrome (BPES), a dominant condition characterised by palpebral malformations associated with POI (type I) or not (type II). Mutations affecting this gene can be detected in about 90% of BPES patients and in 70% of them, the mutation is intragenic (Crisponi *et al.* 2001, Verdin & De Baere 2012). FOXL2 is a single-exon gene encoding a protein of 376 amino acids. The protein contains a forkhead DNA-binding domain and a polyalanine tract of 14 residues, which is expanded to 24 residues in 30% of BPES cases with intragenic mutations (Cocquet *et al.* 2003, Verdin & De Baere 2012). FOXL2 is highly conserved among vertebrates, undergoes strong purifying selection (Cocquet *et al.* 2003) and has emerged as a key factor of ovarian biology.

This review focuses on the transcriptional regulation of GC function, with a particular attention on the role of FOXL2 and its known transcriptional targets (summarised in Table 1). Recent studies have indicated its involvement in virtually all stages of ovarian development and function, as well as its central role in GC-related ovarian pathologies.

Sex determination and early ovarian development

Gonadal sex determination is genetically determined in mammals and involves the mobilisation of sex-specific transcriptional networks in the common precursors of Sertoli cells and GCs. In XY individuals, the first specific event of masculinisation is the expression of the TF SRY, encoded on the Y chromosome. This factor is expressed for a short period of time in the mouse (peaking at 11.5 days post-coitum (d.p.c.)) and is under the control of the TFs NR5A1 (also known as steroidogenic factor 1 (Sf-1)), Wilms tumour 1 (WT1) and the complex Gata-binding protein 4, friend of Gata 2 (GATA4–FOG2) (de Santa Barbara *et al.* 2001, Miyamoto *et al.* 2008). SRY, together with NR5A1, contributes to the activation of a gene encoding another TF from the same family, SRY-box 9 (SOX9), which regulates, in turn, genes essential for testis and genital development such as *Egf9*, *Dhh* or *Amh* (Yao *et al.* 2002, Chaboissier *et al.* 2004, Hiramatsu *et al.* 2010). In mice, *Sox9* is both necessary and sufficient for testis determination, whereas SRY would ‘simply’ upregulate *Sox9* at the right time during development (Vidal *et al.* 2001, Barrionuevo *et al.* 2006).

In the absence of strong SOX9 expression, which is the case in XX individuals, transcriptional networks responsible for female development are activated. Several genes involved in ovarian determination/differentiation encode factors involved in β -catenin signaling. In particular, WNT4 and RSPO1, two secreted positive effectors of β -catenin signaling and of β -catenin expression, are required for the development of functional ovaries (Vainio *et al.* 1999, Chassot *et al.* 2008, Liu *et al.* 2009). Strikingly, β -catenin stabilisation is sufficient to induce femaleness in XY mice (Maatouk *et al.* 2008). This pathway seems to activate the follistatin gene (*Fst*), which encodes a secreted inhibitor of the TGF β pathway, also necessary for ovarian development (Yao *et al.* 2004). Expression of ovarian genes, such as *Wnt4* and *Foxl2*, is decreased in mice where β -catenin has been inactivated in *Nr5a1*-positive cells (including foetal GC precursors), although this may be due to a reduction of the pool of precursors. In addition to secreted inducers, activation of the β -catenin pathway also requires the presence of the GATA4–FOG2 complex. Indeed, in its absence, the inhibitor of the β -catenin signaling pathway DKK1 is highly expressed. *Dkk1* deletion allows reactivation of the pathway, but is not sufficient to promote ovarian expression of *Wnt4* and *Foxl2*. This suggests the existence of a regulatory loop involving GATA4–FOG2 and the canonical pathway to promote an ovarian ‘programme’ (Manuylov *et al.* 2008). Finally, it is important to note that the β -catenin pathway and SRY/SOX9 repress each other, making them mutually exclusive (Bernard *et al.* 2008, Edson *et al.* 2009).

The role of FOXL2 during early ovarian development is less well understood. In the mouse, *Foxl2* is the earliest known ovarian marker, its expression being detectable from 12.0 d.p.c. (Auguste *et al.* 2011). Nevertheless, *Foxl2*^{-/-} mice develop normally until birth, and no perinatal sex reversal is observed (Schmidt *et al.* 2004, Uda *et al.* 2004). It has been suggested that (one of) its function(s) could be to ‘cooperate’ with BMP2 and WNT4 to activate *Fst* expression (Kashimada *et al.* 2011). Although FOXL2 is apparently not required during early ovarian development in mice, several studies suggest that it could play an important role in ovarian determination and development in other mammals. Indeed, a regulatory mutation in goats, responsible for the polled intersex syndrome (PIS^{-/-}), which affects the expression of two transcripts, FOXL2 and polled intersex syndrome regulated transcript 1 (*PISRT1*), during ovarian development was identified in XX sex-reversed animals (Pailhoux *et al.* 2005). Further work showed that *PISRT1* dysregulation could not explain the observed phenotype (Boulanger *et al.* 2008), suggesting that FOXL2

Table 1 Overview of potential direct and indirect FOXL2 transcriptional targets

Gene names	Overall functions	Effects of FOXL2	Context	Methods	References
<i>FST</i>	Ovarian development	Activation	Mouse pituitary cells	Overexpression	Blount <i>et al.</i> (2009) Garcia-Ortiz <i>et al.</i> (2009) and Kashimada <i>et al.</i> (2011)
	Pituitary function	Activation	Mouse foetal ovaries	Knockout	
<i>SOX9</i>	Sex determination	Activation	Mouse adult ovaries	Knockout	Uhlenhaut <i>et al.</i> (2009) Kashimada <i>et al.</i> (2011) Nonis <i>et al.</i> (2013)
		Activation	Mouse granulosa cells (COV434)	Overexpression	
<i>CYP19A1</i>	Oestrogen production	Repression	Human cells (KGN)	Knockout	Uhlenhaut <i>et al.</i> (2009) Rosario <i>et al.</i> (2012) Pannetier <i>et al.</i> (2006)
		Repression	Human cells (KGN)	Knockdown	
<i>KITL</i>	Follicle activation	Activation	Goat promoter, human cells (KGN)	Overexpression	Pannetier <i>et al.</i> (2006) Uhlenhaut <i>et al.</i> (2009) Rosario <i>et al.</i> (2012) Fleming <i>et al.</i> (2010) Bentsi-Barnes <i>et al.</i> (2010) Dai <i>et al.</i> (2013)
		Activation	Mouse adult ovaries	Knockout	
		Repression	Human cells (KGN)	Knockdown	
		Activation	Human cells (KGN)	Overexpression	
		Repression	Hamster cells (CHO)	Overexpression	
<i>CDKN1B</i>	Follicle activation	Repression	Mouse primary granulosa cells	miRNA knockdown	Dai <i>et al.</i> (2013) Garcia-Ortiz <i>et al.</i> (2009)
		Activation	Human cells (KGN)	Overexpression	
<i>SMAD3</i>	Signal transduction	None	Mouse foetal/new-born ovaries	Knockout	Schmidt <i>et al.</i> (2004) Garcia-Ortiz <i>et al.</i> (2009)
		Activation	Mouse newborn/adult ovaries	Knockout	
<i>BMPR1A</i>	Signal transduction	Activation	Mouse foetal/new-born ovaries	Knockout	Garcia-Ortiz <i>et al.</i> (2009)
		Activation	Mouse foetal/new-born ovaries	Knockout	
<i>DKK3</i>	Signal transduction	Activation	Mouse foetal/new-born ovaries	Knockout	Garcia-Ortiz <i>et al.</i> (2009)
		Activation	Mouse foetal/new-born ovaries	Knockout	
<i>STAR</i>	Oestrogen production	Repression	Human promoter/Hamster cells	Overexpression	Pisarska <i>et al.</i> (2004) Rosario <i>et al.</i> (2012)
		Repression	Human cells (KGN)	Knockdown	
<i>CYP17A1</i>	Androgen production	Repression	Human cells (KGN)	Knockdown	Park <i>et al.</i> (2010) Park <i>et al.</i> (2010)
		Repression	Rat granulosa cells	Overexpression	
<i>PTGS2</i>	Ovulation	Activation	Human cells (KGN)	Overexpression	Batista <i>et al.</i> (2007) Kim <i>et al.</i> (2009)
		Repression	Human cells (293FT and MDA-MB-231 cells)	Overexpression	
<i>SERPINE2</i>	Ovulation	Activation	Mouse foetal/adult ovaries	Knockout	Caburet <i>et al.</i> (2012)
<i>HAS2</i>	Ovulation	Repression	Mouse foetal/new-born ovaries	Knockout	Caburet <i>et al.</i> (2012)
<i>PTGER2</i>	Ovulation	Activation	Mouse foetal/new-born ovaries	Knockout	Caburet <i>et al.</i> (2012)
<i>EDNRA</i>	Ovulation	Activation	Mouse adult ovaries	Knockout	Caburet <i>et al.</i> (2012)
<i>FSHB</i>	Follicle growth	Activation	Mouse pituitary	Knockout	Justice <i>et al.</i> (2011) and Tran <i>et al.</i> (2013) Tran <i>et al.</i> (2011) Lamba <i>et al.</i> (2009) and Tran <i>et al.</i> (2011) Corpuz <i>et al.</i> (2010)
		Activation	Porcine promoter, mouse pituitary cells (LbT2)	Overexpression	
		Activation	Mouse pituitary cells	Overexpression	
		Activation	Human promoter, mouse pituitary cells (LbT2)	Overexpression	
<i>TNF-R1</i>	Apoptosis	Activation	Human cells (KGN)	Overexpression	Kim <i>et al.</i> (2011)
<i>FAS</i>	Apoptosis	Activation	Human cells (KGN)	Overexpression	Kim <i>et al.</i> (2011)
<i>TRAIL-R</i>	Apoptosis	Activation	Human cells (KGN)	Overexpression	Kim <i>et al.</i> (2011)

Table 1 Continued

Gene names	Overall functions	Effects of FOXL2	Context	Methods	References
<i>GNRHR</i>	Apoptosis/signaling	Activation Activation Activation	Human cells (KGN) Human luteal cells Mouse granulosa cells (KK1)	Overexpression Knockdown Overexpression	Cheng <i>et al.</i> (2013) Cheng <i>et al.</i> (2013) Escudero <i>et al.</i> (2010)
<i>BCL2A1</i>	Apoptosis	Activation	Human cells (KGN)	Overexpression	Batista <i>et al.</i> (2007)
<i>IEX1</i>	Apoptosis	Activation	Human cells (KGN)	Overexpression	Batista <i>et al.</i> (2007)
<i>FOS</i>	Apoptosis	Activation	Human cells (KGN)	Overexpression	Batista <i>et al.</i> (2007)
<i>CDKN2A</i>	Cell cycle	Activation	Human cells (KGN)	Overexpression	Batista <i>et al.</i> (2007)
<i>SOD2</i>	Stress response	Activation	Human cells (KGN)	Overexpression	Benayoun <i>et al.</i> (2009)
<i>SIRT1</i>	Stress response	Activation	Human cells (KGN)	Overexpression	Benayoun <i>et al.</i> (2009)

misexpression is the responsible event. However, FOXL2 inactivation during foetal development in the goat is still required to formally demonstrate this very plausible hypothesis. This would epitomise dramatic species-specific differences because neither in human nor in mouse are FOXL2 mutations (thus far) known to lead to overt sex reversal.

Recent studies have shown that sexual identity is maintained at the cellular level throughout adulthood in mice. Thus, deletion of the TF *Dmrt1* in Sertoli cells of adult XY mice results in *Foxl2* expression and transdifferentiation of Sertoli cells into granulosa-like cells (Matson *et al.* 2011). Reciprocally, FOXL2 is required to maintain GC identity in females. Indeed, a deletion of *Foxl2* in GCs in adult mice leads to the expression of Sertoli-cell markers and to the appearance of structures reminiscent of seminiferous tubules, in the ovary (Uhlenhaut *et al.* 2009). A similar phenotype has been described in mice where oestrogen receptors *Esr1* and *Esr2* are deleted, or where oestrogen production is impaired, suggesting a functional synergy between oestrogen receptors and FOXL2 in the maintenance of GC identity (Fisher *et al.* 1998, Dupont *et al.* 2000). Interestingly, FOXL2 interacts directly with both oestrogen receptors and, along with ESR1, it directly represses a testis-specific enhancer of *Sox9* in mouse (Uhlenhaut *et al.* 2009).


FOXL2 has been described in various species as a positive regulator of *Cyp19a1*, which encodes the aromatase, responsible for the last step of oestrogen production in GCs (Pannetier *et al.* 2006, Uhlenhaut *et al.* 2009, Fleming *et al.* 2010, Rosario *et al.* 2012). This suggests that FOXL2 can influence directly and indirectly oestrogen signaling. As already noted, in addition to its role in repressing *Sox9*, FOXL2 is also required for the expression of FST, which is involved in follicle assembly (Kashimada *et al.* 2011, Caburet *et al.* 2012). Interestingly, *Fst* deletion in proliferating GCs causes premature depletion of the ovarian reserve and formation of Sertoli-like cells in the

ovary, quite similar to the phenotype observed for the deletion of *Foxl2*, suggesting that *Fst* could be a key target of FOXL2 (Jorgez *et al.* 2004, Yao *et al.* 2004). The role of FOXL2 in the maintenance of GC identity and oestrogen signaling in murine ovaries could explain the differences between the phenotypes of *Foxl2*^{-/-} mice, where ovaries develop apparently normally until birth, and PIS^{-/-} goats, in which decreased FOXL2 expression would induce XX-sex reversal. However, note that despite their normal appearance, *Foxl2*^{-/-} ovaries upregulate male markers such as *Sox9* or *Inhbb* during late developmental stages, suggesting a molecular loss of identity of pre-GCs, without overt macroscopic consequences (Garcia-Ortiz *et al.* 2009).

Follicle activation and GC differentiation

During foetal development, germ cells are organised in clusters (ovigerous cords) surrounded by somatic cells of the GC lineage. A large proportion of the germ cells undergoes apoptosis, while pre-GCs invade the interstices between germ cells to form a single layer around each surviving oocyte. Once formed, these primordial follicles enter a quiescent state in which they can stay for years or decades. After a first wave of folliculogenesis, reactivation of the primordial follicles takes place progressively during mature sexual life. The number of activated follicles is connected to the ovarian reserve and to the mono- or polyovulating capacity of the species in question. Deregulation of primordial follicle activation can thus cause depletion of the ovarian reserve and lead to premature menopause (Adhikari & Liu 2009). This process is finely regulated but the nature of the signal(s) triggering primordial follicle activation has remained elusive for years. Studies on the molecular mechanisms involved suggest that the communication between GCs and the oocyte is a key aspect of this process (Adhikari & Liu 2009).

Several factors are required for primordial follicle activation (Fig. 2). This is the case for the secreted factor

**Figure 2**

Overview of some factors implicated in follicle recruitment for maturation. (A) Summary and representation of principal macroscopic changes between primordial and primary follicles. (B) Representation of the principal factors implied to be involved in primordial follicle activation, and their relations, when they are known. Oocyte-transcribed factors are in black, granulosa-transcribed factors are in red and factors originating from other follicles or of unknown origin are in grey. Activation is represented by an arrow and repression by a blunt arrow.

c-Kit ligand (KITL, also known as stem cell factor), produced by GCs, and its receptor KIT expressed by the oocytes. Indeed, a Kit receptor-blocking antibody strongly prevents activation of primordial murine follicles (Yoshida *et al.* 1997). Conversely, treatment of explanted rat ovaries with KITL increases primordial follicle recruitment (Parrott & Skinner 1999). Mice with mutations of the *Kit* receptor have follicles blocked at the primary stage, suggesting that GCs can start their differentiation but that the oocyte does not exit quiescence (Huang *et al.* 1993). Stimulation of the oocyte by KITL in rodents seems to activate the PI3K/AKT pathway leading to the inactivating phosphorylation of the TF FOXO3A (Reddy *et al.* 2005). This is consistent with the role of FOXO3A and PTEN, an inhibitor of the PI3K/AKT pathway, in the preservation of oocyte quiescence. Oocyte-specific *Pten* knockout results in a constitutive activation of the

PI3K/AKT pathway and in a global activation of primordial follicles in the weeks following birth (John *et al.* 2008, Reddy *et al.* 2008). Other factors signaling through PI3K/AKT could also be involved in follicle activation. For example, the NTRK2 receptor, expressed in the oocytes of primordial follicles, and its ligands BDNF and NT-4 are also necessary for the recruitment of primordial follicles in mice (Paredes *et al.* 2004). Also, other growth factors, such as FGF2, LIF or FGF7, could modulate primordial follicle recruitment by modulating KITL expression (Nilsson *et al.* 2002, Nilsson & Skinner 2004, Kezele *et al.* 2005). Normal follicle activation also requires the expression of several oocyte-specific or -enriched TFs such as SOHL1, SOHL2, LHX8 and NOBOX. Indeed, their absence results in follicles arrested at the primary stage in mice (Rajkovic *et al.* 2004, Pangas *et al.* 2006, Choi *et al.* 2008a,b).

The AMH produced by the GCs of the primary-to-antral follicles is a key factor for the maintenance of the ovarian reserve (Baarends *et al.* 1995). AMH represses primordial follicle activation, suggesting the existence of a negative feedback mechanism avoiding anarchic follicle recruitment. Indeed, *Amh*^{-/-} mice display greater recruitment of primordial follicles and depletion of the follicular reserve (Durlinger *et al.* 1999). Conversely, treatment of explanted human ovaries or rat neonatal ovaries with recombinant AMH *in vitro* leads to a decreased recruitment of primordial follicles (Carlsson *et al.* 2006, Nilsson *et al.* 2007). AMH is thought to induce the activation of the BMP-dependent effectors SMAD1, SMAD5 and SMAD8. Interestingly, however, other BMP factors (i.e. BMP4 or BMP7) play an activating role in follicle recruitment in rats (Lee *et al.* 2001, Nilsson & Skinner 2003), and it has been suggested that AMH may diminish the production of BMP4 and other stimulatory factors (Nilsson *et al.* 2007). Further work is required to identify the precise molecular mechanisms underlying such processes probably involving combinatorial control.

Few determinants of GC differentiation at the transcriptional level have been described so far and FOXL2 is one of them. As previously mentioned, *Foxl2*^{-/-} mice develop normally until birth, at least in appearance. However, examination of their ovaries 2 weeks after birth reveals the presence of follicles with large oocytes surrounded by a single layer of flat GCs, and the absence of normal follicles at later maturation stages (Schmidt *et al.* 2004, Uda *et al.* 2004). In the following weeks, massive 'follicle activation' seems to occur, but no cuboidal GCs are observed. The oocytes of these 'activated' follicles quickly undergo apoptosis, provoking a depletion of the follicular reserve (Schmidt *et al.* 2004, Uda *et al.* 2004).

This mouse phenotype is reminiscent of human POI, suggesting that FOXL2 plays a similar role in humans. Indeed, *in vitro* functional analyses of intragenic FOXL2 mutations suggest that POI (type I BPES) is associated with the loss of FOXL2 transcriptional activity (Dipietromaria *et al.* 2009). FOXL2 haploinsufficiency in humans seems therefore to cause a milder version of the mouse phenotype. A recent analysis used these observations to propose a structural model to predict the outcome (i.e. the occurrence of POI) of FOXL2 mutations when they are located in the DNA-binding domain, thus offering the first tool for genetic counseling of young female BPES patients (Todeschini *et al.* 2011).

The nature of the transcriptional targets of FOXL2 in the pre-GCs has not been characterised in detail. An analysis of the ovarian transcriptome during the foetal period in *Foxl2*^{-/-} mice suggested that *Kitl* may be one of the transcriptional targets of FOXL2 during this period (Garcia-Ortiz *et al.* 2009). This is, however, in contradiction with another work that used *in situ* hybridisation to show that *Kitl* expression was normal in the ovaries of young and adult *Foxl2*^{-/-} mice (Schmidt *et al.* 2004). Other potential targets of FOXL2 in mouse foetal ovaries have been described (Garcia-Ortiz *et al.* 2009), although they may also reflect indirect effects of the absence of FOXL2. One of the most relevant FOXL2 targets suggested by this study may be the cell cycle inhibitor *Cdkn1b*, expressed in both oocytes and pre-GCs. The deletion of *Cdkn1b* in mice leads to an increased recruitment of primordial follicles and to a depletion of the follicular reserve before the age of three months (Rajareddy *et al.* 2007). FOXL2 could help maintaining granulosa quiescence by activating *Cdkn1b*, but it is not clear whether this is related to its role in GC differentiation. Other potentially interesting FOXL2 targets include the TGFβ effector *Smad3*, the BMP/AMH receptor *Bmpr1a* and the β-catenin repressor *Dkk3*, although an in depth analysis is required to characterise the *in vivo* effect of each of these factors, and others, in GC differentiation (Garcia-Ortiz *et al.* 2009).

Putative roles of FOXL2 during folliculogenesis

Owing to the loss of functional GCs in *Foxl2* knockout models, it is hard to determine whether FOXL2 also plays a role in other aspects of folliculogenesis, such as follicular growth, oestrogen production or ovulation (Cocquet *et al.* 2002, Schmidt *et al.* 2004). The examination of FOXL2 transcriptional targets in various contexts suggests that

FOXL2 may be important in many ovarian processes. Several transcriptomic studies on human GCT-derived cells or mice ovaries at different stages of development have described many potential targets of FOXL2 (Batista *et al.* 2007, Garcia-Ortiz *et al.* 2009, Uhlenhaut *et al.* 2009, Escudero *et al.* 2010, Rosario *et al.* 2012). These studies are limited, in the case of GCT-derived cells by the fact that the mutation or extinction of FOXL2 can affect cell identity in the long-term, and 'in the case of mouse ovaries' by the major morphological changes associated with the absence of FOXL2 and by the superposition of direct and indirect effects on multiple cell types (Caburet *et al.* 2012).

FOXL2 regulates several key genes necessary for oestrogen production. As previously mentioned, FOXL2 activates the expression of CYP19A1 in several mammalian models, including goat and mouse ovaries and human cells (Pannetier *et al.* 2006, Uhlenhaut *et al.* 2009, Fleming *et al.* 2010). Some studies have, however, suggested the existence of a repressive effect of FOXL2 on the expression of CYP19A1 in hamster and mouse cells, but an explanation for this difference is lacking (Pisarska *et al.* 2011, Dai *et al.* 2013). FOXL2 also regulates other enzymes of steroidogenesis. One of its first described targets is the gene encoding StAR, involved in intramitochondrial cholesterol transport, whose expression is repressed by FOXL2 in human cells (Pisarska *et al.* 2004, Rosario *et al.* 2012). FOXL2 also represses, in a human GCT-derived cell line, the *CYP17A1* gene, which encodes an enzyme responsible for androgen synthesis in the theca cells (Park *et al.* 2010). This suggests that FOXL2 can repress theca-specific genes in GCs, but *in vivo* confirmation is required.

FOXL2 may also regulate genes playing an essential role in late folliculogenesis. For example, FOXL2 overexpression in GCT-derived cells causes induction of prostaglandin synthase (or COX2), but it has also been showed that it could repress the induction of this gene by ESR1 in breast cancer or embryonic kidney cells (Batista *et al.* 2007, Kim *et al.* 2009), which could be due to a different cellular context. Mouse ovarian transcriptomic studies suggest, in addition, that FOXL2 regulates the expression of genes involved in ovulation such as *Serpine2*, *Has2*, *Ptger2* or *Ednra* (Caburet *et al.* 2012). FOXL2 may thus be required for the acquisition and maintenance of a normal transcriptional programme in GCs of the preovulatory follicles. However, further data are required to determine *in vivo* whether FOXL2 has a role in late folliculogenesis.

In addition to its direct role in GC function in the ovary, FOXL2 may also play a role in folliculogenesis

through its expression in the pituitary. Indeed, recent studies in mice have shown that FOXL2 is required for the production of FSH in gonadotrope cells. FOXL2 coregulates with SMAD3, an effector of activin in gonadotrope cells, the transcription of the *Fshb* gene, encoding the specific subunit of FSH (Lamba *et al.* 2009, 2010, Corpuz *et al.* 2010, Tran *et al.* 2011). Accordingly, the knock-out of *Foxl2*, constitutively or only in gonadotrope cells, leads to a defect in FSH production (Justice *et al.* 2011, Tran *et al.* 2013). Interestingly, cooperation of FOXL2 with SMAD3 is also observed on the promoter of the *Fst* gene in the pituitary and GCs (Blount *et al.* 2009, McTavish *et al.* 2013). Thus, SMAD3 seems to be a key partner of FOXL2 in several contexts.

Altogether, these observations suggest that FOXL2 may act as a global regulator of folliculogenesis, beyond its most obvious role in GC differentiation. It might therefore be interesting to examine its potential role in ovarian pathologies such as PCOS, and isolated POI.

FOXL2: a key player in GC proliferation and tumorigenesis

GCTs are by far the most common type of sex cord tumours and represent up to 5% of ovarian cancers. This type of cancer is also the most common ovarian cancer in other mammals, such as the mare, and appears spontaneously in certain strains of mice (Beamer *et al.* 1985, McCue *et al.* 2006). These tumours can cause hyperoestrogenism, resulting in various symptoms such as disturbance of menstruation or uterine bleeding after menopause (Tavassoli *et al.* 2003). There are two types of GCTs, classified according to their age of onset and histological features. The 'juvenile GCTs' account for 5% of the cases and usually occur before puberty, while the 'adult-type' tumours most often occur during perimenopause or after menopause. The prognosis of these tumours is generally favourable, but aggressive recurrence, usually fatal, occurs in 10–50% of cases. In addition, these tumours are slow growing and can recur several decades after removal of the primary tumour, requiring long-term monitoring of patients (Tavassoli *et al.* 2003).

FOXL2 misexpression has been detected in juvenile GCTs. Indeed, the expression of FOXL2 is reduced or extinct in some juvenile GCTs, correlating with an aggressive phenotype (Kalfa *et al.* 2007). A recent study using NextGen sequencing has identified a recurrent somatic mutation in the coding sequence of FOXL2 in 97% of adult-type tumours (Shah *et al.* 2009). This substitution of one base in the coding sequence of

FOXL2 (c.402>G) results in a p.C134W substitution at the protein level. This mutation was not found in healthy cells of the ovary or other tissues, indicating that the mutation occurs during tumorigenesis (Shah *et al.* 2009). Several studies have confirmed the high prevalence of this mutation and its specificity to adult-type GCTs. Over 94% of GCTs carry this mutation, and it has been suggested that the remaining ones are misdiagnosed sex cord tumours (Jamieson *et al.* 2010, Jamieson & Fuller 2012). The mutation was however identified in 20% of tested thecomas and also in GCTs of the testis (Jamieson & Fuller 2012, Lima *et al.* 2012).


FOXL2 regulates, directly and indirectly, genes involved in cell proliferation and apoptosis. Its overexpression in GCT-derived cells upregulates pro-apoptotic genes, such as *TNF-R1*, *FAS* or *TRAIL-R*, receptors of pro-apoptotic ligands (Kim *et al.* 2011). FOXL2 also activates the expression of the GNRH receptor in human and mouse GCs, which may play a pro-apoptotic role (Escudero *et al.* 2010, Cheng *et al.* 2013). The pro-apoptotic role of FOXL2 has mainly been studied in GCT cells, and has not been demonstrated *in vivo* yet. FOXL2 may also have an anti-proliferative role in regulating the cell cycle as it has been reported to activate the cycle inhibitor CDKN2A and to inhibit cyclin D2 (Batista *et al.* 2007, Bentsi-Barnes *et al.* 2010). Finally, FOXL2 could play a role in the oxidative stress response by regulating the expression of genes such as *SIRT1* and *SOD2* by directly binding to their promoters as shown in GCT-derived cells (Benayoun *et al.* 2009). Several studies have been conducted to elucidate how the C134W mutation is involved in tumorigenesis. The cysteine 134 of FOXL2 is located in the DNA-binding domain, and structural models suggest that the mutation does not directly affect DNA binding, but most probably the interaction with partners. Indeed, the C134W mutation does not cause a general loss of function *in vitro* and the mutated protein can activate normally most target promoters under conditions of overexpression (Benayoun *et al.* 2010). However, FOXL2 C134W has a decreased pro-apoptotic capacity when overexpressed in GCT-derived cultured cells (Kim *et al.* 2011). It has recently been suggested that deregulation of the expression of gonadotrophin-releasing hormone receptor (GNRHR) could contribute to the impairment in induction of apoptosis (Cheng *et al.* 2013). FOXL2-C134W has also been shown to overactivate the *CYP19A1* promoter, compared with the WT version, although the link with tumorigenesis is not clear. A genomic exploration identified an enrichment of FOXL2 transcriptional targets involved in modulating apoptosis and the cell cycle

among genes deregulated in adult GCT samples, compared with healthy (stimulated) GCs (Benayoun *et al.* 2013). A recent study has analysed the transcriptomic differences when FOXL2, WT or mutated, was overexpressed in juvenile GCT-derived cells. This study indicates that the mutation could affect the regulation of the TGF β pathway by FOXL2, promoting the expression of genes of the TGF β /activin/SMAD2/3 pathway, repressing the expression of genes of the BMP/SMAD1/5 pathway and thus promoting GC proliferation (Rosario *et al.* 2012). However, the molecular mechanism(s) by which the mutation induces these effects have not yet been characterised. Interestingly, it has recently been observed that the C134W mutation reduced the ability of FOXL2 to induce the transcription of the *FST* gene in cooperation with SMAD2/3, although it did not prevent the association of FOXL2 with SMAD3 (Nonis *et al.* 2013). We observe a very similar effect on an artificial promoter possessing three repeats of an AP-1/SMAD3/FOXL2 composite element found in the murine promoter of the GNRH receptor (3XGRAS; Ellsworth *et al.* 2003; Fig. 3). Using a constitutively active version of the TGFBR1 (TGFBR1-CA), we stimulated SMAD2 and SMAD3 in presence or absence of FOXL2, in COV434 cells. This led to a moderate activation of the 3XGRAS promoter, which increased in the presence of FOXL2. Overexpression of SMAD7, which binds to the receptor and prevents SMAD2/3 phosphorylation, led to the inhibition of 3XGRAS activation by FOXL2, confirming that the observed effect was SMAD2/3-dependent. Because COV434 cells express SMAD2 around ten times more than SMAD3 at the RNA level, we overexpressed SMAD3 to test the effect of a change in the SMAD2/SMAD3 balance. As expected, this led to a hyperactivation of 3XGRAS promoter, but interestingly mutated FOXL2 had a strongly decreased ability to upregulate 3XGRAS under these conditions. The same experiment in KGN cells, without SMAD3 overexpression, recapitulated the results obtained in COV434 cells with SMAD3 overexpression. This is probably due to the higher level of SMAD3 in KGN cells. This experiment and the one described by Nonis *et al.* (2013) suggest that FOXL2 C134W has a decreased ability to cooperate with SMAD3 in GCs. It would be interesting to quantify the affinity of WT and mutant FOXL2 for SMAD3, and their ability to recognise composite elements to identify the source of this reduced cooperation. A recent study has also revealed results indicative of a lower capacity of mutated FOXL2 to synergise with transcriptional partners such as NR2C1 or GMEB1, although the ability of mutant FOXL2 to interact with these proteins, at least in conditions of overexpression,

was retained (L'Hôte *et al.* 2012). Altogether, these observations suggest that the C134W mutation affects functional interactions of FOXL2 with its partners.

The transcriptomic profile of adult GCTs suggests that they come mainly from proliferating GCs of antral-to-preovulatory follicles. They express the FSH receptor and proliferate in response to FSH stimulation (Jamieson & Fuller 2012). The pathways involved in the proliferation and/or apoptosis of GCs in antral follicles are therefore potentially involved in the genesis of adult GCT. They involve notably the PKA/CREB, PI3K/AKT, MAPK/ERK, β -catenin and TGF β pathways, as epitomised by several mouse models that develop GCTs. For instance, deletion of *Inha*, which causes an absence of inhibin production, induces GCT formation with full penetrance, suggesting that inhibin is required to limit GC proliferation (Matzuk *et al.* 1992). Interestingly, deletion of *Smad1* and *Smad5* in GCs leads to a similar phenotype, suggesting that the BMP/SMAD pathway is required to control GC proliferation (Pangas *et al.* 2008). This is confirmed by the observation that a double *Bmpr1a/Bmpr1b* knockout also leads to GCT formation (Edson *et al.* 2010).

In another model of tumorigenesis, a constitutively active mutant of β -catenin was specifically expressed in GCs. These mice developed abnormal follicles with the presence of tumours in 57% of cases (Boerboom *et al.* 2005). Hypersecretion of LH also leads to GCTs in mice (Risma *et al.* 1995). Secondary inactivation of genes in these models has indicated various factors influencing the progression of GCTs. Thus, deletion of the genes coding the oestrogen receptors *Esr1* and/or *Esr2* in *Inha*^{-/-} mice leads to earlier and more aggressive tumour formation (Burns *et al.* 2003). Inactivation of *Pten* in mice where β -catenin is constitutively active also leads to a worsening of the tumour phenotype, suggesting that the PI3K/AKT pathway can be effectively mobilised in GCTs (Richards *et al.* 2012). More surprisingly, deregulation of the ERK pathway in the same mice, through the use of a *Kras* mutant, also leads to a worsening of the phenotype, while the introduction of this mutation into WT mice leads to cell cycle arrest in GCs (Richards *et al.* 2012). The MAPKs, particularly ERK, could thus play an important role in tumorigenesis of GCs. Several studies suggest that the ERK pathway is constitutively active in cell lines derived from GCTs, and inhibition of ERK1/2 by RNA interference can induce growth arrest of KGN cells (Steinmetz *et al.* 2004, Jamieson & Fuller 2012). However, the mutation of FOXL2 remains the main marker of adult-type GCTs, most of which express the *INHA* gene and do not display constitutive activation

**Figure 3**

Mutant FOXL2 fails to cooperate with SMAD3 on a composite regulatory element. (A) Luciferase activity of the reporter promoter 3XGRAS-luc in COV434 cells. Reporter vector (400 ng) was co-transfected with 200 ng of each of FOXL2 WT or C134W, TGFBR1-CA, SMAD3 and SMAD7, or pcDNA3.1 control vector, as indicated. Error bars represent the s.d. of five replicates. Asterisks represent a Student's *t*-test *P* value $< 10^{-3}$ between the indicated condition and control condition. The # symbol indicates a Student's *t*-test *P* value $< 10^{-3}$ between the FOXL2 WT and mutant conditions. (B) Expression value of SMAD1–9 mRNAs in COV434 cells and KGN cells, determined from microarray data (GEO datasets GSE39890 from

Rosario *et al.* (2012)). Error bars represent the s.d.s of three conditions. (C) Luciferase activity of the reporter promoter 3XGRAS-luc in KGN cells. Reporter vector (400 ng) was co-transfected with 300 ng of each of FOXL2 WT or C134W, TGFBR1-CA or pcDNA3.1 control vector, as indicated. Error bars represent the s.d. of five replicates. Asterisks represent a Student's *t*-test *P* value $< 10^{-3}$, between the indicated condition and control condition. The # symbol indicates a Student's *t*-test *P* value $< 10^{-3}$ between the FOXL2 WT and mutant conditions. Experiments were led as in L'Hôte *et al.* (2012) and Benayoun *et al.* (2013).

of β -catenin (Watson *et al.* 1997, Ohishi *et al.* 2011). This suggests that mouse models might be more representative of the juvenile type of GCTs. Further studies on the functional interactions of FOXL2 with these pathways are required to achieve a better understanding of GC tumorigenesis.

Regulation of FOXL2 expression and function

During mouse development, FOXL2 is expressed from 12 d.p.c. in the female genital ridge. Its expression seems

to require the GATA4–FOG2 complex and activation of the β -catenin pathway. However, factors regulating FOXL2 expression directly are largely unknown and the decreased FOXL2 expression in the various animal models mentioned above could be related to the decrease in the cell population expressing it (Manuylov *et al.* 2008). The developmental expression of FOXL2 is also reduced in *Cited2*^{-/-} mice. It has been suggested that this could be an indirect effect via the effect of CITED2 on *Nr5a1* (Combes *et al.* 2010) required for gonadal and other steroidogenic tissue formation. However, the observed effect on FOXL2

could also result from the reduction of the pool of GC precursors in *Cited2*^{-/-} mice.


Few TFs have been shown to directly regulate the expression of FOXL2. One of them is FOXL2 itself, which binds to its promoter to activate its own transcription, at least *in vitro* (Moumné *et al.* 2008). Another factor potentially regulating FOXL2 expression is retinoblastoma 1 (RB1). In the mouse, depletion of RB in GCs leads to an increase of follicle recruitment followed by massive follicular atresia and early depletion of the ovarian reserve. FOXL2 expression is reduced in the ovaries of *Rb*^{-/-} mice, suggesting a direct or indirect control (Andreu-Vieyra *et al.* 2008). Data from the ENCODE consortium (<http://genome.ucsc.edu/>) show binding of the RB-associated TFs ELF1, E2F1 and E2F6 to the FOXL2 proximal promoter, indicating a potential regulation by these factors. It would be interesting to study in detail the regulation of FOXL2 expression by RB and its partners, and also to determine whether FOXL2 expression is cell-cycle-dependent, as is the case for most RB-E2F targets (Poznic 2009).

Two recent studies have suggested that FOXL2 expression could also be regulated by miRNAs binding to its 3' UTR. Rosario *et al.* proposed that the difference observed in FOXL2 expression in two GC lines (COV434 and KGN) could be due to the action of miR17 family miRNAs. Such miRNAs could negatively regulate FOXL2 expression. The authors show that such miRNAs are expressed in COV434 cells, in which FOXL2 is absent, but are absent in KGN cells, in which FOXL2 is expressed, and that they regulate FOXL2 expression in a 3' UTR-dependent way (Rosario *et al.* 2013). However, no miR17-binding site could be detected in *FOXL2* mRNA, and it was proposed that the observed effect could be indirect. The second study, performed by Dai *et al.*, involves miR133b. Treatment of mouse primary GCs or human KGN cells with miR133b decreased *Foxl2* expression. Moreover, miR133b inhibited the observed *Foxl2*-mediated transcriptional repression of *STAR* and *CYP19A1* in KGN cells (Dai *et al.* 2013). The effect of miR133b on *Foxl2* expression seems to rely on its 3' UTR where putative binding sites were described, but direct binding is yet to be demonstrated.

The regulation of the FOXL2 expression and function can also involve post-translational modifications. The analysis by two-dimensional electrophoresis of the FOXL2 migration pattern suggests that the protein is highly modified post translationally (Benayoun *et al.* 2009). Several modified residues have been identified using mass spectrometry with overexpressed FOXL2.

Serine 33 appears to be one of the principal phosphorylated residues, other phosphorylated sites include serine 211 or tyrosine 215, serine 238 and serine 323 or 326. Interestingly, all *in vitro*-phosphorylated serine residues lie in a similar consensus PxSP, corresponding to the preferred site of MAPKs and structurally related kinases (Georges *et al.* 2011). Acetylation was detected on lysines 96, 114 and 366. The origins and exact roles of these modifications are still poorly understood. However, acetylation has been proposed to be necessary for activation by FOXL2 of a fraction of its target genes, while some targets would be more sensitive to less acetylated FOXL2. The deacetylase SIRT1, which interacts with FOXL2, could thus directly regulate the activity of FOXL2 (Benayoun *et al.* 2009, L'Hôte *et al.* 2012). An enzyme proposed to phosphorylate FOXL2 is the kinase LATS1, which interacts directly with the former. Inactivation of LATS1 with an interfering RNA results in a decreased level of FOXL2 phosphorylation and in a weaker repressive activity of FOXL2 on a reporter promoter (Pisarska *et al.* 2010). LATS1 is a tumour suppressor and a regulator of the HIPPO pathway, which is involved in many types of cancer (Liu *et al.* 2012). *Lats1* depletion in mice results in an almost complete absence of antral follicles, anovulation and in ovarian tumours of stromal origin. It has been suggested that infertility could be a side effect of the tumours (St John *et al.* 1999). FOXL2 interacts directly with the SUMO ligases PIAS1 and UBC9 (Kuo *et al.* 2009). Several potentially SUMOylated FOXL2 residues have been identified and it is not certain that this modification affects any residue in particular. SUMOylation seems to stabilise FOXL2 and consequently increases its transcriptional (activation or repression) activity (Kuo *et al.* 2009, Marongiu *et al.* 2010, Georges *et al.* 2011). Another consequence of SUMOylation under overexpression conditions is an increased recruitment of FOXL2 to PML nuclear bodies. FOXL2 recruitment to these bodies should be necessary to its stabilisation (Georges *et al.* 2011). However, confirmation of these findings *in vivo* is still required.

Beyond the regulation of its expression, FOXL2 function may be regulated through interactions with other proteins. It can form homo-dimers (multimers?), but it is not known whether they are transcriptionally active or not (Lamba *et al.* 2009). The binding sites identified so far suggest a monomeric binding of FOXL2 (Lamba *et al.* 2009). It has been suggested that FOXL2 multimerisation could induce dominant negative effects (Kuo *et al.* 2011). Indeed, it is known that aggregates formed by some FOXL2 mutants can trap the WT protein (Moumné *et al.* 2008). FOXL2 also interacts with the

**Figure 4**

Overview of FOXL2 roles during folliculogenesis. Bold letters represent the physiological roles asserted from observations on mouse models or human pathologies (italicized). Normal letters correspond to potential roles suggested by *in vitro* studies.

pro-apoptotic factor DEADbox helicase DDX20. Interestingly, co-overexpression of DDX20 and FOXL2 in GCs leads to a decreased cell viability, but the targets potentially co-regulated by these factors are yet to be identified (Lee *et al.* 2005a). FOXL2 also directly interacts with several nuclear receptors important for ovarian development and function. This is the case for ESR1 and ESR2, the orphan nuclear receptor NR5A1 and its paralog NR5A2, the retinoic-acid-sensitive receptor NR2C1 and the progesterone receptor PGR (Kim *et al.* 2009, Uhlenhaut *et al.* 2009, Park *et al.* 2010, Ghochani *et al.* 2012, L'Hôte *et al.* 2012). Other forkhead TFs such as FOXO1 and FOXO3 can interact with a variety of nuclear receptors, suggesting a conserved interaction between these two families (van der Vos & Coffey 2008). We have recently reported an interaction of FOXL2 with a coregulator of nuclear receptors (GMEB1) as well as with the TF CREM, involved in cAMP response. Finally, FOXL2 also interacts with chromatin regulators, such as BANF1, the DNA-repair protein XRCC6 and the histone-deacetylase SIRT1, suggesting that FOXL2 could indeed play a role in chromatin organisation (L'Hôte *et al.* 2012). Although an increasing number of FOXL2-interacting partners are known, the precise role of these interactions have been poorly characterised thus far. These interactions have indeed mostly been observed under conditions of over-expression and in non-physiological contexts. Future works clarifying the nature and function of the complexes involving FOXL2 *in vivo* would therefore be greatly informative.

Future directions

Available data show that FOXL2 performs multiple roles throughout folliculogenesis (Fig. 4). Its expression and

activity seem to be finely regulated through several mechanisms and its mutation/dysregulation can be responsible for ovarian pathologies, including cancer. FOXL2 interacts with effectors and modulators of various signaling pathways, and it is therefore interesting to characterise the effect of these pathways on FOXL2 activity and target recognition. Further studies of the influence of FOXL2 on multiple signaling pathways mobilised in GCs (and vice versa) may therefore be of great interest.

The central role of FOXL2 in GC differentiation is not an isolated case within the forkhead family. Other paralogs such as FOXA1, FOXA2, FOXI1, FOXP3 and FOXG1 are also required for the determination of specific cell lineages (Lee *et al.* 2005b, El-Dahr *et al.* 2008, Danesin & Houart 2012, Ohkura *et al.* 2013). It has been proposed that FOXA1 and FOXA2 play the role of pioneer TFs, which are able to remodel chromatin in a lineage-specific way. These factors thus define the subset of potential binding sites occupied by other TFs in differentiated cells, and in particular those of ubiquitous effectors such as SMADs, STATs or oestrogen receptors (Zaret & Carroll 2011). The central position of FOXL2 in determining and maintaining GC identity suggests that it might also be a pioneer factor. Thus, it would be interesting to determine whether FOXL2 is implicated in chromatin remodeling, as suggested by its interaction with some chromatin remodeling factors. Finally, lineage-determining TFs do not act alone but in cooperation within complex regulatory networks. The determination of key signaling pathways and partners regulating or interacting with FOXL2, along with a better definition of its direct transcriptional targets, could therefore allow a better understanding of the role of FOXL2 in the ovary.

A recurrent issue in FOXL2 studies is the lack of a suitable cell model. Primary GCs from mice or rats may

constitute a good model, but they tend to transdifferentiate *in vitro*, making their use rather complex and impractical for many experiments. Cultured cells from human GCTs are the most commonly used models, but they either express a mutant form of FOXL2 (KGN cell line) or do not express it at all (COV434 cell line), therefore being rather unrealistic as models of healthy GC. Recent studies have managed to derive apparently functional GCs from human and mouse embryonic stem cells (Lan *et al.* 2013, Woods *et al.* 2013). Such models could allow a better understanding of FOXL2 function in follicle function and tumorigenesis in the future.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the review.

Funding

This work was supported by the Centre National de la Recherche Scientifique, La Ligue Nationale contre le Cancer (Comité de Paris), Université Paris Diderot-Paris7, Institut Universitaire de France and l'Agence Nationale de la Recherche (ANR).

Author contribution statement

A Auguste, L Bessière and A Vanet contributed equally to this work and are listed alphabetically.

Acknowledgements

We thank two anonymous referees for helpful comments on the manuscript.

References

- Adhikari D & Liu K 2009 Molecular mechanisms underlying the activation of mammalian primordial follicles. *Endocrine Reviews* **30** 438–464. (doi:10.1210/er.2008-0048)
- Andreu-Vieyra C, Chen R & Matzuk MM 2008 Conditional deletion of the retinoblastoma (*Rb*) gene in ovarian granulosa cells leads to premature ovarian failure. *Molecular Endocrinology* **22** 2141–2161. (doi:10.1210/me.2008-0033)
- Auguste A, Chassot A-A, Grégoire EP, Renault L, Pannetier M, Treier M, Pailhoux E & Chaboissier M-C 2011 Loss of *R-spondin1* and *Foxl2* amplifies female-to-male sex reversal in XX mice. *Sexual Development* **5** 304–317. (doi:10.1159/000334517)
- Baarends WM, Uilenbroek JT, Kramer P, Hoogerbrugge JW, van Leeuwen EC, Themmen AP & Grootegoed JA 1995 Anti-Müllerian hormone and anti-Müllerian hormone type II receptor messenger ribonucleic acid expression in rat ovaries during postnatal development, the estrous cycle, and gonadotropin-induced follicle growth. *Endocrinology* **136** 4951–4962. (doi:10.1210/en.136.11.4951)
- Barrionuevo F, Bagheri-Fam S, Klattig J, Kist R, Taketo MM, Englert C & Scherer G 2006 Homozygous inactivation of *Sox9* causes complete XY sex reversal in mice. *Biology of Reproduction* **74** 195–201. (doi:10.1095/biolreprod.105.045930)
- Batista F, Vaiman D, Dausset J, Fellous M & Veitia RA 2007 Potential targets of FOXL2, a transcription factor involved in craniofacial and follicular development, identified by transcriptomics. *PNAS* **104** 3330–3335. (doi:10.1073/pnas.0611326104)
- Beamer WG, Hoppe PC & Whitten WK 1985 Spontaneous malignant granulosa cell tumors in ovaries of young SWR mice. *Cancer Research* **45** 5575–5581.
- Benayoun BA, Batista F, Auer J, Dipietromaria A, L'Hôte D, De Baere E & Veitia RA 2009 Positive and negative feedback regulates the transcription factor FOXL2 in response to cell stress: evidence for a regulatory imbalance induced by disease-causing mutations. *Human Molecular Genetics* **18** 632–644. (doi:10.1093/hmg/ddn389)
- Benayoun BA, Caburet S, Dipietromaria A, Georges A, D'Haene B, Pandaranayaka PJE, L'Hôte D, Todeschini A-L, Krishnaswamy S, Fellous M *et al.* 2010 Functional exploration of the adult ovarian granulosa cell tumor-associated somatic FOXL2 mutation p.Cys134Trp (c.402C>G). *PLoS ONE* **5** e8789. (doi:10.1371/journal.pone.0008789)
- Benayoun BA, Anttonen M, L'Hôte D, Bailly-Bechet M, Andersson N, Heikinheimo M & Veitia RA 2013 Adult ovarian granulosa cell tumor transcriptomics: prevalence of FOXL2 target genes misregulation gives insights into the pathogenic mechanism of the p.Cys134Trp somatic mutation. *Oncogene* **32** 2739–2746. (doi:10.1038/onc.2012.298)
- Bentsi-Barnes IK, Kuo F-T, Barlow GM & Pisarska MD 2010 Human forkhead L2 represses key genes in granulosa cell differentiation including aromatase, P450_{scc}, and cyclin D2. *Fertility and Sterility* **94** 353–356. (doi:10.1016/j.fertnstert.2009.09.050)
- Bernard P, Sim H, Kowner K, Vilain E & Harley V 2008 Human SRY inhibits β -catenin-mediated transcription. *International Journal of Biochemistry & Cell Biology* **40** 2889–2900. (doi:10.1016/j.biocel.2008.06.006)
- Blount AL, Schmidt K, Justice NJ, Vale WW, Fischer WH & Bilezikjian LM 2009 FoxL2 and Smad3 coordinately regulate follistatin gene transcription. *Journal of Biological Chemistry* **284** 7631–7645. (doi:10.1074/jbc.M806676200)
- Boerboom D, Paquet M, Hsieh M, Liu J, Jamin SP, Behringer RR, Sirois J, Taketo MM & Richards JS 2005 Misregulated Wnt/ β -catenin signaling leads to ovarian granulosa cell tumor development. *Cancer Research* **65** 9206–9215. (doi:10.1158/0008-5472.CAN-05-1024)
- Boulanger L, Kocer A, Daniel N, Pannetier M, Chesné P, Heyman Y, Renault L, Mandon-Pépin B, Chavatte-Palmer P, Vignon X *et al.* 2008 Attempt to rescue sex-reversal by transgenic expression of the *PISRT1* gene in XX *PIS^{-/-}* goats. *Sexual Development* **2** 142–151. (doi:10.1159/000143432)
- Burns KH, Agno JE, Chen L, Haupt B, Ogbonna SC, Korach KS & Matzuk MM 2003 Sexually dimorphic roles of steroid hormone receptor signaling in gonadal tumorigenesis. *Molecular Endocrinology* **17** 2039–2052. (doi:10.1210/me.2003-0039)
- Caburet S, Georges A, L'Hôte D, Todeschini A-L, Benayoun BA & Veitia RA 2012 The transcription factor FOXL2: at the crossroads of ovarian physiology and pathology. *Molecular and Cellular Endocrinology* **356** 55–64. (doi:10.1016/j.mce.2011.06.019)
- Carlsson IB, Scott JE, Visser JA, Ritvos O, Themmen APN & Hovatta O 2006 Anti-Müllerian hormone inhibits initiation of growth of human primordial ovarian follicles *in vitro*. *Human Reproduction* **21** 2223–2227. (doi:10.1093/humrep/del165)
- Chaboissier M-C, Kobayashi A, Vidal VIP, Lützkendorf S, van de Kant HJG, Wegner M, de Rooij DG, Behringer RR & Schedl A 2004 Functional analysis of *Sox8* and *Sox9* during sex determination in the mouse. *Development* **131** 1891–1901. (doi:10.1242/dev.01087)
- Chassot A-A, Ranc F, Grégoire EP, Roepers-Gajadien HL, Taketo MM, Camerino G, de Rooij DG, Schedl A & Chaboissier M-C 2008 Activation of β -catenin signaling by *Rspo1* controls differentiation of the mammalian ovary. *Human Molecular Genetics* **17** 1264–1277. (doi:10.1093/hmg/ddn016)
- Cheng J-C, Klausen C & Leung PCK 2013 Overexpression of wild-type but not C134W mutant FOXL2 enhances GnRH-induced cell apoptosis by

- increasing GnRH receptor expression in human granulosa cell tumors. *PLoS ONE* **8** e55099. (doi:10.1371/journal.pone.0055099)
- Choi Y, Ballow DJ, Xin Y & Rajkovic A 2008a Lim homeobox gene, *Lhx8*, is essential for mouse oocyte differentiation and survival. *Biology of Reproduction* **79** 442–449. (doi:10.1095/biolreprod.108.069393)
- Choi Y, Yuan D & Rajkovic A 2008b Germ cell-specific transcriptional regulator *Sohlh2* is essential for early mouse folliculogenesis and oocyte-specific gene expression. *Biology of Reproduction* **79** 1176–1182. (doi:10.1095/biolreprod.108.071217)
- Cocquet J, Pailhoux E, Jaubert F, Servel N, Xia X, Pannetier M, Baere ED, Messiaen L, Cotinot C, Fellous M *et al.* 2002 Evolution and expression of *FOXL2*. *Journal of Medical Genetics* **39** 916–921. (doi:10.1136/jmg.39.12.916)
- Cocquet J, Baere ED, Gareil M, Pannetier M, Xia X, Fellous M & Veitia RA 2003 Structure, evolution and expression of the *FOXL2* transcription unit. *Cytogenetic and Genome Research* **101** 206–211. (doi:10.1159/000074338)
- Combes AN, Spiller CM, Harley VR, Sinclair AH, Dunwoodie SL, Wilhelm D & Koopman P 2010 Gonadal defects in *Cited2*-mutant mice indicate a role for SF1 in both testis and ovary differentiation. *International Journal of Developmental Biology* **54** 683–689. (doi:10.1387/ijdb.092920ac)
- Corpus PS, Lindaman LL, Mellon PL & Coss D 2010 *Foxl2* is required for activin induction of the mouse and human follicle-stimulating hormone β -subunit genes. *Molecular Endocrinology* **24** 1037–1051. (doi:10.1210/me.2009-0425)
- Crisponi L, Deiana M, Loi A, Chiappe F, Uda M, Amati P, Bisceglia L, Zelante L, Nagaraja R, Porcu S *et al.* 2001 The putative forkhead transcription factor *FOXL2* is mutated in blepharophimosis/ptosis/epicanthus inversus syndrome. *Nature Genetics* **27** 159–166. (doi:10.1038/84781)
- Dai A, Sun H, Fang T, Zhang Q, Wu S, Jiang Y, Ding L, Yan G & Hu Y 2013 MicroRNA-133b stimulates ovarian estradiol synthesis by targeting *Foxl2*. *FEBS Letters* **587** 2474–2482. (doi:10.1016/j.febslet.2013.06.023)
- Danesin C & Houart C 2012 A Fox stops the Wnt: implications for forebrain development and diseases. *Current Opinion in Genetics & Development* **22** 323–330. (doi:10.1016/j.gde.2012.05.001)
- De Santa Barbara P, Méjean C, Moniot B, Malclès MH, Berta P & Boizet-Bonhoure B 2001 Steroidogenic factor-1 contributes to the cyclic-adenosine monophosphate down-regulation of human *SRY* gene expression. *Biology of Reproduction* **64** 775–783. (doi:10.1095/biolreprod64.3.775)
- De Vos M, Devroey P & Fauser BC 2011 Primary ovarian insufficiency. *Lancet* **376** 911–921. (doi:10.1016/S0140-6736(10)60355-8)
- Dipietromaria A, Benayoun BA, Todeschini A-L, Rivals I, Bazin C & Veitia RA 2009 Towards a functional classification of pathogenic *FOXL2* mutations using transactivation reporter systems. *Human Molecular Genetics* **18** 3324–3333. (doi:10.1093/hmg/ddp273)
- Dupont S, Krust A, Gansmuller A, Dierich A, Chambon P & Mark M 2000 Effect of single and compound knockouts of estrogen receptors α (*ER α*) and β (*ER β*) on mouse reproductive phenotypes. *Development* **127** 4277–4291.
- Durlinger ALL, Kramer P, Karels B, de Jong FH, Uilenbroek JTT, Grootegoed JA & Themmen APN 1999 Control of primordial follicle recruitment by anti-Müllerian hormone in the mouse ovary. *Endocrinology* **140** 5789–5796. (doi:10.1210/en.140.12.5789)
- Edson MA, Nagaraja AK & Matzuk MM 2009 The mammalian ovary from genesis to revelation. *Endocrine Reviews* **30** 624–712. (doi:10.1210/er.2009-0012)
- Edson MA, Nalam RL, Clementi C, Franco HL, Demayo FJ, Lyons KM, Pangas SA & Matzuk MM 2010 Granulosa cell-expressed *BMPRI1A* and *BMPRI1B* have unique functions in regulating fertility but act redundantly to suppress ovarian tumor development. *Molecular Endocrinology* **24** 1251–1266. (doi:10.1210/me.2009-0461)
- El-Dahr SS, Aboudehen K & Saifudeen Z 2008 Transcriptional control of terminal nephron differentiation. *American Journal of Physiology. Renal Physiology* **294** F1273–F1278. (doi:10.1152/ajprenal.00562.2007)
- Ellsworth BS, Burns AT, Escudero KW, Duval DL, Nelson SE & Clay CM 2003 The gonadotropin releasing hormone (GnRH) receptor activating sequence (GRAS) is a composite regulatory element that interacts with multiple classes of transcription factors including Smads, AP-1 and a forkhead DNA binding protein. *Molecular and Cellular Endocrinology* **206** 93–111. (doi:10.1016/S0303-7207(03)00235-1)
- Escudero JM, Haller JL, Clay CM & Escudero KW 2010 Microarray analysis of *Foxl2* mediated gene regulation in the mouse ovary derived KK1 granulosa cell line: over-expression of *Foxl2* leads to activation of the gonadotropin releasing hormone receptor gene promoter. *Journal of Ovarian Research* **3** 4. (doi:10.1186/1757-2215-3-4)
- Fisher CR, Graves KH, Parlow AF & Simpson ER 1998 Characterization of mice deficient in aromatase (ArKO) because of targeted disruption of the *cyp19* gene. *PNAS* **95** 6965–6970. (doi:10.1073/pnas.95.12.6965)
- Fleming NI, Krowner KC, Lazarus KA, Fuller PJ, Simpson ER & Clyne CD 2010 Aromatase is a direct target of *FOXL2*: C134W in granulosa cell tumors via a single highly conserved binding site in the ovarian specific promoter. *PLoS ONE* **5** e14389. (doi:10.1371/journal.pone.0014389)
- García-Ortiz JE, Pelosi E, Omari S, Nedorezov T, Piao Y, Karmazin J, Uda M, Cao A, Cole SW, Forabosco A *et al.* 2009 *Foxl2* functions in sex determination and histogenesis throughout mouse ovary development. *BMC Developmental Biology* **9** 36. (doi:10.1186/1471-213X-9-36)
- Georges A, Benayoun BA, Marongiu M, Dipietromaria A, L'Hôte D, Todeschini A-L, Auer J, Crisponi L & Veitia RA 2011 SUMOylation of the forkhead transcription factor *FOXL2* promotes its stabilization/activation through transient recruitment to PML bodies. *PLoS ONE* **6** e25463. (doi:10.1371/journal.pone.0025463)
- Ghochani Y, Saini JK, Mellon PL & Thackray VG 2012 *FOXL2* is involved in the synergy between activin and progestins on the follicle-stimulating hormone β -subunit promoter. *Endocrinology* **153** 2023–2033. (doi:10.1210/en.2011-1763)
- Hiramatsu R, Harikae K, Tsunekawa N, Kurohmaru M, Matsuo I & Kanai Y 2010 FGF signaling directs a center-to-pole expansion of tubulogenesis in mouse testis differentiation. *Development* **137** 303–312. (doi:10.1242/dev.040519)
- Huang EJ, Manova K, Packer AI, Sanchez S, Bachvarova RF & Besmer P 1993 The murine steel panda mutation affects Kit ligand expression and growth of early ovarian follicles. *Developmental Biology* **157** 100–109. (doi:10.1006/dbio.1993.1115)
- Jamieson S & Fuller PJ 2012 Molecular pathogenesis of granulosa cell tumors of the ovary. *Endocrine Reviews* **33** 109–144. (doi:10.1210/er.2011-0014)
- Jamieson S, Butzow R, Andersson N, Alexiadis M, Unkila-Kallio L, Heikinheimo M, Fuller PJ & Anttonen M 2010 The *FOXL2* C134W mutation is characteristic of adult granulosa cell tumors of the ovary. *Modern Pathology* **23** 1477–1485. (doi:10.1038/modpathol.2010.145)
- John GB, Gallardo TD, Shirley LJ & Castrillon DH 2008 *Foxo3* is a PI3K-dependent molecular switch controlling the initiation of oocyte growth. *Developmental Biology* **321** 197–204. (doi:10.1016/j.ydbio.2008.06.017)
- Jorgez CJ, Klysiak M, Jamin SP, Behringer RR & Matzuk MM 2004 Granulosa cell-specific inactivation of follistatin causes female fertility defects. *Molecular Endocrinology* **18** 953–967. (doi:10.1210/me.2003-0301)
- Justice NJ, Blount AL, Pelosi E, Schlessinger D, Vale W & Bilezikjian LM 2011 Impaired *FSH β* expression in the pituitaries of *Foxl2* mutant animals. *Molecular Endocrinology* **25** 1404–1415. (doi:10.1210/me.2011-0093)
- Kalfa N, Philibert P, Patte C, Ecochard A, Duviard P, Baldet P, Jaubert F, Fellous M & Sultan C 2007 Extinction of *FOXL2* expression in aggressive ovarian granulosa cell tumors in children. *Fertility and Sterility* **87** 896–901. (doi:10.1016/j.fertnstert.2006.11.016)
- Kashimada K, Pelosi E, Chen H, Schlessinger D, Wilhelm D & Koopman P 2011 *FOXL2* and *BMP2* act cooperatively to regulate follistatin gene

- expression during ovarian development. *Endocrinology* **152** 272–280. (doi:10.1210/en.2010-0636)
- Kezele P, Nilsson EE & Skinner MK 2005 Keratinocyte growth factor acts as a mesenchymal factor that promotes ovarian primordial to primary follicle transition. *Biology of Reproduction* **73** 967–973. (doi:10.1095/biolreprod.105.043117)
- Kim S-Y, Weiss J, Tong M, Laronda MM, Lee E-J & Jameson JL 2009 Foxl2, a forkhead transcription factor, modulates nonclassical activity of the estrogen receptor- α . *Endocrinology* **150** 5085–5093. (doi:10.1210/en.2009-0313)
- Kim J-H, Yoon S, Park M, Park H-O, Ko J-J, Lee K & Bae J 2011 Differential apoptotic activities of wild-type FOXL2 and the adult-type granulosa cell tumor-associated mutant FOXL2 (C134W). *Oncogene* **30** 1653–1663. (doi:10.1038/ncr.2010.541)
- Kuo F-T, Bentsi-Barnes IK, Barlow GM, Bae J & Pisarska MD 2009 Sumoylation of forkhead L2 by Ubc9 is required for its activity as a transcriptional repressor of the steroidogenic acute regulatory gene. *Cellular Signalling* **21** 1935–1944. (doi:10.1016/j.cellsig.2009.09.001)
- Kuo F-T, Bentsi-Barnes IK, Barlow GM & Pisarska MD 2011 Mutant forkhead L2 (FOXL2) proteins associated with premature ovarian failure (POF) dimerize with wild-type FOXL2, leading to altered regulation of genes associated with granulosa cell differentiation. *Endocrinology* **152** 3917–3929. (doi:10.1210/en.2010-0989)
- Lakhal B, Ben-Hadj-Khalifa S, Bouali N, Philipert P, Audran F, Braham R, Hatem E, Sultan C & Saad A 2012 Mutational screening of *SF1* and *WNT4* in Tunisian women with premature ovarian failure. *Gene* **509** 298–301. (doi:10.1016/j.gene.2012.08.007)
- Lamba P, Fortin J, Tran S, Wang Y & Bernard DJ 2009 A novel role for the forkhead transcription factor FOXL2 in activin A-regulated follicle-stimulating hormone β subunit transcription. *Molecular Endocrinology* **23** 1001–1013. (doi:10.1210/me.2008-0324)
- Lamba P, Wang Y, Tran S, Ouspenskaia T, Libasci V, Hébert TE, Miller GJ & Bernard DJ 2010 Activin A regulates porcine follicle-stimulating hormone β -subunit transcription via cooperative actions of SMADs and FOXL2. *Endocrinology* **151** 5456–5467. (doi:10.1210/en.2010-0605)
- Lan C-W, Chen M-J, Jan P-S, Chen H-F & Ho H-N 2013 Differentiation of human embryonic stem cells into functional ovarian granulosa-like cells. *Journal of Clinical Endocrinology and Metabolism* **98** 3713–3723. (doi:10.1210/jc.2012-4302)
- Lee W-S, Otsuka F, Moore RK & Shimasaki S 2001 Effect of bone morphogenetic protein-7 on folliculogenesis and ovulation in the rat. *Biology of Reproduction* **65** 994–999. (doi:10.1095/biolreprod65.4.994)
- Lee MB, Lebedeva LA, Suzawa M, Wadekar SA, Desclozeaux M & Ingraham HA 2005a The DEAD-box protein DP103 (Ddx20 or Gemin-3) represses orphan nuclear receptor activity via SUMO modification. *Molecular and Cellular Biology* **25** 1879–1890. (doi:10.1128/MCB.25.5.1879-1890.2005)
- Lee CS, Friedman JR, Fulmer JT & Kaestner KH 2005b The initiation of liver development is dependent on Foxa transcription factors. *Nature* **435** 944–947. (doi:10.1038/nature03649)
- L'Hôte D, Georges A, Todeschini AL, Kim J-H, Benayoun BA, Bae J & Veitia RA 2012 Discovery of novel protein partners of the transcription factor FOXL2 provides insights into its physiopathological roles. *Human Molecular Genetics* **21** 3264–3274. (doi:10.1093/hmg/dds170)
- Lima JF, Jin L, de Araujo ARC, Erikson-Johnson MR, Oliveira AM, Sebo TJ, Keeney GL & Medeiros F 2012 FOXL2 mutations in granulosa cell tumors occurring in males. *Archives of Pathology & Laboratory Medicine* **136** 825–828. (doi:10.5858/arpa.2011-0355-OA)
- Liu C-F, Bingham N, Parker K & Yao HH-C 2009 Sex-specific roles of β -catenin in mouse gonadal development. *Human Molecular Genetics* **18** 405–417. (doi:10.1093/hmg/ddn362)
- Liu AM, Wong K-F, Jiang X, Qiao Y & Luk JM 2012 Regulators of mammalian Hippo pathway in cancer. *Biochimica et Biophysica Acta* **1826** 357–364. (doi:10.1016/j.bbcan.2012.05.006)
- Maatouk DM, DiNapoli L, Alvers A, Parker KL, Taketo MM & Capel B 2008 Stabilization of β -catenin in XY gonads causes male-to-female sex-reversal. *Human Molecular Genetics* **17** 2949–2955. (doi:10.1093/hmg/ddn193)
- Manuylov NL, Smagulova FO, Leach L & Tevosian SG 2008 Ovarian development in mice requires the GATA4–FOG2 transcription complex. *Development* **135** 3731–3743. (doi:10.1242/dev.024653)
- Marongiu M, Deiana M, Meloni A, Marcia L, Puddu A, Cao A, Schlessinger D & Crisponi L 2010 The forkhead transcription factor Foxl2 is sumoylated in both human and mouse: sumoylation affects its stability, localization, and activity. *PLoS ONE* **5** e9477. (doi:10.1371/journal.pone.0009477)
- Matson CK, Murphy MW, Sarver AL, Griswold MD, Bardwell VJ & Zarkower D 2011 DMRT1 prevents female reprogramming in the postnatal mammalian testis. *Nature* **476** 101–104. (doi:10.1038/nature10239)
- Matzuk MM, Finegold MJ, Su JG, Hsueh AJ & Bradley A 1992 α -Inhibin is a tumour-suppressor gene with gonadal specificity in mice. *Nature* **360** 313–319. (doi:10.1038/360313a0)
- McCue PM, Roser JF, Munro CJ, Liu IKM & Lasley BL 2006 Granulosa cell tumors of the equine ovary. *Veterinary Clinics of North America. Equine Practice* **22** 799–817. (doi:10.1016/j.cveq.2006.08.008)
- McTavish KJ, Nonis D, Hoang YD & Shimasaki S 2013 Granulosa cell tumor mutant FOXL2^{C134W} suppresses GDF-9 and activin A-induced follistatin transcription in primary granulosa cells. *Molecular and Cellular Endocrinology* **372** 57–64. (doi:10.1016/j.mce.2013.03.021)
- Miyamoto Y, Taniguchi H, Hamel F, Silversides DW & Viger RS 2008 A GATA4/WT1 cooperation regulates transcription of genes required for mammalian sex determination and differentiation. *BMC Molecular Biology* **9** 44. (doi:10.1186/1471-2199-9-44)
- Mouméné L, Dipietromaria A, Batista F, Kocer A, Fellous M, Pailhoux E & Veitia RA 2008 Differential aggregation and functional impairment induced by polyaniline expansions in FOXL2, a transcription factor involved in cranio-facial and ovarian development. *Human Molecular Genetics* **17** 1010–1019. (doi:10.1093/hmg/ddm373)
- Nilsson EE & Skinner MK 2003 Bone morphogenetic protein-4 acts as an ovarian follicle survival factor and promotes primordial follicle development. *Biology of Reproduction* **69** 1265–1272. (doi:10.1095/biolreprod.103.018671)
- Nilsson EE & Skinner MK 2004 Kit ligand and basic fibroblast growth factor interactions in the induction of ovarian primordial to primary follicle transition. *Molecular and Cellular Endocrinology* **214** 19–25. (doi:10.1016/j.mce.2003.12.001)
- Nilsson EE, Kezele P & Skinner MK 2002 Leukemia inhibitory factor (LIF) promotes the primordial to primary follicle transition in rat ovaries. *Molecular and Cellular Endocrinology* **188** 65–73. (doi:10.1016/S0303-7207(01)00746-8)
- Nilsson E, Rogers N & Skinner MK 2007 Actions of anti-Müllerian hormone on the ovarian transcriptome to inhibit primordial to primary follicle transition. *Reproduction* **134** 209–221. (doi:10.1530/REP-07-0119)
- Nonis D, McTavish KJ & Shimasaki S 2013 Essential but differential role of FOXL2^{wt} and FOXL2^{C134W} in GDF-9 stimulation of follistatin transcription in co-operation with Smad3 in the human granulosa cell line COV434. *Molecular and Cellular Endocrinology* **372** 42–48. (doi:10.1016/j.mce.2013.02.020)
- Ohishi Y, Oda Y, Kurihara S, Kaku T, Kobayashi H, Wake N & Tsuneyoshi M 2011 Nuclear localization of E-cadherin but not β -catenin in human ovarian granulosa cell tumours and normal ovarian follicles and ovarian stroma. *Histopathology* **58** 423–432. (doi:10.1111/j.1365-2559.2011.03761.x)
- Ohkura N, Kitagawa Y & Sakaguchi S 2013 Development and maintenance of regulatory T cells. *Immunity* **38** 414–423. (doi:10.1016/j.immuni.2013.03.002)
- Pailhoux E, Vigier B, Schibler L, Cribeu EP, Cotinot C & Vaiman D 2005 Positional cloning of the PIS mutation in goats and its impact on understanding mammalian sex-differentiation. *Genetics, Selection, Evolution* **37** (Suppl 1) S55–S64. (doi:10.1186/1297-9686-37-S1-S55)

- Pangas SA, Choi Y, Ballow DJ, Zhao Y, Westphal H, Matzuk MM & Rajkovic A 2006 Oogenesis requires germ cell-specific transcriptional regulators *Sohlh1* and *Lhx8*. *PNAS* **103** 8090–8095. (doi:10.1073/pnas.0601083103)
- Pangas SA, Li X, Umans L, Zwijzen A, Huylebroeck D, Gutierrez C, Wang D, Martin JF, Jamin SP, Behringer RR *et al.* 2008 Conditional deletion of *Smad1* and *Smad5* in somatic cells of male and female gonads leads to metastatic tumor development in mice. *Molecular and Cellular Biology* **28** 248–257. (doi:10.1128/MCB.01404-07)
- Pannetier M, Fabre S, Batista F, Kocer A, Renault L, Jolivet G, Mandon-Pepin B, Cotinot C, Veitia R & Pailhoux E 2006 FOXL2 activates P450 aromatase gene transcription: towards a better characterization of the early steps of mammalian ovarian development. *Journal of Molecular Endocrinology* **36** 399–413. (doi:10.1677/jme.1.01947)
- Paredes A, Romero C, Dissen GA, DeChiara TM, Reichardt L, Cornea A, Ojeda SR & Xu B 2004 TrkB receptors are required for follicular growth and oocyte survival in the mammalian ovary. *Developmental Biology* **267** 430–449. (doi:10.1016/j.ydbio.2003.12.001)
- Park M, Shin E, Won M, Kim J-H, Go H, Kim H-L, Ko J-J, Lee K & Bae J 2010 FOXL2 interacts with steroidogenic factor-1 (SF-1) and represses SF-1-induced *CYP17* transcription in granulosa cells. *Molecular Endocrinology* **24** 1024–1036. (doi:10.1210/me.2009-0375)
- Parrott JA & Skinner MK 1999 Kit-ligand/stem cell factor induces primordial follicle development and initiates folliculogenesis. *Endocrinology* **140** 4262–4271. (doi:10.1210/en.140.9.4262)
- Pisarska MD, Bae J, Klein C & Hsueh AJW 2004 Forkhead L2 is expressed in the ovary and represses the promoter activity of the steroidogenic acute regulatory gene. *Endocrinology* **145** 3424–3433. (doi:10.1210/en.2003-1141)
- Pisarska MD, Kuo F-T, Bentsi-Barnes IK, Khan S & Barlow GM 2010 LATS1 phosphorylates forkhead L2 and regulates its transcriptional activity. *American Journal of Physiology. Endocrinology and Metabolism* **299** E101–E109. (doi:10.1152/ajpendo.00534.2009)
- Pisarska MD, Barlow G & Kuo F-T 2011 Minireview: roles of the forkhead transcription factor FOXL2 in granulosa cell biology and pathology. *Endocrinology* **152** 1199–1208. (doi:10.1210/10.2010-1041)
- Poznic M 2009 Retinoblastoma protein: a central processing unit. *Journal of Biosciences* **34** 305–312. (doi:10.1007/s12038-009-0034-2)
- Qin Y, Choi Y, Zhao H, Simpson JL, Chen Z-J & Rajkovic A 2007 NOBOX homeobox mutation causes premature ovarian failure. *American Journal of Human Genetics* **81** 576–581. (doi:10.1086/519496)
- Rajareddy S, Reddy P, Du C, Liu L, Jagarlamudi K, Tang W, Shen Y, Berthet C, Peng SL, Kaldas P *et al.* 2007 p27^{Kip1} (cyclin-dependent kinase inhibitor 1B) controls ovarian development by suppressing follicle endowment and activation and promoting follicle atresia in mice. *Molecular Endocrinology* **21** 2189–2202. (doi:10.1210/me.2007-0172)
- Rajkovic A, Pangas SA, Ballow D, Suzumori N & Matzuk MM 2004 NOBOX deficiency disrupts early folliculogenesis and oocyte-specific gene expression. *Science* **305** 1157–1159. (doi:10.1126/science.1099755)
- Reddy P, Shen L, Ren C, Boman K, Lundin E, Ottander U, Lindgren P, Liu Y-X, Sun Q-Y & Liu K 2005 Activation of Akt (PKB) and suppression of FKHL1 in mouse and rat oocytes by stem cell factor during follicular activation and development. *Developmental Biology* **281** 160–170. (doi:10.1016/j.ydbio.2005.02.013)
- Reddy P, Liu L, Adhikari D, Jagarlamudi K, Rajareddy S, Shen Y, Du C, Tang W, Hämäläinen T, Peng SL *et al.* 2008 Oocyte-specific deletion of *Pten* causes premature activation of the primordial follicle pool. *Science* **319** 611–613. (doi:10.1126/science.1152257)
- Richards JS, Fan H-Y, Liu Z, Tsoi M, Laguè M-N, Boyer A & Boerboom D 2012 Either *Kras* activation or *Pten* loss similarly enhance the dominant-stable CTNNB1-induced genetic program to promote granulosa cell tumor development in the ovary and testis. *Oncogene* **31** 1504–1520. (doi:10.1038/onc.2011.341)
- Risma KA, Clay CM, Nett TM, Wagner T, Yun J & Nilson JH 1995 Targeted overexpression of luteinizing hormone in transgenic mice leads to infertility, polycystic ovaries, and ovarian tumors. *PNAS* **92** 1322–1326. (doi:10.1073/pnas.92.5.1322)
- Rosario R, Araki H, Print CG & Shelling AN 2012 The transcriptional targets of mutant FOXL2 in granulosa cell tumours. *PLoS ONE* **7** e46270. (doi:10.1371/journal.pone.0046270)
- Rosario R, Blenkiron C & Shelling AN 2013 Comparative study of microRNA regulation on FOXL2 between adult-type and juvenile-type granulosa cell tumours *in vitro*. *Gynecologic Oncology* **129** 209–215. (doi:10.1016/j.ygyno.2012.12.034)
- Schmidt D, Ovitt CE, Anlag K, Fehsenfeld S, Gredsted L, Treier A-C & Treier M 2004 The murine winged-helix transcription factor Foxl2 is required for granulosa cell differentiation and ovary maintenance. *Development* **131** 933–942. (doi:10.1242/dev.00969)
- Shah SP, Köbel M, Senz J, Morin RD, Clarke BA, Wiegand KC, Leung G, Zayed A, Mehl E, Kalloger SE *et al.* 2009 Mutation of FOXL2 in granulosa-cell tumors of the ovary. *New England Journal of Medicine* **360** 2719–2729. (doi:10.1056/NEJMoa0902542)
- Steinmetz R, Wagoner HA, Zeng P, Hammond JR, Hannon TS, Meyers JL & Pescovitz OH 2004 Mechanisms regulating the constitutive activation of the extracellular signal-regulated kinase (ERK) signaling pathway in ovarian cancer and the effect of ribonucleic acid interference for ERK1/2 on cancer cell proliferation. *Molecular Endocrinology* **18** 2570–2582. (doi:10.1210/me.2004-0082)
- St John MA, Tao W, Fei X, Fukumoto R, Carcangiu ML, Brownstein DG, Parlow AF, McGrath J & Xu T 1999 Mice deficient of *Lats1* develop soft-tissue sarcomas, ovarian tumours and pituitary dysfunction. *Nature Genetics* **21** 182–186. (doi:10.1038/5965)
- Tavassoli FA & Devilee P (Eds) 2003 *World Health Organization Classification of Tumours. Pathology and Genetics of Tumours of the Breast and Female Genital Organs*. Lyon: IARC Press.
- Todeschini A-L, Dipietromaria A, L'Hôte D, Boucham FZ, Georges AB, Pandaranayaka PJE, Krishnaswamy S, Rivals I, Bazin C & Veitia RA 2011 Mutational probing of the forkhead domain of the transcription factor FOXL2 provides insights into the pathogenicity of naturally occurring mutations. *Human Molecular Genetics* **20** 3376–3385. (doi:10.1093/hmg/ddr244)
- Tran S, Lamba P, Wang Y & Bernard DJ 2011 SMADs and FOXL2 synergistically regulate murine FSH β transcription via a conserved proximal promoter element. *Molecular Endocrinology* **25** 1170–1183. (doi:10.1210/me.2010-0480)
- Tran S, Zhou X, Lafleur C, Calderon MJ, Ellsworth BS, Kimmins S, Boehm U, Treier M, Boerboom D & Bernard DJ 2013 Impaired fertility and FSH synthesis in gonadotrope-specific *Foxl2* knockout mice. *Molecular Endocrinology* **27** 407–421. (doi:10.1210/me.2012-1286)
- Uda M, Ottolenghi C, Crisponi L, Garcia JE, Deiana M, Kimber W, Forabosco A, Cao A, Schlessinger D & Pilia G 2004 *Foxl2* disruption causes mouse ovarian failure by pervasive blockage of follicle development. *Human Molecular Genetics* **13** 1171–1181. (doi:10.1093/hmg/ddh124)
- Uhlenhaut NH, Jakob S, Anlag K, Eisenberger T, Sekido R, Kress J, Treier A-C, Klugmann C, Klasen C, Holter NI *et al.* 2009 Somatic sex reprogramming of adult ovaries to testes by FOXL2 ablation. *Cell* **139** 1130–1142. (doi:10.1016/j.cell.2009.11.021)
- Vainio S, Heikkilä M, Kispert A, Chin N & McMahon AP 1999 Female development in mammals is regulated by Wnt-4 signalling. *Nature* **397** 405–409. (doi:10.1038/17068)
- Van der Vos KE & Coffey PJ 2008 FOXO-binding partners: it takes two to tango. *Oncogene* **27** 2289–2299. (doi:10.1038/onc.2008.22)
- Verdin H & De Baere E 2012 FOXL2 impairment in human disease. *Hormone Research in Paediatrics* **77** 2–11. (doi:10.1159/000335236)
- Vidal VP, Chaboissier MC, de Rooij DG & Schedl A 2001 *Sox9* induces testis development in XX transgenic mice. *Nature Genetics* **28** 216–217. (doi:10.1038/90046)
- Watson RH, Roy WJ Jr, Davis M, Hitchcock A & Campbell IG 1997 Loss of heterozygosity at the α -inhibin locus on chromosome 2q is not a feature of human granulosa cell tumors. *Gynecologic Oncology* **65** 387–390. (doi:10.1006/gyngo.1997.4682)

- Woods DC, White YAR, Niikura Y, Kiatpongsan S, Lee H-J & Tilly JL 2013 Embryonic stem cell-derived granulosa cells participate in ovarian follicle formation *in vitro* and *in vivo*. *Reproductive Sciences* **20** 524–535. (doi:10.1177/1933719113483017)
- Yao HH-C, Whoriskey W & Capel B 2002 Desert Hedgehog/Patched 1 signaling specifies fetal Leydig cell fate in testis organogenesis. *Genes and Development* **16** 1433–1440. (doi:10.1101/gad.981202)
- Yao HHC, Matzuk MM, Jorgez CJ, Menke DB, Page DC, Swain A & Capel B 2004 *Follistatin* operates downstream of *Wnt4* in mammalian ovary organogenesis. *Developmental Dynamics* **230** 210–215. (doi:10.1002/dvdy.20042)
- Yoshida H, Takakura N, Kataoka H, Kunisada T, Okamura H & Nishikawa SI 1997 Stepwise requirement of c-kit tyrosine kinase in mouse ovarian follicle development. *Developmental Biology* **184** 122–137. (doi:10.1006/dbio.1997.8503)
- Zaret KS & Carroll JS 2011 Pioneer transcription factors: establishing competence for gene expression. *Genes and Development* **25** 2227–2241. (doi:10.1101/gad.176826.111)

Received in final form 26 August 2013

Accepted 18 September 2013

Accepted Preprint published online 18 September 2013