

RESEARCH

Open Access



Fractional differential equations for the generalized Mittag-Leffler function

Praveen Agarwal^{1,2}, Qasem Al-Mdallal^{3*}, Yeol Je Cho^{4,5} and Shilpi Jain⁶

*Correspondence:
q.almdallal@uaeu.ac.ae
³Department of Mathematical Sciences, UAE University, Al Ain, United Arab Emirates
Full list of author information is available at the end of the article

Abstract

In this paper, we establish some (presumably new) differential equation formulas for the extended Mittag-Leffler-type function by using the Saigo-Maeda fractional differential operators involving the Appell function $F_3(\cdot)$ and results in terms of the Wright generalized hypergeometric-type function ${}_{m+1}\Psi_{n+1}^{(\{\kappa_i\}_{i \in \mathbb{N}_0})}(z; \rho)$ recently established by Agarwal. Some interesting special cases are also pointed out.

MSC: Primary 26A33; 33E12; 33C05; 33C15; 33C20; secondary 33C65; 33C90

Keywords: generalized Gamma function; generalized beta functions; generalized Mittag-Leffler function; generalized Wright hypergeometric function; fractional derivative operators

1 Introduction and preliminaries

Fractional calculus (derivative and integrals) is very old as the conventional calculus and has been recently applied in various areas of engineering, science, finance, applied mathematics, and bio engineering (see, e.g., [1, 2]). Many differential equations involving various special functions have found significant importance and applications in various subfields of mathematical analysis. During the last few decades, a number of workers have studied, in depth, the properties, applications, and different extensions of various hypergeometric operators of fractional derivatives. A detailed account of such operators along with their properties and applications have been considered by several authors (see [3–22] and [23]).

A useful generalization of the hypergeometric fractional derivatives, including the Saigo operators [15–17], has been introduced by Marichev [13] as Mellin-type convolution operators with a special function $F_3(\cdot)$ in the kernel (for more details, see Samko et al. [20, p. 194, Eq. (10.47) and Section 10.3]) and later extended and studied by Saigo and Maeda [18, p. 393, Eqs. (4.12) and (4.13)]. Note that the generalized fractional derivative operators (for Saigo-Maeda operators, see [18]) are defined as follows:

$$\begin{aligned} (D_{0+}^{\tau, \tau', \nu, \nu', \sigma} f)(x) &= (I_{0+}^{-\tau', -\tau, -\nu', -\nu, -\sigma} f)(x) \\ &= \left(\frac{d}{dx}\right)^k (I_{0+}^{-\tau', -\tau, -\nu'+k, -\nu, -\sigma+k} f)(x), \end{aligned} \quad (1.1)$$

$$\begin{aligned}
 (D_-^{\tau, \tau', \nu, \nu', \sigma} f)(x) &= (I_-^{-\tau', -\tau, -\nu', -\nu, -\sigma} f)(x) \\
 &= \left(-\frac{d}{dx}\right)^k (I_{0-}^{-\tau', -\tau, -\nu'+k, -\nu, -\sigma+k} f)(x),
 \end{aligned}
 \tag{1.2}$$

where $\Re(\sigma) > 0, k = [\Re(\sigma)] + 1, \tau, \tau', \nu, \nu', \sigma \in \mathbb{C}, \mathbb{C}$ being the set of complex numbers, and $x > 0$.

Following Saigo et al. [18] and Saxena and Saigo [21], the left-hand sided and right-hand sided generalized differentiation for a power function are, respectively, given as follows (see, [24, p. 7, Eqs. (4.1) and (4.2)]):

$$(D_{0+}^{\tau, \tau', \nu, \nu', \sigma} t^{\varrho-1})(x) = \frac{\Gamma(\varrho)\Gamma(\varrho - \sigma + \tau + \tau' + \nu')\Gamma(\varrho + \tau - \nu)}{\Gamma(\varrho - \nu)\Gamma(\varrho - \sigma + \tau + \tau')\Gamma(\varrho - \sigma + \tau + \nu')} x^{\tau+\tau'+\varrho-\sigma-1},
 \tag{1.3}$$

where $\Re(\varrho) > \max\{0, \Re(-\tau + \nu), \Re(-\tau - \tau' - \nu' - \sigma)\}$, and

$$\begin{aligned}
 (D_{0-}^{\tau, \tau', \nu, \nu', \sigma} t^{-\rho})(x) &= \frac{\Gamma(\varrho + \sigma - \tau - \tau')\Gamma(\varrho - \tau' - \nu + \sigma)\Gamma(\varrho + \nu')}{\Gamma(\varrho)\Gamma(\varrho - \tau - \tau' - \nu + \sigma)\Gamma(\varrho - \tau' + \nu')} x^{-\varrho+\tau+\tau'-\sigma},
 \end{aligned}
 \tag{1.4}$$

where $\Re(\varrho) > \max\{\Re(-\nu'), \Re(\tau + \tau' - \sigma), \Re(\tau' + \nu - \sigma) + \Re(\sigma) + 1\}$.

In certain areas of applied mathematics and mathematical physics, special functions and their generalizations are used for finding solutions of the initial or boundary value problems for partial differential equations and fractional differential equations. It is also important to mention here that the special and degenerated cases of hypergeometric functions; in particular, the Bessel, Mittag-Leffler, and Wright hypergeometric functions have an importance due to application point of view.

Recently, Parmar introduced the extended Mittag-Leffler type function in the following form [25, p. 1072, Eq. (16)]:

$$\begin{aligned}
 E_{\xi, \zeta}^{(\{\kappa_l\}_{l \in \mathbb{N}_0}; \lambda)}(z; p) &= \sum_{n=0}^{\infty} \frac{\mathcal{B}^{(\{\kappa_l\}_{l \in \mathbb{N}_0})}(\lambda + n, 1 - \lambda; p)}{B(\lambda, 1 - \lambda)} \frac{z^n}{\Gamma(\xi n + \zeta)} \\
 (z, \zeta, \lambda \in \mathbb{C}, \Re(\xi) > 0, \Re(\zeta) > 0, \Re(\lambda) > 1; p \geq 0),
 \end{aligned}
 \tag{1.5}$$

where $\mathcal{B}^{(\{\kappa_l\}_{l \in \mathbb{N}_0})}(\lambda + n, 1 - \lambda; p)$ is the extended beta function defined by [26]

$$\begin{aligned}
 \mathcal{B}^{(\{\kappa_l\}_{l \in \mathbb{N}_0})}(\alpha, \beta; p) &= \int_0^1 t^{\alpha-1} (1-t)^{\beta-1} \Theta\left(\{\kappa_l\}_{l \in \mathbb{N}_0}; -\frac{p}{t(1-t)}\right) dt \\
 (\min\{\Re(\alpha), \Re(\beta)\} > 0; \Re(p) \geq 0),
 \end{aligned}
 \tag{1.6}$$

and $\Theta(\{\kappa_l\}_{l \in \mathbb{N}_0}; z)$ is a function of an appropriately bounded sequence $\{\kappa_l\}_{l \in \mathbb{N}_0}$ of arbitrary real or complex numbers defined as follows [26, p. 243, Eq. (2.1)]:

$$\begin{aligned}
 \Theta(\{\kappa_l\}_{l \in \mathbb{N}_0}; z) &:= \begin{cases} \sum_{l=0}^{\infty} \{\kappa_l\}_{l \in \mathbb{N}_0} \frac{z^l}{l!}, & |z| < R; 0 < R < \infty; \kappa_0 := 1, \\ \mathcal{M}_0 z^\omega \exp(z) [1 + O(\frac{1}{z})], & \Re(z) \rightarrow \infty; \mathcal{M}_0 > 0; \omega \in \mathbb{C}. \end{cases}
 \end{aligned}
 \tag{1.7}$$

It can be easily seen that different selection of the sequence $\{\kappa_l\}_{l \in \mathbb{N}_0}$ would generate particular cases of (1.5) as explained in the following examples.

Example 1 If we set $\kappa_l = \frac{(\rho)_l}{(\sigma)_l}$ ($l \in \mathbb{N}_0$), then (1.5) results in to the extended generalized Mittag-Leffler function [25, p. 1072, Eq. (17)]

$$E_{\xi, \zeta}^{(\rho, \sigma); \lambda}(z; p) = \sum_{n=0}^{\infty} \frac{\mathcal{B}^{(\rho, \sigma)}(\lambda + n, 1 - \lambda; p)}{B(\lambda, 1 - \lambda)} \frac{z^n}{\Gamma(\xi n + \zeta)}$$

$$(z, \zeta, \lambda \in \mathbb{C}, \Re(\rho) > 0, \Re(\sigma) > 0, \Re(\xi) > 0, \Re(\zeta) > 0, \Re(\lambda) > 1; p \geq 0). \tag{1.8}$$

Example 2 Setting $\kappa_l = 1$ ($l \in \mathbb{N}$) in (1.5) (see [27] with $c = 1$), we get the function

$$E_{\xi, \zeta}^{\lambda}(z; p) = \sum_{n=0}^{\infty} \frac{B(\lambda + n, 1 - \lambda; p)}{B(\lambda, 1 - \lambda)} \frac{z^n}{\Gamma(\xi n + \zeta)}$$

$$(z, \zeta, \lambda \in \mathbb{C}, \Re(\xi) > 0, \Re(\zeta) > 0, \Re(\lambda) > 1; p \geq 0). \tag{1.9}$$

Example 3 Similarly, for $p = 0$, (1.9) immediately reduces to the Prabhakar-type [28] Mittag-Leffler function.

Example 4 For $\xi = \zeta = 1$, (1.5), (1.8), and (1.9) can be expressed, respectively, in terms of the extended confluent hypergeometric functions as follows:

$$E_{1,1}^{(\{\kappa_l\}_{l \in \mathbb{N}_0}; \lambda)}(z; p) = \Phi_p^{(\{\kappa_l\}_{l \in \mathbb{N}_0})}(\lambda; 1; z), \tag{1.10}$$

$$E_{1,1}^{(\rho, \sigma); \lambda}(z; p) = \Phi_p^{(\rho, \sigma)}(\lambda; 1; z) \tag{1.11}$$

and

$$E_{1,1}^{\lambda}(z; p) = \Phi_p(\lambda; 1; z). \tag{1.12}$$

Similarly, by (1.6) and (1.7) of the extended beta function $\mathcal{B}^{(\{\kappa_l\}_{l \in \mathbb{N}_0})}(\lambda + n, 1 - \lambda; p)$ and a function of an appropriately bounded sequence $\{\kappa_l\}_{l \in \mathbb{N}_0}$ of arbitrary real or complex numbers $\Theta(\{\kappa_l\}_{l \in \mathbb{N}_0}; z)$, Agarwal [29] introduced and studied a further potentially useful extension of the Wright hypergeometric function as follows:

$$\begin{aligned} & {}_{m+1}\psi_{n+1}^{(\{\kappa_l\}_{l \in \mathbb{N}_0})}(z; p) \\ &= {}_{m+1}\psi_{n+1}^{(\{\kappa_l\}_{l \in \mathbb{N}_0})} \left[\begin{matrix} (a_i, \alpha_i)_{1,m}, & (\gamma, 1) \\ (b_j, \beta_j)_{1,n}, & (c, 1) \end{matrix} \middle| (z; p) \right] \\ &= \frac{1}{\Gamma(c - \gamma)} \sum_{k=0}^{\infty} \frac{\prod_{i=1}^m \Gamma(a_i + k\alpha_i)}{\prod_{j=1}^n \Gamma(b_j + k\beta_j)} \frac{\mathcal{B}^{(\{\kappa_l\}_{l \in \mathbb{N}_0})}(\gamma + k, c - \gamma; p) z^k}{k!} \\ & (z, \gamma \in \mathbb{C}, \Re(c) > \Re(\gamma) > 0; p \geq 0). \end{aligned} \tag{1.13}$$

In this paper, we aim to establish certain (presumably) new fractional differential equation formulas involving the extended generalized Mittag-Leffler type function (1.9) and

extended Wright Generalized hypergeometric function (1.13) by using the fractional differential operators (1.1) and (1.2), respectively. Some particular cases of our main findings are also pointed out.

2 Main results

In this section, we establish the fractional differential formulas involving the Saigo-Meada fractional derivative operators (1.1) and (1.2). These formulas are given in the following theorems.

Theorem 1 *Let $x > 0, \tau, \tau', \nu, \nu', \sigma, \mu, \xi, \zeta, \lambda, t, p \in \mathbb{C}$ ($\Re(\xi) > 0, \Re(\zeta) > 0, \Re(p) \geq 0$) be such that*

$$\Re(\mu) > \max\{0, \Re(-\tau + \nu), \Re(-\tau - \tau' - \nu' - \sigma)\}. \tag{2.1}$$

Then the following formula holds:

$$\begin{aligned} & (D_{0+}^{\tau, \tau', \nu, \nu', \sigma} t^{\mu-1} E_{\xi, \zeta}^{((k_l)_{l \in \mathbb{N}_0}; \lambda)}(t; p))(x) \\ &= \frac{x^{\mu+\tau+\tau'-\sigma-1}}{\Gamma(\lambda)} \\ & \times {}_4\psi_4^{((k_l)_{l \in \mathbb{N}_0})} \left[\begin{matrix} (\mu, 1), & (\mu - \nu + \tau, 1), & (\mu + \tau + \tau' + \nu' - \sigma, 1), & (\gamma, 1) \\ (\mu - \nu, 1), & (\mu + \tau + \tau' - \sigma, 1), & (\mu - \sigma + \tau + \nu', 1), & (\zeta, \xi) \end{matrix} \middle| (x; p) \right]. \end{aligned} \tag{2.2}$$

Proof Let \mathcal{I} be the left-hand side of (2.2). Using (1.5) and changing the order of integration and summation, which is verified under the conditions of the theorem, we have

$$\begin{aligned} \mathcal{I} &= \sum_{n=0}^{\infty} \frac{\mathcal{B}^{((k_l)_{l \in \mathbb{N}_0})}(\lambda + n, 1 - \lambda; p)}{B(\lambda, 1 - \lambda)} \frac{1}{\Gamma(\xi n + \zeta)} \\ & \times (D_{0+}^{\tau, \tau', \nu, \nu', \sigma} t^{\mu+n-1})(x). \end{aligned} \tag{2.3}$$

Applying (1.1) and (1.3) with ρ replaced by $\eta + n$ yields

$$\begin{aligned} \mathcal{I} &= \sum_{n=0}^{\infty} \frac{\mathcal{B}^{((k_l)_{l \in \mathbb{N}_0})}(\lambda + n, 1 - \lambda; p)}{B(\lambda, 1 - \lambda)} \frac{1}{\Gamma(\xi n + \zeta)} \\ & \times \frac{\Gamma(\mu + n)\Gamma(\mu + n + \tau - \nu)\Gamma(\mu + n + \tau + \tau' + \nu' - \sigma)}{\Gamma(\mu + n - \nu)\Gamma(\mu + n - \sigma + \tau + \tau')\Gamma(\mu + n - \sigma + \tau + \nu')} x^{\mu+n+\tau+\tau'-\sigma-1}, \end{aligned} \tag{2.4}$$

which, in view of (1.13), is equal to the right-hand side of (2.2). This completes the proof. \square

Theorem 2 *Let $x > 0, \tau, \tau', \nu, \nu', \sigma, \mu, \xi, \zeta, \lambda, t, p \in \mathbb{C}$ ($\Re(\xi) > 0, \Re(\zeta) > 0, \Re(\lambda) > 1, \Re(p) \geq 0$) be such that*

$$\Re(\rho) > \max\{\Re(-\nu'), \Re(\tau + \tau' - \sigma), \Re(\tau' + \nu - \sigma) + \Re(\sigma)\}. \tag{2.5}$$

Then the following formula holds:

$$\begin{aligned}
 & (D_{-}^{\tau, \tau', v, v', \sigma} t^{-\mu} E_{\xi, \zeta}^{(\{\kappa_l\}_{l \in \mathbb{N}_0}; \lambda)}(1/t; p))(x) \\
 &= \frac{x^{-\mu + \tau + \tau' - \sigma}}{\Gamma(\lambda)} \\
 & \times {}_4\psi_4^{(\{\kappa_l\}_{l \in \mathbb{N}_0})} \left[\begin{matrix} (\mu + v', 1), & (\mu - v - \tau' + \sigma, 1), & (\mu - \tau - \tau' + \sigma, 1), & (\gamma, 1) \\ (\mu, 1), & (\mu - \tau - \tau' - v + \sigma, 1), & (\mu - \tau' + v, 1), & (\zeta, \xi) \end{matrix} \middle| (x^{-1}; p) \right]. \tag{2.6}
 \end{aligned}$$

Proof We establish the result by a similar argument as in the proof of Theorem 1 using (1.4) instead of (1.3). Therefore, we omit the details. \square

3 Special cases

The results in Theorems 1 and 2 can be easily specialized to yield the corresponding formulas involving simpler functions like Mittag-Leffler-type functions and extended confluent hypergeometric functions given by (1.8)-(1.12) after appropriate selection of the sequence $\{\kappa_l\}_{l \in \mathbb{N}_0}$.

Setting $\kappa_l = \frac{(\phi)_l}{(\varphi)_l}$ ($l \in \mathbb{N}_0$), we obtain the following results from Theorems 1 and 2, respectively.

Corollary 1 *Let $x > 0, \tau, \tau', v, v', \sigma, \mu, \xi, \zeta, \lambda, \phi, \varphi, t, p \in \mathbb{C}$ ($\Re(\xi) > 0, \Re(\zeta) > 0, \Re(p) \geq 0$) be such that*

$$\Re(\mu) > \max\{0, \Re(-\tau + v), \Re(-\tau - \tau' - v' - \sigma)\}. \tag{3.1}$$

Then the following formula holds:

$$\begin{aligned}
 & (D_{0+}^{\tau, \tau', v, v', \sigma} t^{\mu-1} E_{\xi, \zeta}^{(\phi, \varphi); \lambda}(t; p))(x) \\
 &= \frac{x^{\mu + \tau + \tau' - \sigma - 1} \Gamma(\varphi)}{\Gamma(\lambda) \Gamma(\phi)} \\
 & \times {}_5\psi_5 \left[\begin{matrix} (\mu, 1), & (\mu - v + \tau, 1), & (\mu + \tau + \tau' + v' - \sigma, 1), & (\gamma, 1), & (\phi, 1) \\ (\mu - v, 1), & (\mu + \tau + \tau' - \sigma, 1), & (\mu - \sigma + \tau + v', 1), & (\zeta, \xi), & (\varphi, 1) \end{matrix} \middle| (x; p) \right]. \tag{3.2}
 \end{aligned}$$

Corollary 2 *Let $x > 0, \tau, \tau', v, v', \sigma, \mu, \xi, \zeta, \lambda, \phi, \varphi, t, p \in \mathbb{C}$ ($\Re(\xi) > 0, \Re(\zeta) > 0, \Re(\lambda) > 1, \Re(p) \geq 0$) be such that*

$$\Re(\varrho) > \max\{\Re(-v'), \Re(\tau + \tau' - \sigma), \Re(\tau' + v - \sigma) + \Re(\sigma)\}. \tag{3.3}$$

Then the following formula holds:

$$\begin{aligned}
 & (D_{-}^{\tau, \tau', v, v', \sigma} t^{-\mu} E_{\xi, \zeta}^{(\phi, \varphi); \lambda}(1/t; p))(x) \\
 &= \frac{x^{-\mu + \tau + \tau' - \sigma} \Gamma(\varphi)}{\Gamma(\lambda) \Gamma(\phi)} \\
 & \times {}_5\psi_5 \left[\begin{matrix} (\mu + v', 1), & (\mu - v - \tau' + \sigma, 1), & (\mu - \tau - \tau' + \sigma, 1), & (\gamma, 1), & (\phi, 1) \\ (\mu, 1), & (\mu - \tau - \tau' - v + \sigma, 1), & (\mu - \tau' + v, 1), & (\zeta, \xi), & (\varphi, 1) \end{matrix} \middle| (x^{-1}; p) \right]. \tag{3.4}
 \end{aligned}$$

For $\kappa_l = 1$, Theorems 1 and 2 become as follows.

Corollary 3 Let $x > 0, \tau, \tau', v, v', \sigma, \mu, \xi, \zeta, \lambda, t, p \in \mathbb{C}$ ($\Re(\xi) > 0, \Re(\zeta) > 0, \Re(p) \geq 0$) be such that

$$\Re(\mu) > \max\{0, \Re(-\tau + v), \Re(-\tau - \tau' - v' - \sigma)\}. \tag{3.5}$$

Then the following formula holds:

$$\begin{aligned} & (D_{0+}^{\tau, \tau', v, v', \sigma} t^{\mu-1} E_{\xi, \zeta}^{\lambda}(t; p))(x) \\ &= \frac{x^{\mu+\tau+\tau'-\sigma-1}}{\Gamma(\lambda)} \\ & \times {}_4\psi_4 \left[\begin{matrix} (\mu, 1), & (\mu - v + \tau, 1), & (\mu + \tau + \tau' + v' - \sigma, 1), & (\gamma, 1) \\ (\mu - v, 1), & (\mu + \tau + \tau' - \sigma, 1), & (\mu - \sigma + \tau + v', 1), & (\zeta, \xi) \end{matrix} \middle| (x; p) \right]. \end{aligned} \tag{3.6}$$

Corollary 4 Let $x > 0, \tau, \tau', v, v', \sigma, \mu, \xi, \zeta, \lambda, t, p \in \mathbb{C}$ ($\Re(\xi) > 0, \Re(\zeta) > 0, \Re(\lambda) > 1, \Re(p) \geq 0$) be such that

$$\Re(\rho) > \max\{\Re(-v'), \Re(\tau + \tau' - \sigma), \Re(\tau' + v - \sigma) + \Re(\sigma)\}. \tag{3.7}$$

Then the following formula holds:

$$\begin{aligned} & (D_{-}^{\tau, \tau', v, v', \sigma} t^{-\mu} E_{\xi, \zeta}^{\lambda}(1/t; p))(x) \\ &= \frac{x^{-\mu+\tau+\tau'-\sigma} \Gamma(\varphi)}{\Gamma(\lambda)\Gamma(\phi)} \\ & \times {}_4\psi_4 \left[\begin{matrix} (\mu + v', 1), & (\mu - v - \tau' + \sigma, 1), & (\mu - \tau - \tau' + \sigma, 1), & (\gamma, 1) \\ (\mu, 1), & (\mu - \tau - \tau' - v + \sigma, 1), & (\mu - \tau' + v, 1), & (\zeta, \xi) \end{matrix} \middle| (x^{-1}; p) \right]. \end{aligned} \tag{3.8}$$

Similarly, putting $p = 0$ in Corollaries 3 and 4, we get the fractional differential formulas involving the Prabhakar-type [28] Mittag-Leffler function. We omit the details.

Following the same way, setting $\xi = \zeta = 1$, from Theorems 1 and 2 and Corollaries 1 and 2 we obtain the following interesting results.

Corollary 5 Let $x > 0, \tau, \tau', v, v', \sigma, \mu, \lambda, \phi, \varphi, t, p \in \mathbb{C}$ ($\Re(\xi) > 0, \Re(\zeta) > 0, \Re(p) \geq 0$) be such that

$$\Re(\mu) > \max\{0, \Re(-\tau + v), \Re(-\tau - \tau' - v' - \sigma)\}. \tag{3.9}$$

Then the following formula holds:

$$\begin{aligned} & (D_{0+}^{\tau, \tau', v, v', \sigma} t^{\mu-1} \Phi_p^{((k_l)_{l \in \mathbb{N}_0})}(\lambda; 1; t))(x) \\ &= \frac{x^{\mu+\tau+\tau'-\sigma-1}}{\Gamma(\lambda)} \\ & \times {}_4\psi_4^{((k_l)_{l \in \mathbb{N}_0})} \left[\begin{matrix} (\mu, 1), & (\mu - v + \tau, 1), & (\mu + \tau + \tau' + v' - \sigma, 1), & (\gamma, 1) \\ (\mu - v, 1), & (\mu + \tau + \tau' - \sigma, 1), & (\mu - \sigma + \tau + v', 1), & (1, 1) \end{matrix} \middle| (x; p) \right]. \end{aligned} \tag{3.10}$$

Corollary 6 Let $x > 0, \tau, \tau', v, v', \sigma, \mu, \lambda, t, p \in \mathbb{C}$ ($\Re(\xi) > 0, \Re(\zeta) > 0, \Re(\lambda) > 1, \Re(p) \geq 0$) be such that

$$\Re(\rho) > \max\{\Re(-v'), \Re(\tau + \tau' - \sigma), \Re(\tau' + v - \sigma) + \Re(\sigma)\}. \tag{3.11}$$

Then the following formula holds:

$$\begin{aligned} & \left(D_{-}^{\tau, \tau', \nu, \nu', \sigma} t^{-\mu} \Phi_p^{((k_l)_{l \in \mathbb{N}_0})} \left(\lambda; 1; \frac{1}{t} \right) \right) (x) \\ &= \frac{x^{-\mu + \tau + \tau' - \sigma} \Gamma(\varphi)}{\Gamma(\lambda) \Gamma(\phi)} \\ & \quad \times {}_4\psi_4 \left[\begin{matrix} (\mu + \nu', 1), & (\mu - \nu - \tau' + \sigma, 1), & (\mu - \tau - \tau' + \sigma, 1), & (\gamma, 1) \\ (\mu, 1), & (\mu - \tau - \tau' - \nu + \sigma, 1), & (\mu - \tau' + \nu, 1), & (1, 1) \end{matrix} \middle| (x^{-1}; p) \right]. \end{aligned} \tag{3.12}$$

Corollary 7 Let $x > 0, \tau, \tau', \nu, \nu', \sigma, \mu, \lambda, \phi, \varphi, t, p \in \mathbb{C} (\Re(\xi) > 0, \Re(\zeta) > 0, \Re(p) \geq 0)$ be such that

$$\Re(\mu) > \max \{ 0, \Re(-\tau + \nu), \Re(-\tau - \tau' - \nu' - \sigma) \}. \tag{3.13}$$

Then the following formula holds:

$$\begin{aligned} & \left(D_{0+}^{\tau, \tau', \nu, \nu', \sigma} \mu^{-1} \Phi_p^{(\phi, \varphi)}(\lambda; 1; t) \right) (x) \\ &= \frac{x^{\mu + \tau + \tau' - \sigma - 1} \Gamma(\varphi)}{\Gamma(\lambda) \Gamma(\phi)} \\ & \quad \times {}_5\psi_5 \left[\begin{matrix} (\mu, 1), & (\mu - \nu + \tau, 1), & (\mu + \tau + \tau' + \nu' - \sigma, 1), & (\gamma, 1), & (\phi, 1) \\ (\mu - \nu, 1), & (\mu + \tau + \tau' - \sigma, 1), & (\mu - \sigma + \tau + \nu', 1), & (1, 1), & (\varphi, 1) \end{matrix} \middle| (x; p) \right]. \end{aligned} \tag{3.14}$$

Corollary 8 Let $x > 0, \tau, \tau', \nu, \nu', \sigma, \mu, \lambda, \phi, \varphi, t, p \in \mathbb{C} (\Re(\xi) > 0, \Re(\zeta) > 0, \Re(\lambda) > 1, \Re(p) \geq 0)$ be such that

$$\Re(\varrho) > \max \{ \Re(-\nu'), \Re(\tau + \tau' - \sigma), \Re(\tau' + \nu - \sigma) + \Re(\sigma) \}. \tag{3.15}$$

Then the following formula holds:

$$\begin{aligned} & \left(D_{-}^{\tau, \tau', \nu, \nu', \sigma} t^{-\mu} \Phi_p^{((k_l)_{l \in \mathbb{N}_0})}(\lambda; 1; t) \right) (x) \\ &= \frac{x^{-\mu + \tau + \tau' - \sigma} \Gamma(\varphi)}{\Gamma(\lambda) \Gamma(\phi)} \\ & \quad \times {}_5\psi_5 \left[\begin{matrix} (\mu + \nu', 1), & (\mu - \nu - \tau' + \sigma, 1), & (\mu - \tau - \tau' + \sigma, 1), & (\gamma, 1), & (\phi, 1) \\ (\mu, 1), & (\mu - \tau - \tau' - \nu + \sigma, 1), & (\mu - \tau' + \nu, 1), & (1, 1), & (\varphi, 1) \end{matrix} \middle| (x^{-1}; p) \right]. \end{aligned} \tag{3.16}$$

4 Concluding remarks

In this study, we established some fractional differential formulas involving a family of Mittag-Leffler functions. Due to practical importance of the Mittag-Leffler functions, our results are of general character and hence encompass several cases of interest.

Acknowledgements

Authors would like to acknowledge and express their gratitude to the United Arab Emirates University, Al Ain, UAE for providing the financial support with Grant No. 31S240-UPAR (2) 2016.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors read and approved the final manuscript.

Author details

¹Department of Mathematics, Anand International College of Engineering, Jaipur, India. ²International Center for Basic and Applied Sciences, Jaipur, India. ³Department of Mathematical Sciences, UAE University, Al Ain, United Arab Emirates. ⁴School of Mathematical Sciences, University of Electronic Science and Technology of China, Chengdu, People's Republic of China. ⁵Department of Mathematics Education, Gyeongsang National University, Jinju, Korea. ⁶Department of Mathematics, Poornima College of Engineering, Jaipur, India.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 29 July 2017 Accepted: 19 January 2018 Published online: 15 February 2018

References

- Al-Mdallal, QM, Hajji, MA: A convergent algorithm for solving higher-order nonlinear fractional boundary value problems. *Fract. Calc. Appl. Anal.* **18**(6), 1423-1440 (2015)
- Al-Mdallal, QM, Omer, AS: Fractional-order Legendre-collocation method for solving fractional initial value problems. *Appl. Math. Comput.* **321**, 74-84 (2018)
- Agarwal, P: Some inequalities involving Hadamard-type k -fractional integral operators. *Math. Methods Appl. Sci.* **40**(11), 3882-3891 (2017)
- Agarwal, P: Certain properties of the generalized Gauss hypergeometric functions. *Appl. Math. Inf. Sci.* **8**(5), 2315-2320 (2014)
- Agarwal, P: Fractional integration of the product of two multivariables H -function and a general class of polynomials. In: *Advances in Applied Mathematics and Approximation Theory*, Springer Proceedings in Mathematics & Statistics, vol. 41, pp. 359-374 (2013)
- Agarwal, P: Further results on fractional calculus of Saigo operators. *Appl. Appl. Math.* **7**(2), 585-594 (2012)
- Agarwal, P: Generalized fractional integration of the H -function. *Le Matematiche* **LXVII**, 107-118 (2012)
- Agarwal, P, Jain, S: Further results on fractional calculus of Srivastava polynomials. *Bull. Math. Anal. Appl.* **3**(2), 167-174 (2011)
- Agarwal, P, Jain, S, Mansour, T: Further extended Caputo fractional derivative operator and its applications. *Russ. J. Math. Phys.* **24**(4), 415-425 (2017)
- Kim, DS, Kim, T, Mansour, T, Seo, J-J: Degenerate Mittag-Leffler polynomials. *Appl. Math. Comput.* **274**, 258-266 (2016)
- Kim, DS, Kim, T, Rim, S-H: Some identities involving Gegenbauer polynomials. *Adv. Differ. Equ.* **2012**, 219 (2012)
- Kim, T, Kim, DS, Dolgy, DV: Some identities on Bernoulli and Hermite polynomials associated with Jacobi polynomials. *Discrete Dyn. Nat. Soc.* **2012**, Article ID 584643 (2012)
- Marichev, Ol: Volterra equation of Mellin convolution type with a Horn function in the kernel. *Izv. AN BSSR Ser. Fiz.-Mat. Nauk* **1**, 128-129 (1974) (In Russian)
- Miller, KS, Ross, B: *An Introduction to the Fractional Calculus and Differential Equations*. Wiley, New York (1993)
- Saigo, M: A remark on integral operators involving the Gauss hypergeometric functions. *Math. Rep. Kyushu Univ.* **11**, 135-143 (1978)
- Saigo, M: A certain boundary value problem for the Euler-Darboux equation I. *Math. Jpn.* **24**(4), 377-385 (1979)
- Saigo, M: A certain boundary value problem for the Euler-Darboux equation II. *Math. Jpn.* **25**(2), 211-220 (1980)
- Saigo, M, Maeda, N: More generalization of fractional calculus. In: Rusev, P, Dimovski, I, Kiryakova, V (eds.) *Transform Methods and Special Functions*, Varna, 1996. Proc. 2nd Intern. Workshop, pp. 386-400. IMI-BAS, Sofia (1998)
- Saigo, M, Kilbas, AA: Generalized fractional calculus of the H -function. *Fukuoka Univ. Sci. Rep.* **29**, 31-45 (1999)
- Samko, SG, Kilbas, AA, Marichev, Ol: *Fractional Integrals and Derivatives. Theory and Applications*. Gordon & Breach Sci. Publ., New York (1993)
- Saxena, RK, Saigo, M: Generalized fractional calculus of the H -function associated with the Appell function. *J. Fract. Calc.* **19**, 89-104 (2001)
- Srivastava, HM, Tomovski, Z: Fractional calculus with an integral operator containing a generalized Mittag-Leffler function in the kernel. *Appl. Math. Comput.* **211**, 198-210 (2009)
- Ruzhansky, M, Cho, YJ, Agarwal, P, Area, I: *Advances in Real and Complex Analysis with Applications*. Birkhäuser, Basel (2017)
- Kataria, KK, Vellaisamy, P: The generalized k -Wright function and Marichev-Saigo-Maeda fractional operators. arXiv:1408.4762v1 [math.CA] (17 Aug 2014)
- Parmar, RK: A class of extended Mittag-Leffler functions and their properties related to integral transform and fractional calculus. *Matematiche* **3**, 1069-1082 (2015)
- Srivastava, HM, Parmar, RK, Chopra, P: A class of extended fractional derivative operators and associated generating relations involving hypergeometric functions. *Axioms* **1**, 238-258 (2012)
- Özarslan, MA, Yilmaz, B: The extended Mittag-Leffler function and its properties. *J. Inequal. Appl.* **2014**, 85 (2014)
- Prabhakar, TR: A singular integral equation with a generalized Mittag-Leffler function in the kernel. *Yokohama Math. J.* **19**, 7-15 (1971)
- Agarwal, P: Certain properties of extended Wright generalized hypergeometric type function. Submitted