

Pacific Journal of Mathematics

FUNCTION-THEORETIC DEGENERACY CRITERIA FOR RIEMANNIAN MANIFOLDS

MOSES GLASNER AND RICHARD EMANUEL KATZ

FUNCTION-THEORETIC DEGENERACY CRITERIA FOR RIEMANNIAN MANIFOLDS

MOSES GLASNER AND RICHARD KATZ

The modulus of a relatively compact set with border consisting of at least two components is a measure of its magnitude with regard to harmonic functions. A divergent modular sum associated with difference sets obtained from an exhaustion of a Riemannian manifold is characteristic of parabolicity. The existence of a divergent minimum modular sum implies that the manifold carries no nonconstant harmonic functions with finite Dirichlet integral.

The modular criteria presented in this paper are generalizations of criteria established for Riemann surfaces by Noshiro [6] and Sario [8], [9]. In the two-dimensional case function-theoretic degeneracy is completely determined by the conformal structure, whereas it has been shown by Nakai and Sario [5] that the type of a Riemannian manifold varies when the metric is replaced by a conformally equivalent one. The significance of our result stems from this fact.

For the completeness of the presentation it is shown that the various characterizations of parabolicity due to Ahlfors, Brelot, Nevanlinna and Ohtsuka remain equivalent in higher dimensions. This overlaps with the work of Itô [2] and Loeb [3] in different settings. A new proof for Riemannian manifolds of the relation, $O_{UD} = O_{UBD}$, established in [10] is also given.

1. Let R be an orientable noncompact Riemannian manifold. Let $A \subset R$ and denote by $H(A)$ the class of harmonic functions on A and by $H^c(A)$ the functions in $H(A)$ with continuous extensions to \bar{A} . For every parametric region V there exists a Green's function q_x^V with the property $-h(x) = \int_{\partial V} h^* dq_x^V$ for every $h \in H^c(V)$. It is well-known that the sheaf of harmonic functions over R satisfies the axioms of a harmonic space and we shall use the standard facts of the theory freely. These together with Green's formulas will serve as our main tools.

2. Consider a fixed parametric region $V \subset R$ and a point $a \in V$. Let F consist of the constant $+\infty$ and of all nonnegative superharmonic functions s on R such that $s|V - q_a^V$ is bounded. Clearly F is a Perron family on $R - a$ and its lower envelope is either $+\infty$ or a function g_a harmonic on $R - a$. If the function g_a exists it is called the *Green's function* for R .

Let $\{\Omega_n\}$ be an exhaustion of R by regular regions. Let $t_n \in H^c(\Omega_n - \bar{\Omega}_0)$ with $t_n|_{\partial\Omega_0} = 0, t_n|_{\partial\Omega_n} = 1$. The functions t_n form a decreasing sequence and the harmonic function $t = \lim_n t_n$ on $R - \bar{\Omega}_0$ is called the *harmonic measure of the ideal boundary of R with respect to Ω_0* .

3. The above definitions are related as follows.

THEOREM. *Conditions (a), (b) and (c) are equivalent.*

(a) *There exists a nonconstant positive superharmonic function on R .*

(b) *t is not identically zero.*

(c) *R does not belong to the class O_G of manifolds for which g_α does not exist.*

It is obvious that either (b) or (c) implies (a). To show that (a) implies (b), suppose that $t \equiv 0$ and let s be a positive superharmonic function. Set $m = \min_{\partial\Omega_0} s = \min_{\bar{\Omega}_0} s$ and observe that $s \geq m(1 - t_n)$ on $\bar{\Omega}_n - \bar{\Omega}_0$. On letting n tend to ∞ we conclude that $s \geq m$ on R and hence it is a constant. Now suppose that (a) is true. Then (b) holds for all choices of Ω_0 and in particular for the choice

$$\Omega_0 = \{x \mid q_\alpha^V(x) > 1\} \subset V.$$

Set $\alpha = (\min_{\partial V} t)^{-1}$. By virtue of the fact that $q_\alpha^V|_{\partial V} = 0$ we conclude that the function

$$q(x) = \begin{cases} q_\alpha^V(x), & x \in \Omega_0 \\ 1 - 2\alpha t(x), & x \in R - \Omega_0 \end{cases}$$

is superharmonic on R and bounded from below by $1 - 2\alpha$. Thus F contains more than one element and consequently $R \notin O_G$.

The *modulus* τ_n of the region $\Omega_n - \bar{\Omega}_0$ is by definition $(\int_{\partial\Omega_0} *dt_n)^{-1}$ and we can state the following immediate

COROLLARY. *The constants τ_n tend to $+\infty$ if and only if $R \in O_G$.*

4. In addition to the above notions we define the *modulus function* $\omega_i \in H^c(\Omega_i - \bar{\Omega}_{i-1})$ by the conditions $\omega_i|_{\partial\Omega_{i-1}} = 0, \omega_i|_{\partial\Omega_i} = \mu_i$ (const.) and $\int_{\partial\Omega_{i-1}} *d\omega_i = 1$. The constant μ_i is called the *modulus* of $\Omega_i - \bar{\Omega}_{i-1}$.

We now generalize the O_G modular criterion of Noshiro [6] and Sario [9] to Riemannian manifolds.

THEOREM. *There exists an exhaustion $\{\Omega_n\}$ of R such that $\sum \mu_n = +\infty$ if and only if $R \in O_G$.*

Suppose that such an exhaustion exists. The Dirichlet integral of $w_n = \tau_n t_n$ over $\Omega_n - \bar{\Omega}_0$ is

$$D_n(w_n) = \int_{\Omega_n - \bar{\Omega}_0} dw_n \wedge *dw_n = \int_{\partial\Omega_n - \partial\Omega_0} w_n^* dw_n = \tau_n .$$

On the other hand, $D_n(w_n)$ is equal to the sum $\sum_{i=1}^n D_i(w_n)$ of Dirichlet integrals over $\Omega_i - \bar{\Omega}_{i-1}$. Using the Schwarz inequality we obtain

$$D_i(w_n) \geq \frac{D_i^2(w_i, w_n)}{D_i(w_i)} = \frac{\left(\int_{\partial\Omega_i - \partial\Omega_{i-1}} \omega_i^* dw_n \right)^2}{\int_{\partial\Omega_i - \partial\Omega_{i-1}} \omega_i^* d\omega_i} = \frac{\mu_i^2}{\mu_i} .$$

Thus $\tau_n \geq \sum_{i=1}^n \mu_i$, and consequently $\tau_n \rightarrow +\infty$. This implies, by Corollary 3, that $R \in O_G$.

Conversely, suppose that $R \in O_G$ and $\{\Omega_n\}$ is any exhaustion of R . Then there exists an n_1 such that $\tau_{n_1} > 1$. Then $\{\tau'_n\}$ constructed for the exhaustion $\{\Omega_n\}_{n_1}^\infty$ also tend to $+\infty$ and a fortiori there is an n_2 with $\tau'_{n_2} > 1$. Proceeding in this fashion we obtain an exhaustion $\{\Omega_{n_i}\}$ such that the modular sum

$$\sum_{i=1}^\infty \mu_i = \sum_{i=1}^\infty \tau_{n_i}^{(i-1)} = +\infty .$$

The above condition is not necessary in the sense that every manifold has an exhaustion with arbitrarily modular sum (cf. [1]).

5. We now turn to a criterion for membership in the class O_{HD} of manifolds on which every member of the class $HD(R)$ of harmonic functions with finite Dirichlet integral over R is constant. The Dirichlet integral of an $f \in C^1(R)$ over R is, by definition,

$$D(f) = \lim_{n \rightarrow \infty} D_n(f) ,$$

where $D_n(f)$ is taken over Ω_n and $\{\Omega_n\}$ is an exhaustion of R . Clearly $D(f)$ is independent of the choice of the exhaustion $\{\Omega_n\}$ and

$$D(f) < +\infty, D(g) < +\infty$$

imply the existence of the limit $D(f, g) = \lim_{n \rightarrow \infty} D_n(f, g)$.

First we prove the

THEOREM. *There exists a nonconstant element of $HD(R)$ if and only if there exists one which is also bounded. That is, $O_{HD} = O_{HBD}$.*

Let u be a nonconstant element of $HD(R)$ and set

$$u_m = \min(m, \max(u, -m)) .$$

Denote by v_{m_n} the continuous function on R such that $v_{m_n} = u_m$ on $R - \Omega_n$ and v_{m_n} is harmonic on Ω_n . Since $\{v_{m_n}\}_{n=0}^\infty$ is uniformly bounded it has a subsequence also denoted by $\{v_{m_n}\}_{n=0}^\infty$ which converges to a harmonic function v_m uniformly on compact subsets. By Green's formula we obtain $D(u_m - v_{m_n}) = D(u_m) - D(v_{m_n}) \geq 0$ and

$$D(v_{m_n} - v_{m,n+p}) = D(v_{m_n}) - D(v_{m,n+p}) \geq 0 .$$

From this we see that $d = \lim_n D(v_{m_n})$ exists. By Fatou's lemma we also have $D(v_{m_n} - v_m) \leq D(v_{m_n}) - d$ and consequently $\lim_n D(v_{m_n} - v_m) = 0$.

For every m , $v_m \in HBD(R)$ and we shall complete proof by showing that the assumption v_m is a constant for every m leads to a contradiction. Set $g_{m_n} = u_m - v_{m_n}$, $g_n = \lim_n g_{m_n} = u_m - v_m$ and note that $\lim_n D(g_{m_n} - u_m) = 0$. Since $\text{supp } g_{m_n} \subset \bar{\Omega}_n$, we have

$$D(g_{m_n}, u) = \int_{\partial\Omega_n} g_{m_n} * du = 0 .$$

Thus $D(u_m, u) = \lim_n D(g_{m_n}, u) = 0$ and $D(u) = \lim_m D(u_m, u) = 0$, which contradicts the choice of u . Our proof stems in spirit from Royden [7].

6. Sario [8] introduced a sufficient condition for a Riemann surface not to carry nonconstant analytic functions with finite Dirichlet integral. Since analytic, as such, has no meaning in Riemannian manifolds we replace Sario's result with an O_{HD} criterion.

In general the open sets $\Omega_n - \bar{\Omega}_{n-1}$ will consist of components E_{i_n} , $i = 1, \dots, i(n)$. Let ω_{i_n} denote the modulus function of E_{i_n} , i.e., $\omega_{i_n} \in H^c(E_{i_n})$, $\omega_{i_n} | \partial E_{i_n} \cap \partial\Omega_{n-1} = 0$, $\omega_{i_n} | \partial E_{i_n} \cap \partial\Omega_n = \mu_{i_n}$ (const.) such that $\int *d\omega_{i_n} = 1$. Set $\nu_n = \min_i \mu_{i_n}$.

THEOREM. *If $\{\Omega_n\}$ is an exhaustion of R with a divergent minimum modular sum $\sum \nu_n = +\infty$, then $R \in O_{HD}$.*

Since Euclidean n -space ($n \geq 3$) is not in O_G but in O_{HD} , Theorem 4 shows that the divergence of the minimum modular sum is not a necessary condition for a manifold to be in O_{HD} .

For the proof of the theorem let u be a nonconstant harmonic function on R with $D(u) < +\infty$; by Theorem 5 we may assume that $|u| \leq M$. We further normalize u by adding a constant so that

$$\int_{\partial\Omega_0} u * du = 0 .$$

By a theorem of Morse (cf. Milnor [4]) ω_n and its gradient can be uniformly approximated by C^∞ functions h and their gradients, where the h 's have the same boundary values as ω_n and $\text{grad } h = 0$ only at

a finite number of points. We choose such an h with the property that

$$\int_{\beta_{n\lambda}} |\text{grad } h| \, dS \leq 2,$$

where $\beta_{n\lambda}$ is the level surface $h = \lambda$, $0 < \lambda < \mu_n$.

Denote by $D_{n\lambda}(u)$ the Dirichlet integral of u over the open set bounded by $\partial\Omega_0$ and $\beta_{n\lambda}$. By the definition of gradient and the Schwarz inequality we obtain

$$\frac{d}{d\lambda} D_{n\lambda}(u) = \int_{\beta_{n\lambda}} \frac{|\text{grad } u|^2}{|\text{grad } h|} \, dS \geq \frac{\left(\int_{\beta_{n\lambda}} |\text{grad } u| \, dS\right)^2}{\int_{\beta_{n\lambda}} |\text{grad } h| \, dS} \geq \frac{1}{2} \left(\int_{\beta_{n\lambda}} |*du|\right)^2,$$

except for a finite number of values of λ . On the other hand, Green's formula gives

$$D_{n\lambda}(u) = \int_{\beta_{n\lambda}} u * du \leq M \int_{\beta_{n\lambda}} |*du|.$$

We conclude that

$$\frac{d}{d\lambda} D_{n\lambda}(u) \geq \frac{1}{2M^2} D_{n\lambda}^2(u).$$

We now integrate this inequality for $n \geq 2$:

$$\int_0^{\nu_n} \frac{d}{d\lambda} \frac{D_{n\lambda}(u)}{D_{n\lambda}(u)} \, d\lambda \geq \frac{1}{2M^2} \int_0^{\nu_n} D_{n\lambda}(u) \, d\lambda \geq \nu_n a,$$

where $a = (2M^2)^{-1} D_1(u)$. This implies that

$$D_n(u) \geq D_{n,\nu_n}(u) \geq D_{n-1}(u) \exp(\nu_n a)$$

and iteration gives $D_n(u) \geq D_1(u) \exp(\alpha \sum_2^n \nu_n)$. Since $D(u) < +\infty$, we conclude that $\sum_1^\infty \nu_n < +\infty$.

REFERENCES

1. M. Glasner, R. Katz and M. Nakai, *Bisection into small annuli*, Pacific J. Math. **24** (1968), 457-461.
2. S. Itô, *On existence of Green function and positive superharmonic functions for linear elliptic operators of second order*, J. Math. Soc. Japan **16** (1964), 299-306.
3. P. A. Loeb, *An axiomatic treatment of pairs of elliptic differential operators*, Ann. Inst. Fourier Grenoble **16** (1966), 167-208.
4. J. Milnor, *Lectures on the h-cobordism theorem*, Princeton Univ. Press, Princeton, N. J., 1965.
5. M. Nakai and L. Sario, *Classification and deformation of Riemannian spaces*, Math. Scand. **20** (1967), 193-208.

6. K. Noshiro, *Open Riemann surface with null boundary*, Nagoya Math. J. **3** (1951), 73-79.
7. H. L. Royden, *Harmonic functions on open Riemann surfaces*, Trans. Amer. Math. Soc. **73** (1952), 40-94.
8. L. Sario, *Riemannsche Flächen mit hebbarem Rand*, Ann. Acad. Sci. Fenn. Ser. A. I. **50** (1948).
9. ———, *Modular criteria on Riemann surfaces*, Duke Math. J. **20** (1953), 279-286.
10. L. Sario, M. Schiffer and M. Glasner, *The span and principal functions in Riemannian spaces*, J. Analyse Math. **15** (1965), 115-134.

Received February 13, 1968.

CALIFORNIA INSTITUTE OF TECHNOLOGY
PASADENA, CALIFORNIA
CALIFORNIA STATE COLLEGE
LOS ANGELES, CALIFORNIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. R. PHELPS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

Pacific Journal of Mathematics

Vol. 28, No. 2

April, 1969

Richard Arens and Donald George Babbitt, <i>The geometry of relativistic n-particle interactions</i>	243
Kirby Alan Baker, <i>Hypotopological spaces and their embeddings in lattices with Birkhoff interval topology</i>	275
J. Lennart (John) Berggren, <i>Finite groups in which every element is conjugate to its inverse</i>	289
Beverly L. Brechner, <i>Homeomorphism groups of dendrons</i>	295
Robert Ray Colby and Edgar Andrews Rutter, <i>QF - 3 rings with zero singular ideal</i>	303
Stephen Daniel Comer, <i>Classes without the amalgamation property</i>	309
Stephen D. Fisher, <i>Bounded approximation by rational functions</i>	319
Robert Gaines, <i>Continuous dependence for two-point boundary value problems</i>	327
Bernard Russel Gelbaum, <i>Banach algebra bundles</i>	337
Moses Glasner and Richard Emanuel Katz, <i>Function-theoretic degeneracy criteria for Riemannian manifolds</i>	351
Fletcher Gross, <i>Fixed-point-free operator groups of order 8</i>	357
Sav Roman Harasymiv, <i>On approximation by dilations of distributions</i>	363
Cheong Seng Hoo, <i>Nilpotency class of a map and Stasheff's criterion</i>	375
Richard Emanuel Katz, <i>A note on extremal length and modulus</i>	381
H. L. Krall and I. M. Sheffer, <i>Difference equations for some orthogonal polynomials</i>	383
Yu-Lee Lee, <i>On the construction of lower radical properties</i>	393
Robert Phillips, <i>Liouville's theorem</i>	397
Yum-Tong Siu, <i>Analytic sheaf cohomology groups of dimension n of n-dimensional noncompact complex manifolds</i>	407
Michael Samuel Skaff, <i>Vector valued Orlicz spaces. II</i>	413
James DeWitt Stein, <i>Homomorphisms of B^*-algebras</i>	431
Mark Lawrence Teply, <i>Torsionfree injective modules</i>	441
Richard R. Tucker, <i>The δ^2-process and related topics. II</i>	455
David William Walkup and Roger Jean-Baptiste Robert Wets, <i>Lifting projections of convex polyhedra</i>	465
Thomas Paul Whaley, <i>Large sublattices of a lattice</i>	477